



# ANTIBIOTIC ESSENTIALS

FOURTEENTH EDITION

Written by Leading World Experts

- Antibacterial, antifungal, antiviral, antiparasitic therapy
  - Clinical infectious disease syndromes
- Infectious disease differential diagnosis
  - Treatment of HIV infection and AIDS
- Pediatric infectious disease and therapy
  - Antimicrobial drug summaries
  - Immunizations and prophylaxis
  - Antibiotic susceptibility profiles
  - Antibiotic pearls and pitfalls

2015

---

BURKE A. CUNHA, MD, MACP

# ANTIBIOTIC ESSENTIALS

**Fourteenth Edition**

Edited by

**Burke A. Cunha, MD, MACP**

Chief, Infectious Disease Division  
Winthrop-University Hospital  
Mineola, New York  
Professor of Medicine  
State University of New York  
School of Medicine  
Stony Brook, New York



*The Health Sciences Publisher*

New Delhi | London | Philadelphia | Panama



## Jaypee Brothers Medical Publishers Pvt. Ltd.

### Headquarters

Jaypee Brothers Medical Publishers (P) Ltd  
4838/24, Ansari Road, Daryaganj  
New Delhi 110 002, India  
Phone: +91-11-43574357  
Fax: +91-11-43574314  
Email: [jaypee@jaypeebrothers.com](mailto:jaypee@jaypeebrothers.com)

### Overseas Offices

J.P. Medical Ltd  
83 Victoria Street, London  
SW1H 0HW (UK)  
Phone: +44 20 3170 8910  
Fax: +44 (0)20 3008 6180  
Email: [info@jpmepub.com](mailto:info@jpmepub.com)

Jaypee-Highlights Medical Publishers Inc  
City of Knowledge, Bld. 237, Clayton  
Panama City, Panama  
Phone: +1 507-301-0496  
Fax: +1 507-301-0499  
Email: [cSERVICE@jphmedical.com](mailto:cSERVICE@jphmedical.com)

Jaypee Medical Inc  
The Bourse  
111 South Independence Mall East  
Suite 835, Philadelphia, PA 19106, USA  
Phone: +1 267-519-9789  
Email: [jpmed.us@gmail.com](mailto:jpmed.us@gmail.com)

Jaypee Brothers Medical Publishers (P) Ltd  
17/1-B Babar Road, Block-B, Shaymali  
Mohammadpur, Dhaka-1207  
Bangladesh  
Mobile: +08801912003485  
Email: [jaypeedhaka@gmail.com](mailto:jaypeedhaka@gmail.com)

Jaypee Brothers Medical Publishers (P) Ltd  
Bhotahity, Kathmandu, Nepal  
Phone: +977-9741283608  
Email: [kathmandu@jaypeebrothers.com](mailto:kathmandu@jaypeebrothers.com)

Website: [www.jaypeebrothers.com](http://www.jaypeebrothers.com)  
Website: [www.jaypeedigital.com](http://www.jaypeedigital.com)

© 2015, Jaypee Brothers Medical Publishers

The views and opinions expressed in this book are solely those of the original contributor(s)/author(s) and do not necessarily represent those of editor(s) of the book.

All rights reserved. No part of this publication may be reproduced, stored or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior permission in writing of the publishers.

All brand names and product names used in this book are trade names, service marks, trademarks or registered trademarks of their respective owners. The publisher is not associated with any product or vendor mentioned in this book.

Medical knowledge and practice change constantly. This book is designed to provide accurate, authoritative information about the subject matter in question. However, readers are advised to check the most current information available on procedures included and check information from the manufacturer of each product to be administered, to verify the recommended dose, formula, method and duration of administration, adverse effects and contraindications. It is the responsibility of the practitioner to take all appropriate safety precautions. Neither the publisher nor the author(s)/editor(s) assume any liability for any injury and/or damage to persons or property arising from or related to use of material in this book.

This book is sold on the understanding that the publisher is not engaged in providing professional medical services. If such advice or services are required, the services of a competent medical professional should be sought.

Every effort has been made where necessary to contact holders of copyright to obtain permission to reproduce copyright material. If any have been inadvertently overlooked, the publisher will be pleased to make the necessary arrangements at the first opportunity.

**Inquiries for bulk sales may be solicited at:** [jaypee@jaypeebrothers.com](mailto:jaypee@jaypeebrothers.com)

### ***Antibiotic Essentials***

***Fourteenth Edition: 2015***

ISBN 978-93-5152-850-0

*Printed at*

---

## ABOUT THE EDITOR

---

Burke A. Cunha, MD, MACP, is Chief, Infectious Disease Division at Winthrop-University Hospital, Mineola, New York; Professor of Medicine, State University of New York School of Medicine, Stony Brook, New York; and is one of the world's leading authorities on infectious diseases. During his career, he has written/edited over 1250 articles, 200 book chapters, and 30 books on infectious diseases. He has received numerous teaching awards, including the Aesculapius Award for teaching excellence and the Spatz Award for clinical excellence and teaching excellence. Dr. Cunha is a Fellow of the Infectious Disease Society of America, American Academy of Microbiology, American College of Clinical Pharmacology, Surgical Infection Society, American College of Chest Physicians and Society of Hospital Epidemiologists of America. He has had a long interest in clinical syndromic diagnosis, diagnostic reasoning, antimicrobial therapy and resistance, atypical pneumonias, surgical infections, infections in compromised hosts, zoonoses, FUOs, meningitis and encephalitis, endocarditis, and nosocomial infections. Dr. Cunha is a Master of the American College of Physicians, awarded for lifetime achievement as a Master Clinician and Master Teacher of Infectious Diseases.

---

## DEDICATION

---

*for  
Marie*

*"Grace in her steps,  
Heaven in her eye,  
In every gesture, dignity and love"*

**For Downloading Dental Books!** *Milton*  
**Join us on telegram app:**

**<https://t.me/RoyalDentistryLibrary>**

---

---

**TABLE OF CONTENTS**


---

**1. OVERVIEW OF ANTIMICROBIAL****THERAPY**

Factors in Antibiotic Selection .....	1
Spectrum .....	2
Tissue Penetration .....	2
Antibiotic Resistance .....	2
Safety Profile .....	4
Cost .....	4
Factors in Antibiotic Dosing .....	4
Dosing in Renal/Hepatic Insufficiency .....	5
Routes of Elimination (Renal and Hepatic) .....	5
Microbiology and Susceptibility Testing .....	6
In Vitro vs. In Vivo Susceptibility .....	7
PK/PD and Other Considerations in	
Antimicrobial Therapy .....	9
Bactericidal vs. Bacteriostatic Therapy .....	9
Monotherapy vs. Combination Therapy .....	9
Intravenous vs. Oral Switch Therapy .....	10
Bioavailability of Oral Antibiotics .....	11
OPAT .....	11
Duration of Therapy .....	11
Empiric vs. Specific Antibiotic Therapy .....	12
Positive Blood Cultures vs. Bacteremia (MSSA, MRSA, CoNS) .....	13
Drug Fever .....	13
Relative Bradycardia .....	14
Antibiotic Failure .....	14
Pitfalls in Antibiotic Prescribing .....	15

**2. EMPIRIC THERAPY BASED ON CLINICAL SYNDROME****CNS Infections**

Acute Bacterial Meningitis .....	18
Acute Nonbacterial/Chronic Meningitis .....	21
Encephalitis .....	25
Brain Abscess/Subdural Empyema/ Cavernous Vein Thrombosis/Intracranial Suppurative Thrombophlebitis .....	27

**HEENT Infections**

Facial/Periorbital Cellulitis .....	29
Bacterial Sinusitis .....	30
Keratitis .....	31
Conjunctivitis .....	32
Chorioretinitis .....	34
Endophthalmitis .....	35
External Otitis .....	37

Acute Otitis Media .....	38
Mastoiditis .....	39
Suppurative Parotitis .....	40
Pharyngitis .....	40
Thrush (Oropharyngeal Candidiasis) .....	43
Mouth Ulcers/Vesicles .....	44
Deep Neck Infections, Lemierre's Syndrome, Severe Dental Infections .....	46
Epiglottitis .....	47

**Lower Respiratory Tract Infections**

Acute Bacterial Exacerbation of Chronic Bronchitis .....	48
Mediastinitis .....	49
Community-Acquired Pneumonia .....	49
Zoonotic Atypical Pneumonia .....	60
Lung Abscess/Empyema .....	67
Nursing Home-Acquired Pneumonia .....	68
Nosocomial Pneumonia .....	68

**Cardiovascular Infections**

Subacute Bacterial Endocarditis .....	70
Acute Bacterial Endocarditis .....	74
Prosthetic Valve Endocarditis .....	76
Pericarditis/Myocarditis .....	78
Central IV Line/Pacemaker Infections .....	79
Vascular Graft Infections .....	82

**Gastrointestinal Tract Infections**

Esophagitis .....	84
Peptic Ulcer Disease .....	85
Gastric Perforation .....	85
Diarrhea/Typhoid (Enteric) Fever .....	86
Cholecystitis .....	92
Cholangitis .....	93
Gallbladder Wall Abscess/Perforation .....	93
Acute Pancreatitis .....	94
Pancreatic Abscess/Infected Pancreatic Pseudocyst .....	95
Liver Abscess .....	95
Hepatosplenic Candidiasis .....	96
Granulomatous Hepatitis .....	96
Viral Hepatitis .....	97
Intraabdominal or Pelvic Peritonitis/ Abscess (Colitis, Appendicitis, Diverticulitis, Septic Pelvic Thrombophlebitis, Spontaneous Bacterial Peritonitis, CAPD-Associated Peritonitis, TB Peritonitis) .....	100

**Genitourinary Tract Infections**

Dysuria-Pyuria Syndrome .....	103
Cystitis .....	103
Catheter-Associated Bacteriuria/Candiduria .....	104
Epididymitis .....	106
Acute Pyelonephritis .....	107
Chronic Pyelonephritis .....	108
Renal TB .....	108
Renal Abscess (Intrarenal/Perinephric) .....	109
Prostatitis/Prostatic Abscess .....	110
Urosepsis .....	111
Pelvic Inflammatory Disease, Salpingitis, Tuboovarian Abscess, Endometritis/ Endomyometritis, Septic Abortion .....	113

**Sexually Transmitted Diseases**

Urethritis/Cervicitis .....	114
Vaginitis/Balanitis .....	115
Genital Vesicles .....	116
Genital Ulcers .....	117
Suppurating Inguinal Adenopathy .....	118
Genital/Perianal Warts .....	118
Syphilis .....	119

**Bone and Joint Infections**

Septic Arthritis/Bursitis .....	121
Lyme Disease/Lyme Arthritis .....	127
Infected Joint Prosthesis .....	128
Osteomyelitis .....	129

**Skin and Soft Tissue Infections**

Cellulitis, Erysipelas, Mastitis .....	132
Complicated Skin/Skin Structure Infections (cSSSIs) .....	134
Skin Ulcers .....	137
Skin Abscesses/Infected Cysts (Skin Pustules/Boils, Furunculosis) .....	140
Skin Vesicles (Non-Genital) .....	142
Wound Infections .....	144
Superficial Fungal Infections Skin/Nail .....	147
Skin Infestations .....	150
Ischiorectal/Perirectal Abscess .....	151

**Sepsis/Septic Shock**

151

**Febrile Neutropenia**

158

**Transplant Infections**

Bacteremia .....	160
Candidemia .....	161
CMV/HHV-6/BK Virus .....	162
EVB/PTLD .....	163
RSV .....	163
Adenovirus .....	163
Encephalitis/Meningitis .....	164

Brain Abscess/Mass Lesions .....	164
Focal/Segmental Pulmonary Infiltrates .....	165
Diffuse Pulmonary Infiltrates .....	165
Viral Hepatitis .....	166

**Toxin-Mediated Infectious Diseases**

Toxic Shock Syndrome .....	169
Botulism .....	170
Tetanus .....	170
Diphtheria .....	170

**Bioterrorist Agents**

Anthrax .....	173
Tularemia Pneumonia .....	173
Pneumonic Plague .....	174
Botulism .....	174
Smallpox .....	174
Ebola .....	174

**3. ANTIBIOTIC SUSCEPTIBILITY PROFILES AND INITIAL THERAPY OF ISOLATES PENDING SUSCEPTIBILITY RESULTS**

Antibiotic Susceptibility Profiles (Penicillins, Macrolides, Tetracyclines, and Others) .....	185
Antibiotic Susceptibility Profiles (Cephalosporins) .....	191
Antibiotic Susceptibility Profiles (Aminoglycosides, Fluoroquinolones, Carbapenems, and Others) .....	198
Gram Stain Characteristics of Isolates (by Morphology Arrangement, Oxygen Requirements) .....	203
Alphabetical Index of Isolates .....	205
Key Factors in Antibiotic Selection (Isolate Known) .....	207
Antibiotic Resistance Potential .....	207
Aerobic Isolates .....	208
Capnophilic Isolates .....	235
Anaerobic Isolates .....	236
Yeast/Fungi .....	242
Gram Stain/Giemsa Stain Technique .....	246
CSF Gram Stain .....	246
Sputum Gram Stain .....	248
Urine Gram Stain .....	248
Fecal Gram Stain .....	249

**4. PARASITES, FUNGI, UNUSUAL ORGANISMS**

253

**Blood Pathogens**

Microfilaria in Blood .....	254
Trypanosomes in Blood .....	255
Spirochetes in Blood .....	257
Intracellular Inclusion Bodies in Blood .....	258

<b>CSF/Brain Pathogens</b>	
Cysts/Mass Lesions in CSF/Brain.....	261
<b>Lung Pathogens</b>	
Pulmonary Cystic Lesions/Masses.....	265
Pulmonary Coin Lesions.....	266
Pulmonary Infiltrates/Mass Lesions.....	267
<b>Heart Pathogens</b>	
Chagas' Disease.....	273
<b>Liver Pathogens</b>	
Liver Flukes.....	273
Cystic Masses in Liver.....	274
Hepatomegaly.....	275
<b>Stool/Intestines Pathogens</b>	
Intestinal Protozoa.....	276
Intestinal Nematodes (Roundworms).....	278
Intestinal Cestodes (Tapeworms).....	281
Intestinal Trematodes (Flukes/Flatworms).....	282
Other Intestinal Infections.....	282
<b>Skin/Muscle Pathogens</b>	
Infiltrative Skin/Subcutaneous Lesions.....	283
Infiltrative Skin Lesions ± Ulcers/Sinus Tracts/Abscesses.....	284
Skin Vesicles/Bullae.....	288
Subcutaneous Serpiginous Lesions.....	288
Skin Papules/Nodules/Abscesses.....	289
Rickettsia (Fever/Petechial Skin Rash).....	294
Other Skin Lesions.....	297
Myositis.....	298
<b>5. HIV INFECTION</b>	301
Overview.....	302
Stages of HIV Infection.....	303
Acute (Primary) HIV Infection.....	304
Approach to HIV Testing.....	305
Initial Assessment of HIV Patients.....	308
Indications for Treatment of HIV Infection.....	312
Antiretroviral Treatment.....	313
Antiretroviral Treatment Failure.....	316
Opportunistic Infections in HIV Disease.....	319
Treatment of Opportunistic Infections.....	323
Treatment of Other Opportunistic Infections in HIV.....	338
HIV Coinfections (HBV/HCV).....	344
<b>6. PROPHYLAXIS AND IMMUNIZATIONS</b>	353
Surgical Prophylaxis.....	354
Post-Exposure Prophylaxis.....	358
Chronic Medical Prophylaxis/Suppression.....	364
HIV Pre-Exposure Prophylaxis (PrEP).....	365
HIV Post-Exposure Prophylaxis (PEP).....	366
HIV Opportunistic Infections.....	366
Transplant Prophylaxis.....	367
Endocarditis Prophylaxis.....	369
Travel Prophylaxis.....	371
Malaria Prophylaxis.....	375
Tetanus Prophylaxis.....	377
Immunizations.....	378
<b>7. PEDIATRIC INFECTIOUS DISEASES AND PEDIATRIC DRUG SUMMARIES</b>	386
CNS Infections.....	387
HEENT Infections.....	391
Lower Respiratory Tract Infections.....	396
Vascular Infections.....	402
Gastrointestinal Infections.....	403
Bone and Joint Infections.....	407
Skin and Soft Tissue Infections.....	411
Common Pediatric Antimicrobial Drugs.....	414
<b>8. CHEST X-RAY ATLAS</b>	425
Unilateral Focal Segmental/Lobar Infiltrate Without Effusion.....	426
Unilateral Focal Segmental/Lobar Infiltrate With Effusion.....	431
Unilateral Ill-Defined Infiltrates Without Effusion.....	436
Unilateral Ill-Defined Infiltrates With Effusion.....	443
Bilateral Infiltrates Without Effusion.....	448
Bilateral Infiltrates With Effusion.....	458
Cavitary Infiltrates (Thick-Walled).....	463
Cavitary Infiltrates (Thin-Walled).....	470
<b>9. INFECTIOUS DISEASE DIFFERENTIAL DIAGNOSIS</b>	474
<b>10. ANTIBIOTIC PEARLS &amp; PITFALLS</b>	507
<b>11. ANTIMICROBIAL DRUG SUMMARIES</b>	521
<b>APPENDIX</b>	720
<b>INDEX</b>	731
<b>COLOR ATLAS OF CSF, SPUTUM, AND URINE GRAM STAINS &amp; FUNGAL STAINS</b>	

---

**EDITORS**

---

**Burke A. Cunha, MD, MACP**  
*Editor*

**Cheston B. Cunha, MD**  
*Associate Editor*

---

**CONTRIBUTORS**

---

**Burke A. Cunha, MD, MACP**

Chief, Infectious Disease Division  
Winthrop-University Hospital  
Mineola, New York  
Professor of Medicine  
State University of New York  
School of Medicine  
Stony Brook, New York  
*All chapters except HIV Infection &  
Pediatric ID*

**Jeffrey Baron, PharmD**

Clinical Pharmacist  
Winthrop-University Hospital  
Mineola, New York  
*HBV, HCV, and HIV Guidelines  
Antimicrobial Drug Summaries*

**Edward J. Bottone, PhD**

Professor of Medicine  
Professor of Microbiology  
Professor of Pathology  
Mount Sinai  
School of Medicine  
New York, New York  
*Medical Microbiology; Parasites,  
Fungi, Unusual Organisms*

**John L. Bruschi, MD**

Associate Chief of Medicine  
Cambridge Health Alliance  
Infectious Disease Service  
Cambridge Health Alliance  
Medical Director, Somerville Hospital  
Assistant Professor of Medicine  
Harvard Medical School  
Boston, Massachusetts  
*Endocarditis: Therapy & Prophylaxis*

**Cheston B. Cunha, MD**

Director, Antibiotic Stewardship Program  
Rhode Island Hospital and The Miriam Hospital  
Infectious Disease Division  
Assistant Professor of Medicine  
Brown University Alpert School of Medicine  
Providence, Rhode Island  
*Infectious Disease Differential Diagnosis  
Overview of Antimicrobial Therapy  
Empiric Therapy Based of Clinical Syndromes  
Antimicrobial Drug Summaries*

**Daniel Caplivski, MD**

Infectious Disease Division  
Assistant Professor of Medicine  
Mt. Sinai School of Medicine  
New York, New York  
*Fungal Stain Atlas*

**Dennis J. Cleri, MD**

St. Francis Medical Center  
Professor of Medicine  
Seton Hall University  
School of Graduate Medical Education  
Trenton, New Jersey  
*Bioterrorism*

**Staci A. Fischer, MD**

Director, Transplant Infectious Diseases  
Rhode Island Hospital  
Infectious Disease Division  
Associate Professor of Medicine  
Brown University Alpert School of Medicine  
Providence, Rhode Island  
*Transplant Infections: Therapy & Prophylaxis*



**Pierce Gardner, MD**

Senior Advisor, Clinical Research and Training  
National Institutes of Health  
John E. Fogarty International Center for  
Advanced Study in the Health Sciences  
Bethesda, Maryland  
*Prophylaxis and Immunization*

**Arthur Gran, MD**

Infectious Disease Division  
Winthrop-University Hospital  
Mineola, New York  
State University of New York  
School of Medicine  
Stony Brook, New York  
*Antimicrobial Drug Summaries*

**Jean E. Hage, MD**

Infectious Disease Division  
Winthrop-University Hospital  
Mineola, New York  
State University of New York  
School of Medicine  
Stony Brook, New York  
*Empiric Therapy of Clinical Syndromes;  
Prophylaxis & Immunization;  
Parasitic & Fungi; Unusual Organisms;  
Antimicrobial Drug Summaries*

**Mark H. Kaplan, MD**

Professor of Medicine  
Infectious Diseases  
University of Michigan School of Medicine  
Ann Arbor, Michigan  
*HIV Drug Summaries*

**Douglas S. Katz, MD**

Vice Chairman, Clinical Research and Education  
Director, Body CT  
Winthrop-University Hospital  
Mineola, New York  
Professor of Clinical Radiology  
State University of New York School of Medicine  
Stony Brook, New York  
*Chest X-ray Atlas*

**Raymond S. Koff, MD**

Clinical Professor of Medicine  
University of Connecticut School of Medicine  
Farmington, Connecticut  
*Viral Hepatitis: Therapy & Prophylaxis*

**Leonard R. Krilov, MD**

Chief, Pediatric Infectious Disease Division  
Winthrop-University Hospital  
Mineola, New York  
Professor of Pediatrics  
State University of New York  
School of Medicine  
Stony Brook, New York  
*Pediatric Infectious Diseases*

**David W. Kubiak, PharmD**

Infectious Disease Clinical Pharmacist  
Brigham and Women's Hospital  
Boston, Massachusetts  
*Antiretroviral Drug Summaries*

**Brian R Malone, BS, MS, RPh**

Director, Pharmaceutical Services  
Winthrop-University Hospital  
Mineola, New York  
Adjunct Affiliate Professor  
College of Pharmacy St. John's University  
Queens, New York  
*Pediatric Drug Summaries*

**George H. McCracken, Jr., MD**

Distinguished Professor of Pediatric Infectious  
Disease and the Sarah M. and Charles E. Seyay  
Chair in Pediatric Infectious Disease  
University of Texas Southwestern Medical Center  
Dallas, Texas  
*Pediatric Infectious Diseases*

**James H. McGuire, MD**

Master Clinician  
Division of Infectious Diseases  
Brigham and Women's Hospital  
Professor of Medicine  
Harvard Medical School  
Boston, Massachusetts  
*Parasites, Fungi, Unusual Organisms*

**Nardeen Mickail, MD**

Infectious Disease Division  
Winthrop-University Hospital  
Mineola, New York  
State University of New York  
School of Medicine, Stony Brook, New York  
*Antimicrobial Drug Summaries*

**Maria D. Mileno, MD**

Director, Travel Medicine  
The Miriam Hospital  
Infectious Disease Division  
Associate Professor of Medicine  
Brown University Alpert School of Medicine  
Providence, Rhode Island  
*Prophylaxis and Immunizations*

**Robert Moore, MD**

Chairman, Department of Radiology  
Stony Brook University Hospital  
Professor of Radiology  
State University of New York School of Medicine  
Stony Brook, New York  
*Chest X-ray Atlas*

**Sigrith Muñoz-Gomez, MD**

Infectious Disease Division  
Winthrop-University Hospital  
Mineola, New York  
State University of New York  
School of Medicine  
Stony Brook, New York  
*Antimicrobial Drug Summaries*

**Ronald L. Nichols, MD**

William Henderson Professor of Surgery  
Professor of Microbiology and Immunology  
Tulane University School of Medicine  
New Orleans, Louisiana  
*Surgical Prophylaxis and Therapy*

**Genovefa Papanicolaou, MD**

Attending Physician, Infectious Disease Service  
Memorial Sloan Kettering Cancer Center  
Associate Professor of Medicine  
Weill Cornell Medical College  
New York, New York  
*Transplant Infections: Therapy & Prophylaxis*

**Muhammed Raza, MBBS**

Infectious Disease Division  
Winthrop-University Hospital  
Mineola, New York  
State University of New York  
School of Medicine  
Stony Brook, New York  
*Antimicrobial Drug Summaries*

**Michael F. Rein, MD**

Professor of Medicine (Emeritus)  
University of Virginia Health System  
Charlottesville, Virginia  
*Sexually Transmitted Diseases*

**John H. Rex, MD**

Adjunct Professor of Medicine  
University of Texas Medical School  
Houston, Texas  
Vice-President  
Medical Director for Infection  
AstraZeneca Pharmaceuticals  
Macclesfield, UK  
*Antifungal Therapy*

**Paul E. Sax, MD**

Clinical Director  
Division of Infectious Diseases and HIV  
Division of Infectious Diseases  
Brigham and Women's Hospital  
Associate Professor of Medicine  
Harvard Medical School  
Boston, Massachusetts  
*HIV Infection*

**David Schlossberg, MD**

Tuberculosis Control Program  
Philadelphia Department of Health  
Professor of Medicine  
Temple University School of Medicine  
Philadelphia, Pennsylvania  
*Tuberculosis*

**Paul E. Schoch, PhD**

Director  
Clinical Microbiology Laboratory  
Winthrop-University Hospital  
Mineola, New York  
*Medical Microbiology and Gram  
Stains Atlas*

**Daniel S. Siegal, MD**

Department of Radiology  
Mount Auburn Hospital  
Harvard Medical School  
Boston, Massachusetts  
*Chest X-ray Atlas*

**Stephanie Strollo, MD**

Infectious Disease Division  
Winthrop-University Hospital  
Mineola, New York  
State University of New York  
School of Medicine  
Stony Brook, New York  
*Initial Therapy of Isolates Pending  
Susceptibility Testing*

**Uzma Syed, DO**

Infectious Disease Division  
Winthrop-University Hospital  
Mineola, New York  
State University of New York  
School of Medicine  
Stony Brook, New York  
*Initial Therapy of Isolates Pending  
Susceptibility Testing*

**Damary C. Torres, PharmD**

Clinical Pharmacy Specialist  
Winthrop-University Hospital  
Mineola, New York  
Associate Clinical Professor  
College of Pharmacy St. John's University  
Queens, New York  
*Antimicrobial Drug Summaries*

**Kenneth F. Wagner, DO**

Infectious Disease Consultant  
National Naval Medical Center  
Associate Professor of Medicine  
Uniformed Services, University of  
the Health Sciences  
F. Edward Hebert School of Medicine  
Bethesda, Maryland  
*Parasites, Fungi, Unusual Organisms*

---

## ACKNOWLEDGMENTS

---

I would like to thank the many contributors who graciously contributed their time and energy; and especially Mark Freed, MD, past President and Editor-in-Chief of Physicians' Press, for his vision, commitment, and guidance.

Burke A. Cunha, MD

---

## NOTICE

---

The clinical recommendations set forth in this book are those of the authors and are offered as general guidelines, not specific instructions for individual patients. Clinical judgement should always guide the physician in the selection, dosing, and duration of antimicrobial therapy for individual patients.

***Not all recommendations in this book are approved indications by the U.S. Food and Drug Administration, and antimicrobial recommendations are not limited to indications in the package insert. The use of any drug should be preceded by careful review of the package insert, which provides indications and dosing approved by the U.S. Food and Drug Administration.***

The information provided in this book is essential not exhaustive, and the reader is referred to other medical references and the manufacturer's product literature for further information. *Clinical use of the information provided and any consequences that may arise from its use is the responsibility of the prescribing physician. The authors, editors, and publisher do not warrant or guarantee the information herein contained and do not assume and expressly disclaim any liability for errors or omissions or any consequences that may occur from use of this information.*

---

## BASIS OF RECOMMENDATIONS

---

***The therapeutic recommendations in Antibiotic Essentials are based on the contributors' clinical expertise and experience as well as the literature and clinical guidelines.***

---

**ABBREVIATIONS**


---

ABE	acute bacterial endocarditis	DI	diabetes insipidus
ABM	acute bacterial meningitis	DIC	disseminated intravascular coagulation
ADA	adenosine deaminase	DM	diabetes mellitus
AFB	acid fast bacilli	DOT	directly observed therapy
AIH	autoimmune lupoid hepatitis	e.g.	for example
AIHA	autoimmune hemolytic anemia	EBV	Ebstein-Barr virus
AML	acute myelogenous leukemia	EEE	Eastern equine encephalitis
ANA	antinuclear antibody	EEG	electroencephalogram
ARDS	adult respiratory distress syndrome	EIA	enzyme immunoassay
AG	Aspergillus galactomannan	ELISA	enzyme-linked immunosorbent assay
A-V	atrio-ventricular	EM	erythema migrans
$\beta$ -lactams	penicillins, cephalosporins, cephamycins (not monobactams or carbapenems)	EMB	ethambutol
		Enterobacteriaceae:	Citrobacter, Edwardsiella, Enterobacter, E. coli, Klebsiella, Proteus, Providencia, Shigella, Salmonella, Serratia, Hafnia, Morganella, Yersinia
BAL	bronchoalveolar lavage	ESBLs	extended spectrum $\beta$ -lactamases
BG	$\beta$ 1, 3 D-glucan	esp	especially
BMT	bone marrow transplant	ESR	erythrocyte sedimentation rate
BPH	benign prostatic hypertrophy	ESRD	end-stage renal disease
CAB	catheter associated bacteriuria	ET	endotracheal
CABG	coronary artery bypass grafting	EV	enterovirus
CAC	catheter associated candiduria	EVD	external ventricular drain
CAH	chronic active hepatitis	FTA-ABS	fluorescent treponemal antibody absorption test
CA-MRSA	community-acquired MRSA	FUO	fever of unknown origin
CAP	community-acquired pneumonia	G6PD	glucose-6-phosphate dehydrogenase
CD <sub>4</sub>	CD <sub>4</sub> T-cell lymphocyte	GC	gonococcus/gonorrhoea
CE	California encephalitis virus	GCA	giant cell arteritis
CFS	chronic fatigue syndrome	GI	gastrointestinal
CGD	chronic granulomatous disease	gm	gram
CIE	counter-immunoelectrophoresis	GU	genitourinary
CLL	chronic lymphocytic leukemia	GVHD	graft versus host disease
CML	chronic myelogenous leukemia	HA-MRSA	hospital acquired MRSA
CMV	Cytomegalovirus	HAP	hospital acquired pneumonia
CNS	central nervous system	HAV	hepatitis A virus
CO-MRSA	community onset MRSA	HBcAb	hepatitis B core antibody
CoNS	coagulase negative staphylococci	HBoV	human bocavirus
CPH	chronic persistent hepatitis	HBsAg	hepatitis B surface antigen
CPK	creatinine phosphokinase	HBV	hepatitis B virus
CrCl	creatinine clearance	HCV	hepatitis C virus
CRE	carbapenemase resistant Enterobacteriaceae	HD	hemodialysis
CSD	cat scratch disease	HdV	hepatitis D virus
CSF	cerebrospinal fluid	HEV	hepatitis E virus
CT	computerized tomography	HFHD	high flux hemodialysis
CVA	costovertebral angle	HFV	hepatitis F virus
CVC	central venous catheter	HFM	hand foot mouth disease
CVID	common variable immune deficiency		
CWH	continuous veno venous hemo filtration		
CXR	chest x-ray		
DFA	direct fluorescent antibody		

HGA	human granulocytic anaplasmosis	NNRTI	non-nucleoside reverse transcriptase inhibitor
HHV-6,7,8	human herpes virus 6,7,8	NP	nosocomial pneumonia
HME	human monocytic ehrlichiosis	NRTI	nucleoside reverse transcriptase inhibitor
hMPV	human metapneumovirus	NSAIDs	nonsteroidal anti-inflammatory drugs
HPS	Hanta virus pulmonary syndrome	OI	opportunistic infection
HPV	human papilloma virus	OPAT	outpatient parenteral antibiotic therapy
HTLV-1	human T-cell leukemia virus	PAN	polyarteritis nodosa
HRIG	human rabies immune globulin	PBC	primary biliary cirrhosis
HSV	herpes simplex virus	PBS	protected brush specimen
I & D	incision and drainage	PCEC	purified chick embryo cells
IFA	immunofluorescent antibody	PCN	penicillin
IgA	immunoglobulin A	PCP	Pneumocystis (carinii) jiroveci pneumonia
IgG	immunoglobulin G	PCR	polymerase chain reaction
IgM	immunoglobulin M	PD	peritoneal dialysis
IM	intramuscular	PDA	patent ductus arteriosus
INH	isoniazid	PE	Powassan encephalitis
IP	intraoperative	PEP	post-exposure prophylaxis
IT	intrathecal	PI	protease inhibitor
ITP	idiopathic thrombocytopenic purpura	PID	pelvic inflammatory disease
IV	intravenous	PML	progressive multifocal leukoencephalopathy
IV/PO	IV or PO	PMN	polymorphonuclear leukocytes
IVDA	intravenous drug abuser	PO	oral
IVIG	intravenous immunoglobulin	PPD	tuberculin skin test
JE	Japanese encephalitis	PPIs	proton pump inhibitors
kg	kilogram	PPNG	penicillinase-producing N. gonorrhoeae
L	liter	PTBM	partially treated bacterial meningitis
LCM	lymphocytic choriomeningitis	PVE	prosthetic valve endocarditis
LDH	lactate dehydrogenase	PVL	Panton-Valentine leukocidin
LFTs	liver function tests	PZA	pyrazinamide
LGV	lymphogranuloma venereum	q_d	every __ days
MAI	Mycobacterium avium-intracellulare	q_h	every __ hours
MCD	multicentric Castleman's disease	q month	once a month
mcg	microgram	q week	once a week
mcl	microliter	RA	rheumatoid arthritis
MDR	multidrug resistant	RBC	red blood cells
MDRSP	multidrug resistant S. pneumoniae	RE	regional ileitis (Crohn's disease)
MERS	middle east respiratory syndrome	RMSF	rocky mountain spotted fever
mg	milligram	RSV	respiratory syncytial virus
mL	milliliter	RUQ	right upper quadrant
MIC	minimum inhibitory concentration	RVA	rabies vaccine absorbed
min	minute	SARS	severe acute respiratory syndrome
MMR	measles, mumps, rubella	SBE	subacute bacterial endocarditis
MPD	myeloproliferative disorder	SCID	severe combined immune deficiency
MRI	magnetic resonance imaging	SGOT/SGPT	serum transaminases
MRSA	methicillin-resistant S. aureus	SLE	systemic lupus erythematosus
MRSE	methicillin-resistant S. epidermidis	St. LE	Saint Louis encephalitis
MS	multiple sclerosis	SOT	solid organ transplant
MSSA	methicillin-sensitive S. aureus	sp.	species
MSSE	methicillin-sensitive S. epidermidis	SPB	spontaneous bacterial peritonitis
MTT	methyltetrahydrozole	SPEP	serum protein electrophoresis
MU	million units	SQ	subcutaneous
MVP	mitral valve prolapse	STD	sexually transmitted diseases
NDM	New Delhi metallo- $\beta$ -lactamase	TA	temporal arteritis
NHAP	nursing home acquired pneumonia	TAA	teichoic acid antibody titers

TAH/BSO	total abdominal hysterectomy/bilateral salpingoophorectomy	VA	ventriculoatrial
TB	M. tuberculosis	VAP	ventilator associated pneumonia
TDF	tenofovir disoproxil fumarate	VCA	viral capsid antigen
TEE	transesophageal echocardiogram	VEE	Venezuelan equine encephalitis virus
TEN	toxic epidermal necrolysis	VISA	vancomycin intermediate S. aureus
TID	three times per day	VLM	visceral larval migrans
TMP	trimethoprim	VP	ventriculoperitoneal
TMP-SMX	trimethoprim-sulfamethoxazole	VRE	vancomycin resistant enterococci
TPN	total parenteral nutrition	VRSA	vancomycin resistant S. aureus
TRNG	tetracycline-resistant N.gonorrhoeae	VSE	vancomycin-sensitive enterococci
TSS	toxic shock syndrome	VZV	varicella zoster virus
TTE	transthoracic echocardiogram	WBC	white blood cells
TTP	thrombotic thrombocytopenic purpura	WEE	western equine encephalitis
TURP	transurethral resection of prostate	WNE	west nile encephalitis
UC	ulcerative colitis	XMRV	xenotropic murine leukemia related virus
UTI	urinary tract infection	yrs	years

## Chapter 1

**Overview of Antimicrobial Therapy**

**Burke A. Cunha, MD**  
**Cheston B. Cunha, MD**

Factors in Antibiotic Selection .....	2
Spectrum .....	2
Tissue Penetration .....	2
Antibiotic Resistance .....	2
Factors in Antibiotic Dosing .....	4
Dosing in Renal/Hepatic Insufficiency .....	5
Routes of Elimination (Renal and Hepatic) .....	5
Microbiology and Susceptibility Testing .....	6
In Vitro vs. In Vivo Susceptibility .....	7
Susceptibility Breakpoints for <i>S. pneumoniae</i> .....	8
PK/PD and Other Considerations in Antimicrobial Therapy .....	9
Bactericidal vs. Bacteriostatic Antibiotics .....	9
Monotherapy vs. Combination Therapy .....	9
IV to PO Switch Therapy .....	10
Bioavailability of Oral Antibiotics .....	11
OPAT .....	11
Duration of Antibiotic Therapy .....	11
Empiric vs. Specific Antibiotic Therapy .....	12
Positive Blood Cultures vs. Bacteremia (MSSA, MRSA, CoNS) .....	13
Drug Fever .....	13
Relative Bradycardia .....	14
Antibiotic Failure .....	14
Pitfalls in Antibiotic Prescribing .....	15
References and Suggested Readings .....	16



## Overview of Antimicrobial Therapy

---

### FACTORS IN ANTIBIOTIC SELECTION

---

- A. Spectrum.** Antibiotic spectrum refers to the range of microorganisms an antibiotic is usually effective against, and is the basis for empiric antibiotic therapy (Chapter 2). Antibiotic susceptibilities are a guide to predicting antibiotic effectiveness in blood/well vascularized organs. *In vitro* testing does not always predict *in vivo* effectiveness (see p. 6).
- B. Tissue Penetration.** Antibiotics that are effective against a microorganism *in vitro* but unable to reach the site of infection are of little or no benefit to the host. Antibiotic tissue penetration depends on properties of the antibiotic, e.g., lipid solubility, molecular size and tissue, e.g., adequacy of blood supply, presence of inflammation. Antibiotic tissue penetration is rarely problematic in acute infections due to increased microvascular permeability from local release of chemical inflammatory mediators. In contrast, chronic infections, e.g., chronic pyelonephritis, chronic prostatitis, chronic osteomyelitis and infections caused by intracellular pathogens often rely on chemical properties of an antibiotic, e.g., high lipid solubility, small molecular size for adequate tissue penetration. Antibiotics cannot be expected to eradicate organisms from areas that are difficult to penetrate or have impaired blood supply, such as abscesses, which usually require surgical drainage for cure. In addition, implanted foreign materials associated with infection usually need to be removed for cure, since microbes causing infections associated with prosthetic joints, shunts, and intravenous lines produce a slime/biofilm on plastic/metal surfaces that permits organisms to survive despite antimicrobial therapy.
- C. Antibiotic Resistance.** Bacterial resistance to antimicrobial therapy may be classified as natural/intrinsic or acquired relative or absolute. Pathogens not covered by the usual spectrum of an antibiotic are termed *naturally/intrinsically* resistant, e.g., 25% of *S. pneumoniae* are naturally resistant to macrolides; *acquired* resistance refers to a previously susceptible pathogen that is no longer susceptible to an antibiotic, e.g., ampicillin resistant *H. influenzae*. Organisms with *intermediate level (relative)* resistance manifests as an increase in minimum inhibitory concentrations (MICs), but are susceptible if achievable serum/tissue concentrations > MIC, e.g., penicillin-resistant *S. pneumoniae*. In contrast, organisms with *high level (absolute)* resistance manifests as a sudden increase in MICs during therapy, and cannot be overcome by higher-than-usual antibiotic doses, e.g., gentamicin-resistant *P. aeruginosa*. Most acquired antibiotic resistance is *agent-specific*, not a class specific, and is usually limited to one or two species. Resistance is *not* related, per se, to volume or duration of use. Some antibiotics have little resistance potential i.e., "low resistance" potential even when used in high volume; other antibiotics can induce resistance, e.g., "high resistance" potential with little use.

**The Antibiotic Resistance Potential of each antibiotic is included in each Drug Summary (see Chapter 11).**

**Table 1.1. Resistance Potential of Selected Antibiotics**

<b>“High Resistance Potential” Antibiotics</b>	<b>Usual Organism(s) Resistant for Each Antibiotic</b>	<b>Preferred “Low Resistance Potential” Antibiotic Alternatives in Same Class</b>	<b>Preferred “Low Resistance Potential” Antibiotic Alternatives in Different Classes</b>
<b>Aminoglycosides</b>			
Gentamicin or Tobramycin	<i>P. aeruginosa</i>	Amikacin	Levofloxacin or Colistin or Cefepime
<b>Cephalosporins</b>			
Ceftazidime	<i>P. aeruginosa</i>	Cefepime	Levofloxacin or Colistin
<b>Tetracyclines</b>			
Tetracycline	<i>S. pneumoniae</i> <i>S. aureus</i>	Doxycycline or Minocycline	Levofloxacin or Moxifloxacin
<b>Quinolones</b>			
Ciprofloxacin	<i>S. pneumoniae</i>	Levofloxacin or Moxifloxacin	Doxycycline
Ciprofloxacin	<i>P. aeruginosa</i>	Levofloxacin	Amikacin or Colistin or Cefepime
<b>Glycopeptides</b>			
Vancomycin	MSSA MRSA	None	Linezolid or Daptomycin or Minocycline or Tigecycline
<b>Carbapenems</b>			
Imipenem	<i>P. aeruginosa</i>	Meropenem or Doripenem	Amikacin or Cefepime or Colistin

**Table 1.1. Resistance Potential of Selected Antibiotics (Cont'd)**

<b>"High Resistance Potential" Antibiotics</b>	<b>Usual Organism(s) Resistant for Each Antibiotic</b>	<b>Preferred "Low Resistance Potential" Antibiotic Alternatives in Same Class</b>	<b>Preferred "Low Resistance Potential Antibiotic Alternatives in Different Classes</b>
<b>Macrolides</b>			
Azithromycin	<i>S. pneumoniae</i>	None	Doxycycline or Levofloxacin or Moxifloxacin
<b>Dihydrofolate Reductase Inhibitors</b>			
TMP-SMX	<i>S. pneumoniae</i>	None	Doxycycline or Levofloxacin or Moxifloxacin

Adapted from: Cunha BA. Antibiotic Resistance: Effective Control Strategies. *Lancet* 357:1307-1308, 2001; Cunha BA (Ed). *Antibiotic Essentials* (12th ed) Jones & Bartlett. Sudbury, MA 2013. p. 521-719

**D. Safety Profile.** Whenever possible, avoid antibiotics with serious/frequent side effects.

**E. Cost.** Switching early from IV to PO antibiotics is the single most important cost saving strategy in hospitalized patients, as the institutional cost of IV administration (~\$10/dose) may exceed the cost of the antibiotic itself. Antibiotic costs can also be minimized by using antibiotics with long half-lives, and by choosing monotherapy over combination therapy.

## FACTORS IN ANTIBIOTIC DOSING

**A. Renal Insufficiency.** Since most antibiotics eliminated by the kidneys have a wide "toxic-to-therapeutic ratio," dosing strategies are frequently based on formula-derived estimates of creatinine clearance, rather than precise quantitation of glomerular filtration rates. Dosage adjustments are especially important for antibiotics with narrow toxic-to-therapeutic ratios, and for patients who are receiving other nephrotoxic medications or have preexisting renal disease.

- 1. Initial and Maintenance Dosing in Renal Insufficiency.** For drugs eliminated by the kidneys, **the initial dose is unchanged**, and the **maintenance dose/dosing interval are modified in proportion to the degree of renal insufficiency (CrCl)**. Dosing adjustment problems in renal insufficiency can be circumvented by selecting an antibiotic with a similar spectrum that is eliminated by the hepatic route.
- 2. Aminoglycoside Dosing.** Single daily dosing—adjusted for the degree of renal insufficiency after the loading dose is administered—has virtually eliminated the nephrotoxic potential of aminoglycosides, and is recommended for all patients, including the critically ill (a possible exception is enterococcal endocarditis, where gentamicin dosing every 8 hours may be preferable). Aminoglycoside-induced tubular dysfunction is best assessed by quantitative renal tubular cast counts in urine, which more accurately reflect aminoglycoside nephrotoxicity than serum creatinine.

**Table 1.2. Dosing Strategies in Hepatic/Renal Insufficiency\*****Hepatic Insufficiency**

- Decrease total daily dose of hepatically-eliminated antibiotic by 50% in presence of clinically severe liver disease.
- **Alternative: Use antibiotic eliminated/inactivated by the renal route in usual dose.**

**Renal Insufficiency (Examples)**

- If creatinine clearance ~ 40–60 mL/min, decrease dose of renally-eliminated antibiotic by 50% and maintain the usual dosing interval.
- If creatinine clearance ~ 10–40 mL/min, decrease dose of renally-eliminated antibiotic by 50% and double the dosing interval.
- **Alternative: Use antibiotic eliminated/inactivated by the hepatic route in usual dose.**

Major Route of Elimination			
Hepatobiliary		Renal	
Chloramphenicol	Pyrazinamide	Most $\beta$ -lactams	Amantadine
Cefoperazone	Linezolid	$\beta$ -lactam/ $\beta$ -lactamase inhibitors	Rimantadine
Ceftriaxone <sup>t</sup>	Tedizolid	Aminoglycosides	Acyclovir
Doxycycline	Itraconazole	TMP–SMX	Valacyclovir
Minocycline	Isavuconazole	Azthreonam	Famciclovir
Telithromycin	Caspofungin	Carbapenems	Valganciclovir
Moxifloxacin	Micafungin	Polymyxin B	Osetamavir
Macrolides	Anidulafungin	Colistin	Zanamavir
Nafcillin	Ketoconazole	Ciprofloxacin	Peramavir
Clindamycin	Voriconazole	Levofloxacin	Tetracycline
Metronidazole	Posaconazole	Gatifloxacin	Oxacillin
Tigecycline			Daptomycin

**Table 1.2. Dosing Strategies in Hepatic/Renal Insufficiency\* (Cont'd)**

Major Route of Elimination		
Hepatobiliary	Renal	
Quinupristin/dalfopristin	Gemifloxacin	Telavancin
Isoniazid	Flucytosine	Dalbavancin
Ethambutol	Fluconazole	Oritavancin
Rifampin	Amphotericin	Ceftaroline fosamil
	Vancomycin	Fosfomycin
	Cycloserine	Nitrofurantoin

\* CrCl (mL/min) = [(140 – age) × weight (kg)] / [72 × serum creatinine (mg/dL)]. Multiply by 0.85 if female. It is important to recognize that due to age-dependent declines in renal function, elderly patients with “normal” serum creatinines may have CrCls requiring dosage adjustment. For example, a 70-year-old, 50-kg female with a serum creatinine of 1.2 mg/dL has an estimated CrCl of 34 mL/min.

† ~½ eliminated renally.

- B. Hepatic Insufficiency.** Antibiotic dosing for patients with hepatic dysfunction is problematic, since there is no hepatic counterpart to the serum creatinine to accurately assess liver function. In practice, antibiotic dosing is based on clinical assessment of the severity of liver disease. For practical purposes, dosing adjustments are usually not required for mild or moderate hepatic insufficiency. For severe hepatic insufficiency, dosing adjustments are usually made for antibiotics with hepatotoxic potential.
- C. Combined Renal and Hepatic Insufficiency.** There are no good dosing adjustment guidelines for patients with hepatorenal insufficiency. If renal insufficiency is worse than hepatic insufficiency, antibiotics eliminated by the liver are often administered at half the total daily dose. If hepatic insufficiency is more severe than renal insufficiency, renally eliminated antibiotics are usually administered and dosed in proportion to renal function.
- D. Mode of Antibiotic and Excretion/Excretory Organ Toxicity.** The mode of elimination/excretion does not predispose to excretory organ toxicity per se, e.g., nafcillin (hepatically eliminated) is not hepatotoxic, and its main side effect is nephrotoxicity (interstitial nephritis). In contrast, oxacillin (renally eliminated), is not nephrotoxic and its main side effect is hepatotoxicity (hepatitis).

---

## MICROBIOLOGY AND SUSCEPTIBILITY TESTING

---

- A. Overview.** In vitro susceptibility testing provides information about microbial sensitivities of a pathogen to various antibiotics and is useful in guiding therapy.

## B. Limitations of Microbiology Susceptibility Testing

- 1. *In vitro* data do not differentiate between colonizers and pathogens.** Before responding to a culture report from the microbiology laboratory, it is important to determine whether the organism is a pathogen or a colonizer in the clinical context. As a rule, colonization should not be treated.
- 2. *In vitro* data do not necessarily translate into *in vivo* efficacy.** Reports which indicate an organism is “susceptible” or “resistant” to a given antibiotic *in vitro* do not necessarily reflect *in vivo* activity.
- 3. *In vitro* susceptibility testing is dependent on the microbe, methodology, and antibiotic concentration.** *In vitro* susceptibility testing by the microbiology laboratory *assumes* the isolate was recovered from *blood*, and is being exposed to *serum* concentrations of an antibiotic given in the *usual* dose. Since some body sites e.g., bladder urine contains higher antibiotic concentrations than found in serum, and other body sites. CSF contain lower antibiotic concentrations than found in serum, i.e., *in vitro* data may be misleading for non-bloodstream infections. For example, a *Klebsiella pneumoniae* isolate obtained from the CSF may be reported as “sensitive” to cefazolin even though cefazolin does not penetrate the CSF. Likewise, *E. coli* and *Klebsiella* urinary isolates are often reported as “resistant” to ampicillin/sulbactam despite *in vivo* efficacy, due to high antibiotic concentrations in the urinary tract. Antibiotics should be prescribed at the usual recommended doses; attempts to lower cost by reducing dosage may decrease antibiotic efficacy e.g., ceftiofur 2 gm IV inhibits ~ 85% of *B. fragilis* isolates, whereas 1 gm IV inhibits only ~ 20% of strains.

**Table 1.3. Antibiotic-Organism Combinations for Which *In Vitro* Susceptibility Testing Does Not Predict *In Vivo* Effectiveness<sup>1</sup>**

Antibiotic	“Susceptible” Organism
Penicillin	<i>H. influenzae</i> , <i>Yersinia pestis</i>
TMP-SMX	<i>Klebsiella</i> , Enterococci, <i>Bartonella</i>
Polymyxin B	<i>Proteus</i> , <i>Salmonella</i>
Imipenem	<i>Stenotrophomonas maltophilia</i> <sup>2</sup>
Gentamicin	<i>Mycobacterium tuberculosis</i>
Vancomycin	<i>Erysipelothrix rhusiopathiae</i>
Aminoglycosides	Streptococci, <i>Salmonella</i> , <i>Shigella</i>
Clindamycin	Fusobacteria, Clostridia, enterococci, <i>Listeria</i>
Macrolides	<i>P. multocida</i>
1 <sup>st</sup> , 2 <sup>nd</sup> generation cephalosporins	<i>Salmonella</i> , <i>Shigella</i> , <i>Bartonella</i>

**Table 1.3. Antibiotic-Organism Combinations for Which *In Vitro* Susceptibility Testing Does Not Predict *In Vivo* Effectiveness<sup>1</sup> (Cont'd)**

Antibiotic	"Susceptible" Organism
3 <sup>rd</sup> , 4 <sup>th</sup> generation cephalosporins <sup>4</sup>	Enterococci, Listeria, Bartonella
All antibiotics	MRSA <sup>3</sup>

- In vitro* susceptibility does not predict *in vivo* activity; susceptibility data cannot be relied upon to guide therapy for antibiotic-organism combinations in this table.
- Formerly Pseudomonas.
- In spite of apparent *in vitro* susceptibility of antibiotics against MRSA, only vancomycin, minocycline, quinupristin/dalfopristin, linezolid, tedizolid, daptomycin, ceftaroline fosamil, telavancin, dalbavancin, oritavancin, and tigecycline are effective *in vivo*.
- Cefoperazone is the only cephalosporin with clinically useful anti-enterococcal activity against *E. faecalis* (VSE), not *E. faecium* (VRE).

**Table 1.4. CLSI Susceptibility Breakpoints for *Streptococcus pneumoniae***

Antibiotic	MIC (mcg/mL)		
	Sensitive	Intermediate	Resistant
Amoxicillin (non-meningitis)	≤ 2	4	≥ 8
Penicillin (meningitis)	≤ 0.06	–	≥ 0.12
Penicillin (non-meningitis)	≤ 2	4	≥ 8
Doxycycline	≤ 2	4	≥ 8
Cefepime (non-meningitis)	≤ 1	2	≥ 4
Cefepime (meningitis)	≤ 0.5	1	≥ 2
Cefotaxime (non-meningitis)	≤ 1	2	≥ 4
Cefotaxime (meningitis)	≤ 0.5	1	≥ 2
Ceftriaxone (non-meningitis)	≤ 1	2	≥ 4
Ceftriaxone (meningitis)	≤ 0.5	1	≥ 2
Meropenem	≤ 0.25	0.5	≥ 1
Vancomycin	≤ 1	–	–
Moxifloxacin	≤ 1	2	≥ 4
Levofloxacin	≤ 2	4	≥ 8
Chloramphenicol	≤ 4	–	≥ 8
Clindamycin	≤ 0.25	0.5	≥ 1
Linezolid	≤ 2	–	–

CLSI = Clinical and Laboratory Standards Institute (formerly NCCLS = National Committee for Clinical Laboratory Standards) M100–S20 (2010).

**C. Summary.** In vitro susceptibility testing is useful in most situations, but should not be followed blindly. Many factors need to be considered when interpreting in vitro microbiologic data, and infectious disease consultation is recommended for all but the most straightforward susceptibility interpretation problems. IV-to-PO switch changes using antibiotics of the same or other antibiotic class is best made when the oral antibiotic can achieve similar blood/tissue levels as the IV antibiotic.

---

### PK/PD AND OTHER CONSIDERATIONS IN ANTIMICROBIAL THERAPY

---

- A. Bactericidal vs. Bacteriostatic Therapy.** For most infections, bacteriostatic and bactericidal antibiotics inhibit/kill organisms at the same rate, and **should not be a factor in antibiotic selection.** Bactericidal antibiotics have an advantage in certain infections, such as endocarditis, meningitis, and febrile leukopenia, but there are exceptions even in these cases.
- B. Monotherapy vs. Combination Therapy. Monotherapy is preferred to combination therapy for nearly all infections.** In addition to cost savings, monotherapy results in less chance of medication error and fewer missed doses/drug interactions. Combination therapy may be useful for drug synergy or for extending spectrum beyond what can be obtained with a single drug. However, since drug synergy is difficult to assess and the possibility of antagonism always exists, antibiotics should be combined for synergy if synergy is based on actual testing. Combination therapy is not effective in preventing antibiotic resistance, except in very few situations.

#### Table 1.5. Combination Therapy and Antibiotic Resistance

##### Examples of Antibiotic Combinations That Prevent Resistance

Anti-pseudomonal penicillin (carbenicillin) + aminoglycoside (gentamicin, tobramycin, amikacin)  
 Rifampin + other TB drugs (INH, ethambutol, pyrazinamide)  
 5-flucytosine + amphotericin B

##### Examples of Antibiotic Combinations That Do Not Prevent Resistance\*

TMP-SMX  
 Aztreonam + ceftazidime  
 Cefepime + ciprofloxacin  
 Aminoglycoside + imipenem  
 Most other antibiotic combinations

\* These combinations are often prescribed to prevent resistance when, in actuality, they do not.

#### C. Pharmacokinetic (PK)/Pharmacodynamic (PD) Antibiotic Dosing Considerations



**Table 1.6. Antibiotic Dosing: Pharmacokinetic/Pharmacodynamic (PK/PD) Considerations**

Antibiotic-Dependent Antibiotics	Optimal Dosing Strategies
<b>Concentration-Dependent Antibiotics (C<sub>max</sub>: MIC)</b>	
<ul style="list-style-type: none"> <li>• Quinolones</li> <li>• Aminoglycosides</li> <li>• Vancomycin (if MIC ≥ 1 mcg/ml)</li> </ul>	<ul style="list-style-type: none"> <li>• Doxycycline</li> <li>• Tigecycline</li> <li>• Colistin</li> </ul>
<b>Use highest effective dose (without toxicity)</b>	
<b>Time Dependent Antibiotics (T &gt; MIC)</b>	
<ul style="list-style-type: none"> <li>• PCN concentrations &gt; MIC for ≥ 60% of the dosing interval</li> <li>• β-lactam concentrations &gt; MIC for ≥ 75% of the dosing interval</li> <li>• Carbapenems concentrations &gt; MIC for ≥ 40% of the dosing interval</li> <li>• Vancomycin (if MIC ≤ 1 mcg/ml)</li> </ul>	<ul style="list-style-type: none"> <li>• Doxycycline</li> <li>• Tigecycline</li> <li>• Colistin</li> </ul>
<b>Use high doses (which increase serum concentrations and also increases T &gt; MIC for more of the dosing interval)</b>	
<b>Other Antibiotics (C<sub>max</sub>: MIC/T &gt; MIC and/or AUC<sub>0-24</sub>/MIC)</b>	
<ul style="list-style-type: none"> <li>• Quinolones &gt; 125 (effective)</li> <li>• Quinolones &gt; 250 (more effective)</li> </ul>	<ul style="list-style-type: none"> <li>• Doxycycline</li> <li>• Tigecycline</li> <li>• Colistin</li> </ul>
<b>Use highest effective dose (without toxicity)</b>	

Adapted from: Roberts JA, Lipman J. Optimizing use of beta-lactam antibiotics in the critically ill. *Semin Respir Crit Care Med* 28:579-85,2007; Roberts JA, Pharm B, Kruger P, Paterson DL, Lipman J. Antibiotic resistance – What's dosing got to do with it? *Crit Care Med* 36:2433-40,2008; Roberts JA, Pharm B, Lipman J. Pharmacokinetic issues for antibiotics in the critically ill patient. *Crit Care Med* 37:840-851,2009.

**D. Intravenous vs. Oral Switch Therapy.** Patients admitted to the hospital are usually started on IV antibiotic therapy, then switched to equivalent oral therapy after clinical improvement/defervescence (usually within 72 hours). **Advantages of early IV-to-PO switch programs include reduced cost, early hospital discharge, less need for home IV therapy, and virtual elimination of IV line infections.** Drugs well-suited for IV-to-PO switch or for treatment entirely by the oral route include doxycycline, minocycline, clindamycin, metronidazole, chloramphenicol, amoxicillin, trimethoprim-sulfamethoxazole, quinolones, and linezolid.

**Most infectious diseases should be treated orally unless the patient is critically ill, cannot take antibiotics by mouth, or there is no equivalent oral antibiotic.** If the patient is able to take/absorb oral antibiotics, there is no difference in clinical outcome using equivalent IV or PO antibiotics. It is more important to think in terms of antibiotic spectrum, bioavailability and tissue penetration, rather than route of administration. **Nearly all non-critically ill patients should be treated in part or entirely with oral antibiotics.** When

**Table 1.7. Bioavailability of Oral Antimicrobials**

Bioavailability	Antimicrobials		
Excellent (> 90%)	Amoxicillin	TMP	Linezolid
	Cephalexin	TMP-SMX	Tedizolid
	Cefprozil	Doxycycline	Isavuconazole
	Cefadroxil	Minocycline	Voriconazole
	Clindamycin	Fluconazole	Rifampin
	Quinolones	Metronidazole	Isoniazid
	Chloramphenicol	Cycloserine	Pyrazinamide
Good (60–90%)	Cefixime	Valacyclovir	Ethambutol
	Cefpodoxime	Famciclovir	5-Flucytosine
	Ceftibuten	Valganciclovir	Posaconazole
	Cefuroxime	Macrolides	Itraconazole (solution)
		Cefaclor	Nitazoxanide (with food)
Poor (< 60%)	Vancomycin	Cefdinir	Nitazoxanide (without food)
	Acyclovir	Cefditoren	Fosfomycin

switching from IV to PO therapy, **the oral antibiotic chosen should have the same spectrum/degree of activity against the presumed/known pathogen and achieve the same blood and tissue levels as the equivalent IV antibiotic.**

- E. OPAT (outpatient parenteral antibiotic therapy).** OPAT has been used to treat infections IV on an outpatient basis or to complete IV therapy begun during hospitalization. Preferred OPAT antibiotics are those with few adverse effects and those with a long serum half life ( $t_{1/2}$ ). The most frequently used OPAT antibiotics are ceftriaxone and vancomycin. Other agents with long  $t_{1/2}$  ideal for OPAT of Gram positive cSSIs due to MRSA are telavancin 10 mg (IV) q 24 h, dalbavancin 1 gm (IV)  $\times$  1 doses then 500 mg (IV)  $\times$  1 dose 7 days later; tedizolid 200 mg (IV) q 24 h  $\times$  6 days, then 200 mg (PO) q 24 h  $\times$  6 days; and oritavancin 1200 mg (IV)  $\times$  1 dose. The alternative to OPAT is oral antibiotic therapy, e.g., for MRSA, minocycline or linezolid are equally efficacious as OPAT regimens.
- F. Duration of Therapy.** Most bacterial infections in normal hosts are treated with antibiotics for 1–2 weeks. The duration of therapy may need to be extended in patients with impaired immunity e.g., diabetes, SLE, alcoholic liver disease, neutropenia, diminished splenic function, etc., chronic bacterial infections e.g., endocarditis, osteomyelitis, chronic viral and fungal infections, or certain bacterial intracellular pathogens.

**Table 1.8. Infectious Diseases Requiring Prolonged Antimicrobial Therapy**

Therapy	Infectious Diseases
3 weeks	Lymphogranuloma venereum (LGV), syphilis (late latent), <i>H. pylori</i> , chronic prostatitis
4 weeks	Chronic otitis media, chronic sinusitis, acute osteomyelitis, chronic pyelonephritis, brain abscess, SBE
4–6 weeks	Acute bacterial endocarditis ( <i>S. aureus</i> , <i>Listeria</i> , enterococcal), chronic osteomyelitis <sup>4</sup>
3 months	Lung abscess <sup>1</sup> , melioidosis, bartonella
6 months	Pulmonary TB, extrapulmonary TB, actinomycosis <sup>2</sup> , nocardia <sup>3</sup> , prosthetic-related infections <sup>5</sup>
12 months	Whipple's disease
> 12 months	Lepromatous leprosy, HIV, Q fever (SBE/PVE)

1. Treat until resolved or until chest x-ray is normal/nearly normal and remains unchanged.
2. May require longer treatment; treat until resolved.
3. May require longer treatment in compromised hosts.
4. Adequate surgical debridement is required for cure.
5. Implanted foreign materials associated with infection (prosthetic valves, vascular grafts, joint replacements, hemodialysis shunts) should be removed as soon as possible after diagnosis. If removal is not feasible, then chronic suppressive therapy may be attempted, although clinical failure is the rule.

### EMPIRIC VS SPECIFIC ANTIBIOTIC THERAPY

*Always treat the usual pathogens* (related to body site flora) rather than just “covering the cultured organism” (*particularly if the specimen is not representative of the infected tissue*).

#### Examples:

##### Diabetic foot chronic osteomyelitis

**Cover the usual pathogens:** GAS, GBS, common coliforms, *S. aureus*, and *B. fragilis*  
(*not P. aeruginosa*)

**Do not cover surface colonizers cultured:** *P. aeruginosa*, acinetobacter, VSE/VRE, Enterobacter, Burkholderia, Stenotrophomonas

Do not rely on deep ulcer/fistula cultures which represent skin flora (and are not reflective of bone pathogens, i.e., osteomyelitis). If *P. aeruginosa* is cultured from deep ulcer/fistula, do not cover only for *P. aeruginosa*. Over 95% of diabetic foot ulcers/fistulas will be culture positive from *P. aeruginosa* (due to *P. aeruginosa* colonization from wet socks, wet dressings, whirlpool baths). In aseptically collected bone specimens in the OR, *P. aeruginosa* is NOT a bone pathogen in diabetics with chronic osteomyelitis.

##### Sacral decubitus ulcers (stage III/IV) = chronic osteomyelitis

**Cover the usual pathogens:** GAS, GBS, *S. aureus*, common coliforms, and *B. fragilis*.

**Do not cover surface colonizers cultured:** Stenotrophomonas, Acinetobacter, Enterobacter, *P. aeruginosa*, Burkholderia

**Table 1.9. Positive Blood Cultures vs. Bacteremia (MSSA, MRSA, CoNS)**

<b>Factors Favoring + Blood Cultures (not bacteremia)</b>	<b>Factors Favoring Bacteremia (not BC contaminants)</b>
<b>MSSA/MRSA</b> (+ BCs skin contamination likely) with: <ul style="list-style-type: none"> <li>• Intermittently positive BCs</li> <li>• Low level/low grade BC positivity (1/4 – 2/4 BCs +)</li> <li>• TTP = &gt; 2 days</li> <li>• No clinical source of MSSA/MRSA bacteremia (CVC, abscesses, osteomyelitis, ABE)</li> </ul>	<b>MSSA/MRSA Bacteremia</b> with: <ul style="list-style-type: none"> <li>• Persistently positive BCs</li> <li>• High level/high grade bacteremia (3/4 – 4/4 BCs +)</li> <li>• TTP = &lt; 2 days</li> <li>• Clinical source of MSSA/MRSA bacteremia clinically apparent (CVC, abscesses, osteomyelitis, ABE)</li> </ul>
<b>CoNS</b> (skin contamination likely) with: <ul style="list-style-type: none"> <li>• Intermittently positive BCs</li> <li>• Low level/low grade BC positivity (1/4 - 2/4 + BCs)</li> <li>• TTP = &gt; 2 days</li> <li>• No clinical source of CoNS + BCs (CVC, implanted orthopedic/cardiac devices, prosthetic materials, severe/prolonged neutropenia)</li> </ul>	<b>CoNS Bacteremia</b> (infection likely) with: <ul style="list-style-type: none"> <li>• Persistently positive BCs</li> <li>• High level/high grade bacteremia (3/4 – 4/4 BCs +)</li> <li>• TTP = &gt; 2 days</li> <li>• Clinical source of + BCs for CoNS apparent (CVC, implanted orthopedic/cardiac devices, prosthetic materials, severe/prolonged neutropenia)</li> </ul>

Implanted/prosthetic device associated: Dx = gallium or indium scans.

ABE: Dx = cardiac vegetation

Abscess Dx = Gallium scan or CT scan

CVC associated: Dx = SQ removed CVC tip culture with > 15 col of same organism as in BCs not drawn from the CVC

TTP = time to blood culture positivity

**Table 1.10. Clinical Features of Drug Fever**

### History

Many but not all individuals are atopic

Patients have been on a sensitizing medication for days or years “without a problem”

### Physical exam

Relative bradycardia

Fevers may be low- or high-grade, but usually range between 102°–104°F and may exceed 106°F

Patient appears “inappropriately well” for degree of fever

### Laboratory tests

Elevated WBC count (usually with left shift)

Eosinophils almost always present, but eosinophilia is uncommon

Elevated erythrocyte sedimentation rate in majority of cases

Early, transient, mild elevations of serum transaminases (common)

Negative blood cultures (excluding contaminants)

**Relative Bradycardia\*****Temperature-Pulse Relationships**

<b>Temperature</b>	<b>Appropriate Pulse Response (beats/min)</b>	<b>If Relative Bradycardia Pulse (beats/min)</b>
106°F (41.1°C)	150	< 140
105°F (40.6°C)	140	< 130
104°F (40.7°C)	130	< 120
103°F (39.4°C)	120	< 110
102°F (38.9°C)	110	< 100

\* Relative bradycardia refers to heart rates that are inappropriately slow relative to body temperature (pulse must be taken simultaneously with temperature elevation). Applies to adult patients with temperature  $\geq 102^\circ\text{F}$ ; does not apply to patients with second/third-degree heart block, pacemaker-induced rhythms, or those taking beta-blockers, diltiazem, or verapamil.

**ANTIBIOTIC FAILURE****Table 1.11. Causes of Apparent/Actual Antibiotic Failure****Microbiologic Factors**

- *In vitro* susceptibility but ineffective *in vivo*
- Antibiotic "tolerance" with gram-positive cocci
- Treating colonization (not infection)

**Antibiotic Factors**

- Inadequate coverage/spectrum
- Inadequate antibiotic blood tissue levels
- Decreased antibiotic activity in tissue
- Drug-drug interactions (inactivation/antagonism)

**Antibiotic Penetration Problems**

- Undrained abscess
- Foreign body-related infection
- Protected focus e.g., cerebrospinal fluid
- Organ hypoperfusion/diminished blood supply

**Noninfectious Diseases**

- Medical disorders mimicking infection e.g., SLE
- Drug fever

**Antibiotic-unresponsive Infectious Diseases**

- Viral or Fungal infections

---

### PITFALLS IN ANTIBIOTIC PRESCRIBING

---

- **Do not use antibiotics to “treat” colonization or non-infectious/diseases.**
- **Overuse of combination therapy.** *Monotherapy is preferred over combination therapy unless compelling reasons prevail, such as drug synergy or extended spectrum beyond what can be obtained with a single drug. Monotherapy reduces the risk of drug interactions and side effects, and is usually less expensive.*
- **Use of antibiotics for persistent fevers.** For patients with persistent fevers on an antimicrobial regimens that appears to be failing, it is important to reassess the patient rather than add additional antibiotics. Causes of prolonged fevers include undrained septic foci, noninfectious medical disorders, and drug fevers. *Undiagnosed causes of leukocytosis/low-grade fevers should not be treated with prolonged courses of antibiotics.*
- **Inadequate surgical therapy.** Infections involving infected prosthetic materials or fluid collections e.g., abscesses often require surgical therapy for cure. *For infections such as chronic osteomyelitis, surgery is the only way to cure the infection; antibiotics are useful only for suppression or to prevent local infectious complications.*

## REFERENCES AND SUGGESTED READINGS

- Bosso JA, Mauldin PD, Salgado CD. The association between antibiotic use and resistance: the role of secondary antibiotics. *Eur J Clin Microbiol Infect Dis* 29:1125–1192, 2010.
- Cunha BA. Oral Antibiotic Therapy of Serious Systemic Infections. *Med Clin N Am* 90:1197–1222, 2006.
- Cunha BA. *Pseudomonas aeruginosa*: Resistance and therapy. *Semin Respir Infect* 17:231–9, 2002.
- Cunha BA. Clinical relevance of penicillin-resistant *Streptococcus pneumoniae*. *Semin Respir Infect* 17:204–14, 2002.
- Cunha BA. Effective antibiotic resistance and control strategies. *Lancet* 357:1307–1308, 2001.
- Cunha BA. The significance of antibiotic false sensitivity testing with in vitro testing. *J Chemother* 9:25–33, 1997.
- Cunha BA, Ortega A. Antibiotic failure. *Medical Clinics of North America* 79:663–672, 1995.
- Cunha CB, Varughese CA, Mylonakis E. Antimicrobial Stewardship programs (ASPs). The devil is in the details. *Virulence* 4: 147–149, 2013.
- Empey KM, Rapp RP, Evans ME. The effect of an antimicrobial formulary change on hospital resistance patterns. *Pharmacotherapy* 22:81–7, 2002.
- Endimiani A, Perez F, Bajaksouzian S, et al. Evaluation of updated interpretive criteria for categorizing *Klebsiella pneumoniae* with reduced carbapenem susceptibility. *J Clin Microbiol* 48:4417–4425, 2010.
- Hicks LA, Monnet DL, Roberts RM. Increase in pneumococcus macrolide resistance, USA. *Emerg Infect Dis* 16:896–897, 2010.
- Johnson DH, Cunha BA. Drug fever. *Infectious Disease Clinics of North America* 10:85–91, 1996.
- Jorgensen JH, Ferraro MJ. Antimicrobial susceptibility testing: a review of general principles and contemporary practices. *Clin Infect Dis* 49:1749–1755, 2009.
- Leekeha S, Terrell CL, Edson RS. General principles of antimicrobial therapy. *Mayo Clin Proc* 86:156–167, 2011.
- Patel SM, Saravolatz LD. Monotherapy versus combination therapy. *Med Clin N Am* 90:1183–95, 2006.
- Schlossberg D. Clinical approach to antibiotic failure. *Med Clin N Am* 90:1265–77, 2006.
- Tenover FC, Moellering RC. The rationale for revising the clinical and laboratory standards institute vancomycin minimal inhibitory concentration interpretive criteria for *Staphylococcus aureus*. *Clinical Infect Dis* 44:1208–1215, 2007.
- Weinstein MP, Klugman KP, Jones RN. Rationale for revised penicillin susceptibility breakpoints versus *Streptococcus pneumoniae*: coping with antimicrobial susceptibility in an era of resistance. *Clin Infect Dis* 48:1596–1600, 2009.
- Yamagishi Y, Terada M, Ohki E, Miura Y, Umemura T, Mikamo H. Investigation of the clinical breakpoints of piperacillin-tazobactam against infections caused by *Pseudomonas aeruginosa*. *J Chemo* 18: 127–129, 2012.

## TEXTBOOKS

- Anaissie EJ, McGinnis MR, Pfäller MA (eds). *Clinical Mycology*. Churchill Livingstone, New York, 2003.
- Baddour L, Gorbach SL (eds). *Therapy of Infectious Diseases*. Saunders, Philadelphia, 2003.
- Bennett JE, Dolin R, Blaser MJ (eds). *Mandell, Douglas, and Bennett's Principles and Practice of Infectious Diseases* (8th Ed). Philadelphia Elsevier Churchill Livingstone, 2015.
- Bryskier A (ed). *Antimicrobial Agents*. ASM Press, Washington, DC, 2005.
- Cohen J, Powderly WG, Opal S (eds). *Infectious Diseases* (4th Ed). Philadelphia: Elsevier, 2015.
- Finch RG, Greenwood D, Norrby SR, Whitley RJ (eds). *Antibiotic and Chemotherapy* (8th Ed). Churchill Livingstone, Edinburgh, 2003.
- Gorbach SL, Bartlett JG, Blacklow NR (eds). *Infectious Diseases* (3rd Ed). Philadelphia, Lippincott, Williams & Wilkins, 2004.
- Grayson ML (ed). *Kucers' The Use of Antibiotics* (6th Ed), ASM Press, Washington, DC, Blaser MJ 2010.
- O'Grady F, Lambert HP, Finch RG, Greenwood D (eds). *Antibiotic and Chemotherapy* (2nd Ed). Churchill Livingstone, New York, 1997.
- Ristuccia AM, Cunha BA (eds). *Antimicrobial Therapy*. Raven Press, New York, 1984.
- Scholar EM Pratt WB (eds). *The Antimicrobial Drugs* (2nd Ed), Oxford University Press, New York, 2000.
- Schlossberg D (ed). *Clinical Infectious Disease* (2nd Ed), Cambridge University Press, Cambridge, 2015.
- Yoshikawa TT, Rajagopalan S (eds). *Antibiotic Therapy for Geriatric Patients*. Taylor & Francis, New York, 2006.
- Yu V, Edwards G, McKinnon PS, Peloquin C, Morse G (eds). *Antimicrobial Therapy and Vaccines, Volume II: Antimicrobial Agents* (2nd Ed), Esun Technologies, Pittsburgh, 2005.

## Chapter 2

**Empiric Therapy Based on Clinical Syndrome**

**Burke A. Cunha, MD, John L. Bruschi, MD, Ronald L. Nichols, MD**  
**Genovefa Papanicolaou, MD, Jean E. Hage, MD**  
**Raymond S. Koff, MD, John H. Rex, MD, Dennis J. Cleri, MD**  
**Cheston B. Cunha, MD, Staci A. Fischer, MD**  
**David Schlossberg, MD**

Empiric Therapy of CNS Infections .....	18
Empiric Therapy of HEENT Infections .....	29
Empiric Therapy of Lower Respiratory Tract Infections.....	48
Empiric Therapy of Cardiovascular Infections .....	70
Empiric Therapy of GI Tract Infections.....	84
Empiric Therapy of Genitourinary Tract Infections.....	103
Empiric Therapy of Sexually Transmitted Diseases.....	114
Empiric Therapy of Bone and Joint Infections .....	121
Empiric Therapy of Skin and Soft Tissue Infections .....	132
Sepsis/Septic Shock .....	151
Febrile Neutropenia .....	158
Transplant Infections .....	160
Toxin-Mediated Infectious Diseases.....	169
Bioterrorist Agents .....	173
References and Suggested Readings .....	178

This chapter is organized by clinical syndrome, patient subset, and in some cases, specific organism. Clinical summaries immediately follow each treatment grid. Unless otherwise specified, **this chapter pertains to infectious diseases and antimicrobial agents in adults. For pediatric infectious diseases and antimicrobial therapy see Chapter 7.** Therapeutic recommendations are based on antimicrobial effectiveness, reliability, cost, safety, and resistance potential. The antimicrobial dosages in this section represent the usual dosages for normal renal and hepatic function in adults. For any treatment category (i.e., preferred IV therapy, alternate IV therapy, PO therapy), **in any given grid category drugs listed are equivalent alternatives and not ranked.** Dosage adjustments, side effects, drug interactions, and other important prescribing information are



described in the individual drug summaries in Chapter 11. Use of any drug should be preceded by careful review of the package insert, which provides indications and dosing approved by the U.S. Food and Drug Administration. “*IV-to-PO Switch*” in the last column of the shaded title bar in each treatment grid indicates the clinical syndrome should be treated either by IV therapy alone or IV followed by PO therapy, but *not* by PO therapy alone. “*PO Therapy or IV-to-PO Switch*” indicates the clinical syndrome can be treated by IV therapy alone, PO therapy alone, or IV followed by PO therapy (unless otherwise indicated in the footnotes under each treatment grid). Most patients on IV therapy should be switched to PO equivalent therapy after clinical improvement.

## Empiric Therapy of CNS Infections

### Acute Bacterial Meningitis (ABM) (see Color Atlas for CSF Gram stains)

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	IV-to-PO Switch
Normal host	N. meningitidis H. influenzae S. pneumoniae	Ceftriaxone 2 gm (IV) q12h × 2 weeks	Meropenem 2 gm (IV) q8h × 2 weeks <b>or</b> Cefotaxime 3 gm (IV) q6h × 2 weeks <b>or</b> Ceftizoxime 3 gm (IV) q6h × 2 weeks	Chloramphenicol 500 mg (PO) q6h × 2 weeks
Elderly or malignancy	Listeria monocytogenes plus usual meningeal pathogens in normal hosts	<u>Before culture results</u> Ceftriaxone 2 gm (IV) q12h × 3 weeks <b>plus</b> Ampicillin 2 gm (IV) q4h × 3 weeks <u>After culture results</u> <u>Listeria present</u> Ampicillin 2 gm (IV) q4h × 3 weeks <u>Listeria not present</u> Treat as normal host	<u>After culture results</u> <u>Listeria present</u> TMP-SMX 5 mg/kg (IV) q6h × 3 weeks <b>or</b> Meropenem 2 gm (IV) q8h × 3 weeks <b>or</b> Chloramphenicol 500 mg (IV) q6h × 3 weeks <b>or</b> Linezolid 600 mg (IV) q12h × 3 weeks <u>Listeria not present</u> Treat as for normal host, above	<u>For Listeria meningitis only</u> TMP-SMX 5 mg/kg (PO) q6h × 3 weeks <b>or</b> Chloramphenicol 500 mg (PO) q6h × 3 weeks <b>or</b> Linezolid 600 mg (PO) q12h × 3 weeks <u>For usual meningeal pathogens</u> Chloramphenicol 500 mg (PO) q6h × 3 weeks

## Acute Bacterial Meningitis (ABM) (cont'd) (see Color Atlas for CSF Gram stains)

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	IV-to-PO Switch
CNS shunt infections (VA shunts) (Treat initially for MSSA; if later identified as MRSA, MSSE, or MRSE, treat accordingly)	<i>S. aureus</i> <i>S. epidermidis</i> (coagulase-negative staphylococci) CoNS	<u>MSSA/MSSE</u> Meropenem 2 gm (IV) q8h*  <b>or</b> Ceftizoxime 3 gm (IV) q6h*  <u>MRSA/MRSE</u> Linezolid 600 mg (IV) q12h*	<u>MSSA/MSSE</u> Cefepime 2 gm (IV) q8h*  <b>or</b> Cefotaxime 3 gm (IV) q6h*  <u>MRSA/MRSE</u> Vancomycin 2 gm (IV) q12h*  <b>plus</b> 20 mg (IT) q24h until shunt removal	<u>MSSE/MRSE</u> Linezolid 600 mg (PO) q12h*          <u>MSSA/MRSA</u> Minocycline 100mg (PO) q12h*  <b>or</b> Linezolid 600 mg (PO) q12h*
CNS shunt infections (VP shunts)	<i>E. coli</i> <i>K. pneumoniae</i> <i>Enterobacter</i> <i>S. marcescens</i>	Ceftriaxone 2 gm (IV) q12h × 2 weeks (after shunt removal)	TMP-SMX 5 mg/kg (IV) q6h × 2 weeks (after shunt removal)	TMP-SMX 5 mg/kg (PO) q6h × 2 weeks (after shunt removal)
	MDR <i>P. aeruginosa</i>	Meropenem 2 gm (IV) q8h ± Colistin 5 mg/kg (IV) q8h  <b>plus</b> Colistin 10 mg (IT) q24h × 2 weeks (after shunt removal)	Meropenem 2 gm (IV) q8h ± Amikacin 1 gm (IV) q24h  <b>plus</b> Amikacin 10 mg (IT) q24h × 2 weeks (after shunt removal)	
	MDR <i>Acinetobacter baumannii</i>	Meropenem 2 gm (IV) q8h  <b>or</b> Ampicillin/sulbactam 4.5 gm (IV) q6h ± Colistin 5 mg/kg (IV) q8h  <b>plus</b> Colistin 10 mg (IT) q12h × 2 weeks (after shunt removal)	Meropenem 2 gm (IV) q8h  <b>plus</b> Ampicillin/sulbactam 4.5 gm (IV) q6h × 2 weeks (after shunt removal)	

MSSA/MRSA = methicillin-sensitive/resistant *S. aureus*; MSSE/MRSE = methicillin-sensitive/resistant *S. epidermidis*.

\* Treat for 1 week after shunt removal.

**Clinical Presentation:** Abrupt onset of fever, headache, stiff neck.

**Diagnosis:** CSF gram stain/culture.

### **Acute Bacterial Meningitis (Normal Hosts)**

**Diagnostic Considerations:** Gram stain of centrifugated CSF is still the best diagnostic test. CSF antigen/CIE are unhelpful in establishing the diagnosis (many false-negatives). Blood cultures are positive for ABM pathogen in 80–90%. Typical CSF findings include a WBC count of 100–5000 cells/mm<sup>3</sup>, elevated opening pressure, elevated protein and lactic acid levels (> 4–6 mmol/L), and a positive CSF gram stain. If the WBC is extremely high (> 20,000 cells/mm<sup>3</sup>), suspect brain abscess with rupture into the ventricular system, and obtain a CT/MRI to confirm. *S. pneumoniae* meningitis is associated with cranial nerve abnormalities, mental status changes, and neurologic sequelae. With *H. influenzae* or *S. pneumoniae* meningitis, obtain a head CT/MRI to rule out other CNS pathology.

**Pitfalls:** If ABM is suspected, always perform lumbar puncture (LP) *before* obtaining a CT scan, since early antibiotic therapy is critical to prognosis. A CT/MRI should be obtained before LP *only* if a mass lesion/suppurative intracranial process is of primary concern, after blood cultures have been drawn. A stiff neck on physical examination has limited diagnostic value in the elderly, since nuchal rigidity may occur without meningitis (e.g., cervical arthritis) and meningitis may occur without nuchal rigidity. Recurrence of fever during the first week of *H. influenzae* meningitis is commonly due to subdural effusion, which usually resolves spontaneously over several days. Meningococcal meningitis may occur with or without meningococemia. On gram stain, *S. pneumoniae* may be mistaken for *H. influenzae*, and *Listeria* may be mistaken for *S. pneumoniae*. Meningococcal meningitis may occur in those who received the meningococcal vaccine especially serogroup B not in meningococcal vaccine.

**Therapeutic Considerations:** Do not reduce meningeal antibiotic dosing as the patient improves. Repeat LP if the patient is not responding to antibiotics after 48 hours; lack of response may be due to therapeutic failure, relapse, or a Noninfectious CNS disorder. For *S. pneumoniae* meningitis, obtain penicillin MICs on all CSF isolates. Penicillin-resistant strains are susceptible to meningeal doses of beta-lactam antibiotics (e.g., ceftriaxone). All but the most highly penicillin-resistant pneumococci are still effectively treated with meningeal doses of beta-lactams. Highly resistant pneumococcal strains (rare in the CSF) may be treated for 2 weeks with meropenem 2 gm (IV) q8h, or, linezolid 600 mg (IV) q12h. Dexamethasone 0.15 mg/kg (IV) q6h × 4 days may be given to children with ABM to reduce the incidence/severity of neurologic sequelae, although the value of steroids in adult ABM is unclear; if used, give dexamethasone 30 minutes *before* the initial antibiotic dose.

**Prognosis:** Often fatal without treatment. Case-fatality rates in treated adults are 10–20%.

Neurological deficits on presentation are associated with a poor prognosis. Permanent neurological deficits are more frequent with *S. pneumoniae* than *H. influenzae*, even with prompt therapy. In meningococcal meningitis with meningococemia, prognosis is related to the number of petechiae, with few or no neurological deficits in survivors.

### **Acute Bacterial Meningitis (Elderly Malignancy)**

**Diagnostic Considerations:** Diagnosis by CSF gram stain/culture. ABM pathogens include usual pathogens in normal hosts plus *Listeria monocytogenes*, a gram-positive, aerobic, bacillus. *Listeria* is the most common ABM pathogen in patients with malignancies, and is a common pathogen in the elderly. With *Listeria* meningitis, CSF cultures are positive in 100%, but CSF gram stain is negative in 50%. Meningeal carcinomatosis is suggested by multiple cranial nerve abnormalities.

**Pitfalls:** "Diphtheroids" isolated from CSF should be speciated to rule out *Listeria*. *Listeria* are motile and hemolytic on blood agar plate, diphtheroids are not.

**Therapeutic Considerations:** Elderly patients and cancer patients with ABM require empiric coverage of *Listeria* plus other common pathogens in normal hosts (*N. meningitidis*, *H. influenzae*, *S. pneumoniae*). Third-generation cephalosporins are not active against *Listeria*.

**Prognosis:** Related to underlying health of host.

### Acute Bacterial Meningitis (CNS Shunt Infections)

**Diagnostic Considerations:** Diagnosis by CSF gram stain/culture. *S. epidermidis* meningitis usually occurs only with infected prosthetic implant material (e.g., CNS shunt).

**Pitfalls:** Blood cultures are usually negative for shunt pathogens. CSF *P. acnes* cultures usually represent skin contamination. *P. acnes* is rarely a CNS shunt pathogen.

**Therapeutic Considerations:** 15% of *S. epidermidis* strains are resistant to nafcillin/clindamycin. In addition to systemic antibiotics in meningeal doses, adjunctive intraventricular/intrathecal antibiotics are sometimes given to control shunt infections before shunt removal. If vancomycin is used to treat CNS *S. aureus* (MSSA/MRSA) infection, I.T. dosing needed in addition to IV therapy.

**Prognosis:** Good if prosthetic material is removed.

### Acute Nonbacterial Meningitis/Chronic Meningitis

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	PO Therapy or IV-to-PO Switch
Viral (aseptic)	EBV, LCM, Parvo B19, mumps, Enteroviruses, WNE. No effective therapy for most. For HIV (see Chapter 5)			
	HSV-1 HSV-2	<u>IV Therapy:</u> Acyclovir 10 mg/kg (IV) q8h × 10 days <b>or</b> Ganciclovir 5 mg/kg (IV) q12h × 10 days. IV → PO: Valacyclovir 1 gm (PO) q8h × 10 days		
	VZV	Acyclovir 10 mg/kg (IV) q8h × 10 days. IV → PO: Valacyclovir 2 gm (PO) q6h × 10 days		
	HHV-6	<u>Ganciclovir:</u> 5 mg/kg (IV) of 12h × 2 weeks <b>or</b> Foscarnet 90 mg/kg (IV) q12h × 2 weeks		
Primary amebic meningo-encephalitis	<i>Naegleria fowleri</i>	Amphotericin B deoxycholate 1 mg/kg (IV) q24h until cured <b>plus</b> Amphotericin B deoxycholate 1 mg into ventricles via Ommaya reservoir q24h until cured		
Lyme neuro-borreliosis	<i>Borrelia burgdorferi</i>	Ceftriaxone 1 gm (IV) q12h × 2 weeks <b>or</b> Minocycline 100 mg (IV) q12h × 2 weeks <b>or</b> Doxycycline 200 mg (IV) q12h × 3 days, then 100 mg (IV) q12h × 11 days		Minocycline 100 mg (PO) q12h × 2 weeks <b>or</b> Doxycycline 200 mg (PO) q12h × 3 days, then 100 mg (PO) q12h × 11 days

## Acute Nonbacterial Meningitis/Chronic Meningitis (cont'd)

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	PO Therapy or IV-to-PO Switch
Granulomatous amebic meningoencephalitis	Acanthamoeba	No proven treatment (amphotericin B, fluconazole, ketoconazole, itraconazole, flucytosine, rifampin, isoniazid, aminoglycosides, sulfonamides, pentamidine mostly with little success. Success reported in transplant recipient with IV pentamidine followed by itraconazole, and in AIDS patient with ketoconazole plus flucytosine)		
Neurobrucellosis	Brucella sp.	Ceftriaxone 2 gm (IV) q12h <b>plus</b> Doxycycline 100 mg (PO) q12h <b>plus</b> Rifampin 600 mg (PO) q24h × 4 weeks, then Doxycycline <b>plus</b> Rifampin (same doses) × 4 weeks		
TB	M. tuberculosis	INH 300 mg (IV/PO) q24h × 6 months <b>plus</b> Rifampin 600 mg (IV/PO) q24h × 6 months <b>plus</b> PZA 25 mg/kg (PO) q24h × 2 months <b>plus</b> EMB 15 mg/kg (PO) q24h (until susceptibilities known)  Begin with 4 drugs, as for pulmonary TB, but extend treatment for 9–12 months. Corticosteroids tapered over 1–2 months is helpful. If resistance is demonstrated or suspected, ID consultation is advised		
Fungal <i>Non-HIV</i>	Cryptococcus neoformans	Amphotericin B 0.7–1 mg/kg (IV) q24h × 2–6 weeks* <b>plus</b> Flucytosine 25 mg/kg (PO) q6h × 6 weeks*, <b>followed by</b> Fluconazole 800 mg (IV or PO) × 1 dose, then 400 mg (PO) q24h × 10 weeks*	Liposomal amphotericin B (AmBisome) 6 mg/kg (IV) q24h × 6–10 weeks† <b>or</b> Fluconazole 800 mg (IV or PO) × 1 dose, then 400 mg (PO) q24h × 10 weeks*	<u>PO therapy alone</u> Fluconazole may be given PO*
<i>HIV</i>	Cryptococcus neoformans (see p. 329) Blastomyces dermatitidis (see p. 269)			
Chronic meningitis	Coccidioidomycosis (see p. 271), Histoplasmosis (see p. 269), Neurocysticercosis (see p. 262)			Do not treat empirically

\* The 6-week amphotericin B deoxycholate plus flucytosine regimen is the classical approach to achieving a durable cure. Anecdotal data suggest that a shorter course followed by Fluconazole × 10 weeks (or longer) may be successful. ID consultation is advised.

† Other lipid-associated Amphotericin B formulations at 3–5 mg/kg (IV) q24h may also be used (see p. 525).

‡ Levofloxacin 500 mg or Moxifloxacin 400 mg.

### **Viral (Aseptic) Meningitis**

**Clinical Presentation:** Headache, low-grade fever, mild meningismus, photophobia. Natalizumab (NTZ) predisposes to HSV and VZV meningitis or meningoencephalitis.

**Diagnostic Considerations:** Diagnosis by specific serological tests/viral culture. HSV-2 genital infections are often accompanied by mild CNS symptoms, which usually do not require anti-viral therapy. HSV-1, usually more severe, causes meningitis, meningoencephalitis, or encephalitis. HSV HHV-6, and WNE meningitis is indistinguishable clinically from other causes of viral meningitis. EBV meningitis is usually associated with clinical/laboratory features of EBV infectious mononucleosis; suspect the diagnosis in a patient with a positive monospot and unexplained meningoencephalitis. VZV meningitis is typically associated with cutaneous vesicular lesions (H. zoster). VZV meningitis may be later (weeks/months) followed by a CVA. LCM meningitis begins as a “flu-like” illness usually in the fall after hamster contact, and may have low CSF glucose. Enterovirus meningitis is often associated with a maculopapular rash, non-exudative pharyngitis, diarrhea, and rarely low CSF glucose. Aseptic meningitis due to mumps may present without parotid swelling  $\pm$  acute deafness. Drug induced aseptic meningitis may have eosinophils in the CSF and little/no fever.

**Pitfalls:** Consider NSAIDs, TMP-SMX, and IV immunoglobulin as causes of drug induced aseptic meningitis.

**Therapeutic Considerations:** Treat specific pathogen.

**Prognosis:** Without neurological deficits, full recovery is the rule.

### **Primary Amebic Meningoencephalitis (PAM) (Naegleria fowleri)**

**Clinical Presentation:** Acquired by freshwater exposure containing the protozoa, often by jumping into a lake/pool. Affects healthy children/young adults. Organism penetrates cribriform plate and enters CSF. Symptoms occur within 7 days of exposure and may be indistinguishable from fulminant bacterial meningitis, including headache, fever, anorexia, vomiting, signs of meningeal inflammation, altered mental status, coma. May complain of unusual smell/taste sensations early in infection. CSF has RBCs and very low glucose.

**Diagnostic Considerations:** Diagnosis by demonstrating organism in CSF. Worldwide distribution. Free-living freshwater amoeba flourish in warmer climates. Key to diagnosis rests on clinical suspicion based on history of freshwater exposure in previous 1–2 weeks.

**Pitfalls:** CSF findings resemble bacterial meningitis, but RBCs present.

**Therapeutic Considerations:** Often fatal despite early treatment.

**Prognosis:** Frequently fatal.

### **Granulomatous Amebic Meningoencephalitis (Acanthamoeba sp.)**

**Clinical Presentation:** Insidious onset with focal neurologic deficits  $\pm$  mental status changes, seizures, fever, headache, hemiparesis, meningismus, ataxia, visual disturbances. May be associated with Acanthamoeba keratoconjunctivitis, skin ulcers, or disseminated disease. Usually seen only in immunocompromised/debilitated patients.

**Diagnostic Considerations:** Diagnosis by demonstrating organism in brain biopsy specimen. CT/MRI shows mass lesions. “Stellate cysts” characteristic of Acanthamoeba. Worldwide distribution. Strong association with extended wear of contact lenses. Differentiate from *N. fowleri* by culture.

**Pitfalls:** Not associated with freshwater exposure, unlike primary amebic meningoencephalitis (*Naegleria fowleri*). Resembles subacute/chronic meningitis. No trophozoites in CSF. Skin lesions may be present for months before onset of CNS symptoms.

**Therapeutic Considerations:** No proven treatment. Often fatal despite early treatment.

**Prognosis:** Usually fatal.

### **Neuroborreliosis (CNS Lyme Disease)**

**Clinical Presentation:** May have mild neck stiffness not frank nuchal rigidity. Low grade fever < 102°F usually ~ 100–101°F. Recent tick exposure. Often without rash. Sore throat in some, WBC count WNL, mildly ↑ AST/ALT in some.

**Diagnostic Considerations:** Diagnosed by demonstrating *B. burgdorferi* antibody production in the CNS. A CSF-to-serum IgM Lyme titer ratio > 1:1 is diagnostic.

**Pitfalls:** If CNS symptoms present with acute Lyme disease, rash may be absent and serology may be negative early. Neuroborreliosis cannot be reliably diagnosed using CSF-to-serum IgM if antibiotics given before the LP (antibiotics change the ratio and render it useless).

**Therapeutic Considerations:**

**Prognosis:** Related to underlying health of host. Prognosis with neuroborreliosis is good even with delayed treatment.

### **TB Meningitis (Mycobacterium tuberculosis)**

**Clinical Presentation:** Subacute onset of nonspecific symptoms. Fever usually present ± headache, nausea, vomiting. Acute presentation and cranial nerve palsies uncommon.

**Diagnostic Considerations:** Diagnosis by CSF AFB smear/culture; PCR of CSF is specific but not very sensitive. CSF may be normal but usually shows a mild lymphocytic pleocytosis, ↓ glucose, ↑ protein, and few RBCs. Adenosine deaminase (ADA) levels ↑.

**Pitfalls:** CSF may have PMN predominance early, before developing typical lymphocytic predominance. Eosinophils in CSF is not a feature of TB, and should suggest another diagnosis. Chest x-ray, PPD, and CSF smear/culture may be negative. ADA levels more highly elevated in TB than in ABM, listeria, neurobrucellosis.

**Therapeutic Considerations:** Dexamethasone 4 mg (IV or PO) q6h × 4–8 weeks is useful to reduce CSF inflammation if given early. Proteinaceous TB exudates may obstruct ventricles and cause hydrocephalus, which is diagnosed by CT/MRI and may require shunting.

**Prognosis:** Poor prognostic factors include delay in treatment, neurologic deficits, or hydrocephalus.

### **Fungal (Cryptococcal) Meningitis (Cryptococcus neoformans)**

**Clinical Presentation:** Insidious onset of nonspecific symptoms. Headache most common. Chronic cases may have CNS symptoms for weeks to months with intervening asymptomatic periods. Acute manifestations are more common in AIDS, chronic steroid therapy, lymphoreticular malignancies. 50–80% of patients are abnormal hosts.

**Diagnostic Considerations:** *C. neoformans* is the most common cause of fungal meningitis, and the only encapsulated yeast in the CSF to cause meningitis. Diagnosis by CSF India ink preparations showing encapsulated yeasts/cryptococcal latex antigen by latex agglutination/culture. Rule out HIV and other underlying immunosuppressive diseases.

**Pitfalls:** False + cryptococcal antigen may occur with BBL Port-A-Cul transport vials. CSF latex antigen titer may not return to zero. Continue treatment until titers decline/do not decrease further, and until CSF culture is negative for cryptococci. India ink preparations showing encapsulated yeasts of CSF are useful for initial infection, but should not be relied on to diagnose recurrent episodes, since smears may be positive despite negative CSF cultures (dead cryptococci may remain in CSF for years). Diagnosis of recurrences rests on CSF culture.

**Therapeutic Considerations:** Treat until CSF is sterile or CSF latex antigen titer is zero or remains near zero on serial lumbar punctures. After patient defervesces on amphotericin B/5FC, switch to oral fluconazole × 10 weeks. Lipid amphotericin may be used if amphotericin B cannot be tolerated. HIV patients require life-long suppressive therapy with fluconazole 200 mg (PO) q24h. Expert consultation is strongly advised.

**Prognosis:** Good. Poor prognostic factors include no CSF pleocytosis, many organisms in CSF, and altered consciousness on admission.

## Encephalitis

Subset	Usual Pathogens	IV-to-PO Switch		
Herpes simplex	HSV-1 HSV-2	<u>IV Therapy:</u> Acyclovir 10 mg/kg (IV) q8h × 10 days* <b>or</b> Ganciclovir 5 mg/kg (IV) q12h × 10 days.* <u>IV → PO:</u> Valacyclovir 1 gm (PO) q8h × 10 days*		
Herpes zoster (VZV)	VZV	<u>IV Therapy:</u> Acyclovir (IV) q8h 10 mg/kg × 10 days.* <u>IV → PO:</u> Valacyclovir 2 gm (PO) q6h × 10 days*		
	HHV-6	Ganciclovir 5 mg/kg (IV) q12h × 2 weeks <b>or</b> Foscarnet 90 mg/kg (IV) q12h × 2 weeks		
Arbovirus	<u>Mosquito borne:</u> California encephalitis (CE), Western equine encephalitis (WEE), Venezuelan equine encephalitis (VEE), Eastern equine encephalitis (EEE), St. Louis encephalitis (St. LE), Japanese encephalitis (JE), West Nile encephalitis (WNE) <u>Tick borne:</u> Powassan (PE), TBE (RSSE/CCE) <u>IV/PO Therapy:</u> None			
Mycoplasma	M. pneumoniae	Doxycycline 200 mg (IV) q12h × 3 days, then 100 mg (IV) q12h × 2 weeks	Minocycline 100 mg (IV) q12h × 2 weeks	Doxycycline 200 mg (PO) q12h × 3 days, then 100 mg (PO) q12h × 2 weeks <b>or</b> Minocycline 100 mg (PO) q12h × 2 weeks
Listeria	L. monocytogenes	Treat as for Listeria meningitis (see p. 18) but treat for 6–8 weeks		
HIV	see Chapter 5 (p. 304)			
Transplants, HIV	CMV HHV-6 Toxoplasma gondii	Transplants (see p. 160) HIV (see p. 304)		

\* Severe HSV encephalitis may require a longer duration of therapy, i.e., 14–21 days.



### **Herpes Encephalitis (HSV-1)**

**Clinical Presentation:** Acute onset of fever and change in mental status without nuchal rigidity. Natalizumab (NTZ) predisposes to HSV meningitis, meningoencephalitis, or encephalitis.

**Diagnostic Considerations:** EEG is best early (< 72 hours) presumptive test, showing unilateral temporal lobe abnormalities. Brain MRI is abnormal before CT scan, which may require several days before a temporal lobe focus is seen. Definitive diagnosis is by CSF PCR for HSV-1 DNA. Usually presents as encephalitis or meningoencephalitis; RBCs in CSF ~ CNS damage Early/mild HSE may have no RBCs in CSF. CSF may have PMN predominance and low glucose levels, unlike other viral causes of meningitis.

**Pitfalls:** HSV-2 HSE rare in young adults; HSV-2 aseptic meningitis common HSV-2 HSE occurs in the elderly/immunosuppres. In normal hosts, HHV-6 may mimic HSV-1 encephalitis with a frontal/temporal focus on EEG.

**Therapeutic Considerations:** HSV is treatable. Treat as soon as possible, since neurological deficits may be mild and reversible early on, but severe and irreversible later.

**Prognosis:** Related to extent of brain injury and early anti-HSV therapy.

### **Arboviral Encephalitis**

**Clinical Presentation:** Acute onset of fever, headache, change/decrease in mental status days to weeks after insect bite (e.g., mosquito/tick). WNE is suggested by encephalitis ± with highly elevated ferritin levels ± new onset tremors or flaccid paralysis. EEG: diffuse slowing ± ↑ activity basal ganglia/thalamus. Prognosis ~ severity of neurologic deficits and degree/duration of relative lymphopenia/ferritin levels.

**Diagnostic Considerations:** Diagnosis by specific arboviral serology.

**Pitfalls:** Usually occurs in summer/fall. SIADH common, may occur.

**Therapeutic Considerations:** Only supportive therapy is available at present.

**Prognosis:** Permanent neurological deficits are common, but not predictable. May be fatal.

### **Mycoplasma Encephalitis**

**Clinical Presentation:** Acute onset of fever and change in mental status without nuchal rigidity.

**Diagnostic Considerations:** Diagnosis suggested by CAP with sore throat, otitis, E. multiforme, soft stools/diarrhea—with elevated IgM mycoplasma titers, and very high (≥ 1:1024) cold agglutinin titers. CSF shows mild mononucleosis/pleocytosis and normal/low glucose.

**Pitfalls:** CNS findings may overshadow pulmonary findings.

**Therapeutic Considerations:** Macrolides not effective for CNS infection. Doxycycline rapidly ↓ mycoplasma shedding in oropharyngeal secretions (vs. prolonged shedding with macrolides).

**Prognosis:** With early treatment, prognosis is good without neurologic sequelae.

### **Listeria Encephalitis**

**Clinical Presentation:** Fever/mental confusion (encephalitis/cerebritis) ± nuchal rigidity (meningoencephalitis).

**Diagnostic Considerations:** Diagnosis by LP. Suspect Listeria if CSF with “purulent profile” plus RBCs. Listeria seen on Gram stain of CSF in 50%; but culture positive in 100%.

**Pitfalls:** Not to be confused with other gram-positive bacilli (skin contaminants) isolated from CSF (diphtheroids). Listeria are motile and hemolytic on blood agar.

**Therapeutic Considerations:** 3<sup>rd</sup> generation cephalosporins not active against Listeria. In penicillin-tolerant patients, use meningeal dose ampicillin (not penicillin).

**Prognosis:** Good with early/adequate treatment. Related to degree of T lymphocyte dysfunction.

**CMV Encephalitis** (see p. 329)**Toxoplasma Encephalitis** (see p. 331)**Brain Abscess/Subdural Empyema/Cavernous Vein Thrombosis/Intracranial Suppurative Thrombophlebitis**

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	IV-to-PO Switch
<b>Brain Abscess (Single Mass Lesion)</b>				
Open trauma	S. aureus (MSSA) Enterobacteriaceae P. aeruginosa	Meropenem 2 gm (IV) q8h × 2 weeks <b>or</b> Cefepime 2 gm (IV) q8h × 2 weeks		Not applicable
Neurosurgical procedure  (Treat initially for MSSA; if later identified as MRSA, MSSE, or MRSE, treat accordingly)	S. aureus S. epidermidis (CoNS)	<u>MSSA</u> Nafcillin 2 gm (IV) q4h × 2 weeks <b>or</b> <u>MRSA/MSSE/MRSE</u> Linezolid 600 mg (IV) q12h × 2 weeks <b>or</b> Minocycline 100 mg (IV) q12h × 2 weeks	<u>MSSA/MRSA/MSSE/MRSE</u> Linezolid 600 mg (PO) q12h × 2 weeks <b>or</b> Minocycline 100 mg (PO) q12h × 2 weeks	
Mastoid/otitic source	Enterobacter Proteus	Treat the same as for open trauma (see above)		
Dental source	Oral anaerobes Actinomyces	Ceftriaxone 2 gm (IV) q12h × 2 weeks <b>plus</b> Metronidazole 500 mg (IV) q12h × 2 weeks		Ceftizoxime 3 gm (IV) q6h × 2 weeks
Subdural empyema/sinus source	Oral anaerobes H. influenzae	Treat the same as for dental source (see above)		
Cardiac source (ABE; right-to-left shunt)	S. aureus (MSSA) S. pneumoniae H. influenzae	Ceftriaxone 2 gm (IV) q12h × 2 weeks <b>or</b> Meropenem 2 gm (IV) q8h × 2 weeks		
	S. aureus (MRSA)	Linezolid 600 mg (IV) q12h × 2 weeks <b>or</b> Vancomycin 2 gm (IV) q12h × 2 weeks <b>or</b> Minocycline 100 mg (IV) q12h × 2 weeks		Linezolid 600 mg (PO) q12h × 2 weeks <b>or</b> Minocycline 100 mg (PO) q12h × 2 weeks

## Brain Abscess/Subdural Empyema/Cavernous Vein Thrombosis/Intracranial Suppurative Thrombophlebitis (cont'd)

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	IV-to-PO Switch
<b>Brain Abscess (Single Mass Lesion)</b>				
Pulmonary source	Oral anaerobes Actinomyces	Ceftriaxone 2 gm (IV) q12h × 2 weeks <b>plus</b> Metronidazole 500 mg (IV) q12h × 2 weeks		Ceftizoxime 3 gm (IV) q6h × 2 weeks

MSSA/MRSA = methicillin-sensitive/resistant *S. aureus*; MSSE/MRSE = methicillin-sensitive/resistant *S. epidermidis*. Duration of therapy represents total time IV or IV + PO. Most patients on IV therapy able to take PO meds should be switched to PO therapy after clinical improvement.

**Clinical Presentation:** Variable presentation, with fever, change in mental status, cranial nerve abnormalities ± headache.

**Diagnostic Considerations:** Diagnosis by CSF gram stain/culture. If brain abscess is suspected, obtain head CT/MRI. Lumbar puncture may induce herniation.

**Pitfalls:** CSF analysis is negative for bacterial meningitis unless abscess ruptures into ventricular system.

**Therapeutic Considerations:** Treatment with meningeal doses of antibiotics is required. Large single abscesses may be surgically drained; multiple small abscesses are best treated medically.

**Prognosis:** Related to underlying source and health of host.

### **Brain Abscess (Mastoid/Otitic Source)**

**Diagnostic Considerations:** Diagnosis by head CT/MRI demonstrating focus of infection in mastoid.

**Pitfalls:** Rule out associated subdural empyema.

**Therapeutic Considerations:** ENT consult for possible surgical debridement of mastoid.

**Prognosis:** Good. May require mastoid debridement for cure.

### **Brain Abscess (Dental Source)**

**Diagnostic Considerations:** Diagnosis by Panorex x-rays/gallium scan of jaw demonstrating focus in mandible/erosion into sinuses.

**Pitfalls:** Apical root abscess may not be apparent clinically.

**Therapeutic Considerations:** Large single abscess may be surgically drained. Multiple small abscesses are best treated medically. Treat until lesions on CT/MRI resolve or do not become smaller on therapy.

**Prognosis:** Good if dental focus is removed.

### **Brain Abscess (Subdural Empyema/Sinus Source)**

**Diagnostic Considerations:** Diagnosis by sinus films/CT/MRI to confirm presence of sinusitis/bone erosion (cranial osteomyelitis/epidural abscess). Usually from paranasal sinusitis.

**Pitfalls:** Do not overlook underlying sinus infection, which may need surgical drainage.

**Therapeutic Considerations:** Obtain ENT consult for possible surgical debridement of sinuses.

**Prognosis:** Good prognosis if sinus is drained.

### **Brain Abscess (Cardiac Source; Acute Bacterial Endocarditis)**

**Diagnostic Considerations:** Blood cultures often positive if brain abscess due to acute bacterial endocarditis (ABE) pathogen. Head CT/MRI shows multiple mass lesions.

**Pitfalls:** Do not overlook right-to-left cardiac shunt (e.g., patent foramen ovale, atrial septal defect) as source of brain abscess. Cerebral embolization results in aseptic meningitis in SBE, but septic meningitis/brain abscess in ABE (due to high virulence of pathogens).

**Therapeutic Considerations:** Multiple lesions suggest hematogenous spread. Use susceptibility of blood culture isolates to determine coverage. Meningeal doses are the same as endocarditis doses.

**Prognosis:** Related to location/size of CNS lesions and extent of cardiac valvular involvement.

### **Brain Abscess (Pulmonary Source)**

**Diagnostic Considerations:** Diagnosis suggested by underlying bronchiectasis, empyema, cystic fibrosis, or lung abscess in a patient with a brain abscess.

**Pitfalls:** Brain abscesses are most often due to chronic suppurative lung disease (e.g., bronchiectasis, lung abscess/empyema), not chronic bronchitis/AECB. If patient doesn't have a cardiopulmonary disorder predisposing to brain abscess, look for a clinically silent dental (apical root) abscess by gallium scan of jaws/Panorex dental x-rays.

**Therapeutic Considerations:** Lung abscess may need surgical drainage.

**Prognosis:** Related to extent/location of CNS lesions, drainage of lung abscess/empyema, and control of lung infection.

## Empiric Therapy of HEENT Infections

### Facial/Periorbital Cellulitis

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	PO Therapy or IV-to-PO Switch
Facial cellulitis	Group A streptococci H. influenzae	Ceftriaxone 1 gm (IV) q24h × 2 weeks <b>or</b> Cefotaxime 2 gm (IV) q6h × 2 weeks <b>or</b> Ceftizoxime 2 gm (IV) q8h × 2 weeks	Respiratory quinolone* (IV) q24h × 2 weeks	Any oral 2 <sup>nd</sup> or 3 <sup>rd</sup> gen. cephalosporin × 2 weeks <b>or</b> Respiratory quinolone* (PO) q24h × 2 weeks

Duration of therapy represents total time IV, PO, or IV + PO. Most patients on IV therapy able to take PO meds should be switched to PO therapy soon after clinical improvement (usually < 72 hours).

\* Levofloxacin 500 mg or Moxifloxacin 400 mg.

**Clinical Presentation:** Acute onset of warm, painful, facial rash without discharge, swelling, pruritus.

**Diagnostic Considerations:** Diagnosis by clinical appearance. May spread rapidly across face. Purplish hue suggests *H. influenzae*.

**Pitfalls:** If periorbital cellulitis, obtain head CT/MRI to rule out underlying sinusitis/CNS involvement. If secondary to an abrasion after contact with a saliva-contaminated surface, consider *Herpes gladiatorum*; lesions are painful/edematous and do not respond to antibiotic therapy for cellulitis.

**Therapeutic Considerations:** May need to treat × 3 weeks in compromised hosts (chronic steroids, diabetics, SLE, etc.).

**Prognosis:** Good with early treatment; worse if underlying sinusitis/CNS involvement.

## Bacterial Sinusitis

Subset	Usual Pathogens	IV Therapy (Hospitalized)	PO Therapy or IV-to-PO Switch (Ambulatory)
Acute	<i>S. pneumoniae</i> <i>H. influenzae</i> <i>M. catarrhalis</i>	Respiratory quinolone <sup>†</sup> (IV) q24h × 1–2 weeks <b>or</b> Ceftriaxone 1 gm (IV) q24h × 1–2 weeks <b>or</b> Doxycycline 200 mg (IV) q12h × 3 days, then 100 mg (IV) q12h × 11 days	Respiratory quinolone <sup>†</sup> (PO) q24h × 1–2 weeks <b>or</b> Amoxicillin 1gm (PO) q8h × 10 days <b>or</b> Doxycycline 200 mg (PO) q12h × 3 days, then 100 mg (PO) q12h × 11 days* <b>or</b> Cephalosporin <sup>†</sup> (PO) × 2 weeks <b>or</b> Clarithromycin XL 1 gm (PO) q24h × 2 weeks
Chronic	Same as acute + oral anaerobes	Requires prolonged antimicrobial therapy (2–4 weeks)	

Duration of therapy represents total time IV, PO, or IV + PO in adults. Most patients on IV therapy able to take PO meds should be switched to PO therapy soon after clinical improvement (usually < 72 hours). See Chapter 7 for pediatric therapy.

\* Loading dose is not needed PO if given IV with the same drug.

† Moxifloxacin 400 mg × 7 days or Levofloxacin 750 mg × 5 days.

‡ Cefdinir 300 mg q12h or Cefditoren 400 mg q12h or Cefixime 400 mg q12h or Cefpodoxime 200 mg q12h.

## Acute Bacterial Sinusitis

**Clinical Presentation:** Nasal discharge and cough frequently with headache, facial pain, and low-grade fever lasting > 10–14 days. Can also present acutely with high fever (≥ 104°F) and purulent nasal discharge ± intense headache lasting for ≥ 3 days. Other manifestations depend on the affected sinus: maxillary sinus: percussion tenderness of molars; maxillary toothache; local extension may cause osteomyelitis of facial bones with proptosis, retroorbital cellulitis, ophthalmoplegia; direct intracranial extension is rare; frontal sinus: prominent headache; intracranial extension may cause epidural/brain abscess, meningitis, cavernous sinus/superior sagittal sinus thrombosis; orbital extension may cause periorbital cellulitis; ethmoid sinus: eyelid edema and prominent tearing; extension may cause retro-orbital pain/periorbital cellulitis and/or cavernous sinus/superior sagittal sinus thrombosis; sphenoid

**sinus:** severe headache; extension into cavernous sinus may cause meningitis, cranial nerve paralysis (III, IV, VI), temporal lobe abscess, cavernous sinus thrombosis). Cough and nasal discharge are prominent in children.

**Diagnostic Considerations:** Diagnosis by sinus x-rays or CT/MRI showing complete sinus opacification, air-fluid levels, mucosal thickening. Consider sinus aspiration in immunocompromised hosts or treatment failures. In children, acute sinusitis is a clinical diagnosis; imaging studies are not routine.

**Pitfalls:** May present as periorbital cellulitis (obtain head CT/MRI to rule out underlying sinusitis). If CT/MRI demonstrates “post-septal” involvement, treat as acute bacterial meningitis. In children, transillumination, sinus tenderness to percussion, and color of nasal mucus are not reliable indicators of sinusitis.

**Therapeutic Considerations:** Treat for full course to prevent relapses/complications. Macrolides and TMP-SMX may predispose to drug-resistant *S. pneumoniae* (DRSP), and  $\geq 30\%$  of *S. pneumoniae* are naturally resistant to macrolides. Consider local resistance rates before making empiric antibiotic selections.

**Prognosis:** Good if treated for full course. Relapses may occur with suboptimal treatment. For frequent recurrences, consider radiologic studies and ENT consultation.

### **Chronic Bacterial Sinusitis**

**Clinical Presentation:** Generalized headache, fatigue, nasal congestion, post-nasal drip lasting  $> 3$  months with little/no sinus tenderness by percussion. Local symptoms often subtle. Fever is uncommon.

**Diagnostic Considerations:** Sinus films and head CT/MRI are less useful than for acute sinusitis (chronic mucosal abnormalities may persist after infection is treated). Many cases of chronic maxillary sinusitis are due to a dental cause; obtain odontogenic x-rays if suspected.

**Pitfalls:** Clinical presentation is variable/nonspecific. Malaise and irritability may be more prominent than local symptoms. May be mistaken for allergic rhinitis. Head CT/MRI can rule out sinus tumor.

**Therapeutic Considerations:** Therapeutic failure/relapse is usually due to inadequate antibiotic duration, dose, or tissue penetration. Treat for full course. If symptoms persist after 4 weeks of therapy, refer to ENT for surgical drainage procedure.

**Prognosis:** Good if treated for full course. Relapses may occur with suboptimal treatment. For frequent recurrences, consider radiologic studies and ENT consultation.

## **Keratitis**

<b>Subset</b>	<b>Pathogens</b>	<b>Topical Therapy</b>
Bacterial	<i>S. aureus</i> <i>S. pneumoniae</i> <i>Bacillus cereus</i> <i>M. catarrhalis</i> <i>P. aeruginosa</i>	Antibacterial eyedrops (ciprofloxacin, ofloxacin, or tobramycin/bacitracin/polymyxin B) hourly while awake $\times 2$ weeks
Viral	HSV-1	Trifluridine 1% solution 1 drop hourly while awake $\times 2$ days, then 1 drop q6h $\times 14$ –21 days <b>or</b> viral ophthalmic topical ointment (e.g., vidarabine) at bedtime $\times 14$ –21 days

## Keratitis (cont'd)

Subset	Pathogens	Topical Therapy
Amebic	Acanthamoeba	Propamidine (0.1%), neomycin, gramicidin, or polymyxin B eyedrops hourly while awake × 1–2 weeks <b>or</b> polyhexamethylene biguanide (0.02%) eyedrops hourly while awake × 1–2 weeks <b>or</b> chlorhexidine (0.02%) eyedrops hourly while awake × 1–2 weeks

**Clinical Presentation:** Corneal haziness, infiltrates, or ulcers.

**Diagnosis:** Appearance of corneal lesions/culture.

### Bacterial Keratitis

**Diagnostic Considerations:** Usually secondary to eye trauma. Always obtain ophthalmology consult.

**Pitfalls:** Be sure to culture ulcer. Unusual organisms are common in eye trauma.

**Therapeutic Considerations:** Treat until lesions resolve. Ointment easier/lasts longer than solutions. Avoid topical steroids.

**Prognosis:** Related to extent of trauma/organism. *S. aureus*, *B. cereus*, *P. aeruginosa* have worst prognosis.

### Viral Keratitis (HSV-1)

**Diagnostic Considerations:** "Dendritic" corneal ulcers characteristic. Obtain ophthalmology consult.

**Pitfalls:** Small corneal ulcers may be missed without fluorescein staining.

**Therapeutic Considerations:** Treat until lesions resolve. Oral acyclovir is not needed. Avoid ophthalmic steroid ointment.

**Prognosis:** Good if treated early (before eye damage is extensive).

### Amebic Keratitis (Acanthamoeba)

**Diagnostic Considerations:** Usually associated with extended use of soft contact lenses. Corneal scrapings are positive with calcofluor staining. Acanthamoeba keratitis is painful with typical circular, hazy, corneal infiltrate. Always obtain ophthalmology consult.

**Pitfalls:** Do not confuse with HSV-1 dendritic ulcers. Avoid topical steroids.

**Therapeutic Considerations:** If secondary bacterial infection, treat as bacterial keratitis.

**Prognosis:** No good treatment. Poor prognosis.

## Conjunctivitis

Subset	Usual Pathogens	PO/Topical Therapy
Bacterial	M. catarrhalis H. influenzae S. pneumoniae N. meningitidis N. gonorrhoea	Antibacterial eyedrops (ciprofloxacin, ofloxacin, moxifloxacin, or tobramycin/bacitracin/polymyxin B) q12h × 1 week plus antibacterial ointment (same antibiotic) at bedtime × 1 week

## Conjunctivitis (cont'd)

Subset	Usual Pathogens	PO/Topical Therapy
Viral	Adenovirus	Not applicable
	VZV	Valacyclovir 1 gm (PO) q8h × 10–14 days <b>or</b> Famciclovir 500 mg (PO) q8h × 10–14 days
Chlamydial	C. trachomatis (trachoma)	Doxycycline 100 mg (PO) q12h × 1–2 weeks <b>or</b> Azithromycin 1 gm (PO) × 1 dose

### **Bacterial Conjunctivitis**

**Clinical Presentation:** Profuse, purulent exudate from conjunctiva.

**Diagnostic Considerations:** Reddened conjunctiva; culture for specific pathogen.

**Pitfalls:** Do not confuse with allergic conjunctivitis, which itches and has a clear discharge.

**Therapeutic Considerations:** Obtain ophthalmology consult. Ointment lasts longer in eye than solution. Do not use topical steroids without an antibacterial.

**Prognosis:** Excellent when treated early, with no residual visual impairment.

### **Viral Conjunctivitis**

#### **Adenovirus**

**Clinical Presentation:** Reddened conjunctiva, watery discharge, negative bacterial culture.

**Diagnostic Considerations:** Diagnosis by cloudy/steamy cornea with negative bacterial cultures. Clue is punctate infiltrates with a cloudy cornea. Extremely contagious; careful handwashing is essential. Obtain viral culture of conjunctiva for diagnosis.

**Pitfalls:** Pharyngitis a clue to adenoviral etiology (pharyngoconjunctival fever).

**Therapeutic Considerations:** No treatment available. Usually resolves in 1–2 weeks.

**Prognosis:** Related to degree of corneal haziness. Severe cases may take weeks to clear.

#### **VZV Ophthalmicus**

**Diagnostic Considerations:** Vesicles on tip of nose predict eye involvement.

**Pitfalls:** Do not miss vesicular lesions in external auditory canal in patients with facial palsy (Ramsey-Hunt Syndrome).

**Therapeutic Considerations:** Obtain ophthalmology consult. Topical steroids may be used if given with anti-VZV therapy.

**Prognosis:** Good if treated early with systemic antivirals.

### **Trachoma (C. trachomatis) Conjunctivitis**

**Diagnostic Considerations:** Diagnosis by direct fluorescent antibody (DFA)/culture of conjunctiva.

**Pitfalls:** Do not confuse bilateral, upper lid, granular conjunctivitis of Chlamydia with viral/bacterial conjunctivitis, which involves both upper and lower eyelids.

**Therapeutic Considerations:** Ophthalmic erythromycin treatment is useful for neonates. Single-dose azithromycin may be associated with recurrence.

**Prognosis:** Excellent with early treatment.



## Chorioretinitis

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	PO Therapy or IV-to-PO Switch
Viral	CMV	See p. 327		
Fungal†	Candida albicans	Fluconazole 800 mg (IV) × 1 dose, then 400 mg (IV) q24h × 2 weeks <sup>‡</sup> <b>or</b> Voriconazole 6 mg/kg (IV) q12h × 1 day, then 4 mg/kg (IV) q12h × 2 weeks <sup>‡</sup>	Amphotericin B 0.6 mg/kg (IV) q24h for total of 1 gm <b>or</b> Lipid amphotericin (see p. 525) (IV) q24h × 3 weeks <sup>‡</sup>	Fluconazole 800 mg (PO) × 1 dose, then 400 mg (PO) q24h × 2 weeks <sup>**</sup>
Protozoal	Toxoplasma gondii	<u>IV Therapy</u> Not applicable	<u>PO Therapy</u> Pyrimethamine 75 mg (PO) × 1 dose, then 25 mg (PO) q24h × 6 weeks <b>plus either</b> Sulfadiazine 1 gm (PO) q6h × 6 weeks <b>or</b> Clindamycin 300 mg (PO) q8h × 6 weeks	

Duration of therapy represents total time IV, PO, or IV + PO. Most patients on IV therapy able to take PO meds should be switched to PO therapy after clinical improvement.

† Treat IV or IV-to-PO switch.

\* Loading dose is not needed PO if given IV with the same drug.

‡ Duration of therapy is approximate and should be continued until resolution of infection. Close follow-up with an ophthalmologist is required.

### CMV Chorioretinitis (see p. 327)

#### **Candida Chorioretinitis**

**Clinical Presentation:** Small, raised, white, circular lesions on retina.

**Diagnostic Considerations:** Fundus findings similar to white, raised colonies on blood agar plates.

**Pitfalls:** Candida endophthalmitis signifies invasive/disseminated candidiasis.

**Therapeutic Considerations:** Treat as disseminated candidiasis.

**Prognosis:** Good with early treatment.

#### **Toxoplasma Chorioretinitis**

**Clinical Presentation:** Grey/black pigmentation of macula.

**Diagnostic Considerations:** Diagnosis by IgM IFA toxoplasmosis titers.

**Pitfalls:** Unilateral endophthalmitis usually indicates acquired toxoplasmosis; congenital toxoplasmosis is usually bilateral.

**Therapeutic Considerations:** Obtain ophthalmology consult. Treat only acute/active toxoplasmosis with visual symptoms; do not treat chronic chorioretinitis. Add folic acid 10 mg (PO) q24h to prevent folic acid deficiency.

**Prognosis:** Related to degree of immunosuppression.

## Endophthalmitis

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	PO Therapy or IV-to-PO Switch
Bacterial  (Treat initially for MSSA; if later identified as MRSA or S. epidermidis, treat accordingly)	Streptococci H. influenzae S. aureus (MSSA)	<b>Intravitreal injection:</b> Cefepime 2.25 mg/0.1 mL sterile saline × 1 dose; repeat × 1 if needed in 2–3 days <b>plus</b> <b>Subconjunctival injection:</b> Cefepime 100 mg/0.5 mL sterile saline q24h × 1–2 weeks <b>plus</b> Levofloxacin 500 mg (IV/PO) q12h × 1–2 weeks		
	S. aureus (MRSA) S. epidermidis (CoNS)	<b>Intravitreal injection:</b> Vancomycin 1 mg/0.1 mL sterile saline × 1 dose; repeat × 1 if needed in 2–3 days <b>plus</b> <b>Subconjunctival injection:</b> Vancomycin 2.5 mg/0.5 mL sterile saline q24h × 1–2 weeks <b>plus</b> Linezolid 600 mg (IV/PO) q12h × 1–2 weeks		
Fungal <sup>‡</sup>	C. albicans	Fluconazole 800 mg (IV) × 1 dose, then 400 mg (IV) q24h × 2 months	Voriconazole (see “usual dose,” p. 714) × 2 months	Fluconazole (same as IV dose)  <b>or</b> Voriconazole (see p. 714) × 2 months
	A. fumigatus A. flavus	Amphotericin B (intravitreal) 10 mcg (in 0.1 mL of saline ) q2–3 days	Voriconazole (see “usual dose,” p. 714) × 2 months	Voriconazole (see “usual dose,” p. 714) × 2 months
TB	M. tuberculosis	Treat the same as for TB pneumonia (see p. 53)		
Infected lens implant*  (Treat initially for MSSA; if later identified as MRSA, treat accordingly)	Enterobacteriaceae S. aureus (MSSA)	Meropenem 1 gm (IV) q8h × 2 weeks  <b>or</b> Cefepime 2 gm (IV) q8h × 2 weeks	Chloramphenicol 500 mg (IV) q6h × 2 weeks	Chloramphenicol 500 mg (PO) q6h × 2 weeks
	S. aureus (MRSA)	Linezolid 600 mg (IV) q12h × 2 weeks	Minocycline 100 mg (IV) q12h × 2 weeks	Linezolid 600 mg (PO) q12h × 2 weeks  <b>or</b> Minocycline 100 mg (PO) q12h × 2 weeks

MSSA/MRSA = methicillin-sensitive/resistant *S. aureus*. Duration of therapy represents total time IV, PO, or IV + PO. Most patients on IV therapy able to take PO meds should be switched to PO therapy after clinical improvement (usually < 72 hours).

\* Treat only IV or IV-to-PO switch.

‡ Duration of therapy is approximate and should be continued until resolution of disease. Close follow-up with an ophthalmologist is required.

### **Bacterial Endophthalmitis**

**Clinical Presentation:** Ocular pain/sudden vision loss.

**Diagnosis:** Post-op endophthalmitis occurs 1–7 days after surgery. Hypopyon seen in anterior chamber.

**Pitfalls:** Delayed-onset endophthalmitis may occur up to 6 weeks post-op. White intracapsular plaque is characteristic.

**Therapeutic Considerations:** Use steroids (dexamethasone 0.4 mg/0.1 mL intravitreal and 10 mg/1 mL subconjunctival) with antibiotics in post-op endophthalmitis. Also use systemic antibiotics for severe cases. Vitrectomy is usually necessary.

**Prognosis:** Related to pathogen virulence.

### **Fungal Endophthalmitis**

**Clinical Presentation:** Slow deterioration in visual acuity ± eye pain. N fever. Antecedent/concomitant TPN, central IV line, prolonged antibiotic/immunosuppressive therapy, steroids, IVDA, or post-cataract surgery.

**Diagnostic Considerations:** Blood cultures frequently positive in Candida endophthalmitis. Small/round white lesions near retinal vessels. Cultures of aqueous/vitreous humors diagnostic.

**Pitfalls:** Symptoms develop insidiously with little/no eye pain or fever. Blood cultures negative with Aspergillus endophthalmitis.

**Therapeutic Considerations:** Penetration of antibiotic into vitreous humor requires high lipid solubility and inflammation. IV/intravitreal/subconjunctival antimicrobial choice depends on physicochemical properties of drug selected. Vitrectomy preferred by some.

If using intravitreal amphotericin B, there is no benefit in adding systemic IV therapy. Voriconazole penetrates CSF/eye, but clinical experience is limited. Based on limited data, caspofungin appears to have poor penetration into the vitreous and should not be relied upon to treat fungal endophthalmitis.

**Prognosis:** Best results with vitrectomy plus intravitreal antifungal therapy.

### **Tuberculous (TB) Endophthalmitis**

**Clinical Presentation:** Raised retinal punctate lesions ± visual impairment.

**Diagnostic Considerations:** Signs of extraocular TB are usually present. Confirm diagnosis of miliary TB by liver/bone biopsy.

**Pitfalls:** TB endophthalmitis is a sign of disseminated TB.

**Therapeutic Considerations:** Treat as disseminated TB. Systemic steroids may be used if given with anti-TB therapy.

**Prognosis:** Related to degree of immunosuppression.

### **Infected Lens Implant**

**Diagnostic Considerations:** Diagnosis by clinical appearance and culture of anterior chamber. Obtain ophthalmology consult.

**Pitfalls:** Superficial cultures are inadequate. Anterior chamber aspirate may be needed for culture.

**Therapeutic Considerations:** Infected lens must be removed for cure.

**Prognosis:** Good with early lens removal and recommended antibiotics.

## External Otitis

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	Topical Therapy or IV-to-PO Switch
Benign	<i>P. aeruginosa</i>	Use otic solutions only (ofloxacin 0.3%, tobramycin, polymyxin B); apply ear drops q6h × 1 week		
Malignant	<i>P. aeruginosa</i>	<b>One "A" drug ± one "B" drug: "A" drugs:</b> Meropenem 1 gm (IV) q8h <b>or</b> Cefepime 2 gm (IV) q8h <b>or</b> Piperacillin 4 gm (IV) q8h × 4–6 weeks <b>"B" drugs:</b> Ciprofloxacin 400 mg (IV) q8h <b>or</b> Levofloxacin 750 mg (IV) q24h × 4–6 weeks <b>or</b> Amikacin 1 gm (IV) q24h × 4–6 weeks		Ciprofloxacin 750 mg (PO) q12h <b>or</b> Levofloxacin 750 mg (PO) q24h × 4–6 weeks

Duration of therapy represents total time topically (benign), or IV + PO (malignant). Most patients on IV therapy able to take PO meds should be switched to PO therapy soon after clinical improvement (usually < 72 hours).

### **Benign External Otitis (*Pseudomonas aeruginosa*)**

**Clinical Presentation:** Acute external ear canal drainage without perforation of tympanic membrane or bone involvement.

**Diagnostic Considerations:** Diagnosis suggested by external ear drainage after water exposure. Usually acquired from swimming pools ("swimmers ear"). Not an invasive infection.

**Pitfalls:** Be sure external otitis is not associated with perforated tympanic membrane, which requires ENT consultation and systemic antibiotics.

**Therapeutic Considerations:** Treat topically until symptoms/infection resolve.

**Prognosis:** Excellent with topical therapy.

### **Malignant External Otitis (*Pseudomonas aeruginosa*)**

**Clinical Presentation:** External ear canal drainage with bone involvement.

**Diagnostic Considerations:** Diagnosis by demonstrating *P. aeruginosa* in soft tissue culture from ear canal plus bone/cartilage involvement on x-ray. Usually affects diabetics. CT/MRI of head shows bony involvement of external auditory canal.

**Pitfalls:** Rare in non-diabetics.

**Therapeutic Considerations:** Requires surgical debridement plus antibiotic therapy for cure.

**Prognosis:** Related to control of diabetes mellitus.

## Acute Otitis Media

Subset	Usual Pathogens	IM Therapy	PO Therapy
Initial uncomplicated bacterial infection	<i>S. pneumoniae</i> <i>H. influenzae</i> <i>M. catarrhalis</i>	Ceftriaxone 50 mg/ kg (IM) × 1 dose	Amoxicillin 1 gm or 10 mg/kg (PO) q8h × 10 days  <b>or</b> Clarithromycin 7.5 mg/kg (PO) q12h × 10 days  <b>or</b> Azithromycin 10 mg/kg (PO) × 1 dose, then 5 mg/kg (PO) q24h × 4 days
Treatment failure or resistant organism*	MDRSP β-lactamase + <i>H. influenzae</i>	Ceftriaxone 50 mg/kg (IM) q24h × 3 doses	Amoxicillin/clavulanate ES-600 90 mg/kg/day (PO) in 2 divided doses × 10 days <sup>†</sup>  <b>or</b> Cephalosporin <sup>†</sup> (PO) × 10 days

MDRSP = multidrug-resistant *S. pneumoniae*. Pediatric doses are provided; acute otitis media is uncommon in adults. For chronic otitis media, prolonged antimicrobial therapy is required.

\* Treatment failure = persistent symptoms and otoscopy abnormalities 48–72 hours after starting initial antimicrobial therapy. For risk factors for MDRSP, see Therapeutic Considerations.

† ES-600 = 600 mg amoxicillin/5 mL. Cephalosporins: Cefuroxime axetil 15 mg/kg (PO) q12h or Cefdinir 7 mg/kg (PO) q12h or 14 mg/kg (PO) q24h or Cefpodoxime 5 mg/kg (PO) q12h may be used.

### Acute Otitis Media

**Clinical Presentation:** Fever, otalgia, hearing loss. Nonspecific presentation is more common in younger children (irritability, fever). Key to diagnosis is examination of the tympanic membrane. Uncommon in adults.

**Diagnostic Considerations:** Diagnosis is made by finding an opaque, hyperemic, bulging tympanic membrane with loss of landmarks and decreased mobility on pneumatic otoscopy.

**Pitfalls:** Failure to remove cerumen (inadequate visualization of tympanic membrane) and reliance on history of ear tugging/pain are the main factors associated with overdiagnosis of otitis media. Otitis media with effusion (i.e., tympanic membrane retracted or in normal position with decreased mobility or mobility with negative pressure; fluid present behind the drum but normal in color) usually resolves spontaneously and should not be treated with antibiotics.

**Therapeutic Considerations:** Risk factors for infection with MDRSP *S. pneumoniae* include antibiotic therapy in past 30 days, failure to respond within 48–72 hours of therapy, day care attendance, and antimicrobial prophylaxis. Macrolides and TMP–SMX may predispose to DRSP, and 25% of *S. pneumoniae* are naturally resistant to macrolides.

**Prognosis:** Excellent, but tends to recur. Chronic otitis, cholesteatomas, mastoiditis are rare complications. Tympanostomy tubes/adenoidectomy for frequent recurrences of otitis media are the leading surgical procedures in children.

## Mastoiditis

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	PO Therapy or IV-to-PO Switch
Acute	S. pneumoniae H. influenzae S. aureus (MSSA)	Ceftriaxone 1–2 gm (IV) q24h × 2 weeks <b>or</b> Cefotaxime 2 gm (IV) q6h × 2 weeks <b>or</b> Cefepime 2 gm (IV) q12h × 2 weeks	Moxifloxacin 400 mg (IV) q24h × 2 weeks <b>or</b> Levofloxacin 500 mg (IV) q24h × 2 weeks <b>or</b> Doxycycline 200 mg (IV) q12h × 3 days, then 100 mg (PO) q12h × 2 weeks	Moxifloxacin 400 mg (PO) q24h × 2 weeks <b>or</b> Levofloxacin 500 mg (PO) q24h × 2 weeks <b>or</b> Doxycycline 100 mg (PO) q12h × 2 weeks
Chronic	S. pneumoniae H. influenzae P. aeruginosa S. aureus (MSSA) Oral anaerobes	Meropenem 1 gm (IV) q8h × 4 weeks <b>or</b> Cefepime 2 gm (IV) q8h × 4 weeks	Moxifloxacin 400 mg (IV) × 4–6 weeks <b>or</b> Levofloxacin 750 mg (IV) q24h	Moxifloxacin 400 mg (PO) × 4–6 weeks

Duration of therapy represents total time IV, PO, or IV + PO. Most patients on IV therapy able to take PO meds should be switched to PO therapy soon after clinical improvement (usually < 72 hours).

### Acute Mastoiditis

**Clinical Presentation:** Pain/tenderness over mastoid with fever.

**Diagnostic Considerations:** Diagnosis by CT/MRI showing mastoid involvement.

**Pitfalls:** Obtain head CT/MRI to rule out extension into CNS presenting as acute bacterial meningitis.

**Prognosis:** Good if treated early.

### Chronic Mastoiditis

**Clinical Presentation:** Subacute pain/tenderness over mastoid with low-grade fever.

**Diagnostic Considerations:** Diagnosis by CT/MRI showing mastoid involvement. Rarely secondary to TB (diagnose by AFB smear/culture of bone biopsy or debrided bone).

**Pitfalls:** Obtain head CT/MRI to rule out CNS extension.

**Therapeutic Considerations:** Usually requires surgical debridement for cure. Should be viewed as chronic osteomyelitis. If secondary to TB, treat as skeletal TB.

**Prognosis:** Progressive without surgery. Poor prognosis with associated meningitis/brain abscess.

## Suppurative Parotitis

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	PO Therapy or IV-to-PO Switch
Parotitis (bacterial)	S. aureus (MSSA) Enterobacteriaceae Oral anaerobes	Meropenem 1 gm (IV) q8h × 2 weeks <b>or</b> Respiratory quinolone* (IV) q24h × 2 weeks	Ceftriaxone 1 gm (IV) q24h × 2 weeks <b>or</b> Ceftizoxime 2 gm (IV) q8h × 2 weeks	Respiratory quinolone* (PO) q24h × 2 weeks

Duration of therapy represents total time IV, PO, or IV + PO. Most patients on IV therapy able to take PO meds should be switched to PO therapy soon after clinical improvement (usually < 72 hours).

\* Levofloxacin 500 mg or Moxifloxacin 400 mg.

**Clinical Presentation:** Unilateral parotid pain/swelling with discharge from Stensen's duct ± fever.

**Diagnostic Considerations:** Diagnosis by clinical presentation, ↑ amylase, CT/MRI demonstrating stone in parotid duct/gland involvement.

**Pitfalls:** Differentiate from unilateral mumps by purulent discharge from Stensen's duct.

**Therapeutic Considerations:** If duct is obstructed, remove stone.

**Prognosis:** Good with early therapy/hydration.

## Pharyngitis

Subset	Usual Pathogens	PO Therapy
Bacterial	Group A streptococci	Amoxicillin 1 gm (PO) q8h × 7–10 days <b>or</b> Cefprozil 500 mg (PO) q12h × 7–10 days <b>or</b> Clindamycin 300 mg (PO) q8h × 7–10 days <b>or</b> Clarithromycin XL 1 gm (PO) q24h × 7–10 days <b>or</b> Azithromycin 500 mg (PO) × 1 dose, then 250 mg (PO) q24h × 4 days
Membranous	Arcanobacterium (Corynebacterium) hemolyticum	Doxycycline 100 mg (IV or PO) q12h × 1–2 weeks <b>or</b> Clarithromycin XL 1 gm (PO) q24h × 1–2 weeks <b>or</b> Azithromycin 500 mg (IV or PO) × 1 dose then 250 mg (IV or PO) q24h × 1–2 weeks <b>or</b> Cephalosporin (IV or PO) × 1–2 wks
	C. diphtheriae C. ulcerans	Diphtheria antitoxin (see p. 170) ± penicillin or macrolide
Viral	Respiratory viruses, EBV, CMV, HHV-6	Not applicable. EBV/CMV do not cause primary pharyngitis, but pharyngitis part of infectious mononucleosis syndrome, along with hepatitis and lymph node involvement. Viral hepatitis (see pp. 97–98)

## Pharyngitis (cont'd)

Subset	Usual Pathogens	PO Therapy
Other	M. pneumoniae C. pneumoniae	Respiratory quinolone* (PO) q24h × 1 week <b>or</b> Doxycycline 100 mg (PO) q12h × 1 week <b>or</b> Clarithromycin XL 1 gm (PO) q24h × 1 week <b>or</b> Azithromycin 500 mg (PO) × 1 dose, then 250 mg (PO) q24h × 4 days

Chronic fatigue syndrome.<sup>†</sup>

\* Levofloxacin 500 mg or Moxifloxacin 400 mg.

† Not a cause of pharyngitis, pathogen unknown (not EBV) but may be caused by xenotropic murine leukemia related virus (XMRV). For therapeutic considerations (see p. 43).

### Bacterial Pharyngitis

**Clinical Presentation:** Acute sore throat with fever, bilateral anterior cervical adenopathy, and elevated ASO titer. No hoarseness.

**Diagnostic Considerations:** Diagnosis of Group A streptococcal pharyngitis by elevated ASO titer after initial sore throat and positive throat culture. Rapid strep tests unnecessary, since delay in culture results (~1 week) still allows adequate time to initiate therapy and prevent acute rheumatic fever. Group A streptococcal pharyngitis is rare in adults > 30 years.

**Pitfalls:** Gram stain of throat exudate differentiates Group A streptococcal colonization (few or no PMNs) from infection (many PMNs) in patients with a positive throat culture or rapid strep test. Neither throat culture nor rapid strep test alone differentiates colonization from infection.

**Therapeutic Considerations:** Benzathine penicillin 1.2 mu (IM) × 1 dose can be used as an alternative to oral therapy. Penicillin, erythromycin, and ampicillin fail in 15% of cases due to poor penetration into oral secretions or beta-lactamase producing oral organisms.

**Prognosis:** Excellent. Treat within 10 days to prevent acute rheumatic fever.

### Membranous Pharyngitis

**Clinical Presentation:** A. hemolyticum presents with a scarlet fever-like rash and membranous pharyngitis. C. diphtheriae pharyngitis has no fever/rash.

**Diagnostic Considerations:** Diagnosis by throat culture/recovery of organism in patients with scarlatiniform rash.

**Pitfalls:** A. hemolyticum must be differentiated from C. diphtheriae on culture.

**Therapeutic Considerations:** For A. hemolyticum, doxycycline or erythromycin are more active than beta-lactams. Penicillin or macrolides are preferred for C. diphtheriae. C. diphtheriae should be treated with antitoxin ASAP since antibiotic therapy is adjunctive.

**Prognosis:** Related to degree of airway obstruction. Good with early treatment of A. hemolyticum. Diphtheria prognosis related to early antitoxin treatment and to presence/severity of toxic myocarditis.



### **Viral Pharyngitis**

**Clinical Presentation:** Acute sore throat. Other features depend on specific pathogen.

**Diagnostic Considerations:** Most cases of viral pharyngitis are caused by respiratory viruses, and are frequently accompanied by hoarseness, but not high fever, pharyngeal exudates, palatal petechiae, or posterior cervical adenopathy. Other causes of viral pharyngitis (EBV, CMV, HHV-6) are usually associated with posterior cervical adenopathy and ↑ SGOT/SGPT. EBV mono may present with exudative or non-exudative pharyngitis, and is diagnosed by negative ASO titer with a positive mono spot test or elevated EBV IgM viral capsid antigen (VCA) titer. Before mono spot test turns positive (may take up to 8 weeks), a presumptive diagnosis of EBV mono can be made by ESR and SGOT, which are elevated in EBV and normal in Group A streptococcal pharyngitis. If EBV mono spot is negative, retest weekly × 8 weeks; if still negative, obtain IgM CMV/toxoplasmosis titers to diagnose the cause of “mono spot negative” pharyngitis.

**Pitfalls:** 30% of patients with viral pharyngitis have Group A streptococcal colonization. Look for viral features to suggest the correct diagnosis (leukopenia, lymphocytosis, atypical lymphocytes).

**Therapeutic Considerations:** Symptomatic care only. Short-term steroids should only be used in EBV infection if airway obstruction is present/imminent. Since 30% of patients with viral pharyngitis are colonized with Group A streptococci, do not treat throat cultures positive for Group A streptococci if non-streptococcal pharyngitis features are present (e.g., bilateral posterior cervical adenopathy).

**Prognosis:** Related to extent of systemic infection. Post-viral fatigue is common. CMV may remain active in liver for 6–12 months with mildly elevated serum transaminases.

### **Mycoplasma/Chlamydia (Chlamydia) Pharyngitis**

**Clinical Presentation:** Acute sore throat ± laryngitis. Usually non-exudative.

**Diagnostic Considerations:** Diagnosis by elevated IgM M. pneumoniae or C. pneumoniae titers. Consider diagnosis in patients with non-exudative pharyngitis without viral or streptococcal pharyngitis. Mycoplasma pharyngitis is often accompanied by otitis/bullous myringitis.

**Pitfalls:** Patients with C. pneumoniae frequently have laryngitis, which is not a feature of EBV, CMV, Group A streptococcal, or M. pneumoniae pharyngitis.

**Therapeutic Considerations:** Treatment of C. pneumoniae laryngitis results in rapid (~ 3 days) return of normal voice, which does not occur with viral pharyngitis.

**Prognosis:** Excellent.

### **Chronic Fatigue Syndrome (CFS)**

**Clinical Presentation:** Fatigue > 1 year with cognitive impairment and no pharyngitis.

**Diagnostic Considerations:** Rule out other causes of chronic fatigue (cancer, adrenal/thyroid disease, etc.) before diagnosing CFS. HHV-6/Coxsackie B titers are usually elevated. Some have ↓ natural kill (NK) cells/activity. ESR ~ 0. Crimson crescents in the posterior pharynx are common.

**Pitfalls:** ↑ VCA IgG EBV titers is common in CFS, but EBV does not cause CFS. Do not confuse CFS with fibromyalgia, which has muscular “trigger points” and no cognitive impairment. CFS and fibromyalgia may coexist. In presumed CFS, if there is pharyngitis or if the ESR is elevated, consider an alternate diagnosis.

**Therapeutic Considerations:** No specific therapy is available. Patients with ↓ NK cells may benefit from β-carotene 50,000 U (PO) q24h × 3 weeks. Patients with ↑ C. pneumoniae IgG titers may benefit from doxycycline 100 mg (PO) q24h × 3 weeks.

**Prognosis:** Cyclical illness with remissions (precipitated by exertion, stress, or sleep lack). Avoid exercise.

## Thrush (Oropharyngeal Candidiasis)

Subset	Pathogens	PO Therapy
Fungal	C. albicans	Fluconazole 200 mg (PO) × 1 dose, then 100 mg (PO) q24h × <sup>†</sup> weeks <b>or</b> Posaconazole 100 mg (PO) q12h × 1 day, then 100 mg (PO) q24h × 13 days <b>or</b> Clotrimazole 10 mg troches (PO) 5x/day × <sup>†</sup> weeks <b>or</b> Itraconazole 200 mg (PO) q24h × <sup>†</sup> weeks
	Fluconazole-resistant Candida species	Posaconazole 400 mg (PO) q12h × <sup>†</sup> weeks* <b>or</b> Voriconazole 200 mg (PO) q12h × <sup>†</sup> weeks* <b>or</b> Caspofungin 70 mg (IV) × 1, then 50 mg (IV) q24h × <sup>†</sup> weeks* <b>or</b> Itraconazole 200 mg (PO) q12h × <sup>†</sup> weeks*

\* Duration of therapy depends on underlying disease and clinical response.

† See Fluconazole unresponsive esophagitis (see p. 612).

## Thrush (Oropharyngeal Candidiasis)

**Clinical Presentation:** White coated tongue or oropharynx. White adherent plaques may be on any part of the oropharynx.

**Diagnostic Considerations:** Gram stain/culture of white plaques demonstrates yeasts (Candida). Culture and susceptibility testing of causative fungus is useful in analyzing failure to respond to therapy.

**Pitfalls:** Lateral, linear, white, striated tongue lesions may resemble thrush but really represent hairy leukoplakia. Hairy leukoplakia should suggest HIV. Thrush may occur in children, alcoholics, diabetics, those receiving steroids/antibiotic therapy, or HIV.

**Therapeutic Considerations:** Almost all infections are caused by C. albicans, a species that is usually susceptible to fluconazole. Fluconazole-unresponsive infections may be caused by infection with fluconazole-resistant C. albicans (most common explanation), infection with a fluconazole-resistant non-albicans species (rare), noncompliance with therapy (common), or drug interactions (e.g., concomitant usage of rifampin, which markedly reduced azole blood levels). For suspected fluconazole-resistance, a trial with another azole is appropriate as cross-resistance is not universal.

**Prognosis:** Non-HIV patients respond well to therapy, particularly when the predisposing factor is eliminated/decreased (i.e., antibiotics discontinued, steroids reduced, etc.). HIV patients may require longer courses of therapy and should be treated until cured. Relapse is frequent in HIV patients, and institution of effective antiretroviral therapy is the most effective general strategy.

## Mouth Ulcers/Vesicles

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	PO Therapy or IV-to-PO Switch
Vincent's angina	Borrelia Fusobacterium	Clindamycin 600 mg (IV) q8h × 2 weeks	Ceftizoxime 2 gm (IV) q8h × 2 weeks <b>or</b> Any $\beta$ -lactam (IV) × 2 weeks	Clindamycin 300 mg (PO) q8h × 2 weeks <b>or</b> Amoxicillin/clavulanic acid 500/125 mg (PO) q8h × 2 weeks
Ludwig's angina	Group A streptococci	Clindamycin 600 mg (IV) q8h × 2 weeks	Ceftizoxime 2 gm (IV) q8h × 2 weeks <b>or</b> Any $\beta$ -lactam (IV) × 2 weeks	Clindamycin 300 mg (PO) q8h × 2 weeks <b>or</b> Amoxicillin/clavulanic acid 500/125 mg (PO) q8h × 2 weeks
Stomatitis	Normal mouth flora	Not applicable		
Herpangina	Coxsackie A virus	Not applicable		
Herpes gingivo-stomatitis	HSV-1	<u>PO Therapy</u> Valacyclovir 500 mg (PO) q12h × 1 week <b>or</b> Famciclovir 500 mg (PO) q12h × 1 week <b>or</b> Acyclovir 400 mg (PO) 5x/day × 1 week		
Herpes labialis (cold sores/fever blisters)	HSV-1 (recurrent)	<u>PO Therapy</u> Valacyclovir 2 gm (PO) q12h × 1 day (2 doses) started at onset of symptoms (tingling/burning) <b>or</b> Acyclovir 400 mg (PO) 5x/day × 1 week <b>or</b> Famciclovir 500 mg (PO) q12h × 1 week <u>Topical Therapy</u> Penciclovir 1% cream q2h while awake × 4 days <b>or</b> Acyclovir 5% cream 5x/d × 4 days		
Aphthous ulcers	Normal mouth flora	Not applicable		

Duration of therapy represents total time IV, PO, or IV + PO. Most patients on IV therapy able to take PO meds should be switched to PO therapy soon after clinical improvement (usually < 72 hours).

**Vincent's Angina (Borrelia/Fusobacterium)**

**Clinical Presentation:** Painful/bleeding gums without fever.

**Diagnostic Considerations:** Foul breath, poor dental hygiene/pyorrhea.

**Pitfalls:** Do not attribute foul breath to poor dental hygiene without considering other serious causes (e.g., lung abscess, renal failure).

**Therapeutic Considerations:** After control of acute infection, refer to dentist.

**Prognosis:** Excellent with early treatment.

**Ludwig's Angina (Group A streptococci)**

**Clinical Presentation:** Painful floor of mouth with fever.

**Diagnostic Considerations:** Elevated floor of mouth is diagnostic. Massive neck swelling may be present.

**Pitfalls:** C<sub>1q</sub> deficiency has perioral/tongue swelling, but no fever or floor of mouth elevation. Rarely, Ludwig's, angina may be due to F. necrophorum.

**Therapeutic Considerations:** Surgical drainage is not necessary. May need airway emergently; have tracheotomy set at bedside.

**Prognosis:** Early airway obstruction has adverse impact on prognosis.

**Stomatitis (normal mouth flora)**

**Clinical Presentation:** Painful mouth ulcers with fever.

**Diagnostic Considerations:** Diagnosis based on clinical appearance.

**Pitfalls:** Do not miss a systemic cause (e.g., acute leukemia).

**Therapeutic Considerations:** Painful; treat symptomatically.

**Prognosis:** Related to severity of underlying systemic disease.

**Herpangina (Coxsackie A virus)**

**Clinical Presentation:** Painful mouth ulcers/vesicles with fever.

**Diagnostic Considerations:** Ulcers located posteriorly in pharynx. No gum involvement or halitosis.

**Pitfalls:** Do not confuse with anterior vesicular lesions of HSV.

**Therapeutic Considerations:** No good treatment available. Usually resolves spontaneously in 2 weeks.

**Prognosis:** Good, but may be recurrent.

**Herpes Gingivostomatitis (HSV-1)**

**Clinical Presentation:** Painful gums with fever.

**Diagnostic Considerations:** Anterior ulcers in pharynx. Associated with bleeding gums, not halitosis.

**Pitfalls:** Do not miss a systemic disease associated with bleeding gums (e.g., acute myelogenous leukemia). Periodontal disease is not usually associated with oral ulcers.

**Therapeutic Considerations:** Oral analgesic solutions may help in swallowing.

**Prognosis:** Excellent with early treatment.

**Herpes Labialis (Cold Sores/Fever Blisters) (HSV-1)**

**Clinical Presentation:** Painful vesicles along vermilion border of upper or lower lip with fever.

**Diagnostic Considerations:** Caused by recurrent HSV-1 infection, which appears as painful vesicular lesions on/near the vermilion border of lips. Attacks may be triggered by stress, sun exposure, menstruation, and often begin with pain or tingling before vesicles appear. Vesicles crust over and attacks usually resolve by 1 week. "Fever blisters" are not triggered by temperature elevations per se, but may accompany malaria, pneumococcal/meningococcal meningitis. Diagnosis is clinical.

**Pitfalls:** Do not confuse with perioral impetigo; impetigo has crusts (not vesicles), is itchy (not painful), and does not involve the vermilion border of the lips.

**Therapeutic Considerations:** Treatment is not always needed. If valacyclovir is used, it should be started when pain/tingling appear to decrease symptoms/vesicles/duration of attack. Valacyclovir is of no proven value once vesicles have appeared. Non-prescription topical products (docosanol 10%,

tetracaine cream) may decrease pain/itching. Once-daily suppressive therapy may be considered for frequent recurrences or during times of increased risk (e.g., sun exposure). Sun screen may be helpful.

**Prognosis:** Tends to be recurrent in normal hosts. May be severe in compromised hosts.

### **Aphthous Ulcers (normal mouth flora)**

**Clinical Presentation:** Painful mouth ulcers/vesicles without fever.

**Diagnostic Considerations:** Usually an isolated finding. Ulcers are painful.

**Pitfalls:** May be a clue to systemic disorder, e.g., Behçet's syndrome. Mouth ulcers in SLE are painless.

**Therapeutic Considerations:** Usually refractory to all treatment and often recurrent. Steroid ointment (Kenalog) may be helpful.

**Prognosis:** Good, but tends to recur.

## **Deep Neck Infections, Lemierre's Syndrome, Severe Dental Infections**

<b>Subset</b>	<b>Usual Pathogens</b>	<b>Preferred IV Therapy</b>	<b>Alternate IV Therapy</b>	<b>IV-to-PO Switch</b>
Deep neck infections (lateral pharyngeal, retro-pharyngeal, space)	Oral anaerobes Oral streptococci	Meropenem 1 gm (IV) q8h × 2 weeks <b>or</b> Tigecycline 100 mg (IV) × 1 dose, then 100 mg (IV) q24h or 50 mg (IV) q12h × 2 weeks <b>or</b> Piperacillin/tazobactam 3.375 mg (IV) q6h × 2 weeks	Clindamycin 600 mg (IV) q8h × 2 weeks <b>or</b> Ceftizoxime 2 gm (IV) q8h × 2 weeks <b>or</b> Ertapenem 1 gm (IV) q24h × 2 weeks	Clindamycin 300 mg (PO) q8h × 2 weeks <b>or</b> Doxycycline 200 mg (PO) q12h × 3 days, then 100 mg (PO) q12h × 11 days*
Lemierre's Syndrome	Fusobacterium necrophorum	Treat as deep neck infection, see above		
Severe dental infections	Oral anaerobes Oral streptococci	Clindamycin 600 mg (IV) q8h × 2 weeks <b>or</b> Piperacillin/tazobactam 3.375 mg (IV) q6h × 2 weeks	Ertapenem 1 gm (IV) q24h × 2 weeks <b>or</b> Meropenem 1 gm (IV) q8h × 2 weeks	Clindamycin 300 mg (PO) q8h × 2 weeks <b>or</b> Doxycycline 200 mg (PO) q12h × 3 days, then 100 mg (PO) q12h × 11 days*

Duration of therapy represents total time IV or IV + PO. Most patients on IV therapy able to take PO meds should be switched to PO therapy after clinical improvement.

\* Loading dose is not needed PO if given IV with the same drug.

**Deep Neck Infections****(lateral pharyngeal, retropharyngeal)**

**Diagnostic Considerations:** Lateral pharyngeal space infection: Anterior → fever/chills, trismus, dysphagia, swelling or angle of mandible, ± parotid swelling, and bulging of lateral pharyngeal wall. Posterior pharyngeal space infection → bacteremic septic, no trismus/pain, and no pharyngeal wall bulging. Retropharyngeal space infection: fever/chills, dysphagia, dyspnea, ± mucal rigidity, esophageal regurgitation, and bulging of posterior pharyngeal wall.

**Pitfalls:** Retropharyngeal “danger space” infection may extend to mediastinum and present as mediastinitis.

**Therapeutic Considerations:** Obtain ENT consult for surgical drainage.

**Prognosis:** Worst prognosis with posterior lateral and retropharyngeal space infections.

**Lemierre’s Syndrome**

**Clinical Presentation:** Severe sore throat with fever, toxemic appearance, and tenderness over jugular vein. Jugular vein septic thrombophlebitis and septic pulmonary emboli.

**Diagnostic Considerations:** May present as multiple septic pulmonary emboli. Usually follows recent dental infection. Jugular vein tenderness diagnostic. Blood cultures positive for *F. necrophorum* or *F. nucleatum*; rarely MSSA/MRSA.

**Pitfalls:** Suspect Lemierre’s syndrome in patients with concurrent/antecedent dental/oropharyngeal infection with severe sore throat and jugular vein tenderness.

**Therapeutic Considerations:** If unresponsive to antibiotic therapy, may need venotomy.

**Prognosis:** Good with prompt treatment. May be more severe with concurrent *M. pneumoniae* or EBV pharyngitis.

**Severe Dental Infections**

**Diagnostic Considerations:** Obtain CT/MRI of jaws to rule out osteomyelitis or abscess.

**Pitfalls:** Chronic drainage in a patient with an implant is diagnostic of chronic osteomyelitis/abscess until proven otherwise.

**Therapeutic Considerations:** Abscesses should be drained for cure.

**Prognosis:** Poor prognosis and recurrent without adequate surgical drainage.

**Epiglottitis**

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	IV-to-PO Switch
Epiglottitis	<i>S. pneumoniae</i> <i>H. influenzae</i> Respiratory viruses	Ceftriaxone 1 gm (IV) q24h x 2 weeks <b>or</b> Ceftizoxime 2 gm (IV) q8h x 2 weeks	Ertapenem 1 gm (IV) q24h x 2 weeks <b>or</b> Meropenem 1 gm (IV) q8h x 2 weeks <b>or</b> Moxifloxacin 400 mg (IV) q24h x 2 weeks <b>or</b> Levofloxacin 500 mg (IV) q24h x 2 weeks	Moxifloxacin 400 mg (PO) q24h x 2 weeks <b>or</b> Levofloxacin 500 mg (PO) q24h x 2 weeks <b>or</b> Cefprozil 500 mg (PO) q12h x 2 weeks

Duration of therapy represents total time IV or IV + PO. Most patients on IV therapy able to take PO meds should be switched to PO therapy after clinical improvement.

**Clinical Presentation:** Stridor with upper respiratory infection.

**Diagnostic Considerations:** Lateral film of neck shows epiglottic edema. Neck CT/MRI may help if neck films are non-diagnostic.

**Pitfalls:** Do not attempt to culture the epiglottis (may precipitate acute upper airway obstruction).

**Therapeutic Considerations:** Treat empirically as soon as possible. Obtain ENT consult.

**Prognosis:** Early airway obstruction is associated with an adverse prognosis.

## Empiric Therapy of Lower Respiratory Tract Infections

### Acute Bacterial Exacerbation of Chronic Bronchitis (AECB)

Subset	Pathogens	PO Therapy
AECB	S. pneumoniae H. influenzae M. catarrhalis	Respiratory quinolone* (PO) q24h x 5 days <b>or</b> Amoxicillin/clavulanic acid 500/125 mg (PO) q12h x 5 days <b>or</b> Clarithromycin XL 1 gm (PO) q24h x 5 days <b>or</b> Doxycycline 100 mg (PO) q12h x 5 days <b>or</b> Azithromycin 500 mg (PO) x 3 days

\* Moxifloxacin 400 mg or Levofloxacin 500 mg.

**Clinical Presentation:** Productive cough and negative chest x-ray in a patient with chronic bronchitis.

**Diagnostic Considerations:** Diagnosis by productive cough, purulent sputum, and chest x-ray negative for pneumonia. H. influenzae is relatively more common than other pathogens.

**Pitfalls:** Do not obtain sputum cultures in chronic bronchitis; cultures usually reported as normal/mixed flora and should not be used to guide therapy.

**Therapeutic Considerations:** Treated with same antibiotics as for community-acquired pneumonia, since pathogens are the same (even though H. influenzae is relatively more frequent). Respiratory viruses/C. pneumoniae may initiate AECB, but is usually followed by bacterial infection, which is responsible for symptoms and is the aim of therapy. Bronchodilators are helpful for bronchospasm. Macrolide-resistant S. pneumoniae is an important clinical problem (prevalence  $\geq 30\%$ ).

**Prognosis:** Related to underlying cardiopulmonary status.

## Mediastinitis

Subset	Usual Pathogens	IV Therapy	IV-to-PO Switch
Following esophageal perforation or thoracic surgery	Oral anaerobes	<u>Preferred:</u> Piperacillin/tazobactam 3.375 gm (IV) q6h × 2 wks <b>or</b> Ampicillin/sulbactam 3 gm (IV) q6h × 2 wks <u>Alternate:</u> Meropenem 1 gm (IV) q8h × 2 weeks <b>or</b> Ertapenem 1 gm (IV) q24h × 2 weeks	Amoxicillin/clavulanate 500/125 mg (PO) q8h × 2 weeks <b>or</b> Respiratory quinolone* (PO) q24h × 2 weeks

Duration of therapy represents total time IV or IV + PO. Most patients on IV therapy able to take PO meds should be switched to PO therapy after clinical improvement.

\* Moxifloxacin 400 mg or Levofloxacin 500 mg.

**Diagnostic Considerations:** Chest x-ray usually shows perihilar infiltrate in mediastinitis. Pleural effusions from esophageal tears have elevated amylase levels.

**Pitfalls:** Do not overlook esophageal tear in mediastinitis with pleural effusions.

**Therapeutic Considerations:** Obtain surgical consult if esophageal perforation is suspected.

**Prognosis:** Related to extent, location, and duration of esophageal tear/mediastinal infection.

## Community-Acquired Pneumonia (CAP) (see Color Atlas for Sputum Gram stains and Chapter 8 for Differential Diagnosis of CXR patterns)

Subset	Usual Pathogens*	Alternate IV Therapy	PO Therapy or IV-to-PO Switch
Pathogen unknown	<i>S. pneumoniae</i> <sup>‡</sup> <i>H. influenzae</i> <i>M. catarrhalis</i> <i>B. pertussis</i> <i>Legionella</i> sp. <i>Mycoplasma pneumoniae</i>	Respiratory quinolone <sup>†</sup> (IV) q24h <b>or</b> <b>combination therapy with</b> Ceftriaxone 1 gm (IV) q24h × 1–2 week <b>plus either</b> Doxycycline <sup>‡</sup> (IV) × 1–2 weeks <b>or</b> Azithromycin 500 mg (IV) q24h × 1–2 weeks (minimum 2 doses before switching to PO therapy)	Respiratory quinolone <sup>†</sup> (PO) q24h <b>or</b> Doxycycline <sup>‡</sup> (PO) × 1–2 weeks <b>or</b> Macrolide <sup>††</sup> (PO) q24h × 1–2 weeks

Duration of therapy represents total time IV or IV, PO, or IV + PO. Most patients on IV therapy able to take PO meds should be switched to PO therapy after clinical improvement.

\* Compromised hosts may require longer courses of therapy.

† Moxifloxacin 400 mg × 1–2 weeks or Levofloxacin 750 mg × 5 days (or 500 mg × 1–2 weeks).

†† Azithromycin 500 mg or Clarithromycin XL 1 gm.

‡ Macrolide monotherapy should be avoided in areas where *macrolide-resistant S. pneumoniae* (MRSP) or *multidrug-resistant S. pneumoniae* (MDRSP) strains are prevalent.

‡ Doxycycline 200 mg (IV or PO) q12h × 3 days, then 100 mg (IV or PO) q12h × 4–11 days.



## Community-Acquired Pneumonia (CAP) (cont'd)

Subset	Usual Pathogens*	Preferred IV Therapy	Alternate IV Therapy	PO Therapy or IV-to-PO Switch
Typical bacterial pathogens	S. pneumoniae H. influenzae M. catarrhalis	Doxycycline 200 mg (IV) q12h × 3 days, then 100 mg (IV) q12h × 11 days <b>or</b> Respiratory quinolone <sup>†</sup> (IV) q24h × 1–2 weeks <b>or</b> Ceftriaxone 1 gm (IV) q24h × 1–2 wks	Doxycycline <sup>†</sup> (IV) × 1–2 weeks <b>or</b> Ertapenem 1 gm (IV) q24h × 1–2 weeks <b>or</b> Tigecycline** 200 mg (IV) × 1 dose, then 100 mg (IV) q24h × 1–2 weeks	(see unknown pathogen) <b>or</b> Amoxicillin/clavulanate XR 2 tablets (PO) q12h × 7–10 days <b>or</b> Cefprozil 500 mg (PO) q12h × 1–2 weeks
	K. pneumoniae*	Meropenem 1 gm (IV) q8h × 2 weeks <b>or</b> Ertapenem 1 gm (IV) q24h × 2 weeks <b>or</b> Respiratory quinolone <sup>§</sup> (IV) q24h × 2 weeks	Ceftriaxone 1 gm (IV) q24h × 2 weeks <b>or</b> Doripenem 1 gm (IV) q8h	Respiratory quinolone <sup>§</sup> (PO) q24h × 2 weeks <b>or</b> Doxycycline <sup>†</sup> (PO) × 1–2 weeks
	MDR K. pneumoniae CRE	Ceftazidime/avibactam 2.5 gm (IV) q8h × 1–2 weeks <b>or</b> Tigecycline 200 mg (IV) × 1 dose, then 100 mg (IV) q24h × 1–2 weeks**	Colistin 5 mg/kg (IV) q8h <sup>¶</sup> <b>or</b> Polymyxin B 1.25 mg/kg (IV) q12h <sup>¶</sup>	None

\* Compromised hosts may require longer courses of therapy.

† Moxifloxacin 400 mg × 1–2 weeks or Levofloxacin 750 mg × 5 days (or 500 mg × 1–2 weeks).

§ Moxifloxacin 400 mg or Levofloxacin 500 mg.

¶ 1 mg Colistin = 12,500 IU; 1 mg Polymyxin B = 10,000 IU.

‡ Doxycycline 200 mg (IV or PO) q12h × 3 days, then 100 mg (IV or PO) q12h × 4–11 days.

\*\* Depending on MIC, higher doses of tigecycline (LD: 400 mg (IV) × 1 dose, then MD: 200 mg (IV) q24h) may be necessary for susceptible MDR GNBS.

## Community-Acquired Pneumonia (CAP) (cont'd)

Subset	Usual Pathogens*	Preferred IV Therapy	Alternate IV Therapy	PO Therapy or IV-to-PO Switch
Atypical pathogens <i>Zoonotic</i>	<i>C. psittaci</i> (psittacosis) <i>Coxiella burnetii</i> (Q fever) <i>Francisella tularensis</i> (tularemia)	Doxycycline 200 mg (IV) q12h × 3 days, then 100 mg (IV) q12h × 2 weeks	Respiratory quinolone <sup>†</sup> (IV) q24h × 2 weeks	Doxycycline 200 mg (PO) q12h × 3 days, then 100 mg (PO) q12h × 11 days <b>or</b> Quinolone <sup>†</sup> (PO) q24h × 2 weeks
<i>Non-zoonotic</i>	<i>Legionella</i> sp. <sup>‡</sup> <i>Mycoplasma pneumoniae</i> <sup>‡</sup> <i>C. pneumoniae</i> <sup>‡</sup>	Moxifloxacin 400 mg (IV) q24h × 1–2 weeks <b>or</b> Levofloxacin 500 mg (IV) q24h × 1–2 weeks <b>or</b> Doxycycline 200 mg (IV) q12h × 3 days, then 100 mg (IV) q12h × 4–11 days	Tigecycline 200 mg (IV) × 1 dose, then 100 mg (IV) q24h × 1–2 weeks <b>or</b> Azithromycin 500 mg (IV) q24h × 1–2 weeks (minimum of 2 doses before switching to PO)	Respiratory quinolone <sup>†</sup> (PO) q24h × 1–2 weeks <b>or</b> Doxycycline 200 mg (PO) q12h × 3 days, then 100 mg (PO) q12h × 4–11 days <b>or</b> Azithromycin 500 mg (PO) q24h × 1–2 weeks
	CMV	Valganciclovir 900 mg (PO) q12h until cured		
Influenza ( <i>mild or moderate</i> ) without bacterial CAP	Influenza A/B	Oseltamivir (Tamiflu) 75 mg (PO) q24h × 5 days	Zanamivir (Relenza) 10 mg (2 puffs via oral inhaler) q12h × 5 days	Oseltamivir may be ineffective against avian influenza (H <sub>5</sub> N <sub>1</sub> ), but effective against swine influenza (H <sub>1</sub> N <sub>1</sub> ) and swine-like influenza (H <sub>3</sub> N <sub>2</sub> )

\* Compromised hosts may require longer courses of therapy.

† Moxifloxacin 400 mg or Levofloxacin 500 mg.

‡ May require prolonged therapy: *Legionella* (2–3 weeks); *Mycoplasma* (2 weeks); *Chlamydia* (2 weeks).

## Community-Acquired Pneumonia (CAP) (cont'd)

Subset	Usual Pathogens*	Preferred IV Therapy	Alternate IV Therapy	PO Therapy or IV-to-PO Switch
Influenza (severe with <b>simultaneous</b> CAP)*	Influenza A pneumonia	Peramivir 600 mg (IV) q24h × 1 days <b>or</b> Oseltamivir 75 mg (PO) q12h × 5 days <b>plus</b> Amantadine 200 mg (PO) q24h × 7–10 days**		Start treatment as soon as possible after onset of symptoms, preferably within 3 days. Rimantadine/ amantadine inactive against Influenza B.
	with MSSA/MRSA CAP	<u>MSSA</u> Nafcillin 2 gm (IV) q4h × 2 weeks <b>or</b> Cefazolin 1 gm (IV) q8h × 2 weeks <b>or</b> Meropenem 1 gm (IV) q8h × 2 weeks <u>MRSA</u> Vancomycin 1 gm (IV) q12h × 2 weeks <b>or</b> Tigecycline 200 mg (IV) × 1 dose then 100 mg (IV) q24h × 2 weeks <b>or</b> Linezolid 600 mg (IV) q12h × 2 weeks		
Post-influenza (with <b>subsequent</b> bacterial CAP). Recovering from influenza/treated previously	S. pneumoniae H. influenzae	Respiratory quinolone† (IV) q24h × 2 weeks <b>or</b> Ceftriaxone 1 gm (IV) q24h × 2 wks	Doxycycline 200 mg (IV) × 3 days, then 100 mg (IV) q12h × 11 days	Respiratory quinolone‡ (PO) q24h × 2 weeks
		Plus influenza therapy (see preceding page)		
Chickenpox pneumonia	VZV	Acyclovir 5–10 mg/kg (IV) q8h × 10 days		Valacyclovir 1–2 gm (PO) q8h × 10 days

Duration of therapy represents total time IV or IV, PO, or IV + PO.

\* Compromised hosts predisposed to organisms listed, but may be infected by usual pathogens in normal hosts.

‡ May require prolonged therapy: Legionella (2–3 weeks); Mycoplasma (2 weeks); Chlamydia (2 weeks).

¶ If CXR shows multiple rapidly cavitating infiltrates < 72 hours, begin empiric anti-MSSA/CA-MRSA therapy with anti-influenza therapy.

¶¶ Although resistance common, may improve oxygenation in severe influenza A.

§ Dose for CrCl > 30 ml/min = 30 mg (PO) q12h; for CrCl > 10–30 ml/min = 30 mg (PO) q24h.

## Community-Acquired Pneumonia (CAP) (cont'd)

Subset	Usual Pathogens*	Preferred IV Therapy	Alternate IV Therapy	PO Therapy or IV-to-PO Switch
Aspiration	Oral anaerobes S. pneumoniae H. influenzae M. catarrhalis	Ceftriaxone 1 gm (IV) q24h x 2 weeks  <b>or</b> Respiratory quinolone <sup>‡</sup> (IV) q24h x 2 weeks	Doxycycline 200 mg (IV) q12h x 3 days, then 100 mg (IV) q12h x 11 days	Respiratory quinolone <sup>‡</sup> (PO) q24h x 2 weeks  <b>or</b> Doxycycline 200 mg (PO) q12h x 3 days, then 100 mg (PO) q12h x 4–11 days**  <b>or</b> Amoxicillin 1 gm (PO) q8h x 2 weeks
Tuberculosis (TB)	M. tuberculosis	INH 300 mg (PO) q24h (and pyridoxine 50 mg (PO) q24h) x 6 months <b>plus</b> RIF 600 mg (PO) q24h x 6 months <b>plus</b> PZA 25 mg/kg (PO) q24h x 2 months <b>plus</b> EMB 15 mg/kg (PO) q24h (until susceptibilities known) <sup>††</sup>		
MDR TB	M. tuberculosis	Bedaquiline (Sirturo) 400 mg (PO) q24h (D.O.T.) x 2 weeks, then 200 mg (PO) 3 x per week x 22 weeks*.		
Non-tuberculous (atypical tuberculosis) Mycobacteria		<u>Treat for 12 months after sputum negative for MAI</u> ; EMB 15 mg/kg (PO) q24h <b>plus either</b> Clarithromycin 500 mg (PO) q12h <b>or</b> Azithromycin 500 mg (PO) q24h <b>plus either</b> (Rifampin or Rifabutin). May substitute a quinolone (PO) <sup>‡</sup> q24h for rifamycin		
	M. kansasii	<u>Preferred therapy</u> RIF 600 mg (PO) q24h <b>plus</b> INH 300 mg (PO) q24h <b>plus</b> pyridoxine 50 mg (PO) q24h <b>plus</b> EMB 15 mg/kg (PO) q24h. Treat x 12 months after negative sputum <u>Alternate therapy</u> RIF 600 mg (PO) thrice weekly <b>or</b> daily <b>plus</b> EMB 15 mg/kg (PO) thrice weekly <b>or</b> daily <b>plus</b> clarithromycin 500–1000 mg (PO) thrice weekly <b>or</b> daily. Treat x 12 months after negative sputum		

\* May be used in adults as part of a multidrug TB regimen (>3 drugs). Take with food. Side effects include nausea, headache, arthralgias, QT<sub>c</sub> prolongation.

‡ Moxifloxacin 400 mg or Levofloxacin 500 mg.

†† If isolate is sensitive, discontinue EMB and continue as above to complete 6 months. If INH-resistant, continue EMB, RIF and PZA to complete 6 months. If any other resistance is present (MDR-TB), obtain infectious disease or pulmonary consult.

## Community-Acquired Pneumonia (CAP) (cont'd)

Subset	Usual Pathogens*	Preferred IV Therapy	Alternate IV Therapy	PO Therapy or IV-to-PO Switch
Chronic alcoholics	K. pneumoniae S. pneumoniae H. influenzae M. catarrhalis	Meropenem 1 gm (IV) q24h × 2 weeks <b>or</b> Ertapenem 1 gm (IV) q24h × 2 weeks <b>or</b> Tigecycline 200 mg (IV) × 1, then 100 mg (IV) q24h	Ceftriaxone 1 gm (IV) q24h × 2 weeks <b>or</b> Respiratory quinolone <sup>‡</sup> (IV) q24h × 2 weeks	Respiratory quinolone (PO) <sup>‡</sup> q24h × 2 weeks
	MDR K. pneumoniae CRE	Ceftazidime/avibactam 2.5 gm (IV) q8h × 1-2 weeks <b>or</b> Tigecycline 200 mg (IV) × 1 dose, 100 mg (IV) q24h × 1-2 weeks	Colistin 5 mg/kg (IV) q8h × 1-2 weeks <sup>¥</sup> <b>or</b> Polymyxin B 1.25 mg/kg (IV) q12h × 1-2 weeks <sup>¥</sup>	None
Bronchiectasis, cystic fibrosis	S. maltophilia B. cepacia Alcaligenes xyloxidans	<u>Preferred:</u> TMP-SMX 2.5 mg/kg (IV) q6h <sup>¶</sup> <b>or</b> Minocycline 100 mg (IV) q12h <sup>¶</sup> <u>Alternate:</u> Doxycycline 100 mg (IV) q12h <sup>¶</sup>		TMP-SMX 1 SS tablet (PO) q6h <sup>¶</sup> <b>or</b> Minocycline 100 mg (PO) q12h <sup>¶</sup> <b>or</b> Respiratory quinolone <sup>‡</sup> (PO) q24h <sup>¶</sup>
	P. aeruginosa	Meropenem 2 gm (IV) q8h <sup>¶</sup>	Levofloxacin 750 mg (IV) q24h <sup>¶</sup> <b>or</b> Cefepime 2 gm (IV) q8h <sup>¶</sup> ± Amikacin 1 gm (IV) q24h <sup>¶</sup>	Levofloxacin 750 mg (PO) q24h <sup>¶</sup> <b>or</b> Ciprofloxacin 750 mg (PO) q12h
	MDR P. aeruginosa	Ceftazidime/avibactam 2.5 gm (IV) q8h × 1-2 weeks <b>or</b> Doripenem 1 gm (IV) q8h <sup>¶</sup>	Polymyxin B 1.25 mg/kg (IV) q12h <sup>¥</sup> <b>or</b> Colistin 5 mg/kg (IV) q8h <sup>¶</sup>	None

\* Compromised hosts predisposed to organisms listed but may be infected by usual pathogens in normal hosts.

¶ Treat until cured.

‡ Moxifloxacin 400 mg or Levofloxacin 500 mg.

¥ Colistin 1 mg = 12,500 IU; Polymyxin B 1 mg = 10,000 IU.

## Community-Acquired Pneumonia (CAP) (cont'd)

Subset	Usual Pathogens*	Preferred IV Therapy	Alternate IV Therapy	PO Therapy or IV-to-PO Switch
Chronic steroid therapy†	Aspergillus sp.	Voriconazole (see "usual dose," p. 714)¶	Lipid amphotericin (see p. 525)¶ <b>or</b> Caspofungin 70 mg (IV) × 1 then 50 mg (IV) q24h¶ <b>or</b> amphotericin B 1–1.5 mg/kg (IV) q24h¶	Voriconazole (see "usual dose," p. 714)¶ <b>or</b> Posaconazole 200 mg (PO) q6h initially, then 400 mg (PO) q12h¶ <b>or</b> Itraconazole 200 mg (IV) q12h × 2 days then 200 mg (PO) q12h***
	P. (carinii) jiroveci (PCP)	TMP–SMX 5 mg/kg (IV) q6h × 3 weeks‡	Pentamidine 4 mg/kg (IV) q24h × 3 weeks‡	TMP–SMX 5 mg/kg (PO) q6h × 3 weeks <b>or</b> Atovaquone 750 mg (PO) of q12h <b>or</b> Dapsone 100 mg (PO) of q24h
	CMV	Valganciclovir 900 mg (PO) q12h until cured		
HIV	Typical/atypical CAP pathogens	see pp. 50–51		
	P. (carinii) jiroveci (PCP)	see p. 325		
Other pathogens	For <i>Cryptococcus neoformans</i> , <i>Blastomyces</i> , <i>Histoplasma</i> , <i>Coccidioides</i> , <i>Paracoccidioides</i> , <i>Actinomyces</i> , <i>Nocardia</i> , <i>Pseudallescheria boydii</i> , <i>Sporothrix</i> , <i>Mucor</i> (see pp. 267–269).			

¶ Treat until cured.

† Treat IV or IV-to-PO switch.

\* Compromised hosts predisposed to organisms listed, but may be infected by usual pathogens of normal hosts.

\*\* Loading dose is not needed PO if given IV with the same drug.

‡ Steroid dosage: prednisone 40 mg (PO) q12h on days 1–5, then 40 mg (PO) q24h on days 6–10, then 20 mg (PO) q24h on days 11–21. Methylprednisolone (IV) can be substituted at 75% of prednisone dose.

†† Moxifloxacin 400 mg or Levofloxacin 500 mg.

**Clinical Presentation:** Fever, cough, respiratory symptoms, chest x-ray consistent with pneumonia. Typical CAPs present only with pneumonia without extrapulmonary findings.

**Diagnosis:** Identification of organism on sputum gram stain/culture. Same organism is found in blood if blood cultures are positive. Also see Chapter 8 for typical chest x-ray patterns.

### **Community-Acquired Pneumonia (Typical Bacterial Pathogens)**

**Diagnostic Considerations:** Sputum is useful if a single organism predominates and is not contaminated by saliva. Purulent sputum, pleuritic chest pain, pleural effusion favor typical pathogens.

**Pitfalls:** Obtain chest x-ray to verify the diagnosis and rule out noninfectious mimics (e.g., heart failure).

**Therapeutic Considerations:** Do not switch to narrow-spectrum antibiotic after organism is identified on gram stain/blood culture. Pathogen identification is important for prognostic and public health reasons, not for therapy. Severity of CAP is related to the degree of cardiopulmonary/immune dysfunction and impacts the length of hospital stay, not the therapeutic approach or antibiotic choice.

**Prognosis:** Related to cardiopulmonary status and splenic function.

### **Community-Acquired Pneumonia (Pertussis)**

**Clinical Presentation:** Rhinorrhea over 1–2 weeks (catarrhal stage) progressing to paroxysms of cough (paroxysmal stage) lasting 2–4 weeks, often without a characteristic inspiratory whoop, followed by a convalescent stage lasting 1–2 weeks during which cough paroxysms decrease in frequency/severity. Fever is low grade or absent. In children < 6 months, whoop is frequently absent and apnea may occur. Older children/adults may present with persistent cough (without whoop) lasting 2–6 weeks.

**Diagnostic Considerations:** A positive PCR or DFA for *Bordetella pertussis* from a nasopharyngeal swab (NP). May be cultured by bedside inoculation from NP swab of Bordet-Gengou media. "Shaggy heart" on chest x-ray is characteristic. The only CAP with lymphocytosis > 60%.

**Pitfalls:** Consider pertussis in older children and adults with prolonged coughing. Hoarseness uncommon, but may be misleading by mimicking viral laryngitis or *C. pneumoniae*.

**Therapeutic Considerations:** By the paroxysmal stage, antibiotics have minimal effect on the course of the illness but are indicated to decrease transmission.

**Prognosis:** Good, despite the prolonged course.

### **Community-Acquired Pneumonia (Atypical Pathogens)**

**Clinical Presentation:** CAP with extrapulmonary symptoms, signs, or laboratory abnormalities.

**Diagnosis:** Confirm by specific serological tests.

### **Legionnaire's Disease (*Legionella* sp.)**

*Legionella* organisms live in a fresh water environment and survive in a symbiotic relationship with freshwater amoeba. *Legionella* infection may be introduced into humans via droplet aerosolization of *Legionella*-contaminated water supplies (e.g., cooling towers, water systems, whirlpools, showers, air conditioners, respiratory therapy devices). LD can occur in isolated cases or in outbreaks (community-acquired or nosocomial). *Legionella* has a seasonal predisposition and is most common in the late summer/early fall. LD is rare in young children, uncommon in young adults, but is most common in adult/elderly patients. Individuals with impaired cell-mediated immunity ( $\downarrow$  T-lymphocyte function) e.g., TNF- $\alpha$  antagonists are predisposed to LD, but normal hosts can be infected as well.

**Clinical Presentation:** LD is more severe than *M. pneumoniae* or *C. pneumoniae* CAP. Onset is usually acute, with high fever ( $\geq 102^{\circ}\text{F}$ ) and relative bradycardia. Myalgias and chills are not uncommon. Like other atypical pneumonias, LD has a characteristic pattern of extrapulmonary

manifestations, which is the key to presumptive clinical diagnosis. On chest x-ray, LD is suggested not by the appearance of the infiltrate, but by its rapid, asymmetric progression. With LD, bilateral involvement is usual, consolidation/pleural effusion are not uncommon, and cavitation is rare. Extrapulmonary LD findings include, otherwise unexplained, mental confusion, watery diarrhea, abdominal pain, relative lymphopenia  $\uparrow$  ESR/CRP, mild/transient  $\uparrow$  SGOT/SGPT, early/transient  $\downarrow$  serum phosphorus,  $\uparrow$  ferritin levels ( $> 2 \times n$ )  $\uparrow$  CPK, and early microscopic hematuria.

**Diagnostic Considerations:** Lack of response to beta-lactam therapy in a CAP patient suggests the possibility of LD and should prompt specific testing. Diagnosis is confirmed by Legionella DFA of sputum/respiratory secretions, Legionella IgM/IgG titers (a single titer  $\geq 1:256$  or  $\geq 4$  fold increase between acute and convalescent titers) is diagnostic. Legionella may also be cultured on BCYE agar from sputum/respiratory secretions, pleural effusion, or lung tissue. Legionella pneumophila (serotypes 01-06) may also be diagnosed by urinary antigen test.

**Pitfalls:** As no individual finding is pathognomonic the constellation of signs/symptoms of LD are often overlooked. The characteristic pattern of extrapulmonary organ involvement differentiates LD from other CAPs. Hyponatremia is common but nonspecific, but serum phosphorus is a more specific for LD. Findings suggestive of an alternate (non-Legionella) cause of CAP include  $\uparrow$  cold agglutinin titer, normal SGOT/SGPT, normal serum phosphorus, normal CPK level, upper respiratory tract involvement (non-exudative pharyngitis, laryngitis, otitis) or splenomegaly. Legionella DFA of sputum/respiratory secretions is positive early, but rapidly becomes negative with treatment. Legionella titers, may be negative if ordered early (usually at the time of presentation). Antimicrobial therapy may eliminate/blunt a titer rise. Legionella urinary antigen testing is useful but has limitations, as it is not always positive early when the patient presents, and detects only Legionella pneumophila (serotypes 01-06), but not other Legionella species.

**Therapeutic Considerations:** Antimicrobial agents with a high degree of anti-Legionella activity are quinolones, doxycycline, tigecycline, and macrolides. Rifampin has in vitro anti-Legionella activity, but as part of combination therapy (erythromycin + rifampin) has not been shown to offer any advantage over well selected monotherapy (quinolone, doxycycline). LD does not respond to beta-lactams and only somewhat to erythromycin. Patients treated with a quinolone, doxycycline, or tigecycline typically defervesce slowly (5–7 days). Therapy is usually continued for 2–3 weeks depending upon clinical severity/host factors.

**Prognosis:** In immunocompetent hosts with good cardiopulmonary function, prognosis is good. Prognosis is guarded in the elderly and those with impaired CMI (HIV, organ transplants, corticosteroid therapy) and in those severe cardiopulmonary disease (CAD, valvular heart disease, COPD, chronic bronchiectasis).

## **Mycoplasma pneumoniae**

**Epidemiology:** M. pneumoniae primarily affects young adults and the elderly. M. pneumoniae is spread from person-to-person via aerosolized droplets.

**Clinical Presentation:** Onset is subacute with a characteristic dry/nonproductive cough that lasts for weeks, low-grade fever ( $< 102^{\circ}\text{F}$ ) /chills, and mild myalgias. As with other atypical CAPs, M. pneumoniae is characterized by its pattern of extrapulmonary organ involvement, which may include sore throat (non-exudative pharyngitis), bullous myringitis, otitis media, and/or watery loose stools/diarrhea (Table 2.1). Features that argue against the diagnosis of M. pneumoniae CAP include high fevers ( $>102^{\circ}\text{F}$ ), relative bradycardia, laryngitis, abdominal pain,  $\uparrow$  SGOT/SGPT, low serum phosphorus, or renal abnormalities. Chest x-ray typically shows ill-defined unilateral infiltrates without pleural effusion, consolidation or cavitation. M. pneumoniae may present as severe CAP in compromised hosts, the elderly, and those with advanced COPD.



**Diagnostic Considerations:** *M. pneumoniae* testing is recommended for patients with otherwise unexplained protracted nonproductive cough, low-grade fevers, loose stools/watery diarrhea, and an ill-defined infiltrate on chest x-ray. Cold agglutinin titers are elevated early/transiently (75%) and provide a rapid presumptive diagnosis; titers  $\geq 1:64$  are due to *M. pneumoniae*. An elevated *M. pneumoniae* IgM ELISA titer suggests acute infection, but an elevated IgG titer indicates only past exposure; *M. pneumoniae* may be cultured from respiratory secretions on viral media (requires 1–2 weeks for growth).

**Pitfalls:** A dry, nonproductive cough also suggests an ILI or *C. pneumoniae*. Before ascribing *E. multiforme* to *M. pneumoniae*, be sure that it is not due to other causes (e.g., drug induced). Features that argue against the diagnosis of *M. pneumoniae* include temperature  $> 102^{\circ}\text{F}$ , relative bradycardia, moderate/large pleural effusion, consolidation/cavitation, otherwise unexplained abdominal pain,  $\uparrow$  SGOT/SGPT or low serum phosphorus. Suspect *M. pneumoniae* in chronic dry cough/new onset asthma.

**Therapeutic Considerations:** Doxycycline, macrolides, quinolones, or tigecycline result in rapid clinical response: in  $< 72$  hours, but dry cough usually persists for weeks. Because *Mycoplasma* organisms reside on/in the bronchial epithelium, antimicrobial therapy should be continued for 2 weeks to eliminate carriage from oropharyngeal secretions and reduce the risk of transmission via aerosolized droplets.

**Prognosis:** *M. pneumoniae* is usually a mild, self-limiting illness in immunocompetent adults. *M. pneumoniae* may be severe in compromised hosts, the elderly, and those with advanced lung disease. Patients with *M. pneumoniae* and meningoencephalitis (headaches/stiff neck, mental confusion, very high cold agglutinin titers [often  $\geq 1:1024$ ]) have a good prognosis. *M. pneumoniae* may be complicated by transverse myelitis, brain stem/cerebellar ataxia, or Guillain-Barré syndrome after *M. pneumoniae* some patients develop permanent asthma.

### **Chlamydophilia (Chlamydia) pneumoniae**

**Epidemiology:** *C. pneumoniae* may occur as sporadic cases part of an outbreak NHAP. *C. pneumoniae* is spread from person-to-person via aerosolized droplets. *C. pneumoniae* has no seasonal predisposition. Some patients with *C. pneumoniae* CAP develop permanent asthma.

**Clinical Presentation:** *C. pneumoniae* presents as a “mycoplasma-like” illness. Like other atypical CAPs, *C. pneumoniae* CAP is characterized by extrapulmonary findings, particularly nasal discharge, sore throat, and hoarseness (Table 2.1). Temperature is usually  $\leq 102^{\circ}\text{F}$ . Chest x-ray findings typically show a unilateral ill-defined infiltrate without consolidation, cavitation, or pleural effusion. *C. pneumoniae* testing is recommended in a patient with a “mycoplasma-like” illness and hoarseness.

**Diagnostic Considerations:** The diagnosis of *C. pneumoniae* CAP requires a single IgM titer of  $\geq 1:16$  or an IgG titer  $> 1:512$  (ELISA or MIF). *C. pneumoniae* may be cultured from respiratory secretions using viral culture media.

**Pitfalls:** IgG titers indicate past exposure, not current infection.  $\uparrow$  IgM and  $\uparrow$  IgG titers may be delayed for more than 3 and 6 weeks, respectively. False negatives may occur if IgM titers are obtained too early. Recovery of *C. pneumoniae* from respiratory secretions does not differentiate carriage from infection. In a CAP patient with otherwise unexplained wheezing should suggest the possibility of *C. pneumoniae*.

**Therapeutic Considerations:** Doxycycline, or quinolones are effective. *C. pneumoniae* is sensitive to erythromycin in vitro, but erythromycin is often ineffective in vivo. *C. pneumoniae* also responds to newer macrolides.

**Prognosis:** *C. pneumoniae* CAP is usually a mild/moderate pneumonia with an excellent prognosis. Severe CAP may occur in compromised hosts, the elderly, or those with severe lung disease some develop permanent asthma.

**Clinical Presentation:** CAP with extrapulmonary symptoms, signs, or laboratory abnormalities (Table 2.2).

**Table 2.1. Diagnostic Features of Non-zoonotic Atypical Pneumonias**

Key Characteristics	Mycoplasma pneumoniae	Legionnaire's Disease	C. pneumoniae
<b>Symptoms</b>			
Mental confusion	- <sup>†</sup>	+	-
Prominent headache	-	±	-
Meningismus	-	-	-
Myalgias	±	±	±
Ear pain	+	-	±
Pleuritic pain	-	±	-
Abdominal pain	-	±	-
Loose stools/watery diarrhea	±	±	-
<b>Signs</b>			
Rash	± <sup>a</sup>	-	-
Non-exudative pharyngitis	+	-	+
Hemoptysis	-	±	-
Wheezing	-	-	+
Lobar consolidation	-	±	-
Cardiac involvement	± <sup>b</sup>	- <sup>c</sup>	-
Splenomegaly	-	-	-
Relative bradycardia	-	+	-
<b>Laboratory Abnormalities</b>			
WBC count	↑/N	↑	N
Acute thrombocytosis	±	-	-
Hyponatremia	-	+	-
Hypophosphatemia (early)	-	+	-
↑ AST/ALT (SGOT/SGPT)	-	+	-
↑ CPK	-	±	-
↑ ESR (> 100 mm/h)	-	+	-
↑ Ferritin (> 2 x n)	-	+	-
↑ Cold agglutinins (≥ 1:64)	+	-	-
Microscopic hematuria (early)	-	±	-
<b>Chest X-Ray</b>			
Infiltrate	Patchy	Patchy	Circumscribed lesions
Bilateral hilar adenopathy	-	-	-
Pleural effusion	± (small)	±	-

ALT = alanine aminotransferase; AST = aspartate aminotransferase; N = normal; WBC = white blood cell.

+ = usually present; ± = sometimes present; - = usually absent.

↑ = increased; ↓ = decreased; ↑↑↑ = markedly increased.

a = erythema multiforme; b = myocarditis, heart block, or pericarditis; c = unless endocarditis.

† = mental confusion only if meningoencephalitis.

†† = often not be positive early, but antigenuria persists for weeks. Useful only to diagnose *L. pneumophila* (serogroups 1-6), not other species/serogroups.

## Zoonotic Atypical Pneumonia (see Chapter 8 for Differential Diagnosis of Chest X-Ray patterns)

Subset	Usual Pathogens*	Preferred IV Therapy	Alternate IV Therapy	PO Therapy or IV-to-PO Switch
Zoonotic pathogens (zoonotic vector)	C. psittaci (psittacosis) Coxiella burnetii (Q fever) Francisella tularensis (tularemia)	Doxycycline 200 mg (IV) q12h x 3 days, then 100 mg (IV) q12h x 11 days	Levofloxacin 500 mg (IV) x 2 weeks <b>or</b> Moxifloxacin 400 mg (IV) x 2 weeks	Doxycycline 200 mg (PO) q12h x 3 days, then 100 mg (PO) q12h x 11 days <sup>†</sup> <b>or</b> Levofloxacin 500 mg (PO) x 2 weeks <b>or</b> Moxifloxacin 400 mg (PO) x 2 weeks

Duration of therapy represents total time IV, PO, or IV + PO.

\* *Compromised hosts may require longer courses of therapy.*

† *Loading dose is not needed PO if give IV with the same drug.*

**Diagnostic Considerations:** Zoonotic contact history is key to presumptive diagnosis: psittacosis (parrots and psittacine birds); Q fever (sheep, parturient cats); tularemia (rabbit, deer, deer fly bite). The diagnosis is confirmed by specific serological testing. Zoonotic CAP with splenomegaly/relative bradycardia suggests Q fever or psittacosis.

**Pitfalls:** Organisms are difficult/dangerous to grow. Do not culture. Use serological tests for diagnosis. The diagnosis of acute Q fever is by demonstrating elevated IgG (phase II) C. burnetii titers. Most patients with acute Q fever do not become chronic e.g., Q fever SBE. The diagnosis of chronic Q fever is made by demonstrating elevated IgG (phase I) titers  $\geq 1:1024$  at 6 months following acute Q fever. C. burnetii (Q fever) titers may cross react with Bartonella titers.

**Therapeutic Considerations:** Q fever endocarditis requires prolonged therapy.

**Prognosis:** Good except for Q fever with complications, e.g., myocarditis, SBE.

**Table 2.2. Diagnostic Features of Zoonotic Atypical Pneumonias**

Key Characteristics	Psittacosis	Q Fever	Tularemia
<b>Symptoms</b>			
Mental confusion	–	±	–
Prominent headache	+	+	+
Meningismus	–	–	–
Myalgias	+	+	+
Ear pain	–	–	–
Pleuritic pain	±	±	+
Abdominal pain	–	–	–
Diarrhea	–	–	–
<b>Signs</b>			
Rash	± a	–	–
Nonexudative pharyngitis	–	–	±
Hemoptysis	–	–	±
Lobar consolidation	+	+	+
Cardiac involvement	± b	± c	–
Splenomegaly	±	+	–
Relative bradycardia	±	±	–
<b>Chest X-Ray</b>			
Infiltrate	Patchy consolidation	"Round" infiltrates or consolidation	"Ovoid infiltrates"
Bilateral hilar adenopathy	–	–	±
Pleural effusion	–	–	Bloody
<b>Laboratory Abnormalities</b>			
WBC count	↓	↑/N	↑/N
Thrombocytosis	–	+	–
Hyponatremia	±	±	±
Hypophosphatemia	–	–	–
↑ AST/ALT (SGOT/SGPT)	+	+	–
↑ Cold agglutinins	–	±	–
Anti-smooth muscle antibodies (anti-SMA)	–	±	–
Microscopic hematuria	–	–	–

ALT = alanine aminotransferase; AST = aspartate aminotransferase; N = normal; WBC = white blood cell.

+ = usually present; ± = sometimes present; – = usually absent.

↑ = increased; ↓ = decreased; ↑↑↑ = markedly increased.

a = Horder's spots (facial spots, resemble the abdominal rash of typhoid fever (rose spots); b = Myocarditis; c = endocarditis.

### Severe Acute Respiratory Syndrome (SARS)

**Presentation:** Fever, dry cough, myalgias, diarrhea in some. Auscultation of the lungs resembles viral influenza (i.e., quiet, no rales). Biphasic infection: fever decreases after few days, patient improves, then fever recurs in a few days and patients become short of breath/hypoxic. Chest x-ray shows bilateral interstitial (diffuse) infiltrates. WBC and platelet counts are usually normal or slightly decreased. Relative lymphopenia present early. Mild increases in SGOT/SGPT, LDH, CPK are common.

**Diagnostic Considerations:** Diagnosis by viral isolation or specific SARS serology. Important to exclude influenza A, Legionnaires' disease and tularemic pneumonia.

**Pitfalls:** Should not be confused with viral influenza (a 3-day illness). Patients with SARS deteriorate during week 2 when influenza patients are recovering. Early, chest x-ray in SARS has discrete infiltrates (unlike influenza unless superimposed CAP) that may be ovoid. Late, ARDS on chest x-ray.

**Therapeutic Considerations:** Most patients are severely hypoxemic and require oxygen/ventilatory support. In a preliminary study, some patients benefitted from corticosteroids (pulse-dosed methylprednisolone 500 mg [IV] q24h x 3 days followed by taper/step down with prednisone [PO] to complete 20 days) plus Interferon alfa-con-1 (9 mcg [SQ] q24h x at least 2 days, increased to 15 mcg/d if no response) x 8–13 days. Ribavirin of no benefit.

**Prognosis:** Related to underlying cardiopulmonary/immune status/ARDS. Frequently fatal.

**Pitfalls:** Failure to respond to beta-lactams should suggest diagnosis of atypical CAP.

**Therapeutic Considerations:** Treat Legionella x 4 weeks. Treat Mycoplasma or Chlamydia x 2 weeks.

**Prognosis:** Related to severity of underlying cardiopulmonary disease.

### Middle East Respiratory Syndrome (MERS)

**Clinical Presentation:** Presents as an influenza like illness (ILI) with acute onset of fever, chills, dry cough/SOB, myalgias  $\pm$  N/V/D. Headache/sore throat common. Hemoptysis in some. Rapidly progressive respiratory failure/ARDS is typical (~ 1 week). WBC counts usually normal, thrombocytopenia,  $\uparrow$ LFTs,  $\uparrow$ LDH, relative lymphopenia, but lymphocytosis in some. CXR: basilar unilateral (initially) or bilateral dense interstitial/nodular infiltrates (later). Consolidation common  $\pm$  pleural effusions.

**Diagnostic Considerations:** MERS resembles an ILI clinically (unlike SARS-CoA). Contact with dromedary camels (MERS reservoir). Dx is by RT-PCR of MERS-CoA virus in respiratory secretions (lower > upper). MERS, in respiratory secretions (lower > upper), but not in stools/urine (unlike SARS-CoA).

**Pitfalls:** Resembles severe influenza A. Not a biphasic illness (like SARS-CoA).

**Therapeutic Considerations:** Ventilatory support, but no effective treatment available. No bacterial co-infections, i.e., no antibiotics for superimposed bacterial CAP. IC measures important to prevent person-to-person/nosocomial spread.

**Prognosis:** High mortality rate with rapidly progressive severe respiratory failure/ARDS especially with comorbidities.

### Avian Influenza (Influenza A (H<sub>5</sub>N<sub>1</sub>, H<sub>7</sub>N<sub>9</sub>) Pneumonia)

**Clinical Presentation:** Influenza following close contact with infected poultry. Recent outbreaks in humans in Asia. Human-to-human transmission reported. Often fulminant respiratory illness rapidly followed by ARDS/death.

**Diagnostic Considerations:** Often acute onset of severe influenza illness  $\pm$  diarrhea/conjunctival suffusion with leukopenia, lymphopenia, mildly  $\uparrow$  serum transaminases and  $\uparrow$  LDH. Diagnosis by hemagglutinin-specific RT-PCR for avian influenza or culture of respiratory secretions. Not complicated by bacterial CAP. No need for empiric antibiotic therapy.

**Pitfalls:** Although avian influenza is caused by influenza A virus, the hemagglutinin inhibition serological test used to diagnose influenza A is insensitive to avian hemagglutinin, resulting in negative testing.

**Therapeutic Considerations:** Antivirals must be given early to be effective. Avian influenza strains may be resistant to oseltamivir and amantadine/rimantadine. Oseltamivir 150 mg dose may be more effective than 75 mg dose. Even if resistant, amantadine/rimantadine should be given to increase peripheral airway dilatation/oxygenation.

**Prognosis:** Often fulminant with ARDS/death. Unlike human/swine influenza, avian influenza is often rapidly/highly lethal but *not* complicated by MSSA/MRSA CAP.

### Influenza A/B Pneumonia

**Clinical Presentation:** Acute onset of fever, headache, myalgias/artralgias, sore throat, prostration, dry cough. Myalgias most pronounced in lower back/legs. Eye pain is common. Rapid high fever initially, which decreases in

2–3 days. Severity ranges from mild flu to life-threatening pneumonia. Chest x-ray in early influenza is normal/near normal without focal/segmental infiltrates or pleural effusion.

**Diagnostic Considerations:** Mild cases with headache, sore throat, and rhinorrhea resemble the common cold/respiratory viruses (influenza-like illnesses) and can be caused by type A or B. Severe flu is usually due to type A. Rapid diagnosis by DFA of respiratory secretions (nasopharyngeal swabs). Influenza virus may be cultured in viral media from respiratory secretions and typed.

**Pitfalls:** Influenza pneumonia has no chest auscultatory findings and the chest x-ray is normal/near normal. Severe influenza pneumonia is accompanied by an oxygen diffusion defect ( $\uparrow$  A-a gradient), and patients are hypoxemic/cyanotic. Pleuritic chest pain may be present. Influenza can invade the intercostal muscles to mimic pleuritic chest pain. Labor/segmental infiltrates on CXR with influenza pneumonia indicate simultaneous or subsequent bacterial CAP. If CAP presents with fulminant/severe necrotizing pneumonia in a normal host look for antecedent or concomitant influenza pneumonia.

**Therapeutic Considerations:** Mild/moderate influenza can be treated with neuraminidase inhibitors (Tamiflu/Relenza), which reduce symptoms by 1–2 days. Start treatment within 2 days of symptom onset. Reduce the dose of Tamiflu to 75 mg (PO) q48h for CrCl 10–30 ml/min. Relenza is generally not recommended for patients with underlying COPD/asthma (increased risk of bronchospasm) and should be discontinued if bronchospasm or a decline in respiratory function occurs. For severe influenza pneumonia, amantadine may increase peripheral airway dilatation/oxygenation. Reduce the dose of rimantadine to 100 mg (PO) q24h in the elderly, severe liver dysfunction, or CrCl < 10 ml/min. Peramivir may be given in severe cases. Flu with simultaneous bacterial CAP is due to MSSA/MRSA and post-influenza CAP usually due to *S. pneumoniae* or *H. influenzae* (Table 2.3).

**Prognosis:** Good for mild/moderate flu. Severe flu may be fatal due to influenza pneumonia with profound hypoxemia. Prognosis is worse if presents with simultaneous MSSA/MRSA. Influenza with necrotizing community-acquired (CA) MSSA/MRSA CAP is frequently fatal. Prognosis is worst for CA-MRSA Panton-Valentine Leukocidin (PVL)-positive strains. Dry cough/fatigue may persist for weeks after influenza.

**Table 2.3. The 3 Clinical Presentations of Influenza A Pneumonia in Adults**

Clinical features at time of onset	Severity	Usual Pathogens
• Influenza pneumonia ( <b>no focal/lobar infiltrates</b> )	Mild/moderate $\rightarrow$ rarely fatal	Influenza (human, avian, swine <sup>†</sup> )
• Influenza pneumonia with simultaneous bacterial CAP ( <b>with multiple rapidly cavitating infiltrates</b> )	Severe $\rightarrow$ often rapidly fatal	Influenza (human, swine <i>not</i> avian) <b>plus</b> MSSA/CA-MRSA*
• Influenza pneumonia with sequential bacterial CAP ( <b>followed by non-cavitating focal/lobar infiltrates</b> (after ~1 week of improvement))	Mild/moderate $\rightarrow$ rarely fatal (unless underlying cardiopulmonary disease/Immunodeficiency). Same prognosis as in <i>S. pneumoniae</i> /H. influenzae CAP without influenza.	<i>S. pneumoniae</i> <i>H. influenzae</i>

† Avian influenza ( $H_5N_1$ ) often rapidly fatal in young/healthy adults.

\* CA-MRSA PVL + strains highly virulent.

### **Swine Influenza (H<sub>1</sub>N<sub>1</sub>) Pneumonia**

**Clinical Presentation:** Swine influenza (H<sub>1</sub>N<sub>1</sub>) may present with a mild illness with fever, cough, and loose stools/diarrhea to severe viral pneumonia requiring ventilation that may rapidly progress to ARDS/death. Because the clinical presentation resembles that of human seasonal influenza, patients with swine influenza (H<sub>1</sub>N<sub>1</sub>) pneumonia as well as other viral pneumonias typically present as influenza-like illnesses (ILIs). Unlike human seasonal influenza, swine influenza (H<sub>1</sub>N<sub>1</sub>) pneumonia affects primarily young healthy adults. As with human seasonal influenza, swine influenza (H<sub>1</sub>N<sub>1</sub>) may affect children, pregnant females, and the elderly with co-morbid conditions. In admitted adults, swine influenza (H<sub>1</sub>N<sub>1</sub>) pneumonia typically presents with high fevers > 102°F often with shaking chills and myalgias. Patients may also complain of headache or sore throat. Dry cough is characteristic but a mildly productive cough with thin mucoid sputum is also common. Patients are variably short of breath. Conjunctival suffusion is rare. Lungs are clear to auscultation. Patients often have loose stools/diarrhea but not abdominal pain. Swine influenza (H<sub>1</sub>N<sub>1</sub>) pneumonia clinically resembles human seasonal influenza, avian influenza, CMV or adenoviral CAP. CXR early (< 48 hours) typically is clear or may show accentuated bibasilar lung markings resembling basilar atelectasis. CXR later (> 48 hours) typically shows bilateral patchy interstitial infiltrates. Consolidation may occur later, but cavitation or moderate/large pleural effusions are not features. Degree of hypoxemia is related to the severity of pneumonia. In adults, nonspecific laboratory clues include otherwise unexplained relative lymphopenia, thrombocytopenia, elevated CPKs or mildly elevated serum transaminases (SGOT/SGPT). Atypical lymphocytes are not present in adults. Unlike in human seasonal and avian influenza (H<sub>2</sub>N<sub>2</sub>), leukopenia is uncommon in swine influenza (H<sub>1</sub>N<sub>1</sub>) pneumonia in adults. Serum LDH is variably elevated. Elevated cold agglutinin titers are not elevated. Nosocomial transmission may occur.

**Diagnostic Considerations:** Definitive diagnosis is by RT-PCR of oropharyngeal/respiratory secretions/lung. Rapid influenza diagnostic screening tests (RIDTs), if positive, correlate well with RT-PCR positivity. However, ~30% of (RIDTs) are false negative. Discordant testing results with rapid influenza A, respiratory FA viral panel, and RT-PCR may be due to variability of specimens rather than differences in test sensitivity/specificity. All respiratory secretion diagnostic tests may be negative in autopsy proven cases of swine influenza (H<sub>1</sub>N<sub>1</sub>) pneumonia whose lungs are RT-PCR positive.

**Pitfalls:** Since swine influenza (H<sub>1</sub>N<sub>1</sub>) pneumonia presents as an ILI, care must be taken to rule out other viral pneumonias as well as mimics of viral pneumonia. The CXR is important in identifying possible mimics of swine flu. While human seasonal influenza may present simultaneously with MSSA/MRSA CAP or subsequently after an interval (5–7 days) of improvement with *S. pneumoniae* or *H. influenzae* CAP, this and has been uncommon with swine influenza (H<sub>1</sub>N<sub>1</sub>) pneumonia and has not been a complication of avian influenza (H<sub>3</sub>N<sub>2</sub>) pneumonia. Swine influenza (H<sub>1</sub>N<sub>1</sub>) pneumonia on CXR appears initially with clear lung fields and with subsequent bilateral patchy infiltrates. Cold agglutinins and atypical lymphocytes may occur in pediatric cases but in adults should suggest an alternate diagnosis, e.g., adenoviral or CMV pneumonia.

**Therapeutic Considerations:** Swine influenza (H<sub>1</sub>N<sub>1</sub>) appears to respond to oseltamivir. There are no benefits to steroids. Although amantadine or rimantadine are ineffective against swine influenza (H<sub>1</sub>N<sub>1</sub>), patients who are severely hypoxemic may benefit from amantadine or rimantadine by increasing peripheral airway dilatation/oxygenation. If the CXR in swine influenza (H<sub>1</sub>N<sub>1</sub>) pneumonia has no lobar/focal infiltrates, there is no need to treat with empiric antimicrobial therapy. Swine influenza (H<sub>1</sub>N<sub>1</sub>) pneumonia with multiple lobar/focal infiltrates that rapidly cavitate (< 72 hours) empiric anti-MSSA/CA-MRSA therapy should be given with oseltamivir. For patients with swine influenza (H<sub>1</sub>N<sub>1</sub>) pneumonia that

improves and after 5–7 days re-present with focal segmental/lobar (non-cavitating) infiltrates, treat as for typical CAP to cover *S. pneumoniae* or *H. influenzae*.

In patients unable to take oral oseltamivir, IV peramivir may be given. (See p. 665 for dosing recommendations.)

**Prognosis:** In immunocompetent adults, prognosis is related to severity of swine influenza ( $H_1N_1$ ) pneumonia. With swine influenza ( $H_1N_1$ ), prognosis is directly related to the severity of pneumonia (degree/duration of severe hypoxemia) and degree/persistence of relative lymphopenia. Prognosis worst with simultaneous MSSA/MRSA CAP, but is not worse with subsequent *S. pneumoniae* or *H. influenzae* CAP.

### ***S. aureus* (MSSA/MRSA) Pneumonia**

**Clinical Presentation:** Only occurs with an antecedent ILI or concurrent influenza pneumonia. Does not occur in DM or those on chronic steroids/immunosuppressive therapy. Presents as a fulminant necrotizing CAP with high fevers, hemoptysis, cyanosis, and hypotension. CXR shows unilateral/bilateral rapid cavitation of pulmonary infiltrates < 72 hrs.

**Diagnostic Considerations:** Patient severely hypoxic with  $\uparrow$  A-a gradient ( $>35$ ) due to underlying influenza pneumonia. Blood/sputum cultures + for MSSA/MRSA.

**Pitfalls:** Culture of MSSA/MRSA from sputum in CAP without rapid cavitation < 72 hrs on CXR, represents colonization, not diagnostic of MSSA/MRSA CAP. Patients CAP due to MSSA/MRSA are critically ill with high spiking fevers, hemoptysis, cyanosis, and hypotension. Diagnosis not supported by MSSA/MRSA in sputum/ blood cultures unless accompanied by the clinical features of MSSA/MRSA pneumonia.

**Therapeutic Considerations:** Treat underlying influenza as well as the superimposed MSSA/MRSA pneumonia with an antibiotic with a high degree of anti- MSSA/MRSA activity, e.g., linezolid (avoid daptomycin). Resection of necrotic lung segments may be life saving.

**Prognosis:** Poor (independent of host factors). Worst prognosis is with PVL + strains.

### **Aspiration Pneumonia**

**Diagnostic Considerations:** Sputum not diagnostic. No need for transtracheal aspirate culture.

**Pitfalls:** Lobar location varies with patient position during aspiration.

**Therapeutic Considerations:** Oral anaerobes are sensitive to all beta-lactams and most antibiotics used to treat CAP. Additional anaerobic (*B. fragilis*) coverage is not needed.

**Prognosis:** Related to severity of CNS/esophageal disease.

### **HIV with CAP (*Sputum Negative for AFB*) (PCP see p. 325)**

**Clinical Presentation:** Bacterial CAP in HIV patient with focal infiltrate(s) and normal/slightly depressed  $CD_4$ .

**Diagnostic Considerations:** Diagnosis by sputum gram stain/culture  $\pm$  positive blood cultures (bacterial pathogens) or Legionella/Chlamydia serology (atypical pathogens). *S. pneumoniae* and *H. influenzae* are particularly common.

**Pitfalls:** Atypical chest x-ray appearance is not uncommon. Treat syndrome of CAP, not chest x-ray. Bacterial CAP (focal/segmental infiltrates) in HIV does not resemble PCP (bilateral patchy interstitial infiltrates). PCP presents with profound hypoxemia and  $\uparrow$  LDH.  $\beta$  1,3 D-glucan +, aspergillus galactomannan –.

**Therapeutic Considerations:** R/O TB/MAI with negative sputum AFB stain. If AFB negative, treat same as bacterial CAP in normal hosts.

**Prognosis:** Same as bacterial CAP in normal hosts.



### **Tuberculous (TB) Pneumonia**

**Clinical Presentation:** Community-acquired pneumonia with single/multiple infiltrates.

**Diagnostic Considerations:** Diagnosis by sputum AFB smear/culture. Primary TB is lower lobe usually with a pleural effusion. Reactivation TB is usually bilateral/apical  $\pm$  old, healed Ghon complex; cavitation/fibrosis are common, but adenopathy or pleural effusions uncommon. TB reactivation  $\uparrow$  with steroids, immunosuppressive drugs, HIV, gastrectomy, jejunioileal bypass, silicosis, DM, CRI, advanced age.

**Pitfalls:** Primary TB may present as CAP and improve transiently with quinolone therapy. Primary TB patients with large pleural effusion usually anergic.

**Therapeutic Considerations:** Usually sputum becomes smear negative in 1–2 weeks of therapy.

**Prognosis:** Related to underlying health status.

### **Mycobacterium avium-intracellulare (MAI) Pneumonia**

**Clinical Presentation:** Community-acquired pneumonia in normal hosts or immunosuppressed/HIV patient with focal single/multiple infiltrates indistinguishable from TB.

**Diagnostic Considerations:** Diagnosis by AFB culture. In HIV, MAI may disseminate, resembling military TB. MAI typically lingular with “tree in bud” appearance on chest CT. BHA more frequent than with TB.

**Pitfalls:** May mimic reactivation TB. Must differentiate TB from MAI by AFB culture, since therapy for MAI differs from TB.

**Therapeutic Considerations:** Should be treated until sputum cultures are negative  $\times$  1 year.

**Prognosis:** Good in normal hosts. In HIV related to degree of immunosuppression/CD<sub>4</sub> count.

### **Mycobacterium kansasii Pneumonia**

**Clinical Presentation:** Subacute CAP resembling reactivation TB.

**Diagnostic Considerations:** Chest x-ray infiltrates/lung disease plus *M. kansasii* in a single sputum specimen. *M. kansasii* can cause disseminated infection, like TB, and is diagnosed by culturing *M. kansasii* from sputum, blood, liver, or bone marrow.

**Pitfalls:** *M. kansasii* in sputum with a normal chest x-ray and no symptoms does not indicate infection.

**Therapeutic Considerations:** Should be treated until sputum cultures are negative  $\times$  1 year.

**Prognosis:** Good in normal hosts. May be rapidly progressive/fatal without treatment in HIV patients.

### **Pneumonia in Chronic Alcoholics**

**Diagnostic Considerations:** *Klebsiella pneumoniae* usually occurs only in chronic alcoholics, and is characterized by blood-flecked “currant jelly” sputum and cavitation (typically in 3–5 days).

**Pitfalls:** Suspect *Klebsiella* in “pneumococcal” pneumonia that cavitates. Empyema is more common than pleural effusion.

**Therapeutic Considerations:** Monotherapy with newer anti-*Klebsiella* agents, e.g., 3<sup>rd</sup> GC, carbapenems, colistin, tigecycline, are as effective or superior to “double-drug” therapy with older agents.

**Prognosis:** Related to degree of hepatic/splenic dysfunction.

### **Bronchiectasis/Cystic Fibrosis**

**Diagnostic Considerations:** Cystic fibrosis/bronchiectasis is characterized by viscous secretions  $\pm$  low grade fevers; less commonly may present as lung abscess. Onset of pneumonia/lung abscess heralded by cough/decrease in pulmonary function.

**Pitfalls:** Sputum colonization is common (e.g., *S. maltophilia*, *B. cepacia*, *P. aeruginosa*); may not reflect pathogens.

**Therapeutic Considerations:** Important to select antibiotics with low resistance potential and good penetration into respiratory secretions (e.g., quinolones, meropenem).

**Prognosis:** Related to extent of underlying lung disease/severity of infection.

### **Pneumonia in Organ Transplants**

**Clinical Presentation:** CAP with perihilar infiltrates and hypoxemia.

**Diagnostic Considerations:** CMV is diagnosed by stain/culture of lung biopsy.

**Therapeutic Considerations:** Treat as CMV pneumonia if CMV is predominant pathogen on lung biopsy. CMV may progress despite ganciclovir therapy.

**Prognosis:** Related to degree of immunosuppression.

### **Pneumonia in Chronic Steroid Therapy**

If fungal infection is suspected, obtain lung biopsy to confirm diagnosis/identify causative organism. Non-responsiveness to appropriate antibiotics should suggest fungal infection. Avoid empirically treating fungi; due to the required duration of therapy, it is advantageous to confirm the diagnosis by lung biopsy first. Prognosis related to degree of immunosuppression.

### **Acute Aspergillus Pneumonia**

**Clinical Presentation:** Chest x-ray shows progressive necrotizing pneumonia (bilateral in half) unresponsive to antibiotics. Characteristic of early Aspergillus pneumonia is the "halo sign". A few days later the halo decreases in size. After a week after the appearance of the halo sign, the "air crescent sign" is typically present. Usually seen only in compromised hosts.

**Diagnostic Considerations:** Diagnosis by lung biopsy (not bronchoalveolar lavage) demonstrating hyphae invading lung parenchyma/blood vessels. Usually occurs only in patients on chronic steroids, cancer chemotherapy, organ transplants, leukopenic compromised hosts, or with chronic granulomatous disease (CGD).  $\beta$  1,3 D-glucan +, aspergillus galactomannan +. Aspergillus galactomannan (GM) may XR with Geotrichum, Blastomyces, Penicillium, or Alternaria.  $\uparrow$  GM levels with cyclophosphamide and if enteritis/colitis ( $\uparrow$  absorption of GM).

**Pitfalls:** Invasive Aspergillus pneumonia does not occur in normal/non-immunosuppressed hosts. PCP has  $\uparrow$  LDH levels and  $\beta$  1,3 D-glucan +, aspergillus galactomannan –.

**Prognosis:** Cavitation is a good prognostic sign. Prognosis related to degree of immunosuppression.

## **Lung Abscess/Empyema**

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	PO Therapy or IV-to-PO Switch
Lung abscess/empyema	Oral anaerobes <i>S. aureus</i> <i>K. pneumoniae</i> <i>S. pneumoniae</i>	Clindamycin 600 mg (IV) q8h* <b>or</b> Piperacillin/tazobactam 3.375 gm (IV) q8h*	Meropenem 1 gm (IV) q8h* <b>or</b> Ertapenem 1 gm (IV) q24h*	Clindamycin 300 mg (PO) q8h* <b>or</b> Quinolone <sup>†</sup> (PO) q24h*

Bronchiectasis, cystic fibrosis (*P. aeruginosa*): see p. 54

\* Treat until resolved. Duration of therapy represents total time IV, PO, or IV + PO. Most patients on IV therapy able to take PO meds should be switched to PO therapy soon after clinical improvement (usually < 72 hours).

† Moxifloxacin 400 mg or Levofloxacin 500 mg.

**Clinical Presentation:** Lung abscess presents as single/multiple cavitory lung lesion(s) with fever. Empyema presents as persistent fever/pleural effusion without layering on lateral decubitus chest x-ray.

**Diagnostic Considerations:** In lung abscess, plain film/CT scan demonstrates cavitory lung lesions appearing > 1 week after pneumonia. Most CAPs are not associated with pleural effusion, and few develop empyema. In empyema, pleural fluid pH is  $\leq 7.2$ ; culture purulent exudate for pathogen.

**Pitfalls:** Pleural effusions secondary to CAP usually resolve rapidly with treatment. Suspect empyema in patients with persistent pleural effusions with fever.

**Therapeutic Considerations:** Chest tube/surgical drainage needed for empyema. Treat lung abscess until it resolves (usually 3–12 months).

**Prognosis:** Good if adequately drained.

## Nursing Home-Acquired Pneumonia (NHAP)

Subset	Usual Pathogens*	Preferred IV Therapy	Alternate IV Therapy	PO Therapy or IV-to-PO Switch
Nursing home-acquired pneumonia (NHAP)	H. influenzae S. pneumoniae M. catarrhalis	Ceftriaxone 1 gm (IV) q24h × 2 weeks <b>or</b> Respiratory quinolone <sup>††</sup> (IV) q24h × 2 weeks <b>or</b> Doxycycline 200 mg (IV) q12h × 3 days; then 100 mg (IV) q12h × 2 weeks	Ertapenem 1 gm (IV) q24h × 2 weeks <b>or</b> Cefepime 2 gm (IV) q12h × 2 weeks	Respiratory quinolone <sup>††</sup> (PO) q24h × 2 weeks <b>or</b> Doxycycline 200 mg (PO) q12h × 3 days, then 100 mg (PO) q12h × 11 days

**Diagnostic Considerations:** H. influenzae is common.

**Pitfalls:** Resembles community-acquired pneumonia in terms of pathogens and length of stay, not nosocomial pneumonia. NHAP outbreaks may be due to H. influenzae, C. pneumoniae, hMPV, or Legionnaire's disease.

**Therapeutic Considerations:** Treat as community-acquired pneumonia, not nosocomial pneumonia. No need to cover P. aeruginosa or GNBs which are common colonizers of nursing home patients, but not causes of NHAP.

**Prognosis:** Related to underlying cardiopulmonary status.

## Nosocomial Pneumonia (NP) / Hospital-Acquired Pneumonia (HAP) / Ventilator-Associated Pneumonia (VAP)<sup>§</sup>

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	IV-to-PO Switch
Empiric therapy	P. aeruginosa* E. coli K. pneumoniae S. marcescens (S. aureus) <sup>†</sup>	Meropenem 1 gm (IV) q8h × 1–2 weeks <b>or</b> Doripenem 1 gm (IV) q8h × 2 weeks <b>or</b> Levofloxacin 750 mg (IV) q24h × 1–2 weeks <b>or</b> Piperacillin/tazobactam 4.5 gm (IV) q6h <b>plus</b> amikacin 1 gm (IV) q24h × 1–2 wks		Levofloxacin 750 mg (PO) q24h × 1–2 weeks <b>or</b> Ciprofloxacin 750 mg (PO) q12h × 1–2 weeks

\* For confirmed P. aeruginosa NP/VAP, combination therapy preferred.

† MSSA/MRSA are common colonizers of respiratory secretions. In ventilated patients with fever, leukocytosis, and infiltrates on CXR, are not diagnostic of MSSA/MRSA NP/HAP/VAP. MSSA/MRSA NP/HAP/VAP is clinically distinctive and rare.

§ Treat aspiration NP same as NP/VAP. Anaerobes not pathogens in aspiration NP.

## Nosocomial Pneumonia (NP) / Hospital-Acquired Pneumonia (HAP) / Ventilator-Associated Pneumonia (VAP) (cont'd)

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	IV-to-PO Switch
Specific therapy	<i>P. aeruginosa</i> **	Meropenem 1 gm (IV) q8h × 2 weeks <b>plus either</b> Levofloxacin 750 mg (IV) q24h × 2 weeks <b>or</b> Amikacin 1 gm (IV) q24h × 2 weeks		Levofloxacin 750 mg (PO) q24h × 2 weeks <b>or</b> Ciprofloxacin 750 mg (PO) q12h × 2 weeks
	MDR Klebsiella or MDR Acinetobacter	Meropenem 1 gm (IV) q8h × 2 weeks <b>or</b> Tigecycline 200 mg (IV) × 1 dose, then 100 mg (IV) q24h × 2 weeks* <b>or</b> Ceftolozane/tazobactam 1.5 gm (IV) q8h × 2 weeks <b>or</b> Colistin 5 mg/kg (IV) q8h × 2 weeks		
	MDR <i>P. aeruginosa</i>	Doripenem 1 gm (IV) q8h × 2 weeks† <b>or</b> Meropenem 1 gm (IV) q8h × 2 weeks‡ <b>or</b> Ceftolozane/tazobactam 1.5 gm (IV) q8h × 2 weeks		
	CRE	Ceftazidime/avibactam 2.5 gm (IV) q8h × 2 weeks <b>or</b> Colistin 5 mg/kg (IV) q8h × 2 weeks		
	HSV-1**	Acyclovir 10 mg/kg (IV) q8h × 10 days		Valacyclovir 1 gm (PO) q8h × 10 days

\* **For empiric *P. aeruginosa* coverage, monotherapy sufficient.** For proven *P. aeruginosa* NP/VAP, combination therapy preferred.

‡ *P. aeruginosa* NP/VAP shows multiple infiltrates with rapid cavitation in <72 h on CXR ± otherwise unexplained blood cultures for *P. aeruginosa* (BCs negative with inhalation acquired *P. aeruginosa* VAP and BCs + with in hematogenously acquired *P. aeruginosa* NP/VAP).

† Give as a 4 hour infusion.

\*\* Presents late in course of VAP as "failure to wean", does *not* present as early VAP.

§ If susceptible.

¶ Higher than usual doses of tigecycline may be necessary for non-susceptible/relatively resistant MDR GNBs.

**Clinical Presentation:** Pulmonary infiltrate compatible with a bacterial pneumonia occurring ≥ 1 week in-hospital ± fever/leukocytosis.

**Diagnostic Considerations:** CXR infiltrates with leukocytosis and fever are nonspecific/nondiagnostic of NP/VAP. Definitive diagnosis by lung biopsy. *P. aeruginosa* is a common colonizer in ventilated patients. *P. aeruginosa* VAP manifests as a necrotizing pneumonia with rapid cavitation (< 72 hours), microabscesses, and blood vessel invasion. *S. aureus* (MSSA/MRSA) NP/VAP remains rare. MSSA/MRSA common colonizers of respiratory secretions. *Acinetobacter*/*Legionella* NP/VAP usually occur in clusters/outbreaks.

**Pitfalls:** No rationale for “covering” non-pulmonary pathogens colonizing respiratory secretions in ventilated patients (Enterobacter, B. cepacia, S. maltophilia, Citrobacter, Flavobacterium, Enterococci); these organisms rarely if ever cause NP/VAP. Semi-quantitative BAL/protected brush specimens often reflect airway colonization. Characteristic clinical presentation (necrotizing pneumonia with rapid cavitation < 72 hrs) or tissue biopsy is needed for diagnosis of P. aeruginosa NP/VAP. MSSA/MRSA are common colonizers of respiratory secretions and are not uncommon causes of tracheobronchitis in ventilated patients. While MSSA/MRSA colonization of respiratory secretions is common, MSSA/MRSA necrotizing/rapidly cavitating NP/VAP is rare.

**Therapeutic Considerations:** NP (HAP/VAP) is defined as pneumonia acquired > 5 hospital days. So called “early” NP occurring < 5 days of hospitalization, usually due to S. pneumoniae or H. influenzae, represents CAP that manifests early after hospital admission. Empiric therapy recommendations cover usual NP/VAP pathogens. Monotherapy is as effective as combination therapy for non-P. aeruginosa NP/VAP, but 2-drug therapy is recommended for confirmed P. aeruginosa NP/VAP. If MSSA/MRSA necrotizing/rapidly cavitating NP/VAP present, may be treated with vancomycin, quinupristin/dalfopristin, tigecycline, or linezolid. After 2 weeks of appropriate antibiotic therapy, nonprogressive/stable pulmonary infiltrates with fever and leukocytosis are usually due to a noninfectious cause, rather than persistent infection. Treat aspiration NP/VAP the same as NP/VAP since anaerobes are not important pathogens in NP/VAP.

**Prognosis:** Related to underlying cardiopulmonary status.

## Empiric Therapy of Cardiovascular Infections

### Native Valve Subacute Bacterial Endocarditis (SBE)

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	PO Therapy or IV-to-PO Switch
No obvious source or oral	Viridans streptococci*† Groups B, C, G streptococci, S. bovis (PCN MIC < 0.12 mcg/ml)	Ceftriaxone 2 gm (IV) q24h × 2 weeks*‡ <b>plus</b> Gentamicin 240 mg or 3 mg/kg (IV) q24h × 2 weeks* <b>or</b> <b>monotherapy</b> <b>with</b> Ceftriaxone 2 gm (IV) q24h × 4 weeks	Penicillin G 3 mu (IV) q4h × 2 weeks*‡ <b>plus</b> Gentamicin 240 mg or 3 mg/kg (IV) q24h × 2 weeks* <b>or</b> Vancomycin 1 gm (IV) q12h × 2 weeks <b>or</b> Linezolid 600 mg (IV) q12h × 4 weeks	Linezolid <sup>§</sup> 600 mg (PO) q12h × 4–6 weeks <b>or</b> Amoxicillin <sup>§</sup> 1 gm (PO) q8h × 4–6 weeks

\* If relatively PCN resistant (MIC 0.12–0.5 mcg/ml) or if no intra/extra cardiac complications treat × 4 weeks.

† S. mitis (S. mitior), S. oralis, S. gordonii, S. parasanguis, S. sanguis, S. mutans, S. sobrinus, S. thermophilus, S. angiosus, S. milleri, S. intermedius, S. constellatus.

‡ In PCN allergic patients, Vancomycin 1 gm (IV) q12h × 2 weeks.

§ IV therapy preferred. PO therapy if no IV access or if patient not a surgical candidate.

## Native Valve Subacute Bacterial Endocarditis (SBE) (cont'd)

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	PO Therapy or IV-to-PO Switch
	Nutritionally-variant streptococci (NVS) pyridoxal B <sub>6</sub> dependent streptococci <sup>‡</sup>	As above, but treat × 6 weeks	As above, but treat × 6 weeks	As above <sup>‡</sup> , but treat × 6 weeks
	Listeria monocytogenes	Ampicillin 2 gm (IV) q4h × 4–6 weeks <b>or</b> TMP-SMX 5 mg/kg (IV) q6h	Meropenem 1 gm (IV) q8h × 4–6 weeks	Amoxicillin <sup>§</sup> 1 gm (PO) q8h × 4–6 weeks <b>or</b> Linezolid <sup>§</sup> 600 mg (PO) q12h × 4–6 weeks
GI/GU source likely  (Treat initially for E. faecalis (VSE); if later identified as E. faecium (VRE), treat accordingly)	E. faecalis (VSE) <sup>†</sup>	Ampicillin 2 gm (IV) q4h × 4–6 weeks <b>plus</b> Gentamicin* 120 mg or 1 mg/kg (IV) or 80 mg (IV) q8h × 4–6 weeks	Vancomycin 1 gm or 15 mg/kg (IV) q12h × 4–6 weeks <b>plus</b> Gentamicin* 120 mg or 1 mg/kg (IV) q24h × 2 weeks	Amoxicillin <sup>§</sup> 1 gm (PO) q8h × 4–6 weeks <b>or</b> Linezolid <sup>§</sup> 600 mg (PO) q12h × 4–6 weeks
	E. faecium (VRE) <sup>†</sup>	Linezolid 600 mg (IV) q12h × ≥ 8 weeks <b>or</b> Daptomycin 12 mg/kg (IV) q24h × 6–8 weeks		Linezolid <sup>§</sup> 600 mg (PO) q12h × ≥ 8 weeks
GI source	S. (bovis) gallolyticus (Non-enterococcal group D streptococci)	Treat the same as “no obvious source” subset (see p. 70)		

VRE = vancomycin-resistant enterococci. Duration of therapy represents total time IV, PO, or IV + PO.

† Symptoms < 3 months → treat × 4 weeks.

Symptoms > 3 months → treat × 6 weeks.

‡ NVS (*nutritionally variant streptococci*) include: Abiotrophia defectiva, Granulicata adjacens/elegans.

§ IV therapy preferred. PO therapy if no IV access or if patient not a surgical candidate.

\* If gentamicin resistant, give streptomycin 7.5 mg/kg (IV/1M) q12h.

## Native Valve Subacute Bacterial Endocarditis (SBE) (cont'd)

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	PO Therapy or IV-to-PO Switch
"Culture negative" SBE*	Pathogen unknown (Abiotrophia, enterococci) PCN resistant organisms	Ampicillin/ Sulbactam 3 gm (IV) q6h x 4–6 weeks  <b>plus</b> Gentamicin <sup>§</sup> 120 mg or 1 mg/kg (IV) q24h x 2 weeks	Quinolone <sup>†</sup> (IV) q24h x 4–6 weeks	Quinolone <sup>15</sup> (PO) q24h x 4–6 weeks
	Hemophilus sp. Actinobacillus actinomycetem-comitans Cardiobacterium hominis Eikenella corrodens Kingella kingae (HACEK organisms)	Ceftriaxone 2 gm (IV) q24h x 4–6 weeks  <b>or</b> Any 3 <sup>rd</sup> generation cephalosporin (IV) x 4–6 weeks  <b>or</b> Cefepime 2 gm (IV) q12h x 4–6 weeks	Ampicillin 2 gm (IV) q4h x 4–6 weeks <b>plus</b> Gentamicin <sup>§</sup> 120 mg or 1 mg/kg (IV) q24h x 2 weeks  <b>or</b> <b>monotherapy with</b> Ampicillin/ Sulbactam 3 gm (IV) q6h x 4–6 weeks  <b>or</b> Quinolone <sup>†</sup> (IV) q24h x 4–6 weeks	Quinolone <sup>15</sup> (PO) q24h x 4–6 weeks
	Legionella Chlamydia (Chlamydia) psittaci	Doxycycline <sup>**</sup> 200 mg (IV) q12h x 3 days, then 100 mg (IV) q12h x 3 months	Quinolone <sup>†</sup> (IV) q24h x 3 months	Doxycycline <sup>**§</sup> 200 mg (PO) q12h x 3 days, then 100 mg (PO) q12h x 3 months
	Coxiella burnetii (Q fever)	Doxycycline <sup>**</sup> 200 mg (IV/PO) q12h x 3 days, then 100 mg (PO) q12h x 24 months  <b>plus</b> Hydroxchloroquin 200 mg (PO) q8h x 24 months		

\* Slow growing/fastidious organisms may require ↑ CO<sub>2</sub> and prolonged incubation.

† Levofloxacin 500 mg or Moxifloxacin 400 mg.

\*\* Loading dose is not needed PO if given IV with the same drug.

§ IV therapy preferred. PO therapy if no IV access or if patient not a surgical candidate.

## Native Valve Subacute Bacterial Endocarditis (SBE) (cont'd)

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	PO Therapy or IV-to-PO Switch
Brucella SBE†	Brucella melitensis, abortus, canis, suis	Doxycycline 200 mg** (IV) q12h × 3 days, then 100 mg (PO) q12h × ≥ 6 weeks		
			<b>plus</b>	
			Rifampin 300 mg (PO) q8h × ≥ 6 weeks	
			<b>plus</b>	
			Gentamicin <sup>§§</sup> 3 mg/kg (IV) q24h × ≥ 6 weeks	

† Levofloxacin 500 mg or Moxifloxacin 400 mg.

\*\* Loading dose is not needed PO if given IV with the same drug or if patient not a surgical candidate.

§§ May be given as a once daily dose or given as divided doses q8h.

### Native Valve SBE

**Clinical Presentation:** Subacute febrile illness ± localizing symptoms/signs in a patient with a heart murmur. Peripheral manifestations are commonly absent with early diagnosis/treatment.

**Diagnosis:** High grade/continuous bacteremia (due to a endocarditis pathogen) plus vegetation on TTE/TEE is diagnostic.

### SBE (No Obvious Source)

**Diagnostic Considerations:** Most common pathogen is viridans streptococci. Source is usually the mouth, although oral/dental infection is usually inapparent clinically.

**Pitfalls:** Vegetations without positive blood cultures or peripheral manifestations of SBE are not diagnostic of endocarditis. SBE vegetations may persist after antibiotic therapy, but are sterile.

**Therapeutic Considerations:** In penicillin-allergic (anaphylactic) patients, vancomycin may be used alone or in combination with gentamicin. Follow ESR weekly to monitor antibiotic response. No need to repeat blood cultures unless patient has persistent fever or is not responding clinically. Two-week treatment is acceptable for uncomplicated viridans streptococcal SBE. Treat nutritionally-variant streptococci (NVS) (*B<sub>g</sub>*/pyridoxal dependent streptococci) same as viridans streptococcal SBE.

**Prognosis:** Related to extent of embolization/severity of heart failure.

### SBE (GI/GU Source Likely)

**Diagnostic Considerations:** Commonest pathogens from GI/GU source are Enterococci (especially *E. faecalis*). If *S. bovis*, look for GI polyp, tumor source. Enterococcal SBE commonly follows GI/GU instrumentation.

**Pitfalls:** Low back pain prominent with enterococcal/*S. bovis* SBE, *S. bovis* associated with CNS embolization, Vertebral osteomyelitis, and large valvular vegetations.

**Therapeutic Considerations:** For penicillin-allergic patients, use vancomycin plus gentamicin. Vancomycin alone is inadequate for enterococcal (*E. faecalis*) SBE. Treat enterococcal PVE the same as for native valve enterococcal SBE. Treat *S. bovis* SBE the same as *S. viridans* SBE.

**Prognosis:** Related to extent of embolization.



**“Culture Negative” SBE (Culturable Pathogens)**

**Diagnostic Considerations:** Culture of organisms may require enhanced CO<sub>2</sub>/special media (Castaneda vented bottles) and prolonged incubation (2–4 weeks). True “culture negative” SBE is rare, and is characterized by peripheral signs of SBE, a murmur, vegetation (by TTE/TEE), and negative blood cultures.

**Pitfalls:** Most cases of “culture negative” SBE are not, in fact, culture negative SBE, but are due to slow growing organisms. Vegetations may not be visible by TTE/TEE with Q fever SBE.

**Therapeutic Considerations:** Follow clinical improvement with serial ESRs, which should return to pretreatment levels with therapy. Serial TTEs/TEEs show decreasing vegetation size during effective therapy. Sterile vegetations may persist after antibiotic therapy.

**Prognosis:** Related to extent of embolization.

**“Culture Negative” SBE Non-Culturable/(Serologically Diagnosed Pathogens)**

**Diagnostic Considerations:** Diagnosis by specific serology. Large vessel emboli suggests culture negative SBE in patients with negative blood cultures but signs of SBE, e.g., splenomegaly/peripheral manifestation’s.

**Pitfalls:** Do not diagnose culture negative SBE in patients with a heart murmur and negative blood cultures if peripheral SBE manifestations are absent. Vegetations may not be visible on TTE/TEE in Q fever SBE. The diagnosis of chronic Q fever is made by demonstrating elevated IgG (phase I) titers  $\geq 1:1024$  at 6 months following acute Q fever. Since Bartonella titers may cross react with C. burnetii (Q fever) titers and vice versa, an elevation of either titer may be a clue to the other and should prompt testing for both.

**Therapeutic Considerations:** Treatment is based on specific organism identified by diagnostic tests.

**Prognosis:** Related to extent of embolization/severity of heart failure.

**Native Valve Acute Bacterial Endocarditis (ABE)**

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	IV-to-PO Switch Therapy
Normal hosts*  (Treat initially for MRSA; if later identified as MSSA, treat accordingly)	S. aureus (MRSA)	Daptomycin 12 mg/kg (IV) q24h × 4–6 weeks  <b>or</b> Linezolid 600 mg (IV) q24h × 4–6 weeks  <b>or</b> Vancomycin 1 gm or 15 mg/kg (IV) q12h × 4–6 weeks  <b>or</b> Minocycline 100 mg (IV) q12h × 4–6 weeks		Linezolid 600 mg (PO) q12h × 4–6 wks  <b>or</b> Minocycline 100 mg (PO) q12h × 4–6 wks
	S. aureus (MSSA)	Nafcillin 2 gm (IV) q4h × 4–6 weeks  <b>or</b> Cefazolin 1 gm (IV) q8h × 4–6 weeks		Linezolid 600 mg (PO) q12h × 4–6 wks  <b>or</b> Cephalexin 1 gm (PO) q6h × 4–6 weeks

MRSA/MSSA = methicillin-resistant/sensitive S. aureus. Duration of therapy represents total time IV, PO, or IV + PO.

\* Treat IV or with IV-to-PO switch therapy or if patient not a surgical candidate.

## Native Valve Acute Bacterial Endocarditis (ABE) (cont'd)

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	IV-to-PO Switch Therapy
IV drug abusers (IVDAs) <sup>†</sup>  (Treat as MRSA before culture results; treat according to pathogen after culture results)	S. aureus (MRSA)	<u>Before culture results</u> Vancomycin 1 gm or 15 mg/kg (IV) q12h  <b>or</b> Daptomycin 12 mg/kg (IV) q24h × 4–6 weeks	<u>After culture results</u> Linezolid 600 mg (IV) q12h × 4–6 weeks  <b>or</b> Vancomycin 1 gm or 15 mg/kg (IV) q12h × 4–6 weeks  <b>or</b> Minocycline 100 mg (IV) q12h × 4–6 weeks	<u>After culture results</u> Linezolid 600 mg (PO) q12h × 4–6 weeks
	S. aureus (MSSA)	<u>Before culture results</u> Meropenem 1 gm (IV) q8h	<u>After culture results</u> Nafcillin 2 gm (IV) q4h × 4–6 weeks <b>or</b> Cefazolin 1 gm (IV) q8h × 4–6 weeks <b>or</b> Daptomycin 6 mg/kg (IV) q24h × 4–6 weeks <b>or</b> Linezolid 600 mg (IV) q12h × 4–6 weeks <b>or</b> Meropenem 1 gm (IV) q8h × 4–6 weeks	<u>After culture results</u> Linezolid 600 mg (PO) q12h × 4–6 weeks  <b>or</b> Cephalexin 1 gm (PO) q6h × 4–6 weeks  <b>or</b> Minocycline <sup>†</sup> 100 mg (PO) q12h × 4–6 weeks
	P. aeruginosa* S. marcescens Aerobic GNBS	<u>Before culture results</u> Meropenem 1 gm (IV) q8h	<u>After culture results</u> <b>One "A" drug ± one "B" drug</b> <b>"A" Drugs</b> Meropenem 1 gm (IV) q8h × 4–6 weeks <b>or</b> Doripenem 1 gm (IV) q8h × 4–6 weeks <b>"B" Drugs</b> Amikacin 1 gm or 15 mg/kg (IV) q24h × 4–6 weeks <b>or</b> Aztreonam 2 gm (IV) q8h × 4–6 weeks	None

MRSA/MSSA = methicillin-resistant/sensitive *S. aureus*. Duration of therapy represents total time IV, PO, or IV + PO.

\* Treat IV or with IV-to-PO switch therapy or if patient not a surgical candidate.

<sup>†</sup> MSSA/MRSA TV ABE may be treated IV/PO × 2 weeks if no intra-cardiac or CNS complications. Pulmonary septic emboli are a feature of TV ABE e.g., and are *not*, per se, an extracardiac complication.

### **Acute Bacterial Endocarditis (ABE)**

**Diagnostic Considerations:** Clinical criteria for MRSA/MSSA ABE: continuous/high-grade bacteremia (repeatedly 3/4 or 4/4 positive blood cultures), fever (temperature usually  $\geq 102^{\circ}\text{F}$ ), no, new or changing murmur, and vegetation by transesophageal/trans thoracic echocardiogram.

**Pitfalls:** Obtain a baseline TTE for comparative purposes should ABE be complicated by valve destruction, heart failure or ring/perivalvular abscess. On TTE/TEE a vegetation may not be visible until  $> 1$  week of ABE. In MSSA/MRSA ABE, resistance to daptomycin may occur during therapy particularly in patients initially treated with vancomycin.

**Therapeutic Considerations:** Treat for 4–6 weeks. Follow teichoic acid antibody titers (TAA) weekly in MSSA/MRSA ABE, which  $\downarrow$  (along with the ESR) with effective therapy. If MSSA/MRSA ABE unresponsive (persistent high grade bacteremia) to appropriate MSSA/MRSA therapy or if a myocardial/paravalvular abscess present that cannot be surgically drained, “high dose” daptomycin 12 mg/kg (IV) q24h may be effective. With daptomycin resistant MSSA/MRSA strains, Quinupristin/dalfopristin, linezolid or minocycline may be effective.

**Prognosis:** Related to extent of embolization/severity of valve destruction/heart failure.

### **Acute Bacterial Endocarditis (IVDAs)**

**Diagnostic Considerations:** Clinically IVDAs with *S. aureus* usually have relatively mild ABE, permitting oral treatment.

**Pitfalls:** IVDAs with new aortic or tricuspid regurgitation should be treated IV  $\pm$  valve replacement.

**Therapeutic Considerations:** After pathogen is isolated, may switch from IV to PO regimen to complete treatment course.

**Prognosis:** Prognosis is better than ABE in normal hosts if not complicated by abscess, valve regurgitation, or heart failure.

**Clinical Presentation:** Prolonged fevers and chills following prosthetic valve replacement (PVR).

**Diagnosis:** High-grade blood culture positivity (3/4 or 4/4) with endocarditis pathogen and no other source of infection.

### **Prosthetic Valve Endocarditis (PVE)**

#### **Early PVE (< 60 days post-PVR)**

**Diagnostic Considerations:** Blood cultures persistently positive. Temperature usually  $\leq 102^{\circ}\text{F}$ .

**Pitfalls:** Obtain baseline TTE/TEE. Premature closure of mitral leaflet is early sign of impending aortic valve regurgitation. Rifampin should be given 2 days after strain shown to be susceptible.

**Therapeutic Considerations:** Patients improve clinically on treatment, but prosthetic valve removal may be necessary for cure.

**Prognosis:** Related to extent of embolization/severity of heart failure.

#### **Late PVE (> 60 days post-PVR)**

**Pitfalls:** Culture of removed valve may be negative, but valve gram stain will be positive.

**Therapeutic Considerations:** Clinically, late PVE resembles viridans streptococcal SBE. Patients improve clinically on treatment, but prosthetic valve removal may be necessary for cure.

**Prognosis:** Related to extent of embolization/severity of heart failure.

## Prosthetic Valve Endocarditis (PVE)

Subset	Usual Pathogens	Before Culture Results	After Culture Results
Early PVE (< 60 days post-PVR) <sup>§</sup>	<i>S. aureus</i> (MSSA/MRSA) <i>S. epidermidis</i> (CoNS) Enterobacteriaceae	Vancomycin 1 gm or 15 mg/kg (IV) q12h <b>plus</b> Gentamicin 120 mg or 1 mg/kg (IV) q24h	<u>Enterobacteriaceae</u> Meropenem 1 gm (IV) q8h × 6 weeks <b>or</b> <u>MSSA</u> Cefazolin 1 gm (IV) q8h × 6 weeks <b>or</b> Meropenem 1 gm (IV) q8h × 6 weeks <u>MRSA</u> <sup>†</sup> Daptomycin 12 mg/kg (IV) q24h × 6 weeks <b>or</b> Linezolid 600 mg (IV or PO) q12h × 6 weeks <b>or</b> Vancomycin 1 gm or 15 mg/kg (IV) q12h × 6 weeks <b>plus</b> Gentamicin 120 mg (IV) q24h × 2 weeks <b>plus</b> Rifampin 300 mg (IV) q24h × 6 weeks*
	<i>Candida</i> species <sup>§</sup>	Mycafungin 150 mg or 3 mg/kg (IV) q24h	Lipid amphotericin 3–5 mg/kg (IV) q24h × 6 weeks <b>plus</b> 5-FC 25 mg/kg (PO) q6h × 6 weeks
Late PVE (> 60 days post-PVR) <sup>§</sup>	Viridans streptococci <i>S. epidermidis</i> (CoNS)	Vancomycin 1 gm or 15 mg/kg (IV) q12h <b>plus</b> Gentamicin 120 mg or 1 mg/kg (IV) q24h	<u>Viridans streptococci</u> Ceftriaxone 2 gm (IV) q24h × 6 weeks <u>CoNS</u> Linezolid 600 mg (IV or PO) q12h × 6 weeks <b>or</b> Vancomycin 1 gm or 15 mg/kg (IV) q12h × 6 weeks <b>plus</b> Rifampin 300 mg (PO) q8h × 6 weeks

MSSA/MRSA = methicillin-sensitive/resistant *S. aureus*; MSSE/MRSE = methicillin-sensitive/resistant *S. epidermidis*. Duration of therapy represents total time IV or IV + PO.

† MRSA drugs also effective against MSSA.

§ Valve removal usually needed for cure.

\* Begin 2 days after pathogen known to be susceptible.

## Pericarditis/Myocarditis

Subset	Usual Pathogens	Preferred Therapy
Diphtheritic myocarditis	C. diphtheriae	Treat same as diphtheriae (see p. 170)
Viral	Coxsackie, influenza	No treatment for Coxsackie, for influenza therapy
Lyme myocarditis	B. burgdorferi	Ceftriaxone 2 gm (IV) q24h x 2 weeks  <b>or</b> Doxycycline 100 mg (IV) q12h x 2 weeks
TB pericarditis	M. tuberculosis	Treat same as pulmonary TB (see p. 53). Add a tapering dose of corticosteroids x 4–8 weeks
Suppurative pericarditis	S. pneumoniae S. aureus	Treat same as lung abscess/empyema (see p. 67)

**Clinical Presentation:** Viral pericarditis presents with acute onset of fever/chest pain (made worse by sitting up) following a viral illness. TB pericarditis is indolent in presentation, with ↑ jugular venous distension (JVD), pericardial friction rub (40%), paradoxical pulse (25%), and chest x-ray with cardiomegaly ± left-sided pleural effusion. Suppurative pericarditis presents as acute pericarditis (patients are critically ill). Develops from contiguous (e.g., pneumonia) or hematogenous spread e.g., S. aureus bacteremia. Viral myocarditis presents with heart failure, arrhythmias ± emboli.

**Diagnostic Considerations:** Pericarditis/effusion manifests cardiomegaly with decreased heart sounds ± tamponade. Diagnosis by culture/biopsy of pericardial fluid or pericardium for viruses, bacteria, or acid-fast bacilli (AFB). Diagnosis of myocarditis is clinical ± myocardial biopsy.

**Pitfalls:** Consider other causes of pericardial effusion (malignancy, esp. if bloody effusion, uremia, etc). Otherwise unexplained tachycardia should suggest myocarditis until proven otherwise. Rule out treatable non-viral causes of myocarditis e.g., RMSF, Lyme, diphtheria. Suspect Lyme disease in any person without heart disease with exposure to endemic area who presents with otherwise unexplained heart block. Lyme IgM titers may be negative early when patients present with heart block.

**Therapeutic Considerations:** No specific treatment for viral myocarditis/pericarditis. TB pericarditis is treated the same as pulmonary TB ± pericardiectomy. Suppurative pericarditis is treated the same as lung abscess plus surgical drainage (pericardial window). Heart block in Lyme disease rapidly reverses with therapy, but a temporary pacemaker may be needed until heart block is reversed.

**Prognosis:** For viral pericarditis, the prognosis is good, but viral myocarditis may be fatal. For TB pericarditis, the prognosis is good if treated before constrictive pericarditis/adhesions develop. Suppurative pericarditis is often fatal without early pericardial window/antibiotic therapy. With Lyme myocarditis/heart block, prognosis is good if treatment is started early when diagnosis is suspected.

## Central Venous Catheter (CVC) and Pacemaker/Lead Infections

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	IV-to-PO Switch
<b>CVC (temporary) Infections*</b>  <i>Bacterial</i>  (Treat initially for MSSA; if later identified as MRSA, treat accordingly)	S. aureus (MSSA) Aerobic GNBS	Meropenem 1 gm (IV) q8h**  <b>or</b> Cefepime 2 gm (IV) q12h**	Ceftriaxone 2 gm (IV) q24h**	Quinolone <sup>†</sup> (PO) q24h***
	S. aureus (MRSA/MSSA <sup>§</sup> )	Daptomycin 12 mg/kg (IV) q24h**  <b>or</b> Linezolid 600 mg (IV) q12h**  <b>or</b> Vancomycin 1 gm or 15 mg/kg (IV) q12h**		Linezolid 600 mg (PO) q12h***  <b>or</b> Minocycline 100 mg (PO) q12h***
	<i>Candida</i> <sup>*</sup>	C. albicans	Treat the same as for Candidemia (see p. 153)	
	(Treat initially for C. albicans; if later identified as non-albicans Candida, treat accordingly).	Non-albicans Candida	Treat the same as for Candidemia (see p. 153)	
<b>CVC (semi-permanent Hickman/Broviac/Tessio) Infections*</b>  <i>Bacterial</i>	S. aureus (MSSA/MRSA)	Daptomycin 12 mg/kg (IV) q24h**  <b>or</b> Linezolid 600 mg (IV) q12h**  <b>or</b> Vancomycin 1 gm or 15 mg/kg (IV) q12h**  <b>or</b> Minocycline 100 mg (IV) q12h**		Linezolid 600 mg (PO) q12h***  <b>or</b> Minocycline 100 mg (PO) q12h***

MSSA/MRSA = methicillin-sensitive/resistant *S. aureus*.

† IV therapy preferred. PO therapy if no IV access

‡ Levofloxacin 500 mg or Moxifloxacin 400 mg.

\* If clinically possible, CVC should be removed if CVC suspected source of bacteremia in a septic patient. If high grade/sustained bacteremia obtain TTE/TEE to rule out ABE.

\*\* Treat × 2 weeks after CVC removal if no ABE.

¶ For candidemia not associated with CVCs, (see pp. 153–154).

§ MRSA drugs also effective against MSSA.

## Central Venous Catheter (CVC) and Pacemaker/Lead Infections (cont'd)

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	IV-to-PO Switch
	<i>S. epidermidis</i> (CoNS)	Daptomycin 6 mg/kg (IV) q24h** <b>or</b> Linezolid 600 mg (IV) q12h** <b>or</b> Vancomycin 1 gm or 15 mg/kg (IV) q12h**		Linezolid 600 mg (PO) q12h**†
Pacemaker wire/generator infection, LVAD, ICD (Treat initially for <i>S. aureus</i> ; if later identified as <i>S. epidermidis</i> , treat accordingly)	<i>S. aureus</i> (MSSA/MRSA)	Daptomycin 6 mg/kg (IV) q24h <sup>†</sup> <b>or</b> Linezolid 600 mg (IV) q12h <sup>†</sup> <b>or</b> Vancomycin 1 gm or 15 mg/kg (IV) q12h <sup>†</sup>		Linezolid 600 mg (PO) q12h <sup>†</sup> ** <b>or</b> Minocycline 100 mg (PO) q12h <sup>†</sup> **
	<i>S. epidermidis</i> (CoNS)	Daptomycin 6 mg/kg (IV) q24h <sup>†</sup> <b>or</b> Linezolid 600 mg (IV) q12h <sup>†</sup> <b>or</b> Vancomycin 1 gm or 15 mg/kg (IV) q12h <sup>†</sup>		Linezolid 600 mg (PO) q12h <sup>†</sup> *

MSSA/MRSA = methicillin-sensitive/resistant *S. aureus*; MSSE/MRSE = methicillin-sensitive/resistant *S. epidermidis*.

† IV therapy preferred. PO therapy if no IV access.

\*\* Treat × 2 weeks after CVC removal if no IE.

\* Treat × 2 weeks after wire/generator removal if no IE.

§ Obtain teichoic acid antibody titers (TAA) after 2 weeks. If titers are < 1:4, 2 weeks of therapy is sufficient. If titers are ≥ 1:4 rule out endocarditis and complete 4–6 weeks of therapy.

## Central Venous Catheter (CVC) and Pacemaker/Lead Infections (cont'd)

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	IV-to-PO Switch
Septic thrombophlebitis  (Treat initially for MSSA; if later identified as MRSA, treat accordingly)	S. aureus (MSSA)	Nafcillin 2 gm (IV) q4h x 2 weeks* <b>or</b> Linezolid 600 mg (IV) q12h x 2 weeks* <b>or</b> Meropenem 1 gm (IV) q8h x 2 weeks*	Cefazolin 1 gm (IV) q8h x 2 weeks* <b>or</b> Ceftriaxone 1 gm (IV) q12h x 2 weeks* <b>or</b> Minocycline 100 mg (IV) q12h x 2 weeks	Linezolid 600 mg (PO) q12h x 2 weeks*† <b>or</b> Clindamycin 300 mg (PO) q8h x 2 weeks*† <b>or</b> Cephalexin 1 gm (PO) q6h x 2 weeks*†
	S. aureus (MRSA)	Linezolid 600 mg (IV) q12h x 2 weeks* <b>or</b> Vancomycin 1 gm or 15 mg/kg (IV) q12h x 2 weeks* <b>or</b> Minocycline 100 mg (IV) q12h x 2 weeks*		Linezolid 600 mg (PO) q12h x 2 weeks*† <b>or</b> Minocycline 100 mg (PO) q12h x 2 weeks*†

MSSA/MRSA = methicillin-sensitive/resistant *S. aureus*. Duration of therapy represents total time IV or IV + PO. Most patients on IV therapy able to take PO meds should be switched to PO therapy after clinical improvement.

\* Obtain teichoic acid antibody titers after 2 weeks. If titers are < 1:2, 2 weeks of therapy is sufficient. If titers are ≥ 1:2, TTE/TEE to rule out ABE and complete 4–6 weeks of therapy.

† IV therapy preferred. PO therapy if no IV access.

### CVC (Temporary) Infections

**Clinical Presentation:** Temperature ≥ 102°F ± IV site erythema.

**Diagnostic Considerations:** Diagnosis by semi-quantitative catheter tip culture with ≥ 15 colonies plus blood cultures with same pathogen. If no other explanation for fever and line has been in place ≥ 7 days, remove line and obtain semi-quantitative catheter tip culture. Suppurative thrombophlebitis presents with hectic/septic fevers and pus at IV site ± palpable venous cord.

**Pitfalls:** Temperature ≥ 102°F with IV line infection, in contrast to phlebitis.

**Therapeutic Considerations:** Line removal is usually curative, but antibiotic therapy is usually given for 1 week after IV line removal for gram-negative bacilli or 2 weeks after IV line removal for *S. aureus* (MSSA/MRSA). Antifungal therapy is usually given for 2 weeks after IV line removal for Candidemia.

**Prognosis:** Good if line is removed before endocarditis/metastatic spread.



### **CVC (Semi-Permanent) Hickman/Broviac Infections**

**Clinical Presentation:** Fever  $\pm$  IV site erythema.

**Diagnostic Considerations:** Positive blood cultures plus gallium scan pickup on catheter is diagnostic.

**Pitfalls:** Antibiotics will lower temperature, but patient will usually not be afebrile without line removal.

**Therapeutic Considerations:** Lines usually need to be removed for cure. Rifampin 600 mg (PO) q24h may be added to IV/PO regimen if pathogen is *S. aureus*.

**Prognosis:** Good with organisms of low virulence.

### **Pacemaker/Lead Infections**

**Clinical Presentation:** Persistently positive blood cultures without endocarditis in a pacemaker patient.

**Diagnostic Considerations:** Positive blood cultures with gallium scan pickup on wire/pacemaker generator is diagnostic. Differentiate wire from pacemaker pocket infection by chest CT/MRI.

**Pitfalls:** Positive blood cultures are more common in wire infections than pocket infections. Blood cultures may be negative in both, but more so with pocket infections.

**Therapeutic Considerations:** Wire alone may be replaced if infection does not involve pacemaker generator. Replace pacemaker generator if involved; wire if uninvolved can usually be left in place.

**Prognosis:** Good if pacemaker wire/generator replaced before septic complications develop.

### **Septic Thrombophlebitis**

**Clinical Presentation:** Temperature  $\geq 102^\circ\text{F}$  with local erythema and signs of sepsis.

**Diagnostic Considerations:** Palpable venous cord and pus at IV site when IV line is removed.

**Pitfalls:** Suspect diagnosis if persistent bacteremia and no other source of infection in a patient with a peripheral IV.

**Therapeutic Considerations:** Remove IV catheter. Surgical venotomy is usually needed for cure.

**Prognosis:** Good if removed early before septic complications develop.

## **Vascular Graft Infections**

<b>Subset</b>	<b>Usual Pathogens</b>	<b>Preferred IV Therapy</b>	<b>Alternate IV Therapy</b>	<b>IV-to-PO Switch</b>
AV graft/shunt infection  (Treat initially for MRSA, etc.; if later identified as MSSA, treat accordingly)	<i>S. aureus</i> (MRSA)	Daptomycin* <sup>§</sup> 12 mg/kg (IV) q24h <b>plus</b> Rifampin 300 mg (PO) q8h <b>or</b> Linezolid <sup>†§</sup> 600 mg (IV) q12h <b>plus</b> Rifampin <sup>†§</sup> 300 mg (PO) q8h	Vancomycin* <sup>§</sup> 1 gm or 15 mg/kg (IV) q12h <b>plus</b> Rifampin 300 mg (PO) q8h <b>or</b> Minocycline <sup>†§</sup> 100 mg (IV) q12h <b>plus</b> Rifampin <sup>†§</sup> 300 mg (PO) q8h	Linezolid <sup>†§</sup> 600 mg (PO) q12h <b>plus</b> Rifampin 300 mg (PO) q8h <b>or</b> Minocycline <sup>†§</sup> 100 mg (PO) q12h <b>plus</b> Rifampin <sup>†§</sup> 300 mg (PO) q8h

MRSA/MSSA = methicillin-resistant/sensitive *S. aureus*. Duration of therapy represents total time IV or IV + PO. Most patients on IV therapy able to take PO meds should be switched to PO therapy after clinical improvement.

\* Follow with maintenance dosing for renal failure (CrCl < 10 mL/min) and type of dialysis.

† If ABE not present, treat for 2 weeks after graft is removed/replaced. If ABE present, treat for 6 weeks.

§ IV therapy preferred. PO therapy if no IV access.

## Vascular Graft Infections (cont'd)

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	IV-to-PO Switch
	<i>S. aureus</i> (MSSA)	Nafcillin 3 gm (IV) q12h <sup>†</sup>	Vancomycin 1 gm or 15 mg/kg (IV) q12h <sup>*†</sup>	Moxifloxacin 400 mg (IV or PO) q24h <sup>*†§</sup>
	<i>E. faecalis</i> (VSE) Aerobic GNBs	Ampicillin 2 gm (IV) q4h <sup>*†</sup> <b>plus</b> Gentamicin 240 mg or 3 mg/kg (IV) q24h <sup>*†</sup>	Vancomycin 1 gm or 15 mg/kg (IV) q24h <sup>†</sup> <b>plus</b> Gentamicin 120 mg or 3 mg/kg (IV) q24h <sup>*†</sup>	
Aortic graft infection	<i>S. aureus</i> (MSSA) Aerobic GNBs <i>P. aeruginosa</i>	Meropenem 1 gm (IV) q8h <sup>†**</sup> <b>or</b> Cefepime 2 gm (IV) q12h <sup>†**</sup>	Levofloxacin 750 mg (IV) q24h <sup>†</sup>	Levofloxacin 750 mg (PO) q24h <sup>†</sup> <b>or</b> Ciprofloxacin 750 mg (PO) q12h <sup>†</sup>

MRSA/MSSA = methicillin-resistant/sensitive *S. aureus*. Duration of therapy represents total time IV or IV

\* Follow with maintenance dosing for renal failure (CrCl < 10 mL/min) and type of dialysis.

\*\* Depending on susceptibilities.

† If ABE not present, treat for 2 weeks after graft is removed/replaced. If graft not removable/replaceable, treat for 6 weeks.

§ IV therapy preferred. PO therapy if no IV access.

### AV Graft Infection

**Clinical Presentation:** Persistent fever/bacteremia without endocarditis in a patient with an AV graft on hemodialysis.

**Diagnostic Considerations:** Diagnosis by persistently positive blood cultures and gallium scan pickup over infected AV graft. Gallium scan will detect deep AV graft infection not apparent on exam.

**Pitfalls:** Antibiotics will lower temperature, but patient will usually not become afebrile without AV graft replacement.

**Therapeutic Considerations:** Graft usually must be removed for cure. MRSA is a rare cause of AV graft infection; if present, treat with linezolid 600 mg (IV or PO) q12h until graft is removed/replaced.

**Prognosis:** Good if new graft does not become infected at same site.

### Aortic Graft Infection

**Clinical Presentation:** Persistently positive blood cultures without endocarditis in a patient with an aortic graft.

**Diagnostic Considerations:** Diagnosis by positive blood cultures plus gallium scan pickup over infected aortic graft or abdominal CT/MRI scan.

**Pitfalls:** Infection typically occurs at anastomotic sites.

**Therapeutic Considerations:** Graft must be removed for cure. Operate as soon as diagnosis is confirmed (no value in waiting for surgery). MRSA is a rare cause of AV graft infection; if present, treat with linezolid 600 mg (IV or PO) q12h until graft is replaced.

**Prognosis:** Good if infected graft is removed before septic complications develop.

## Empiric Therapy of GI Tract Infections

### Esophagitis

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	PO Therapy or IV-to-PO Switch
Fungal	Candida albicans	Fluconazole 200 mg (IV or PO) × 1 dose, then 100 mg (IV or PO) q24h × 2–3 weeks	Micafungin 150 mg (IV) q24h × 2–3 weeks <b>or</b> Caspofungin 50 mg (IV) q24h × 2–3 weeks <b>or</b> Anidulafungin 100 mg (IV or PO) × 1 dose then 50 mg (IV) q24h × 2–3 weeks <b>or</b> Amphotericin B deoxycholate 0.5 mg/kg (IV) q24h × 2–3 weeks <b>or</b> Itraconazole 200 mg (IV) q12h × 2 days, then 200 mg (IV) q24h × 2–3 weeks	Fluconazole 200 mg (PO) × 1 dose, then 100 mg (PO) q24h × 2–3 weeks* <b>or</b> Posaconazole 100 mg (PO) q12h × 1 day, then 100 mg (PO) q24h × 2 weeks <b>or</b> Itraconazole 200 mg (PO) solution q24h × 2–3 weeks
Viral	HSV-1	Acyclovir 5 mg/kg (IV) q8h × 3 weeks	Valacyclovir 500 mg (PO) q12h × 3 weeks <b>or</b> Famciclovir 500 mg (PO) q12h × 3 weeks	
	CMV	Ganciclovir 5 mg/kg (IV) q12h × 3 weeks	Valganciclovir 900 mg (PO) q12h × 3 weeks	

Duration of therapy represents total time IV, PO, or IV + PO. Most patients on IV therapy able to take PO meds should be switched to PO therapy soon after clinical improvement (usually < 72 hours).

\* Loading dose is not needed PO if given IV with the same drug.

**Clinical Presentation:** Pain on swallowing.

**Diagnosis:** Stain/culture for fungi/HSV/CMV on biopsy specimen.

### Fungal (Candida) Esophagitis

**Diagnostic Considerations:** Rarely if ever in normal hosts. Often (but not always) associated with Candida in mouth. If patient is not alcoholic or diabetic and is not receiving antibiotics, test for HIV.

**Pitfalls:** Therapy as a diagnostic trial is appropriate. Suspect CMV-related disease and proceed to endoscopy if a patient with a typical symptom complex fails to respond to antifungal therapy.

**Therapeutic Considerations:** In normal hosts, treat for 1 week after clinical resolution. HIV patients respond more slowly than normal hosts and may need higher doses/treatment for 2–3 weeks after clinical resolution (see p. 336).

**Prognosis:** Related to degree of immunosuppression.

### Viral Esophagitis

**Diagnostic Considerations:** Rarely in normal hosts. May occur in the immunosuppressed.

**Pitfalls:** Viral and non-viral esophageal ulcers look similar; need biopsy for specific viral diagnosis.

**Therapeutic Considerations:** In normal hosts, treat for 2–3 weeks after clinical resolution. HIV patients respond more slowly and may need treatment for weeks after clinical resolution.

**Prognosis:** Related to degree of immunosuppression.

## Peptic Ulcer Disease (*H. pylori*)

Triple Therapy	Quadruple Therapy	Sequential Therapy
PPI + amoxicillin 1 gm (PO) q12h <b>plus either</b> clarithromycin 500 mg (PO) q12h <b>or</b> tinidazole (or metronidazole) 500 mg (PO) q12h all × 2 weeks	PPI + metronidazole 500 mg (PO) q12h + doxycycline 100 mg (PO) q12h + bismuth subsalicylate 525 mg tabs (PO) q6h all × 2 weeks	<b>5 days:</b> PPI + amoxicillin 1 gm (PO) q12h; <b>next 5 days:</b> PPI <b>plus</b> either clarithromycin 500 mg (PO) q12h <b>or</b> levofloxacin 500 mg (PO) q24h

**Diagnostic Considerations:** **Invasive:** rapid urease test, histology, culture. **Noninvasive:** serum ELISA test, urea breath test, stool (monoclonal antibody) antigen test.

**Pitfalls:** *False negative H. pylori tests with antibiotics, bismuth, PPIs.*

**Therapeutic Considerations:** See above grid. **Therapeutic failure:** *substitute* bismuth (for amoxicillin) *or substitute* nitazoxanide 1 gm (PO) q12h (for clarithromycin, metronidazole or tinidazole) all × 2 weeks. **1 week of therapy often fails.**

**Tests of Cure:** Urea breath test → 4–6 weeks post-therapy. Stool antigen test → 6–8 weeks post-therapy. *Stop PPIs 2 weeks before re-testing.*

## Gastric Perforation

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	IV-to-PO Switch
Gastric perforation	Oral anaerobes	Cefazolin 1 gm (IV) q8h × 1–3 days	Any beta-lactam (IV) × 1–3 days	Amoxicillin 1 gm (PO) q8h × 1–3 days <b>or</b> Cephalexin 500 mg (PO) q6h × 1–3 days <b>or</b> Quinolone* (PO) q24h × 1–3 days

\* Levofloxacin 500 mg or Moxifloxacin 400 mg.

**Clinical Presentation:** Presents acutely with fever and peritonitis.

**Diagnostic Considerations:** Obtain CT/MRI of abdomen for perforation/fluid collection.

**Pitfalls:** No need to cover *B. fragilis* with perforation of stomach/small intestine.

**Therapeutic Considerations:** Obtain surgical consult for possible repair.

**Prognosis:** Good if repaired.

## Infectious Diarrhea/Typhoid (Enteric) Fever

Subset	Usual Pathogens	Preferred Therapy	Alternate Therapy
Acute watery diarrhea	E. coli (ETEC, EHEC) Campylobacter Yersinia Salmonella Vibrio sp.	Quinolone <sup>†</sup> (IV or PO) × 5 days	Doxycycline 100 mg (IV or PO) q12h × 5 days <b>or</b> TMP-SMX 1 DS tablet (PO) q12h × 5 days
C. difficile diarrhea/colitis	Clostridium difficile	<b>Diarrhea</b> <sup>†§</sup> <b>Initial episode:</b> Vancomycin 250 mg (PO) q6h × 7–10 days* (If no improvement in 3 days, ↑ dose to 500 mg (PO) q6h × 7 days)** <b>Relapse:</b> Vancomycin 500 mg (PO) q6h × 14 days** <b>Recurrence:</b> <sup>††</sup> Vancomycin 500 mg (PO) q6h × 1 month ( <b>do not taper vancomycin dose</b> ). If another recurrence, <i>re-treat</i> with Vancomycin 500 mg (PO) q6h × 2 months. If another recurrence, <i>re-treat</i> with Vancomycin 500 mg (PO) q6h × 3 months. <i>If dose not tapered and if no colitis, this regimen will not fail!</i> If diarrhea continues, rule out C. difficile colitis or look for alternate diagnosis.	<b>Diarrhea</b> <b>Initial episode:</b> Nitazoxanide 500 mg (PO) q12h × 7–10 days <b>or</b> Metronidazole 250 mg (PO) q6h × 7–10 days* <b>Relapse:</b> Nitazoxanide 500 mg (PO) q12h × 7–10 days <b>or</b> Rifaximin 400 mg (PO) q8h × 10 days <b>or</b> Fidoxamicin 200 mg (PO) q12h × 10 days

\* **Treatment failure common with Vancomycin 125 mg (PO) q6h and with Metronidazole at any dose, (Flagyl frequently fails!).**

\*\* *If no improvement after 3 days with Vancomycin 500 mg (PO) q6h, rule out colitis with abdominal CT scan.* If abdominal CT scan shows **colitis** treat as C. difficile **colitis**.

† When treating C. difficile diarrhea or colitis, **discontinue antibiotics with a high C. difficile potential**, e.g., **cilindamycin, ciprofloxacin, α-lactams** (excluding ceftriaxone).

Patients being treated with levofloxacin or moxifloxacin *should not concurrently be taking PPIs*. Either discontinue PPI or switch to a H<sub>2</sub> blocker during quinolone therapy.

†† **Avoid C. difficile “prophylaxis.” Do not treat “history of C. difficile.”** Instead, *repeat C. difficile stool PCR to verify diagnosis.* Treat C. difficile diarrhea **only if PCR + for C. difficile.**

§ **Avoid anti-spasmodics in C. difficile diarrhea;** use may result in C. difficile **colitis**.

§§ Colectomy may be lifesaving in severe C. difficile pancolitis.

**Infectious Diarrhea/Typhoid (Enteric) Fever (cont'd)**

Subset	Usual Pathogens	Preferred Therapy	Alternate Therapy
		<p><b>Colitis</b>  <b>Initial episode:</b>            Metronidazole 1 gm (IV) q24h until cured  <b>plus</b>            Ertapenem 1 gm (IV) 124h (until associated peritonitis component resolves)</p> <p><b>Severe Pancolitis:<sup>55</sup></b>            Metronidazole 500 mg (IV/PO) q6h–8h until cured  <b>plus</b>            Nitazoxanide 500 mg (PO) q12h x until cured  <b>plus</b>            Tigecycline 200 mg (IV) x 1 dose, then, 100 mg (IV) q24h until cured</p> <p><b>Relapse or Recurrence:</b>            Re-treat with Metronidazole 500 mg (IV/PO) q6–8h until cured</p>	<p><b>Colitis</b>  <b>Initial Episode:</b>            Metronidazole 250 mg (PO) q6h until cured  <b>plus</b>            Tigecycline 200 mg (IV) x 1 dose, then 100 mg (IV) q24h until cured  <b>or</b>            Nitazoxanide 500 mg (PO) q12h until cured</p> <p><b>Relapse or Recurrence:</b>            Nitazoxanide 500 mg (PO) q12h until cured</p>
Cytomegalovirus colitis	CMV	Valganciclovir 900 mg (PO) q12h x 21 days; for HIV, see p. 334	

<sup>55</sup> Colectomy may be lifesaving in severe *C. difficile* pancolitis.

## Infectious Diarrhea/Typhoid (Enteric) Fever (cont'd)

Subset	Usual Pathogens	Preferred Therapy	Alternate Therapy
Typhoid (enteric) fever	Salmonella typhi/paratyphi	Quinolone <sup>†</sup> (IV or PO) × 10–14 days <b>or</b> TMP–SMX 5 mg/kg (IV or PO) q6h × 10–14 days	Chloramphenicol 500 mg (IV or PO) q6h × 10–14 days <b>or</b> any 3 <sup>rd</sup> gen cephalosporin (IV or PO) × 10–14 days <b>or</b> Azithromycin 1 gm (PO) q24h × 5 days
Chronic watery diarrhea	Giardia lamblia*	Tinidazole 2 gm (PO) × 1 dose <b>or</b> Nitazoxanide 500 mg (PO) q12h × 3 days <b>or</b> Metronidazole 2 gm (PO) q24h × 3 days	Albendazole 400 mg (PO) q24h × 5 days <b>or</b> Quinacrine 100 mg (PO) q8h × 5 days
	Cryptosporidia*	Nitazoxanide 500 mg (PO) q12h × 5 days <sup>§</sup>	Paromomycin 500–750 mg (PO) q8h until response <b>or</b> Azithromycin 600 mg (PO) q24h × 4 weeks
	Isospora* Cyclospora*	TMP–SMX 1 DS tablet (PO) q12h × 10 days <sup>‡</sup>	
Acute dysentery	Entamoeba histolytica	<u>Preferred therapy</u> : Metronidazole 750 mg (PO) q8h × 10 days <b>followed by either</b> Iodoquinol 650 mg (PO) q8h × 20 days <b>or</b> Paromomycin 500 mg (PO) q8h × 7 days <u>Alternate therapy</u> : Tinidazole 1 gm (PO) q12h × 3 days	
	Shigella	Quinolone <sup>†</sup> (IV or PO) × 3 days	TMP–SMX 1 DS tablet (PO) q12h × 3 days <b>or</b> Azithromycin 500 mg (IV or PO) q24h × 3 days

Duration of therapy represents total time IV, PO, or IV + PO. Most patients on IV therapy able to take PO meds should be switched to PO therapy soon after clinical improvement (usually < 72 hours).

\* May also present as acute watery diarrhea.

† Ciprofloxacin 400 mg (IV) or 500 mg (PO) q12h or Levofloxacin 500 mg (IV or PO) q24h or Moxifloxacin 400 mg (IV or PO) q24h.

§ Longer duration of therapy may be needed in immunosuppressed patients (treat until cured).

### **Acute Watery Diarrhea**

**Clinical Presentation:** Acute onset of watery diarrhea without blood/mucus.

**Diagnostic Considerations:** Diagnosis by culture of organism from stool specimens.

**Pitfalls:** Recommended antibiotics are active against most susceptible enterotoxigenic bacterial pathogens causing diarrhea, but not viruses/parasites. Concomitant transient lactase deficiency may prolong diarrhea if dairy products are taken during an infectious diarrhea.

**Therapeutic Considerations:** Avoid norfloxacin and ciprofloxacin due to resistance potential. *V. cholerae* may be treated with a single dose of any oral respiratory quinolone or doxycycline.

**Prognosis:** Excellent. Most recover with supportive treatment.

### **Clostridium difficile Diarrhea/Colitis**

**Clinical Presentation:** Voluminous watery diarrhea following exposure to *C. difficile* contaminated fomites, exposure to patients with *C. difficile*, recent cancer chemotherapy or antibiotic therapy with some, *but not most*, antibiotics. Most often associated with clindamycin, ciprofloxacin, and  $\beta$ -lactams (excluding ceftriaxone).

**Diagnostic Considerations:** Watery diarrhea with positive *C. difficile* stool toxin. A single positive PCR *C. difficile* stool toxin test is sufficiently sensitive/specific for diagnosis (endpoint is end of diarrhea, not stool toxin negativity); *If negative, no need to retest*. If *C. difficile* colitis suspected in *C. difficile* positive patients with fever/prominent leukocytosis/abdominal pain, confirm diagnosis by abdominal CT scan.

**Pitfalls:** *C. difficile* colitis is suggested by the presence of otherwise unexplained leukocytosis,  $\uparrow$  ESR, abdominal pain and often temperature  $> 102^{\circ}\text{F}$ ; confirm diagnosis with CT/MRI of abdomen. Virulent strains of *C. difficile* may present with colitis with temperature  $\leq 102^{\circ}\text{F}$ , leukocytosis (often very high, i.e., 25–50  $\text{K}/\text{mm}^3$ ), and little/no abdominal pain; confirm diagnosis with CT/MRI of abdomen. Radiographically *C. difficile* colitis typically is a pancolitis. Segmental colitis suggests a non-*C. difficile* etiology, i.e., ischemic colitis. *C. difficile* toxin test may remain positive in stools following resolution of diarrhea; In patients receiving enteral feeds, diarrhea is likely due to enteral feeds (high infusion rates/high osmotic loads) rather than *C. difficile*. Norovirus diarrhea may mimic *C. difficile* diarrhea or concurrent outbreaks may occur. Vancomycin may be *ineffective* in *C. difficile colitis*.

**Therapeutic Considerations:** For ***C. difficile* diarrhea** oral vancomycin or nitazoxanide are preferred; *Flagyl frequently fails*. Rifaximin and Fidaxomylin have no advantage over vancomycin. With effective therapy, *C. difficile* diarrhea begins to improve ( $\leq 3$  days) and usually resolves by 5–7 days, although some patients require 10 days of therapy. If no improvement with vancomycin 250 mg (PO) q6h,  $\uparrow$  dose to 500 mg (PO) q6h or use nitazoxanide. *Do not treat C. difficile negative diarrhea with oral vancomycin or metronidazole*. For ***C. difficile* colitis**, treat until colitis resolves (follow with serial ESRs/abdominal and or CT scans). Add aerobic GNB coverage for microscopic/gross peritonitis with ESRs ertapenem. Tigecycline highly effective monotherapy for *C. difficile* colitis; and also provides anti-B fragilis coverage (for associated microscopic/clinical peritonitis). **Avoid anti-motility agents**, e.g., loperamide with *C. difficile* diarrhea which may result in *C. difficile colitis/toxic megacolon*.

**Prognosis:** Prognosis with *C. difficile* diarrhea is excellent. *C. difficile* colitis prognosis is related to strain virulence/and extent of the colitis.

### **Typhoid (Enteric) Fever (Salmonella typhi/Paratyphi)**

**Clinical Presentation:** High fevers ( $> 102^{\circ}\text{F}$ ) increasing in a stepwise fashion accompanied by relative bradycardia (without chills) with watery diarrhea/constipation, headache, anorexia, abdominal pain, dry cough/sore throat, tender hepatomegaly,  $\pm$  Rose spots. Dull mental affect ("apathetic facies") peculiar to typhoid fever. Thrombocytopenia should suggest another diagnosis. e.g. malaria or viral coinfection, e.g. dengue.



**Diagnostic Considerations:** Most community-acquired watery diarrheas are not accompanied by temperatures  $> 102^{\circ}\text{F}$  and relative bradycardia. Diagnosis is confirmed by demonstrating *Salmonella* in blood, bone marrow, Rose spots, or stool cultures. Culture of bone marrow is the quickest/most reliable method of diagnosis. WBC count is usually low/low normal. Leukocytosis should suggest another diagnosis or bowel perforation, which may occur during 2<sup>nd</sup> week of typhoid fever. Eosinopenia characteristic of typhoid/enteric fever. In patients with suspected typhoid fever, eosinophilia should suggest an alternate diagnosis or parasitic coinfection.

**Pitfalls:** Rose spots are few/difficult to see and not present in all cases. Typhoid fever usually presents with constipation, not diarrhea. Severe rigors not a feature of typhoid fever.

**Therapeutic Considerations:** 2<sup>nd</sup> generation cephalosporins, aztreonam, and aminoglycosides are ineffective. Since *Salmonella* strains causing enteric fever are intracellular pathogens, treat for a full 2 weeks to maximize cure rates/minimize relapses. Treat relapses with the suggested antibiotics  $\times$  2–3 weeks. *Salmonella* excretion into feces usually persists  $< 3$  months. Persistent excretion  $> 3$  months suggests a carrier state—rule out hepatobiliary/urinary calculi.

**Prognosis:** Good if treated early. Poor with late treatment/bowel perforation.

### **Chronic Watery Diarrhea**

**Clinical Presentation:** Watery diarrhea without blood/mucus lasting  $> 1$  month.

**Diagnostic Considerations:** Diagnosis by demonstrating organisms/cysts in stool specimens. Multiple fresh daily stool samples often needed for diagnosis especially for protozoan parasites.

**Pitfalls:** Concomitant transient lactase deficiency may prolong diarrhea if dairy products are consumed during an infectious diarrhea.

**Therapeutic Considerations:** Cryptosporidia and *Isospora* are being recognized increasingly in acute/chronic diarrhea in normal hosts.

**Prognosis:** Excellent in well-nourished patients. Untreated patients may develop malabsorption.

### **Giardia lamblia**

**Clinical Presentation:** Acute/subacute onset of diarrhea, abdominal cramps, bloating, flatulence. Incubation period 1–2 weeks. Malabsorption may occur in chronic cases. No eosinophilia.

**Diagnostic Considerations:** Diagnosis by demonstrating trophozoites or cysts in stool/antigen detection assay. If stool exam and antigen test are negative and Giardiasis is suspected, perform “string test”/duodenal aspirate and biopsy.

**Pitfalls:** Cysts intermittently excreted into stool. Usually need multiple stool samples for diagnosis. Often accompanied by transient lactose intolerance.

**Therapeutic Considerations:** Nitazoxanide effective therapy. Diarrhea may be prolonged if milk (lactose-containing) products are ingested after treatment/cure. May need repeat courses of therapy.

**Prognosis:** Related to severity of malabsorption and health of host.

### **Cryptosporidia**

**Clinical Presentation:** Acute/subacute onset of diarrhea. Usually occurs in HIV patients with CD<sub>4</sub> counts  $< 200$ . Biliary cryptosporidiosis is seen only in HIV; may present as acalculous cholecystitis or sclerosing cholangitis with RUQ pain, fever,  $\uparrow$  alkaline phosphatase, but bilirubin is normal.

**Diagnostic Considerations:** Diagnosis by demonstrating organism in stool/intestinal biopsy specimen. Cholera-like illness in normal hosts. Chronic watery diarrhea in compromised hosts.

**Pitfalls:** Smaller than *Cyclospora*. Oocyst walls are smooth (not wrinkled) on acid fast staining.

**Therapeutic Considerations:** Nitazoxanide effective therapy.

**Prognosis:** Related to adequacy of fluid replacement/underlying health of host.

### **Cyclospora**

**Clinical Presentation:** Acute/subacute onset of diarrhea. Incubation period 1–14 days.

**Diagnostic Considerations:** Diagnosis by demonstrating organism in stool/intestinal biopsy specimen. Clinically indistinguishable from cryptosporidial diarrhea (intermittent watery diarrhea without blood or mucus). Fatigue/weight loss common.

**Pitfalls:** Oocysts only form seen in stool and are best identified with modified Kinyoun acid fast staining. Acid fast fat globules stain pink with acid fast staining. "Wrinkled wall" oocysts are characteristic of Cyclospora, not Cryptosporidia. Oocysts are twice the size of similar appearing Cryptosporidia (~ 10 µm vs. 5 µm).

**Therapeutic Considerations:** Nitazoxanide effective therapy.

**Prognosis:** Related to adequacy of fluid replacement/underlying health of host.

### **Acute Dysentery**

#### **Entamoeba histolytica**

**Clinical Presentation:** Acute/subacute onset of bloody diarrhea/mucus. Fecal WBC/RBCs due to mucosal invasion. *E. histolytica* may also cause chronic diarrhea. Colonic ulcers secondary to *E. histolytica* are round and may form "collar stud" abscesses.

**Diagnostic Considerations:** Diagnosis by demonstrating organism/trophozoites in stool/intestinal biopsy specimen. Serology is negative with amebic dysentery, but positive with extra-intestinal forms. Test to separate *E. histolytica* from non-pathogenic *E. dispar* cyst passers. On sigmoidoscopy, ulcers due to *E. histolytica* are round with normal mucosa in between, and may form "collar stud" abscesses. In contrast, ulcers due to *Shigella* are linear and serpiginous without normal intervening mucosa. Bloody dysentery is more subacute with *E. histolytica* compared to *Shigella*.

**Pitfalls:** Intestinal perforation/abscess may complicate amebic colitis. Rule out infectious causes of bloody diarrhea with mucus before diagnosing/treating inflammatory bowel disease (IBD). Obtain multiple stool cultures for bacterial pathogens/parasites. Do not confuse *E. histolytica* in stool specimens with *E. hartmanni*, a non-pathogen protozoa similar in appearance but smaller in size.

**Therapeutic Considerations:** *E. histolytica* cyst passers should be treated, but metronidazole is ineffective against cysts. Recommended antibiotics treat both luminal and hepatic *E. histolytica*. Use paromomycin 500 mg (PO) q8h × 7 days for asymptomatic cysts.

**Prognosis:** Good if treated early. Related to severity of dysentery/ulcers/extra-intestinal amebiasis.

#### **Shigella**

**Clinical Presentation:** Acute onset of bloody diarrhea/mucus.

**Diagnostic Considerations:** Diagnosis by demonstrating organism in stool specimens. *Shigella* ulcers in colon are linear, serpiginous, and rarely lead to perforation.

**Therapeutic Considerations:** *Shigella* dysentery is more acute/fulminating than amebic dysentery. *Shigella* has no carrier state, unlike *Entamoeba*.

**Prognosis:** Good if treated early. Severity of illness related to *Shigella* species: *S. dysenteriae* (most severe) > *S. flexneri* > *S. boydii*/*S. sonnei* (mildest).

## Cholecystitis

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	PO Therapy or IV-to-PO Switch
Normal host	E. coli Klebsiella E. faecalis (VSE)	Meropenem 1 gm (IV) q8h* <b>or</b> Piperacillin/tazobactam 3.375 gm (IV) q6h* <b>or</b> Tigecycline 200 mg (IV) × 1 dose, then 100 mg (IV) q24h*	Cefazolin 1 gm (IV) q8h* ± Ampicillin 1 gm (IV) q6h* <b>or</b> Quinolone <sup>†</sup> (IV)*	Quinolone <sup>†</sup> (PO)*
Emphysematous cholecystitis <sup>‡</sup>	Clostridium perfringens E. coli	Meropenem 1 gm (IV) q8h <sup>‡</sup> <b>or</b> Piperacillin/tazobactam 3.375 gm (IV) q6h <sup>‡</sup>	Ertapenem 1 gm (IV) q24h <sup>‡</sup> <b>or</b> Ticarcillin/clavulanate 3.1 gm (IV) q6h <sup>‡</sup>	Clindamycin 300 mg (PO) q8h <sup>‡</sup>

Duration of therapy represents total time IV, PO, or IV + PO. Most patients on IV therapy able to take PO meds should be switched to PO therapy after clinical improvement.

† Treat only IV or IV-to-PO switch.

‡ Ciprofloxacin 400 mg (IV) or Levofloxacin 500 mg (IV or PO) q24h or Moxifloxacin 400 mg (IV or PO) q24h.

\* If no cholecystectomy, treat × 5–7 days. If cholecystectomy is performed, treat × 3–4 days postoperatively.

¶ Treat × 4–7 days after cholecystectomy.

### Cholecystitis

**Clinical Presentation:** RUQ pain, fever usually  $\leq 102^{\circ}\text{F}$ , positive Murphy's sign, no percussion tenderness over right lower ribs.

**Diagnostic Considerations:** Diagnosis by RUQ ultrasound/positive HIDA scan.

**Pitfalls:** No need to cover *B. fragilis*.

**Therapeutic Considerations:** Obtain surgical consult for possible cholecystectomy.

**Prognosis:** Related to cardiopulmonary status.

### Emphysematous Cholecystitis

**Clinical Presentation:** Clinically presents as cholecystitis. Usually in diabetics.

**Diagnostic Considerations:** RUQ/gallbladder gas on flat plate of abdomen.

**Pitfalls:** Requires immediate cholecystectomy.

**Therapeutic Considerations:** Usually a difficult/prolonged post-op course.

**Prognosis:** Related to speed of gallbladder removal.

## Cholangitis

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	IV-to-PO Switch
Normal host	E. coli Klebsiella E. faecalis (VSE)	Meropenem 1 gm (IV) q8h*  <b>or</b> Tigecycline 200 mg (IV) x 1 dose, then 100 mg (IV) q24h*  <b>or</b> Piperacillin/tazobactam 3.375 gm (IV) q6h*	Ampicillin/sulbactam 3 gm (IV) q6h*  <b>or</b> Cefoperazone 2 gm (IV) q12h*  <b>or</b> Doripenem 1 gm (IV) q8h	Ciprofloxacin 500 mg (PO) q12h*  <b>or</b> Levofloxacin 500 mg (PO) q24h*  <b>or</b> Moxifloxacin 400 mg (PO) q24h*

Duration of therapy represents total time IV or IV + PO. Most patients on IV therapy able to take PO meds should be switched to PO therapy after clinical improvement.

\* Treat until resolved (usually 5–7 days).

**Clinical Presentation:** RUQ pain, fever > 102°F, positive Murphy's sign, percussion tenderness over right lower ribs.

**Diagnostic Considerations:** Obstructed common bile duct on ultrasound/CT/MRI of abdomen.

**Pitfalls:** Charcot's triad (fever, RUQ pain, jaundice) is present in only 50%.

**Therapeutic Considerations:** Obtain surgical consult to relieve obstruction. Continue antibiotics for 4–7 days after obstruction is relieved.

**Prognosis:** Related to speed of surgical relief of obstruction.

## Gallbladder Wall Abscess/Perforation

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	IV-to-PO Switch
Gallbladder wall abscess/perforation	E. coli Klebsiella E. faecalis (VSE)	Piperacillin/ tazobactam 3.375 gm (IV) q6h*  <b>or</b> Tigecycline 200 mg (IV) x 1 dose, then 100 mg (IV) q24h*  <b>or</b> Meropenem 1 gm (IV) q8h*	Ampicillin/sulbactam 3 gm (IV) q6h*  <b>or</b> Cefoperazone 2 gm (IV) q12h*  <b>or</b> Doripenem 1 gm (IV) q8h	Ciprofloxacin 500 mg (PO) q12h*  <b>or</b> Levofloxacin 500 mg (PO) q24h*  <b>or</b> Moxifloxacin 400 mg (PO) q24h*

Duration of therapy represents total time IV or IV + PO.

\* Treat until resolved (usually 1–2 weeks).

**Clinical Presentation:** RUQ pain, fever ≤ 102°F, positive Murphy's sign, no percussion tenderness over right lower ribs.

**Diagnostic Considerations:** Diagnosis by CT/MRI of abdomen. Bile peritonitis is common.

**Pitfalls:** Bacterial peritonitis may be present.

**Therapeutic Considerations:** Obtain surgical consult for possible gallbladder removal. Usually a difficult and prolonged post-op course.

**Prognosis:** Related to removal of gallbladder/repair of perforation.

## Acute Pancreatitis

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	IV-to-PO Switch
Edematous pancreatitis	None	Not applicable	Not applicable	Not applicable
Hemorrhagic/necrotizing pancreatitis	Aerobic GNBs B. fragilis	Meropenem 1 gm (IV) q8h* <b>or</b> Ertapenem 1 gm (IV) q24h*	Piperacillin/tazobactam 3.375 gm (IV) q6h* <b>or</b> Ampicillin/sulbactam 1.5 gm (IV) q6h* <b>or</b> Ticarcillin/clavulanate 3.1 gm (IV) q6h*	Clindamycin 300 mg (PO) q8h* <b>plus</b> Levofloxacin 500 mg (PO) q24h* <b>or monotherapy with</b> Moxifloxacin 400 mg (PO) q24h*

Duration of therapy represents total time IV or IV + PO and varies depending on the clinical response. Most patients on IV therapy able to take PO meds should be switched to PO therapy after clinical improvement.

\* Treat until resolved (usually 1–2 weeks).

### Edematous Pancreatitis

**Clinical Presentation:** Sharp abdominal pain with fever  $\leq 102^{\circ}\text{F}$   $\pm$  hypotension.

**Diagnostic Considerations:** Diagnosis by elevated serum amylase and lipase levels with normal meth-hemalbumin levels. May be drug-induced (e.g., steroids).

**Pitfalls:** Amylase elevation alone is not diagnostic of acute pancreatitis.

**Therapeutic Considerations:** NG tube is not needed. Aggressively replace fluids.

**Prognosis:** Good with adequate fluid replacement.

### Hemorrhagic/Necrotizing Pancreatitis

**Clinical Presentation:** Sharp abdominal pain with fever  $\leq 102^{\circ}\text{F}$   $\pm$  hypotension. Grey-Turner/Cullen's sign present in some.

**Diagnostic Considerations:** Mildly elevated serum amylase and lipase levels with high meth-hemalbumin levels.

**Pitfalls:** With elevated lipase, amylase level is inversely related to severity of disease.

**Therapeutic Considerations:** Obtain surgical consult for possible peritoneal lavage as adjunct to antibiotics. Serum albumin/dextran are preferred volume expanders.

**Prognosis:** Poor with hypocalcemia or shock.

## Pancreatic Abscess/Infected Pancreatic Pseudocyst

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	IV-to-PO Switch
Infected pancreatic pseudocyst/ pancreatic abscess	Aerobic GNBs B. fragilis	Meropenem 1 gm (IV) q8h* <b>or</b> Piperacillin/tazobactam 3.375 gm (IV) q6h* <b>or</b> Ertapenem 1 gm (IV) q24h*	Ampicillin/sulbactam 1.5 gm (IV) q6h* <b>or</b> Ticarcillin/clavulanate 3.1 gm (IV) q6h* <b>or</b> Doripenem 1 gm (IV) q8h	Moxifloxacin 400 mg (PO) q24h* <b>or combination therapy with</b> Clindamycin 300 mg (PO) q8h* <b>plus</b> Quinolone <sup>†</sup> (PO)*

Duration of therapy represents total time IV or IV + PO.

† Ciprofloxacin 500 mg q12h or Levofloxacin 500 mg q24h.

\* Treat until resolved.

**Clinical Presentation:** Follows acute pancreatitis or develops in a pancreatic pseudocyst. An infected pancreatic pseudocyst is an abscess equivalent. Fevers usually  $\geq 102^{\circ}\text{F}$ .

**Diagnostic Considerations:** CT/MRI of abdomen demonstrates pancreatic abscess.

**Pitfalls:** Peritoneal signs are typically absent.

**Prognosis:** Related to size/extent of abscess and adequacy of drainage.

## Liver Abscess

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	PO Therapy or IV-to-PO Switch
Liver abscess	Aerobic GNBs Enterococci (VSE) B. fragilis	Piperacillin/tazobactam 3.375 gm (IV) q6h* <b>or</b> Tigecycline 100 mg (IV) $\times$ 1 dose, then 50 mg (IV) q12h* <b>or</b> Meropenem 1 gm (IV) q8h* <b>or</b>	Quinolone <sup>†</sup> (IV)* <b>plus either</b> Metronidazole 1 gm (IV) q24h* <b>or</b> Clindamycin 600 mg (IV) q8h*  Moxifloxacin 400 mg (IV) q24h* <b>or</b> Sulbactam/ampicillin 3 gm (IV) q6h	Amoxicillin/clavulanic acid 875/125 mg (PO) q12h* <b>or</b> Moxifloxacin 400 mg (PO) q24h* <b>or combination therapy with</b>  Quinolone <sup>†</sup> (PO)* <b>plus either</b> Metronidazole 500 mg (PO) q12h* <b>or</b> Clindamycin 300 mg (PO) q8h*
	E. histolytica	See p. 274		

Duration of therapy represents total time IV, PO, or IV + PO.

\* Treat until abscess(es) are no longer present or stop decreasing in size on serial CT scans.

† Ciprofloxacin 400 mg (IV) or 500 mg (PO) q12h or Levofloxacin 500 mg (IV or PO) q24h.

## Liver Abscess (cont'd)

**Clinical Presentation:** Fever, RUQ tenderness, negative Murphy's sign, and negative right lower rib percussion tenderness.

**Diagnostic Considerations:** Diagnosis by CT/MRI scan of liver and aspiration of abscess. CT shows multiple lesions in liver. Source is usually either the colon (diverticulitis or diverticular abscess with portal pyemia) or retrograde infection from the gallbladder (cholecystitis or gallbladder wall abscess).

**Pitfalls:** Bacterial abscesses are usually multiple and involve multiple lobes of liver; amebic abscesses are usually solitary and involve the right lobe of liver.

**Therapeutic Considerations:** Liver laceration/trauma usually requires ~ 2 weeks of antibiotics.

**Prognosis:** Good if treated early.

## Hepatosplenic Candidiasis

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	IV-to-PO Switch
Hepato-splenic candidiasis	Candida albicans	Fluconazole 800 mg (IV) × 1 dose, then 400 mg (IV) q24h × 2–4 weeks <b>or</b> Micafungin 100 mg (IV) q24h × 2–4 weeks <b>or</b> Caspofungin 70 mg (IV) × 1 dose, then 50 mg (IV) q24h × 2–4 weeks	Ambisome (L-Amb) (see p. 540) (IV) q24h × 2–4 weeks <b>or</b> Amphotericin B 0.7 mg/kg (IV) q24h × 2–4 weeks	Fluconazole 800 mg (PO) × 1 dose, then 400 mg (PO) q24h × 2–4 weeks* <b>or</b> Itraconazole 200 mg (PO) solution q12h × 2–4 weeks

Duration of therapy represents total time IV or IV + PO. Most patients on IV therapy able to take PO meds should be switched to PO therapy after clinical improvement.

\* Loading dose is not needed PO if given IV with the same drug.

**Clinical Presentation:** New high spiking fevers with RUQ/LUQ pain after 2 weeks in a patient with febrile leukopenia.

**Diagnostic Considerations:** Diagnosis by abdominal CT/MRI showing mass lesions in liver/spleen.

**Pitfalls:** Do not overlook RUQ tenderness and elevated alkaline phosphatase in leukopenic cancer patients as a clue to the diagnosis.

**Therapeutic Considerations:** Treat until liver/spleen lesions resolve. Should be viewed as a form of disseminated disease.

**Prognosis:** Related to degree/duration of leukopenia.

## Granulomatous Hepatitis (BCG)

Subset	Pathogen	Preferred Therapy
BCG hepatitis	Bacille Calmette-Guérin (BCG)	INH 300 mg (PO) q24h × 6 months + rifampin 600 mg (PO) q24h × 6 months

**Clinical Presentation:** Fever, chills, anorexia, weight loss, hepatomegaly  $\pm$  RUQ pain days to weeks after intravesicular BCG for bladder cancer.

**Diagnostic Considerations:**  $\uparrow$  alkaline phosphatase  $>$   $\uparrow$  SGOT/SGPT. Liver biopsy is negative for AFB/positive for granulomas.

**Pitfalls:** Exclude other causes of hepatomegaly.

**Therapeutic Considerations:** INH plus rifampin  $\times$  6 months is curative.

**Prognosis:** Excellent with early treatment.

## Leptospirosis

**Clinical Presentation:** After incubation period of 1–2 weeks, abrupt onset of fever, no chills, severe headache, conjunctival suffusion, meningismus, dry cough, intense myalgias (gastroc myalgias/tenderness)  $\pm$  abdominal pain. 1<sup>st</sup> (leptospiremic) phase (4–9 days); 2<sup>nd</sup> (immune) phase (6–12 days). Leukocytosis with no thrombocytopenia usual. Hepatic ( $\uparrow$  bilirubin/transaminases) and renal (azotemia, proteinuria, sterile, pyuria) involvement. Meningitis (early)  $\rightarrow$  ABM with CSF + for leptospires; meningitis (late)  $\rightarrow$  aseptic meningitis CSF—for leptospires. Progressive vomiting, epistaxis, jaundice/hepatosplenomegaly only with Weil's syndrome (icteric leptospirosis). Intense myalgias with hepatic and renal involvement should suggest leptospirosis.

**Diagnostic Considerations:** Recent occupational or contaminated water (rat urine) exposure. Leptospiuria (after 2 weeks) seen in dark-field of urine. BCs + early (– late); UCs + late. Leptospires die rapidly in acid urine (add buffered saline before culture). Diagnosis by blood/urine culture or  $\uparrow$  MAT IgM titers (after 1 week). Antibiotic therapy may blunt, delay or abort titer rise.

**Pitfalls:** Like influenza (headache, sore throat, myalgias), patients can recall exact hour of onset. Unlike influenza, no chills but conjunctival suffusion. CXR—in influenza (early) but CXR often + in leptospirosis. Differentiate from viral hepatitis ( $<$  102°F, no conjunctival suffusion, leukopenia, highly  $\uparrow$  transaminases, no renal involvement). Differentiate from EBV, CMV, HSV hepatitis by + atypical lymphocytes (– with leptospirosis) and specific serologies. Like aseptic (viral) meningitis (normal glucose, WBCs  $<$  500/hpf). Only ID with CSF bilirubin  $>$  serum. Clinically irrelevant serologic XR with Borrelia/treponeme titer.

**Therapeutic Considerations:** Penicillin or doxycycline ( $\pm$  Jarisch-Herxheimer reaction).

**Prognosis:** Excellent if treated early, 5–40% mortality for Weil's syndrome and elderly.

## Viral Hepatitis

Subset	Pathogens	Therapy
Acute	HAV (none); HBV (consider nucleos(t)ides for HBV-acute liver failure); HCV (consider pegylated interferon +/- ribavirin if no viral clearance after 12–16 weeks); HDV (none); HEV (consider ribavirin fir severe hepatitis)	
Chronic	HBV*	<p><u>Preferred:</u></p> <p>Tenofovir 300 mg (PO) <b>or</b> Entecavir 0.5 mg (PO) q24h <math>\times</math> <math>\geq</math> 12 months</p> <p><u>For lamivudine resistant strains use either</u></p> <p>Entecavir 1.0 mg q24h <b>or</b> Pegylated interferon alfa-2a 180 mcg (SQ)/week <i>without</i> ribavirin <math>\times</math> 12 months</p> <p><u>Alternate:</u> Telbivudine 600 mg (PO) q24h <b>or</b> Adefovir 10 mg (PO) q24h <b>or</b> Lamivudine 100 mg (PO) for <math>\times</math> <math>\geq</math> 12 months</p>

\* HBeAg+ with ALT  $>$  2  $\times$  n and HBV DNA  $>$  20,000 IU/ml or alternately, chronic HBV with moderate/severe inflammation or significant fibrosis on liver biopsy. Primary therapeutic response ( $<$  2 log  $\downarrow$ ) in a HBV DNA levels after  $\geq$  6 months of therapy. Treatment failures should be treated with an alternate regimen or additional treatment.



## Viral Hepatitis (cont'd)

Subset	Pathogens	Therapy
	HCV	<p><b>Genotype 1 (includes 1a &amp; 1b)</b> Preferred therapy (similar efficacy): Ledipasvir 90 mg/sofosbuvir 400 mg (Harvoni) (PO) q24h x 12 weeks Paritaprevir 150 mg/ritonavir 100 mg/ombitasvir 25 mg (PO) q24h + dasabuvir 250 mg (Viekira Pak) (PO) q12h + ribavirin<sup>1,2</sup> x weeks<sup>3</sup> Sofosbuvir 400 mg (Sovaldi) (PO) q24h + simeprevir<sup>4</sup> 150 mg (Olysio) (PO) q24h ± ribavirin<sup>1</sup> x weeks<sup>3</sup></p> <p><b>Genotype 2</b> Preferred therapy: Sofosbuvir 400 mg (Sovaldi) (PO) q24h + ribavirin<sup>1</sup> x 12 weeks (16 weeks in cirrhosis)</p> <p><b>Genotype 3</b> Preferred therapies: Sofosbuvir 400 mg (Sovaldi) (PO) q24h + ribavirin<sup>1</sup> x 24 weeks Alternate therapy: Sofosbuvir 400 mg (Sovaldi) (PO) q24h + pegylated IFN alfa-2a/2b<sup>5</sup> ribavirin<sup>1</sup> x 12 weeks</p> <p><b>Genotype 4</b> Preferred therapies (similar efficacy): Ledipasvir 90 mg/sofosbuvir 400 mg (Harvoni) (PO) q24h x 12 weeks Paritaprevir 150 mg/ritonavir 100 mg/ombitasvir 25 mg (Viekira Pak minus dasabuvir) (PO) q24h + ribavirin<sup>1</sup> x 12 weeks Sofosbuvir 400 mg (Sovaldi) (PO) q24h + ribavirin<sup>1</sup> x 24 weeks Alternate therapy: Sofosbuvir 400 mg (Sovaldi) (PO) q24h + pegylated IFN alfa-2a/2b<sup>5</sup> + ribavirin<sup>1</sup> x 12 weeks, Sofosbuvir 400 mg (Sovaldi) (PO) q24h + simeprevir 150 mg (Olysio) (PO) q24h ± ribavirin<sup>1</sup> x 12 weeks</p> <p><b>Genotype 5</b> Preferred therapy: Sofosbuvir 400 mg (Sovaldi) (PO) q24h + pegylated IFN alfa-2a/2b<sup>5</sup> + ribavirin<sup>1</sup> x 12 weeks Alternate therapy: Pegylated IFN alfa-2a/2b<sup>5</sup> + ribavirin<sup>1</sup> x 48 weeks</p> <p><b>Genotype 6</b> Preferred therapy: Ledipasvir 90 mg/sofosbuvir 400 mg (Harvoni) (PO) q24h x 12 weeks Alternate therapy: Sofosbuvir 400 mg (Sovaldi) (PO) q24h + pegylated IFN alfa-2a/2b<sup>5</sup> + ribavirin<sup>1</sup> x 12 weeks</p>
	HDV	Prolonged therapy with pegylated interferon alfa-2a <i>without</i> ribavirin as for chronic hepatitis B (see above)
	HEV	Ribavirin 600–1200 mg daily x 12 weeks or pegylated interferon weekly x 12 weeks +/- ribavirin

<sup>1</sup>Ribavirin should be dosed as following:

Weight < 75 kg: Ribavirin 1000 mg (PO) daily in 2 divided doses (400 mg in am, 600 mg in pm)

Weight > 75 kg: Ribavirin 1200 mg (PO) daily in 2 divided doses (600 mg q12h)

<sup>2</sup>In Genotype 1b, ribavirin is only Indicated in cirrhosis

<sup>3</sup>12 week therapy without cirrhosis; 24 week therapy with cirrhosis

<sup>4</sup>Simeprevir may be ineffective in patients with HCV genotype 1a Q80K polymorphism

<sup>5</sup>Pegylated IFN alfa-2a should be dosed 180 µg (SC) weekly. Pegylated IFN alfa-2b should be dosed 1.5 µg/kg (SC) weekly. Both types of pegylated IFN alfa are approved to treat genotype 1

Adapted from: American Association for the Study of Liver Disease and Infectious Disease Society of America. Recommendations for testing, managing, and treating hepatitis C. <http://www.hcvguidelines.org>, 2014

### **Acute Viral Hepatitis**

**Clinical Presentation:** Asymptomatic or anorexia, malaise, nausea, serum-sickness like syndrome, ± jaundice, dark urine.

**Diagnostic Considerations:** Diagnosis by presence in HAV of IgM anti-HAV or in HBV of HBsAg and IgM anti-HBc or in HCV of anti-HCV and HCV RNA or in HDV of HBsAg, anti-HDV and HDV RNA or in HEV of anti-HEV and HEV RNA, with markedly elevated serum aminotransferases (ALT > 400). Serum alkaline phosphatase normal/mildly elevated; albumin normal or mildly depressed; INR normal or minimally elevated.

**Pitfalls:** Rule out other hepatitis-causing agents (EBV, CMV, HSV, etc) and drug-induced liver injury if serologic tests for HAV, HBV, HCV, HDV, and HEV are negative.

**Therapeutic Considerations:** For patients who develop encephalopathy and/or a prolonged INR (acute liver failure) admit to liver transplantation center.

**Prognosis:** Hepatitis A generally self-limited; Hepatitis B, C, D and rarely E may progress to chronic hepatitis/cirrhosis/hepatocellular carcinoma.

### **EBV/CMV Hepatitis**

**Clinical Presentation:** Similar to acute viral hepatitis plus bilateral posterior cervical adenopathy and fatigue.

**Diagnostic Considerations:** EBV hepatitis is part of infectious mononucleosis and may be the presenting sign in adults. CMV hepatitis also may appear as a “mono-like” infection in normal hosts. In compromised hosts, hepatitis may be the primary manifestation of CMV infection. Diagnosis of EBV/CMV hepatitis can be made by serology (positive mono spot test, high EBV VCA IgM titer, high CMV IgM titer, CMV early antigen detection (via shell vial cultures) or molecular amplification techniques.

**Pitfalls:** In normal and immunocompromised hosts with unexplained aminotransferase (ALT/AST) elevations, consider CMV hepatitis and order appropriate tests.

**Treatment:** None effective in EBV hepatitis; CMV hepatitis in normal hosts may be treated with Valganciclovir 900 mg (PO) q12h × 12 days.

**Prognosis:** In normal hosts EBV and CMV hepatitis are usually self-limited. In immunocompromised hosts, the prognosis of CMV hepatitis is related to the degree of Immunosuppression and the rapidity of onset of treatment.

### **Chronic Viral Hepatitis**

**Clinical Presentation:** Often asymptomatic with persistently or intermittently elevated serum aminotransferases but in some aminotransferases are normal. Some present with signs and symptoms of chronic liver disease.

**Diagnostic Considerations:** Chronic hepatitis B is diagnosed by presence of HBsAg and HBV DNA. Chronic hepatitis C is diagnosed by positive tests for anti-HCV and HCV RNA. Signal cutoff (SCO) >1.0 = HCV infection. SCO < 1.0 = false + test. Most HCV infections have SCOs > 5-10. Chronic hepatitis D is diagnosed by positive tests for HBsAg and HDV RNA. Chronic hepatitis E is diagnosed by presence of anti-HEV and HEV RNA. Liver biopsy may be used to grade inflammation and stage fibrosis.

**Pitfalls:** Autoimmune hepatitis may present as an acute or chronic hepatitis but with elevated IgG levels, elevated ANAs, and anti-smooth muscle antibodies. For both HCV and HBV, rule out co-infection with HIV.

**Therapeutic Considerations:** In chronic hepatitis B (see p. 97) treat until HBV DNA becomes undetectable, ALT normalizes, and for HBeAg-positive patients, HBeAg seroconversion occurs (then continue for 6–12 additional months). For chronic hepatitis C therapy is based on genotype (see p. 98).

**Prognosis:** Sustained clearance of HCV RNA (measured 24 weeks after discontinuation of treatment) results in a cure in 99% of chronic hepatitis C patients. Sustained clearance of HBV DNA is linked with an improved prognosis in chronic hepatitis B.

## Intraabdominal or Pelvic Peritonitis/Abscess

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	PO Therapy or IV-to-PO Switch
Mild/moderate peritonitis	Entero-bacteriaceae B. fragilis	Moxifloxacin 400 mg (IV) q24h <sup>‡</sup> <b>or</b> Cefoxitin 2 gm (IV) q6h <sup>‡</sup> <b>or</b> Piperacillin/tazobactam 3.375 gm (IV) q6h <sup>‡</sup>	Ampicillin/sulbactam 1.5 gm (IV) q6h <sup>‡</sup> <b>or</b> Ceftazidime/avibactam 2.5 gm (IV) q8h <b>plus</b> Metronidazole 500 mg (IV) q8h	Moxifloxacin 400 mg (PO) q24h <sup>‡</sup> <b>or</b> Amoxicillin/clavulanate 875/125 mg (PO) q12h <sup>‡</sup> <b>or combination therapy with</b> Levofloxacin 500 mg (PO) q24h <sup>‡</sup> <b>plus</b> Clindamycin 300 mg (PO) q8h <sup>‡</sup>
Severe peritonitis <sup>‡</sup>	Entero-bacteriaceae B. fragilis	Ertapenem 1 gm (IV) q24h <sup>‡</sup> <b>or</b> Ceftazidime/avibactam 2.5 gm (IV) q8h <b>plus</b> Metronidazole 1 gm (IV) q24h <sup>‡</sup> <b>or</b> Tigecycline 200 mg (IV) × 1 dose, then 100 mg (IV) q24h <sup>‡</sup> <b>or</b> Meropenem 1 gm (IV) q8h <sup>‡</sup>	Ampicillin/sulbactam 3 gm (IV) q6h <sup>‡</sup> <b>or</b> Doripenem 1 gm (IV) q8h <b>or combination therapy with</b> Metronidazole 1 gm (IV) q24h <sup>‡</sup> <b>plus either</b> Ceftriaxone 1 gm (IV) q24h <sup>‡</sup> <b>or</b> Levofloxacin 500 mg (IV) q24h	Moxifloxacin 400 mg (PO) q24h <sup>‡</sup> <b>or</b> Amoxicillin/clavulanate 875/125 mg (PO) q12h <sup>‡</sup>
	CRE	Ceftazidime/avibactam 2.5 gm (IV) q8h <sup>‡</sup> <b>plus</b> Metronidazole 1 gm (IV) q24h <sup>‡</sup>		

Duration of therapy represents total time IV, PO, or IV + PO.

‡ Duration of therapy as clinically indicated or for 5–7 days following corrective surgery.

**Intraabdominal or Pelvic Peritonitis/Abscess (cont'd)**

<b>Subset</b>	<b>Usual Pathogens</b>	<b>Preferred IV Therapy</b>	<b>Alternate IV Therapy</b>	<b>PO Therapy or IV-to-PO Switch</b>
Spontaneous bacterial peritonitis (SBP) <sup>‡</sup>	Enterobacteriaceae S. pneumoniae (children)	Ceftriaxone 1 gm (IV) q24h × 1–2 wks <b>or</b> Levofloxacin 500 mg (IV) × 1–2 weeks	Aztreonam 2 gm (IV) q8h × 1–2 weeks <b>or</b> Any aminoglycoside (IV) q24h × 1–2 wks	Quinolone* (PO) × 1–2 weeks <b>or</b> Amoxicillin/clavulanate 875/125 mg (PO) q12h × 1–2 weeks
Chronic TB peritonitis	M. tuberculosis	Not applicable		Treat the same as pulmonary TB (see p. 53)
CAPD-associated peritonitis <sup>‡</sup>	S. epidermidis (CoNS) S. aureus (MSSA) Enterobacteriaceae Non-fermentative GNBS	<u>Before culture results</u> Vancomycin 1 gm (IV) initial dose* <b>plus</b> Gentamicin 5 mg/kg or 240 mg (IV) initial dose*	<u>After culture results</u> <u>MSSA/Enterobacteriaceae</u> Meropenem 1 gm (IV) initial dose* <u>MRSA</u> Vancomycin 1 gm (IV) initial dose* <b>or</b> Linezolid 600 mg (IV or PO)	

MSSA/MRSA = methicillin-sensitive/resistant *S. aureus*. Duration of therapy represents total time IV, PO, or IV + PO.

<sup>‡</sup> Treat only IV or IV-to-PO switch.

\* Follow with maintenance dosing × 2 weeks after culture results are available. For maintenance dosing, use renal failure (CrCl < 10 mL/min) and post-peritoneal dialysis dosing (see Drug Summaries).

**Intraabdominal or Pelvic Peritonitis/Abscess****(Appendicitis/Diverticulitis/Septic Pelvic Thrombophlebitis)**

**Clinical Presentation:** Spiking fevers with acute abdominal pain and peritoneal signs. In diverticulitis, the pain is localized over the involved segment of colon. Appendicitis ± perforation presents as RLQ pain/rebound tenderness or mass. Peri-diverticular abscess presents the same as intraabdominal/pelvic abscess, most commonly in the LLQ. Septic pelvic thrombophlebitis (SPT) presents as high spiking fevers unresponsive to antibiotic therapy following delivery/pelvic surgery.

**Diagnostic Considerations:** Diagnosis by CT/MRI scan of abdomen/pelvis.

**Pitfalls:** Tympany over liver suggests abdominal/visceral perforation. Pelvic peritonitis/abscess presents the same as intraabdominal abscess/peritonitis, but peritoneal signs are often absent.

**Therapeutic Considerations:** Patients with ischemic/inflammatory colitis should be treated the same as peritonitis, depending on severity. Obtain surgical consult for repair/lavage or abscess drainage. In SPT, fever rapidly falls when heparin is added to antibiotics.

**Prognosis:** Related to degree/duration of peritoneal spillage and rapidity/completeness of lavage. Prognosis for SPT is good if treated early and clots remain limited to pelvic veins.

### **Spontaneous Bacterial Peritonitis (SBP)**

**Clinical Presentation:** Acute or subacute onset of fever  $\pm$  abdominal pain.

**Diagnostic Considerations:** Diagnosis by positive blood cultures of SBP pathogens. For patients with abdominal pain, ascites, and a negative CT/MRI, paracentesis ascitic fluid with  $> 500$  WBCs and  $> 100$  PMNs predicts a positive ascitic fluid culture and is diagnostic of SBP. Some degree of splenic dysfunction usually exists, predisposing to infection with encapsulated organisms.

**Pitfalls:** Do not overlook GI source of peritonitis (e.g. appendicitis, diverticulitis); obtain CT/MRI.

**Therapeutic Considerations:** *B. fragilis*/anaerobes are not common pathogens in SBP, and *B. fragilis* coverage is unnecessary.

**Prognosis:** Related to degree of hepatic/splenic dysfunction.

### **Chronic Tuberculous Peritonitis (Mycobacterium tuberculosis)**

**Clinical Presentation:** Abdominal pain with fevers, weight loss, ascites over 1–3 months.

**Diagnostic Considerations:** “Doughy consistency” on abdominal palpation. Diagnosis by AFB on peritoneal biopsy/culture.

**Pitfalls:** Chest x-ray is normal in  $\sim 70\%$ . Increased incidence in alcoholic cirrhosis.

**Therapeutic Considerations:** Treated the same as pulmonary TB.

**Prognosis:** Good if treated early.

### **CAPD Associated Peritonitis**

**Clinical Presentation:** Abdominal pain  $\pm$  fever in a CAPD patient.

**Diagnostic Considerations:** Diagnosis by gram stain/culture and  $\uparrow$  WBC count in peritoneal fluid. Lymphocytic predominance may suggest TB or fungi. Unlike SBP, there are not specific diagnostic criteria in peritoneal fluid.

**Pitfalls:** Fever is often absent.

**Therapeutic Considerations:** Treat with systemic antibiotics  $\pm$  antibiotics in dialysate.

**Prognosis:** Good with early therapy and removal of peritoneal catheter.

## Empiric Therapy of Genitourinary Tract Infections

### Dysuria-Pyuria Syndrome (Acute Urethral Syndrome)

Subset	Usual Pathogens	IV Therapy	PO Therapy
Acute urethral syndrome	S. saprophyticus C. trachomatis E. coli (< 10 <sup>5</sup> cfu/mL)	Not applicable	Doxycycline 100 mg (PO) q12h × 10 days <b>or</b> Quinolone (PO)* × 7 days

\* Ciprofloxacin 500 mg q12h or Levofloxacin 500 mg q24h.

**Clinical Presentation:** Dysuria, frequency, urgency, lower abdominal discomfort, fevers < 102°F.

**Diagnostic Considerations:** Diagnosis by symptoms of cystitis with pyuria and no growth or low concentration of E. coli ( $\leq 10^3$  colonies/mL) by urine culture. Clue to S. saprophyticus is alkaline urinary pH and RBCs in urine.

**Pitfalls:** Resembles "culture negative" cystitis.

**Therapeutic Considerations:** S. saprophyticus is susceptible to most antibiotics used to treat UTIs.

**Prognosis:** Excellent.

### Cystitis (see Color Atlas for Urine Gram stains)

Subset	Usual Pathogens	Therapy
Bacterial	Enterobacteriaceae E. faecalis (VSE) S. agalactiae (group B streptococci) S. saprophyticus	Amoxicillin 500 mg (PO) × q12h × 3 days <b>or</b> TMP-SMX 1 SS tablet (PO) × q12h × 3 days <b>or</b> Levofloxacin 500 mg (PO) q24h × 3 days <b>or</b> Nitrofurantoin 100 mg (PO) q12h × 3 days
	MDR GNBs	Fosfomycin 3 gm (PO) q24h × 3 days
Fungal	C. albicans <sup>†</sup>	Fluconazole 200 mg (PO) × 1 dose, then 100 mg (PO) q24h × 4 days
	Fluconazole-resistant <sup>‡</sup> Candida isolates or fluconazole-refractory disease	Amphotericin B 0.3 mg/kg (IV) × 1 dose

<sup>†</sup> C. albicans cystitis (see p. 104)

<sup>‡</sup> Fluconazole-resistant (see p. 104)

### Bacterial Cystitis

**Clinical Presentation:** Dysuria, frequency, urgency, lower abdominal discomfort, fevers < 102°F.

**Diagnostic Considerations:** Pyuria plus bacteriuria.

**Pitfalls:** Compromised hosts (chronic steroids, diabetes, SLE, cirrhosis, multiple myeloma) may require 3–5 days of therapy. A single dose of amoxicillin or TMP–SMX may be sufficient in acute uncomplicated cystitis in normal hosts.

**Therapeutic Considerations:** Pyridium 200 mg (PO) q8h after meals × 24–48h is useful to decrease dysuria (inform patients urine will turn orange).

**Prognosis:** Excellent in normal hosts.

### Candida Cystitis

**Diagnostic Considerations:** Marked pyuria, urine nitrate negative ± RBCs. Speciate if not *C. albicans*.

**Pitfalls:** Lack of response suggests renal candidiasis or a “fungus ball” in the renal collecting system.

**Therapeutic Considerations:** If fluconazole fails, use amphotericin. For chronic renal failure/dialysis patients with candiduria, use amphotericin B deoxycholate bladder irrigation (as for catheter-associated candiduria, below). Removal of devices and correction of anatomic abnormalities are critical to success.

**Prognosis:** Patients with impaired host defenses, abnormal collecting systems, cysts, renal disease or stones are prone to recurrent UTIs/uropsepsis.

## Catheter Associated Bacteriuria (CAB)

Subset	Usual Pathogens	Therapy
Catheter associated bacteriuria (CAB) <sup>†*</sup>	<i>E. coli</i> <i>E. faecalis</i> (VSE)	Nitrofurantoin 100 mg (PO) q12h × 3 days <b>or</b> Amoxicillin 500 mg (PO) q12h × 3 days
	<i>E. faecium</i> (VRE)	Nitrofurantoin 100 mg (PO) q12h × 3 days <b>or</b> Fosomycin 3 gm (PO) q24h × 3 days <sup>††</sup>
	MDR <i>Klebsiella</i> <sup>††</sup> MDR <i>Acinetobacter</i> <sup>††</sup> MDR <i>P. aeruginosa</i> <sup>††</sup>	Nitrofurantoin 100 mg (PO) q12h × 3 days <sup>††</sup> <b>or</b> Fosomycin 3 gm (PO) q24h × 3 days <sup>††</sup> <b>or</b> Meropenem 1 gm (IV) q8h × 3 day <sup>††§</sup> <b>or</b> Doripenem 1 gm (IV) q8h × 3 days <sup>††§</sup>

VSE/VRE = vancomycin-sensitive/resistant enterococci, MDR = multidrug resistant

\* **Remove/replace Foley catheter before initiating antibiotic therapy.**

† No need to treat CAB in normal hosts, pre-emptive therapy suggested in compromised hosts, e.g., cirrhosis, SLE, DM, myeloma, steroids, immunosuppressives, and those with renal insufficiency.

†† Longer courses of therapy may be needed in compromised hosts. If pyuria substantially decreased after 2 days of therapy, complete 3 days of therapy. If not, continue therapy for 7 days.

§ Only if oral therapy not possible.

## Catheter Associated Candiduria (CAC)

Subset	Usual Pathogens	Therapy
Catheter associated candiduria (CAC)*	<i>C. albicans</i>	Fluconazole 200 mg (PO) × 1 dose, then 100 mg (PO) q24h × 2 weeks
	Fluconazole-resistant <i>C. albicans</i> /non- <i>albicans</i> <i>Candida</i>	Amphotericin B 0.3–0.6 mg/kg q24h × 1–7 days <b>or</b> Flucytosine 25 mg/kg QID (PO) × 7–10 days <b>or</b> Amphotericin B bladder irrigation (continuous: 50 mg in 1 liter sterile water over 24h × 1–2 days; intermittent: 50 mg in 200–300 ml sterile water q6-8h × 1–2 days) <sup>§</sup>

§ Limited efficacy.

**Clinical Presentation:** Indwelling urinary (Foley) catheter with bacteriuria and pyuria; no symptoms.

**Diagnostic Considerations:** Pyuria plus bacteriuria/candiduria. Usually afebrile or temperature < 101°F.

**Pitfalls:** Bacteriuria/candiduria often represent colonization, not infection. Persistent candiduria after amphotericin B deoxycholate bladder irrigation suggests renal candidiasis.

**Therapeutic Considerations:** Avoid treating catheter-associated bacteriuria in normal hosts without GU tract abnormalities/disease. Compromised hosts (diabetes, SLE, chronic steroids, multiple myeloma, cirrhosis) may require therapy for duration of catheterization. If bacteriuria/candiduria does not clear with appropriate therapy, change the catheter. For chronic renal failure/dialysis patients with candiduria, use amphotericin B deoxycholate bladder irrigation. Efficacy of therapy of catheter-associated candiduria is limited and relapse is frequent unless the catheter can be replaced or (preferably) removed.

**Prognosis:** Excellent in normal hosts. Untreated bacteriuria/candiduria in compromised hosts may result in ascending infection (e.g., pyelonephritis) or bacteremia/candidemia.



## Epididymitis

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	PO Therapy or IV-to-PO Switch
Acute <i>Young males</i>	<i>C. trachomatis</i>	Doxycycline 200 mg (IV) q12h × 3 days, then 100 mg (IV) q12h × 4 days	Levofloxacin 500 mg (IV) q24h × 7 days	Doxycycline 200 mg (PO) q12h × 3 days, then 100 mg (PO) q12h × 7 days* <b>or</b> Levofloxacin 500 mg (PO) q24h × 10 days <b>or</b> Ofloxacin 300 mg (PO) q12h × 10 days
<i>Elderly males</i>	<i>P. aeruginosa</i>	Cefepime 2 gm (IV) q8h × 10 days <b>or</b> Meropenem 1 gm (IV) q8h × 10 days	Ciprofloxacin 400 mg (IV) q8h × 10 days <b>or</b> Levofloxacin 750 mg (IV) q24h × 10 days	Ciprofloxacin 750 mg (PO) q12h × 10 days
Chronic	<i>M. tuberculosis</i> <i>Blastomyces dermatitidis</i>	Treat the same as pulmonary TB (see p. 53) or pulmonary blastomycosis (see p. 267)		

Duration of therapy represents total time IV, PO, or IV + PO. Most patients on IV therapy able to take PO meds should be switched to PO therapy soon after clinical improvement (usually < 72 hours).

\* Loading dose is not needed PO if given IV with the same drug.

### **Acute Epididymitis (*Chlamydia trachomatis*/*Pseudomonas aeruginosa*)**

**Clinical Presentation:** Acute unilateral testicular pain ± fever.

**Diagnostic Considerations:** Ultrasound to rule out torsion or tumor.

**Pitfalls:** Rule out torsion by absence of fever and ultrasound.

**Therapeutic Considerations:** Young males respond to treatment slowly over 1 week. Elderly males respond to anti-Pseudomonal therapy within 72 hours.

**Prognosis:** Excellent in young males. Related to health of host in elderly.

### **Chronic Epididymitis (*Mycobacterium tuberculosis*/*Blastomyces dermatitidis*)**

**Clinical Presentation:** Chronic epididymo-orchitis with epididymal nodules.

**Diagnostic Considerations:** Diagnosis by AFB on biopsy/culture of epididymis. TB epididymitis is always associated with renal TB. Blastomyces epididymitis is a manifestation of systemic infection.

**Pitfalls:** Vasculitis (e.g., polyarteritis nodosum) and lymphomas may present the same way.

**Therapeutic Considerations:** Treated the same as pulmonary TB/blastomycosis.

**Prognosis:** Good.

**Acute Pyelonephritis** (see Color Atlas for Urine Gram stains)

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	PO Therapy or IV-to-PO Switch
Acute pyelonephritis  (Treat initially based on urine gram stain; see therapeutic considerations, below)	Enterobacteriaceae	Ceftriaxone 1 gm (IV) q24h × 2 weeks <b>or</b> Levofloxacin 500 mg (IV) q24h × 2 weeks	Meropenem 1 gm (IV) q8h × 2 weeks <b>or</b> Aztreonam 2 gm (IV) q8h × 2 weeks <b>or</b> Gentamicin 240 mg (IV) q24h × 2 weeks	Levofloxacin 500 mg (PO) q24h × 2 weeks <b>or</b> Amoxicillin 1 gm (PO) q8h × 2 weeks
	Enterococcus faecalis (VSE) <sup>§</sup>	Ampicillin 1 gm (IV) q4h × 2 weeks <b>or</b> Linezolid 600 mg (IV) q12h × 2 weeks <b>or</b> Meropenem 1 gm (IV) q8h × 2 weeks	Quinolone (IV) × 2 weeks	Amoxicillin 1 gm (PO) q8h × 2 weeks <b>or</b> Linezolid 600 mg (PO) q12h × 2 weeks <b>or</b> Levofloxacin 500 mg (PO) q24h × 2 weeks
	Enterococcus faecium (VRE)	Linezolid 600 mg (IV) q12h × 2 weeks	Quinupristin/ dalfopristin 7.5 mg/ kg (IV) q8h × 2 weeks <b>or</b> Doxycycline 200 mg (IV) q12h × 3 days, then 100 mg q12h × 2 weeks	Linezolid 600 mg (PO) q12h × 2 weeks <b>or</b> Doxycycline 200 mg (PO) q12h × 3 days, then 100 mg (PO) q12h × 2 weeks
	MDR GNB	Meropenem 1 gm (IV) q8h × 2 weeks <b>or</b> Ceftolozane/ tazobactam 1.5 gm (IV) q8h × 2 weeks	Colistin 5 mg/kg (IV) q8h × 2 weeks	
	CRE	Ceftazidime/ avibactam 2.5 gm (IV) q8h × 2 weeks	Tigecycline 200 mg (IV) × 1 dose then 100 mg (IV) q24h × 2 weeks* <b>or</b> Colistin 5 mg/kg (IV) q8h × 2 weeks	

VSE/VRE = vancomycin-sensitive/resistant enterococci.

\* Depending on MICs, higher doses may be necessary LD: 400 mg (IV) × 1 dose, then MD: 200 mg (IV) q24h.

§ Treat vancomycin resistant *E. faecalis* as VRE.

## Chronic Pyelonephritis/Renal TB

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	PO Therapy or IV-to-PO Switch
Chronic pyelonephritis	Enterobacteriaceae	IV Therapy Not applicable		Quinolone <sup>†</sup> (PO) × 4–6 weeks <b>or</b> TMP-SMX 1 DS tab (PO) q12h × 4–6 weeks <b>or</b> Doxycycline 200 mg (PO) q12h × 3 days, then 100 mg (PO) q12h × 4–6 weeks total
Renal TB	M. tuberculosis	IV Therapy Not applicable		Treated the same as pulmonary TB (see p. 53)

Duration of therapy represents total time IV, PO, or IV + PO. Most patients on IV therapy able to take PO meds should be switched to PO therapy after clinical improvement (usually < 72 hours).

† Ciprofloxacin XR 1000 mg (PO) q24h or Ciprofloxacin 400 mg (IV) q12h or Levofloxacin 500 mg (IV or PO) q24h.

### **Acute Bacterial Pyelonephritis (Enterobacteriaceae, E. faecalis/faecium)**

**Clinical Presentation:** Unilateral CVA tenderness with fevers  $\geq 102^{\circ}\text{F}$ .

**Diagnostic Considerations:** Bacteriuria plus pyuria with unilateral CVA tenderness and temperature  $\geq 102^{\circ}\text{F}$ . Bacteremia usually accompanies acute pyelonephritis; obtain blood and urine cultures.

**Pitfalls:** Temperature decreases in 72 hours with or without antibiotic treatment. If temperature does not fall after 72 hours of antibiotic therapy, suspect renal/perinephric abscess.

**Therapeutic Considerations:** Initial treatment is based on the urinary gram stain: If gram-negative bacilli, treat as Enterobacteriaceae. If gram-positive cocci in chains (enterococcus), treat as E. faecalis; if enterococcus is subsequently identified as E. faecium, treat accordingly. Acute pyelonephritis is usually treated initially for 1–3 days IV, then switched to PO to complete 4 weeks of antibiotics to minimize progression to chronic pyelonephritis. Obtain a CT/MRI in persistently febrile patients after 72 hours of antibiotics to rule out renal calculi, obstruction, abscess, or xanthomatous pyelonephritis.

**Prognosis:** Excellent if first episode is adequately treated with antibiotics for 4 weeks.

### **Chronic Bacterial Pyelonephritis (Enterobacteriaceae)**

**Clinical Presentation:** Previous history of acute pyelonephritis with same symptoms as acute pyelonephritis but less CVA tenderness/fever.

**Diagnostic Considerations:** Diagnosis by CT/MRI showing changes of chronic pyelonephritis plus bacteriuria/pyuria. Urine cultures may be intermittently negative before treatment. Chronic pyelonephritis is bilateral pathologically, but unilateral clinically.

**Pitfalls:** Urine culture may be intermittently positive after treatment; repeat weekly × 4 to confirm urine remains culture-negative.

**Therapeutic Considerations:** Treat × 4–6 weeks. Impaired medullary vascular blood supply/renal anatomical distortion makes eradication of pathogen difficult.

**Prognosis:** Related to extent of renal damage.

**Renal TB (*Mycobacterium tuberculosis*)**

**Clinical Presentation:** Renal mass lesion with ureteral abnormalities (pipestem, corkscrew, or spiral ureters) microscopic hematuria/sterile pyuria. Painless unless complicated by ureteral obstruction.

**Diagnostic Considerations:** Combined upper/lower urinary tract abnormalities  $\pm$  microscopic hematuria/urinary pH  $\leq$  5.5. Diagnosis by culture of TB from urine. Urine TB PCR is specific, but not very sensitive.

**Pitfalls:** Chest x-ray is normal in 50%, but most patients are PPD positive. Rule out other infectious/inflammatory causes of sterile pyuria (e.g., *Trichomonas*, interstitial nephritis).

**Therapeutic Considerations:** Treat the same as pulmonary TB.

**Prognosis:** Good if treated before renal parenchymal destruction/ureteral obstruction occur.

**Renal Abscess (Intrarenal/Perinephric)**

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	PO Therapy or IV-to-PO Switch
Cortical (Treat initially for MSSA; if later identified as MRSA, treat accordingly)	<i>S. aureus</i>	MSSA: Nafcillin 2 gm (IV) q4h* <b>or</b> Ceftriaxone 1 gm (IV) q24h* <b>or</b> Clindamycin 600 mg (IV) q8h*	MSSA Meropenem 1 gm (IV) q8h* <b>or</b> Ertapenem 1 gm (IV) q24h*	MSSA/MRSA Linezolid 600 mg (PO) q12h* <b>or</b> Minocycline 100 mg (PO) q12h*
		MRSA: Linezolid 600 mg (IV) q12h* <b>or</b> Minocycline 100 mg (IV) q12h*	MRSA Vancomycin 1 gm (IV) q12h*	
Medullary	Entero-bacteriaceae	Quinolone (IV)**	TMP-SMX 2.5 mg/kg (IV) q6h*	Quinolone (PO)**

MSSA/MRSA = methicillin-sensitive/resistant *S. aureus*. Duration of therapy represents total time IV, PO, or IV + PO. Most patients on IV therapy able to take PO meds should be switched to PO therapy soon after clinical improvement.

\* Treat until renal abscess resolves completely or is no longer decreasing in size on CT/MRI.

† Levofloxacin 500 mg (IV/PO) q24h.

**Clinical Presentation:** Similar to pyelonephritis but fever remains elevated after 72 hours of antibiotics.

**Diagnostic Considerations:** Obtain CT/MRI to diagnose perinephric/intra-renal abscess and rule out mass lesion. Cortical abscesses are usually secondary to hematogenous/contiguous spread. Medullary abscesses are usually due to extension of intrarenal infection.

**Pitfalls:** Urine cultures may be negative with cortical abscesses.

**Therapeutic Considerations:** Most large abscesses need to be drained. Multiple small abscesses are managed medically. Obtain urology consult.

**Prognosis:** Related to degree of baseline renal dysfunction.

## Prostatitis/Prostatic Abscess

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	PO Therapy or IV-to-PO Switch
Acute prostatitis/ acute prostatic abscess	Entero- bacteriaceae	Quinolone* (IV) × 2 weeks <b>or</b> Ceftriaxone 1 gm (IV) q24h × 2 weeks	TMP-SMX 2.5 mg/kg (IV) q6h × 2 weeks <b>or</b> Aztreonam 2 gm (IV) q8h × 2 weeks	Quinolone* (PO) × 2 weeks <b>or</b> Doxycycline 200 mg (PO) q12h × 3 days, then 100 mg (PO) q24h × 11 days <b>or</b> TMP-SMX 1 SS tablet (PO) q12h × 2 weeks
Chronic prostatitis	Entero- bacteriaceae		Quinolone* (PO) × 1–3 months <b>or</b> TMP-SMX 1 DS tablet (PO) q12h × 1–3 months	
	MDR GNB		Fosfomycin 3 gm (PO) q48h × 30 days <b>±</b> Doxycycline 100 mg (PO) q24h × 1–3 months	

Duration of therapy represents total time IV, PO, or IV + PO. Most patients on IV therapy able to take PO meds should be switched to PO therapy soon after clinical improvement (usually < 72 hours).

\* Ciprofloxacin XR 1000 mg (PO) q24h or Ciprofloxacin 400 mg (IV) q12h or Levofloxacin 500 mg (IV or PO) q24h.

### **Acute Prostatitis/Acute Prostatic Abscess (Enterobacteriaceae)**

**Clinical Presentation:** Acute prostatitis presents as an acute febrile illness in males with dysuria and no CVA tenderness. Prostatic abscess presents with hectic/septic fevers without localizing signs.

**Diagnostic Considerations:** Acute prostatitis is diagnosed by bacteriuria, pyuria plus mucus threads, with exquisite prostate tenderness, and is seen primarily in young males. Positive urine culture is due to contamination of urine as it passes through infected prostate. Prostatic abscess is diagnosed by transrectal ultrasound or CT/MRI of prostate.

**Pitfalls:** Do not overlook acute prostatitis in males with bacteriuria without localizing signs, or prostatic abscess in patients with a history of prostatitis.

**Therapeutic Considerations:** Treat acute prostatitis for 2 full weeks to decrease progression to chronic prostatitis. Prostatic abscess is treated the same as acute prostatitis plus surgical drainage.

**Prognosis:** Excellent if treated early with full course of antibiotics (plus drainage for prostatic abscess).

### Chronic Prostatitis (Enterobacteriaceae/MDR GNB)

**Clinical Presentation:** Vague urinary symptoms (mild dysuria  $\pm$  low back pain), history of acute prostatitis, and little or no fever.

**Diagnostic Considerations:** Diagnosis by bacteriuria plus pyuria with mucus threads  $\pm$  mild prostate tenderness. Urine, semen, or prostate expressate are culture positive.

**Pitfalls:** Commonest cause of treatment failure is inadequate duration of therapy. Chronic prostatitis with prostatic calcifications (transrectal ultrasound) will not clear with antibiotics; transurethral resection of prostate (TURP) with removal of all calcifications curative.

**Therapeutic Considerations:** In sulfa-allergic patients, TMP alone may be used in place of TMP-SMX.

**Prognosis:** Excellent if treated  $\times$  1–3 months. Prostatic abscess is a rare but serious complication (may cause urosepsis).

### Urosepsis (see Color Atlas for Urine Gram stains)

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	IV-to-PO Switch
Community acquired  (Treat initially based on urine gram stain)	Entero-bacteriaceae (ESBL-)	Ceftriaxone 1 gm (IV) q24h $\times$ 7 days*  <b>or</b> Levofloxacin 500 mg (IV) q24h $\times$ 7 days*	Amikacin 1 gm (IV) q24h $\times$ 7 days*  <b>or</b> Aztreonam 2 gm (IV) q8h $\times$ 7 days*	Levofloxacin 500 mg (PO) q24h $\times$ 7 days*  <b>or</b> TMP-SMX 1 SS tablet (PO) q12h $\times$ 7 days*
	(ESBL +)	Meropenem 1 gm (IV) of q8h $\times$ 7 days*	Doripenem 1 gm (IV) of q8h $\times$ 7 days*	Fosfomycin 3 gm (PO) q 3 days $\times$ 7 days*
	E. faecalis (VSE) Group B streptococci	Ampicillin 2 gm (IV) q4h $\times$ 7 days*	Meropenem 1 gm (IV) q8h $\times$ 7 days*	Amoxicillin 1 gm (PO) q8h $\times$ 7 days*  <b>or</b> Levofloxacin 500 mg (PO) q24h $\times$ 7 days*
(No urine gram stain)	Entero-bacteriaceae E. faecalis (VSE) Group B streptococci	Meropenem 1 gm (IV) q8h $\times$ 7 days*	Piperacillin/tazobactam 3.375 mg (IV) q6h $\times$ 7 days*	Levofloxacin 500 mg (PO) q24h $\times$ 7 days*

\* Longer duration needed if urologic/renal abnormalities present.

**Urosepsis (cont'd)** (see Color Atlas of Urine Gram stains)

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	IV-to-PO Switch
Related to urological procedure  (Treat initially for <i>P. aeruginosa</i> , etc; if later identified as non- <i>aeruginosa Pseudomonas</i> , treat accordingly)	<i>P. aeruginosa</i> Enterobacter Klebsiella Serratia	Meropenem 1 gm (IV) q8h × 7 days <b>or</b> Levofloxacin 750 mg (IV) q24h × 7 days <b>or</b> Cefepime 2 gm (IV) q8h × 7 days	Doripenem 1 gm (IV) q8h × 7 days <b>or</b> Aztreonam 2 gm (IV) q8h × 7 days <b>or</b> Amikacin 1 gm (IV) q24h × 7 days <b>or</b> Ciprofloxacin 400 mg (IV) q8h × 7 days	Ciprofloxacin 750 mg (PO) q12h × 7 days <b>or</b> Levofloxacin 750 mg (PO) q24h × 7 days
	MDR GNB	Meropenem 1 gm (IV) q8h × 2 weeks <b>or</b> Ceftolozane/ tazobactam 1.5 gm (IV) q8h × 2 weeks	Colistin 5 mg/kg (IV) q8h × 2 weeks	
	CRE	Ceftazidime/ avibactam 2.5 gm (IV) q8h × 2 weeks <b>or</b> Tigecycline 200 mg (IV) × 1 dose then 100 mg (IV) q24h × 2 weeks*	Colistin 5 mg/kg (IV) q8h × 2 weeks	

\* Depending on MICs, higher doses may be necessary LD: 400 mg (IV) × 1 dose, then MD: 200 mg (IV) q24h.

**Community-Acquired Urosepsis**

**Clinical Presentation:** Sepsis from urinary tract source.

**Diagnostic Considerations:** Blood and urine cultures positive for same uropathogen. If patient does not have diabetes, SLE, cirrhosis, myeloma, steroids, pre-existing renal disease or obstruction, obtain CT/MRI of GU tract to rule out abscess/obstruction. Prostatic abscess is rarely a cause of urosepsis.

**Pitfalls:** Mixed gram-positive/negative urine cultures suggest specimen contamination or enterovesicular fistula.

**Therapeutic Considerations:** Empiric treatment is based on urine gram stain. If urine gram stain shows pyuria and gram-positive cocci, treat as group D enterococci (*E. faecalis*-VSE). If gram-negative bacilli, treat as Enterobacteriaceae. *S. aureus*/*S. pneumoniae* are not uropathogens.

**Prognosis:** Related to severity of underlying condition causing urosepsis and health of host.

**Urosepsis Following Urological Procedures**

**Clinical Presentation:** Sepsis within 24 hours after GU procedure.

**Diagnostic Considerations:** Blood and urine cultures positive for same uropathogen. Use pre-procedural urine culture to identify uropathogen and guide therapy.

**Pitfalls:** If non-*aeruginosa Pseudomonas* in urine/blood, switch to TMP-SMX pending susceptibilities.

**Therapeutic Considerations:** Empiric *P. aeruginosa* monotherapy will cover most other uropathogens.

**Prognosis:** Related to severity of underlying condition causing urosepsis and health of host.

## Pelvic Inflammatory Disease (PID), Salpingitis, Tuboovarian Abscess, Endometritis/Endomyometritis, Septic Abortion

Subset	Usual Pathogens	IV Therapy	PO Therapy or IV-to-PO Switch
Hospitalized patients†	B. fragilis Enterobacteriaceae N. gonorrhoeae C. trachomatis C. sordelli‡ (septic abortion)	<b>Monotherapy with</b> Moxifloxacin 400 mg (IV) q24h × 2 weeks <b>or combination therapy with</b> Doxycycline 200 mg (IV) q12h × 3 days, then 100 mg (IV) q12h × 11 days <b>plus either</b> Piperacillin/tazobactam 4.5 gm (IV) q8h × 2 weeks <b>or</b> Ertapenem 1 gm (IV) q24h × 3–10 days <b>or</b> Cefoxitin 2 gm (IV) q6h × 2 weeks <b>or</b> Cefotetan 2 gm (IV) q12h × 2 weeks <b>Alternate combination therapy</b> Doxycycline 200 mg (IV) q12h × 3 days, then 100 mg (IV) q12h × 11 days <b>plus</b> Ampicillin/sulbactam 3 gm (IV) q6h × 2 weeks <b>or</b> Quinolone‡ (IV) q24h × 2 weeks <b>plus</b> Metronidazole 1 gm (IV) q24h × 2 weeks	<b>Monotherapy with</b> Moxifloxacin 400 mg (PO) q24h × 2 weeks
Outpatients (mild PID only)	N. gonorrhoeae C. trachomatis B. fragilis Enterobacteriaceae	Moxifloxacin 400 mg (PO) q24h × 2 weeks** <b>or</b> Doxycycline 100 mg (PO) q12h × 2 weeks	

Duration of therapy represents total time IV, PO, or IV + PO. Most patients on IV therapy able to take PO meds should be switched to PO therapy after clinical improvement.

† Treat only IV or IV-to-PO switch for salpingitis, tuboovarian abscess, endometritis, endomyometritis, septic abortion, or severe PID.

‡ Levofloxacin 500 mg (IV or PO) q24h or Ofloxacin 400 mg (IV or PO) q12h.

\*\* Recent gonococcal resistance to quinolones requires careful follow-up during/after therapy.

§ Antibiotic therapy of septic abortion same as salpingitis/endometritis *plus* evacuation of uterine contents.

**Clinical Presentation:** PID/salpingitis presents with cervical motion/adnexal tenderness, lower quadrant abdominal pain, and fever. Endometritis/endomyometritis presents with uterine tenderness ± cervical discharge/fever. Endomyometritis is the most common postpartum infection.

**Diagnostic Considerations:** Unilateral lower abdominal pain in a female without a non-pelvic cause suggests PID/salpingitis.

**Pitfalls:** Obtain CT/MRI of abdomen/pelvis to confirm diagnosis and rule out other pathology or tuboovarian abscess.

**Therapeutic Considerations:** Tuboovarian abscess usually requires drainage/removal ± TAH/BSO, plus antibiotics (see p. 113) × 1–2 weeks after drainage/removal. Septic abortion is treated the same as endometritis/endomyometritis plus uterine evacuation.



**Prognosis:** Related to promptness of treatment/adequacy of drainage if tuboovarian abscess. Late complications of PID/salpingitis include tubal scarring/infertility.

## Empiric Therapy of Sexually Transmitted Diseases

### Urethritis/Cervicitis

Subset	Usual Pathogens	IM Therapy	PO Therapy
Gonococcal (GC)	<i>N. gonorrhoeae</i>	Ceftriaxone 250 mg (IM) × 1 dose <b>plus either</b> Azithromycin 1–2 gm (PO) × 1 dose <b>or</b> Doxycycline 100 mg (PO) of q12h × days	Cefixime <sup>†</sup> 400 mg (PO) × 1 dose <sup>†</sup> <b>or</b> Cefpodoxime <sup>†</sup> 400 mg (PO) × 1 dose <sup>†</sup> <b>or</b> Azithromycin <sup>†</sup> 2 gm (PO) × 1 dose <sup>†</sup>
Non-gonococcal (NGU)	<i>C. trachomatis</i> <i>U. urealyticum</i> <i>M. genitalium</i>	Not applicable	Doxycycline 100 mg (PO) q12h × 7 days <sup>‡</sup> <b>or</b> Quinolone* <sup>†</sup> (PO) × 7 days <b>or</b> Azithromycin 1 gm (PO) × 1 dose <b>or</b> Erythromycin 500 mg (PO) q6h × 7 days
	<i>Trichomonas vaginalis</i>	Not applicable	Tinidazole 2 gm (PO) × 1 dose <b>or</b> Metronidazole 2 gm (PO) × 1 dose

\* Levofloxacin 500 mg q24h or Moxifloxacin 400 mg q24h or Ofloxacin 300 mg q12h.

† Increased resistance oral regimens should be considered alternate (second line) therapy and test of cure essential.

‡ (Doxycycline may be given as a 200 mg (PO) of q24h × 7 days.

### Gonococcal Urethritis/Cervicitis (*Neisseria gonorrhoeae*)

**Clinical Presentation:** Purulent penile/cervical discharge with burning/dysuria 3–5 days after contact.

**Diagnostic Considerations:** Rapid diagnosis in males by Gram stain of urethral discharge showing gram-negative diplococci; urethral cultures also positive. In females, diagnosis requires identification of organism by culture or DNA probe, not Gram stain. Rapid diagnosis in males/females by DNA probe. Obtain throat/rectal culture for *N. gonorrhoeae*. Co-infections are common; obtain syphilis and HIV serologies.

**Pitfalls:** Gram stain of cervical discharge showing gram-negative diplococci is not diagnostic of *N. gonorrhoeae*; must confirm by culture or nucleic acid amplification. *N. gonorrhoeae* infections are asymptomatic in 10% of men and 70% of women.

**Therapeutic Considerations:** Failure to respond suggests re-infection or relapse. Treat all GC with ceftriaxone. Because of frequent coinfection with agents of NGU, add oral azithromycin. Because of increasing resistance, consider using azithromycin 2 gm for dual therapy in patients with GC.

**Prognosis:** Good even with disseminated infection.

### **Non-Gonococcal Urethritis/Cervicitis (Chlamydia/Ureaplasma/Mycoplasma)**

**Clinical Presentation:** Mucopurulent penile/cervical discharge  $\pm$  dysuria  $\sim$  1 week after contact.

**Diagnostic Considerations:** Diagnosis by positive chlamydial NAAT/Ureaplasma or Mycoplasma NAAT of urethral/cervical discharge. Evaluate urethral/cervical discharge to rule out *N. gonorrhoeae*. Co-infections are common; obtain syphilis and HIV serologies.

**Pitfalls:** *C. trachomatis* infections are asymptomatic in 25%.

**Therapeutic Considerations:** Failure to respond to doxycycline therapy suggests re-infection or *Trichomonas/Ureaplasma/Mycoplasma* infection. Failure to respond to azithromycin suggests trichomoniasis, or infection due to a resistant *Mycoplasma* (consider quinolone therapy).

**Prognosis:** Tubal scarring/infertility in chronic infection.

### **Trichomonas Urethritis/Cervicitis (Trichomonas vaginalis)**

**Clinical Presentation:** Frothy, pruritic vaginal discharge.

**Diagnostic Considerations:** *Trichomonas* by wet mount/culture on special media.

**Pitfalls:** Classic "strawberry cervix" is infrequently seen.

**Therapeutic Considerations:** Use week-long regimen if single dose fails. Resistance now recognized as a cause of treatment failure.

**Prognosis:** Excellent if partner is also treated.

## **Vaginitis/Vaginosis**

Subset	Usual Pathogens	PO Therapy
Bacterial vaginosis/vaginitis	Polymicrobial ( <i>Gardnerella vaginalis</i> , <i>Mobiluncus</i> , <i>Prevotella</i> , <i>M. hominis</i> , etc.)	Tinidazole 1 gm (PO) $\times$ 5 days <b>or</b> 2 gm (PO) $\times$ 2 days <b>or</b> Clindamycin 300 mg (PO) q12h $\times$ 7 days <b>or</b> Metronidazole 500 mg (PO) q12h $\times$ 7 days
Candida vaginitis/balanitis	<i>Candida</i>	Fluconazole 150 mg (PO) $\times$ 1 dose <sup>†</sup>

<sup>†</sup> Those failing to respond should be treated with Fluconazole 200 mg (PO)  $\times$  1 dose then 100 mg (PO) q24h  $\times$  1 week.

### **Bacterial Vaginosis/Vaginitis**

**Clinical Presentation:** Non-pruritic vaginal discharge with "fishy" odor.

**Diagnostic Considerations:** Diagnosis by "clue cells" in vaginal fluid wet mount. Vaginal pH  $\geq$  4.5.

**Pitfalls:** "Fishy" odor from smear of vaginal secretions intensified when 10% KOH solution is added (positive "whiff test").

**Therapeutic Considerations:** As an alternative to oral therapy, clindamycin cream 2% intravaginally qHS  $\times$  7 days (avoid in pregnancy) or metronidazole gel 0.075% 1 application intravaginally q12h  $\times$  5 days can be used.

**Prognosis:** Complications include premature rupture of membranes, premature delivery, increased risk of PID. Recurrences very common.

**Candida Vaginitis/Balanitis**

**Clinical Presentation:** Pruritic white plaques in vagina/erythema of glans penis.

**Diagnostic Considerations:** Diagnosis by gram stain/culture of whitish plaques.

**Pitfalls:** Rule out Trichomonas, which also presents with pruritus in females.

**Therapeutic Considerations:** Uncomplicated vaginitis (mild sporadic infections in healthy individuals) responds readily to single-dose therapy. Complicated vaginitis (severe, recurrent, or in difficult-to-control diabetes) often requires  $\geq 7$  days of therapy (daily topical therapy or 2 doses of fluconazole 150 mg given 72h apart). Non-albicans infections respond poorly to azoles. Topical boric acid (600 mg/d in a gelatin capsule  $\times$  14 days) is often effective in this setting.

**Prognosis:** Good with systemic therapy. Diabetics/uncircumcised males may need prolonged therapy.

**Genital Vesicles (Genital Herpes) (HSV-2/HSV-1)**

Subset	PO Therapy
Initial therapy	Acyclovir 200 mg (PO) 5 $\times$ /day $\times$ 10 days <b>or</b> Famciclovir 500 mg (PO) q12h $\times$ 7–10 days <b>or</b> Valacyclovir 1 gm (PO) q12h $\times$ 3 days
Recurrent/ intermittent therapy ( $< 6$ episodes/year)	Acyclovir 200 mg (PO) 5 $\times$ /day $\times$ 5 days <b>or</b> Valacyclovir ( <u>normal host</u> : 500 mg [PO] q24h $\times$ 5 days; <u>HIV positive</u> : 1 gm [PO] q12h $\times$ 7–10 days)** <b>or</b> Famciclovir ( <u>normal host</u> : 125 mg [PO] q12h $\times$ 5 days <b>or</b> 1 gm [PO] q12h $\times$ 1 day*; <u>HIV-positive</u> : 500 mg [PO] q12h $\times$ 7 days)
Chronic suppressive therapy ( $> 6$ episodes/year)	Acyclovir 400 mg (PO) q12h $\times$ 1 year <b>or</b> Valacyclovir ( <u>normal host</u> : 1 gm [PO] q24h $\times$ 1 year; <u>HIV-positive</u> : 500 mg [PO] q12h $\times$ 1 year) <b>or</b> Famciclovir 250 mg (PO) q12h $\times$ 1 year

\* Patient initiated therapy to be started immediately when recurrence begins.

\*\* Short-course therapy with Valacyclovir 500 mg (PO) q12h  $\times$  3 days or Acyclovir 800 mg (PO) q8h  $\times$  2 days also effective.

**Clinical Presentation:** Painful vesicles/ulcers on genitals with painful bilateral regional adenopathy  $\pm$  low-grade fever.

**Diagnostic Considerations:** Diagnosis by clinical presentation may be misleading.

**Pitfalls:** 70% of newly acquired genital herpes is due to HSV-1. Elevated IgG HSV-2 titer indicates past exposure, not acute infection. HSV-2 IgM titers may be negative.

**Therapeutic Considerations:** If concomitant rectal herpes, increase acyclovir to 800 mg (PO) q8h  $\times$  7 days. For recurrent genital herpes, use acyclovir or valacyclovir (dose same as primary infection) for 7 days after each relapse. Recurrent episodes of HSV-2 are less painful than primary infection, and inguinal adenopathy is less prominent/painful.

**Prognosis:** HSV-2 tends to recur, especially during the first year. HSV-1 recurrences less frequent.

## Genital Ulcers

Subset	Usual Pathogens	IM Therapy	PO Therapy
Primary syphilis	<i>Treponema pallidum</i>	Benzathine penicillin 2.4 mu (IM) × 1 dose	Doxycycline 100 mg (PO) q12h × 2 weeks <b>or</b> Azithromycin 2 gm (PO) × 1 dose**
Chancroid	<i>Hemophilus ducreyi</i>	Ceftriaxone 250 mg (IM) × 1 dose <b>or</b> Any 3 <sup>rd</sup> generation cephalosporin 250–500 mg (IM) × 1 dose	Azithromycin 1 gm (PO) × 1 dose <b>or</b> Quinolone* (PO) × 3 days <b>or</b> Erythromycin base 500 mg (PO) q8h × 7 days

\* Ciprofloxacin 500 mg q12h or Levofloxacin 500 mg or Moxifloxacin 400 mg q24h.

\*\* Resistance increasing (careful followup essential).

### **Primary Syphilis (*Treponema pallidum*)**

**Clinical Presentation:** Painless, indurated ulcers (chancres) with bilateral painless inguinal adenopathy. Syphilitic chancres are elevated, clean and indurated, but not undermined.

**Diagnostic Considerations:** Diagnosis by spirochetes on darkfield examination of ulcer exudate. Elevated non-treponemal (VDRL/RPR) titers after 1 week.

**Pitfalls:** Non-treponemal (VDRL/RPR) titers fall slowly within 1 year; failure to decline suggests treatment failure/HIV. Even after effective treatment some patients remain VDRL/RPR positive for life (serofast).

**Therapeutic Considerations:** Parenteral penicillin is the preferred antibiotic for all stages of syphilis. If treatment fails and VDRL/RPR titers do not decline, obtain HIV serology.

**Prognosis:** Good with early treatment.

### **Chancroid (*Hemophilus ducreyi*)**

**Clinical Presentation:** Ragged, undermined, painful ulcer(s) + painful unilateral inguinal adenopathy.

**Diagnostic Considerations:** Diagnosis by streptobacilli in “school of fish” configuration on gram-stained smear of ulcer exudate/culture of *H. ducreyi*/NAAT.

**Pitfalls:** Co-infection is common; obtain Syphilis and HIV serologies.

**Therapeutic Considerations:** In HIV, multiple dose regimens or azithromycin is preferred. Resistance to erythromycin/ciprofloxacin has been reported.

**Prognosis:** Good with early treatment.

## Suppurating Inguinal Adenopathy

Subset	Pathogens	IV Therapy	PO Therapy
Lympho-granuloma venereum (LGV)	Chlamydia trachomatis (L <sub>1-3</sub> serotypes)	Not applicable	Doxycycline 100 mg (PO) q12h × 3 wks <b>or</b> Erythromycin base 500 mg (PO) q6h × 3 weeks
Granuloma inguinale (Donovanosis)	Klebsiella (Calymmatobacterium) granulomatis	Not applicable	Azithromycin 1 gm (PO) q week until cured. Doxycycline 100 mg (PO) q12h until cured <b>or</b> Erythromycin 500 mg (PO) q6h until cured <b>or</b> TMP-SMX 1 DS (PO) q12h until cured <b>or</b> Ciprofloxacin 750 mg (PO) q12h until cured

### Lymphogranuloma Venereum (Chlamydia trachomatis) LGV

**Clinical Presentation:** Unilateral inguinal adenopathy ± discharge/sinus tract.

**Diagnostic Consideration:** Diagnosis by very high Chlamydia trachomatis L<sub>1-3</sub> titers. Do not biopsy site (often does not heal and may form a fistula). May present as FUO.

**Pitfalls:** Initial papule not visible at clinical presentation. Biopsy shows granulomas; may be confused with perianal Crohn's disease.

**Therapeutic Considerations:** Rectal LGV may require additional courses of treatment.

**Prognosis:** Fibrotic perirectal/pelvic damage does not reverse with therapy.

### Granuloma Inguinale (Klebsiella [Calymmatobacterium] granulomatis)

#### Donovanosis

**Clinical Presentation:** Pseudolymphadenopathy with painless inguinal ulcers.

**Diagnostic Considerations:** Donovan bodies ("puffed-wheat" appearance) in tissue biopsy.

**Pitfalls:** No true inguinal adenopathy, as opposed to LGV infection.

**Therapeutic Considerations:** Doxycycline or azithromycin preferred. Continue therapy until lesions are healed.

**Prognosis:** Good if treated early.

## Genital/Perianal Warts (Condylomata acuminata)

Subset	Pathogens	Therapy
Genital/perianal warts	Human papilloma virus (HPV)	Podophyllin 10–25% in tincture of benzoin or podofilox or imiquimod (patient applies) <b>or</b> surgical/laser removal/cryotherapy with liquid nitrogen <b>or</b> cidofovir gel (1%) QHS × 5 days every other week for 6 cycles <b>or</b> trichloroacetic acid (TCA)/bichloroacetic acid (BCA) <b>or</b> intralesional interferon. Sinecatechins (15% ointment) q8h × 4 months

**Clinical Presentation:** Single/multiple verrucous genital lesions ± pigmentation, without inguinal adenopathy.

**Diagnostic Considerations:** Diagnosis by clinical appearance. Genital warts are usually caused by HPV types 6, 11. Anogenital warts caused by HPV types 16,18,31,33,35 and others are associated with cervical neoplasia. Females with anogenital warts need serial cervical PAP smears to detect cervical dysplasia/neoplasia.

**Pitfalls:** Most HPV infections are asymptomatic.

**Therapeutic Considerations:** Cidofovir cures/halts HPV progression in 50% of cases.

**Prognosis:** Related to HPV serotypes with malignant potential (HPV types 16,18,31,33,35). Preventative (not therapeutic) vaccines now available.

## Syphilis

Subset	Pathogen	IV/IM Therapy	PO Therapy
Primary, secondary, or early latent (duration < 1 year) syphilis	<i>Treponema pallidum</i>	Benzathine penicillin 2.4 mu (IM) × 1 dose	Doxycycline 100 mg (PO) q12h × 2 weeks <b>or</b> Azithromycin 2 gm (PO) × 1 dose*
Late latent (duration > 1 year) or tertiary syphilis	<i>Treponema pallidum</i>	Benzathine penicillin 2.4 mu (IM) weekly × 3 weeks	Doxycycline 100 mg (PO) q12h × 4 weeks
Neurosyphilis	<i>Treponema pallidum</i>	Penicillin G 3–4 mu (IV) q4h or continuous infusion × 10–14 days <b>or</b> <u>Alternate:</u> Procaine penicillin 2.4 mu (IM) q24h × 10–14 days plus probenecid 500 mg (PO) q6h × 10–14 days <b>or</b> Ceftriaxone 2 gm (IV) q24h × 10–14 days	Doxycycline 100 mg (PO) q12h × 4 weeks <b>or</b> Minocycline 100 mg (PO) q12h × 4 weeks

\* Resistance increasing (careful follow up essential).

Duration of therapy represents total time IV, IM, or PO. All stages of syphilis in HIV patients usually respond to therapeutic regimens recommended for normal hosts. Syphilis in pregnancy should be treated according to the stage of syphilis; penicillin-allergic pregnant patients should be desensitized and treated with penicillin.

### Primary Syphilis (*Treponema pallidum*)

**Clinical Presentation:** Painless, indurated ulcer(s) (chancere) with bilateral painless inguinal adenopathy.

**Diagnostic Considerations:** Diagnosis by spirochetes on darkfield or DFA examination of ulcer exudate. Reactive non-treponemal (VDRL/RPR) or treponemal (TPPA, others) test after 1 week. Patients with ↑ low titers of VDRL/RPR and – treponemal test may be BFP, e.g., SLE.

**Pitfalls:** VDRL/RPR titers fall slowly within 1 year; failure to decline suggests treatment failure.

**Therapeutic Considerations:** Parenteral penicillin is the preferred antibiotic for all stages of syphilis; if treatment fails and VDRL/RPR titers do not decline, obtain HIV serology. Some effectively treated patients remain VDRL/RPR positive in low titers eg 1:4 (serofast) for years.

**Prognosis:** Good with early treatment.

### **Secondary Syphilis (*Treponema pallidum*)**

**Clinical Presentation:** Facial/truncal macular, papular, papulosquamous, non-pruritic, non-tender, symmetrical rash which may involve the palms/soles. Usually accompanied by generalized adenopathy. Typically appears 4–10 weeks after primary chancre, although stages may overlap. Alopecia, condyloma lata, mucous patches, iritis/uveitis may be present. Renal involvement ranges from mild proteinuria to nephrotic syndrome. Without treatment, spontaneous resolution occurs after 3–12 weeks.

**Diagnostic Considerations:** Diagnosis by clinical findings and VDRL/RPR in high titers ( $\geq 1:64$ ). After treatment, RPR/VDRL titers usually  $\downarrow \times 4 < 6$  months.

**Pitfalls:** If only undiluted serum is tested, prozone phenomenon may render VDRL/RPR falsely negative.

**Therapeutic Considerations:** Parenteral penicillin is the preferred antibiotic for all stages of syphilis.

**Prognosis:** Excellent with early treatment.

### **Latent Syphilis (*Treponema pallidum*)**

**Clinical Presentation:** Patients are asymptomatic with elevated non-treponemal titers and reactive treponemal tests.

**Diagnostic Considerations:** Diagnosis by positive serology  $\pm$  prior history, but no signs/symptoms of syphilis. Asymptomatic syphilis  $< 1$  year in duration is termed “early” latent syphilis; asymptomatic syphilis  $> 1$  year/unknown duration is termed “late” latent syphilis. Secondary syphilis may relapse in up to 25% of patients with early latent syphilis, but relapse is rare in late latent syphilis. Evaluate patients for neurosyphilis.

**Pitfalls:** Treponemal tests (FTA-ABS, MHA-TP, TPPA, others) usually remain positive for life, even after adequate treatment.

**Therapeutic Considerations:** Parenteral penicillin is the preferred antibiotic for all stages of syphilis. Repeat VDRL/RPR titers at 6, 12, and 24 months; therapeutic response is defined as a 4-fold reduction in RPR/VDRL titers (2 tube dilutions).

**Prognosis:** Excellent even if treated late.

### **Tertiary Syphilis (*Treponema pallidum*)**

**Clinical Presentation:** May present with aortitis, neurosyphilis, iritis, or gummata 5–30 years after initial infection.

**Diagnostic Considerations:** Diagnosis by history of syphilis plus positive serological tests with signs/symptoms of late syphilis.

**Pitfalls:** Treat for signs of neurosyphilis on clinical exam or LP, even if VDRL/RPR are non-reactive.

**Therapeutic Considerations:** Parenteral penicillin is the preferred antibiotic for all stages of syphilis.

**Prognosis:** Related to extent of end-organ damage.

### **Neurosyphilis (*Treponema pallidum*)**

**Clinical Presentation:** Patients are often asymptomatic, but may have ophthalmic/auditory symptoms, cranial nerve abnormalities, tabes dorsalis, paresis, psychosis, or signs of meningitis/dementia.

**Diagnostic Considerations:** Diagnosis by elevated CSF VDRL titers; no need to obtain CSF FTA-ABS titers. Diagnosis confirmed if CSF has pleocytosis ( $> 5$  WBCs/hpf) or increased protein ( $> 50$  mg/dL), and positive VDRL.

**Pitfalls:** Persistent CSF abnormalities suggest treatment failure. CSF VDRL (60% sensitive) may be negative in neurosyphilis.

**Therapeutic Considerations:** Parenteral penicillin is the preferred antibiotic for all stages of syphilis. CSF abnormalities should decrease in 6 months and return to normal after 2 years; repeat lumbar puncture 6 months after treatment. Failure rate with ceftriaxone is 20%.

**Prognosis:** Related to extent of end-organ damage.

## Empiric Therapy of Bone and Joint Infections

### Septic Arthritis/Bursitis

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	PO Therapy or IV-to-PO Switch
Acute  (Treat initially based on gram stain of synovial fluid. If gram-positive cocci in clusters, treat initially for MRSA; if later identified as MSSA, treat accordingly)	S. aureus (MSSA)	Cefazolin 1 gm (IV) q8h x 3 weeks  <b>or</b> Ceftriaxone 1 gm (IV) q24h x 3 weeks  <b>or</b> Clindamycin 600 mg (IV) q8h x 3 weeks	Nafcillin 2 gm (IV) q4h x 3 weeks  <b>or</b> Meropenem 1 gm (IV) q8h x 3 weeks  <b>or</b> Ertapenem 1 gm (IV) q24h x 3 weeks	Cephalexin 1 gm (PO) q6h x 3 weeks  <b>or</b> Clindamycin 300 mg (PO) q8h x 3 weeks  <b>or</b> Quinolone* (PO) q24h x 3 weeks
	S. aureus (MRSA)	Linezolid 600 mg (IV) q12h x 3 weeks  <b>or</b> Quinupristin/dalfopristin 7.5 mg/kg (IV) q8h x 3 weeks  <b>or</b> Minocycline 100 mg (IV) q12h x 3 weeks		Linezolid 600 mg (PO) q12h x 3 weeks  <b>or</b> Minocycline 100 mg (PO) q12h x 3 weeks
	Group A,B,C,G streptococci	Ceftriaxone 1 gm (IV) q24h x 3 weeks  <b>or</b> Clindamycin 600 mg (IV) q8h x 3 weeks	Cefazolin 1 gm (IV) q8h x 3 weeks  <b>or</b> Quinolone* (IV) q24h x 3 weeks	Clindamycin 300 mg (PO) q8h x 3 weeks  <b>or</b> Cephalexin 500 mg (PO) q6h x 3 weeks <b>or</b> Quinolone* (PO) q24h x 3 weeks

\* Moxifloxacin 400 mg or Levofloxacin 500 mg.



## Septic Arthritis/Bursitis (cont'd)

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	PO Therapy or IV-to-PO Switch
Acute (cont'd)	Entero-bacteriaceae	Ceftriaxone 1 gm (IV) q24h x 3 weeks <b>or</b> Cefepime 2 gm (IV) q12h x 3 weeks <b>or</b> Cefotaxime 2 gm (IV) q6h x 3 weeks <b>or</b> Ceftizoxime 2 gm (IV) q8h x 3 weeks	Aztreonam 2 gm (IV) q8h x 3 weeks <b>or</b> Respiratory quinolone <sup>†</sup> (IV) x 3 weeks	Respiratory quinolone <sup>†</sup> (PO) x 3 weeks
	P. aeruginosa	Meropenem 1 gm (IV) q8h x 3 weeks <b>or</b> Ciprofloxacin 400 mg (IV) q8h x 3 weeks	Aztreonam 2 gm (IV) q8h x 3 weeks <b>or</b> Cefepime 2 gm (IV) q8h x 3 weeks	Ciprofloxacin 750 mg (PO) q12h x 3 weeks <b>or</b> Levofloxacin 750 mg (PO) q24h x 3 weeks
	N. gonorrhoea (PSNG/PPNG)	Ceftriaxone 1 gm (IV) q24h x 2 weeks <b>or</b> Ceftizoxime 2 gm (IV) q8h x 2 weeks	Levofloxacin 500 mg (IV) q24h x 2 weeks <b>or</b> Moxifloxacin 400 mg (IV) q24h x 2 weeks	Levofloxacin 500 mg (PO) q24h x 2 weeks <b>or</b> Moxifloxacin 400 mg (PO) q24h x 2 weeks
	Salmonella	Ceftriaxone 2 gm (IV) q24h x 2–3 weeks <b>or</b> Respiratory quinolone* (IV) x 2–3 weeks	Aztreonam 2 gm (IV) q8h x 2–3 weeks <b>or</b> TMP–SMX 2.5 mg/kg (IV) q6h x 2–3 weeks	Respiratory quinolone* (PO) x 2–3 weeks <b>or</b> TMP–SMX 1 DS tablet (PO) q12h x 2–3 weeks

‡ Moxifloxacin 400 mg (IV/PO) q24h or Levofloxacin 500 mg (IV/PO) q24h.

† Loading dose is not needed PO if given IV with the same drug.

\* Levofloxacin 500 mg (IV or PO) q24h or Moxifloxacin 400 mg (IV or PO) q24h.

**Septic Arthritis/Bursitis (cont'd)**

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	PO Therapy or IV-to-PO Switch
Secondary to animal bite wound	<i>Pasteurella multocida</i> <i>Streptobacillus moniliformis</i> <i>Eikenella corrodens</i>	Piperacillin/tazobactam 3.375 gm (IV) q6h x 2 weeks <b>or</b> Ampicillin/sulbactam 3 gm (IV) q6h x 2 weeks <b>or</b> Ticarcillin/clavulanate 3.1 gm (IV) q6h x 2 weeks	Meropenem 1 gm (IV) q8h x 2 weeks <b>or</b> Ertapenem 1 gm (IV) q24h x 2 weeks <b>or</b> Doxycycline 200 mg (IV) q12h x 3 days, then 100 mg (IV) q12h x 11 days	Amoxicillin/clavulanic acid 875/125 mg (PO) q12h x 2 weeks <b>or</b> Doxycycline 200 mg (PO) q12h x 3 days, then 100 mg (PO) q12h x 11 days <sup>†</sup> <b>or</b> Moxifloxacin 400 mg (PO) q24h x 2 weeks
Fungal arthritis	<i>Coccidioides immitis</i>	Not applicable	Itraconazole 200 mg (PO) solution q12h x 12 months or until cured* <b>or</b> Fluconazole 800 mg (PO) q24h until cured*	
	<i>Sporothrix schenckii</i>	Not applicable	Itraconazole 200 mg (PO) q12h until cured*	
TB arthritis	<i>M. tuberculosis</i>	Not applicable	Treat the same as for pulmonary TB (p. 53) except treat for 6–9 months	

MSSA/MRSA = methicillin-sensitive/resistant *S. aureus*. Duration of therapy represents total time IV, PO, or IV + PO. Most patients on IV therapy able to take PO meds should be switched to PO therapy after clinical improvement.

\* Itraconazole solution provides more reliable absorption than capsules.

**Acute Septic Arthritis/Bursitis**

**Clinical Presentation:** Acute joint pain with fever. Septic joint unable to bear weight. Septic bursitis presents with pain on joint motion, but patient is able to bear weight.

**Diagnostic Considerations:** Diagnosis by demonstrating organisms in synovial fluid by stain/culture. In septic bursitis (knee most common), there is pain on joint flexion (although the joint can bear weight), and synovial fluid findings are negative for septic arthritis. Except for *N. gonorrhoeae*, polyarthritis is not usually due to bacterial pathogens. Post-infectious polyarthritis is usually viral in origin, most commonly due to parvovirus B19, rubella, or HBV.

**Pitfalls:** Reactive arthritis may follow *C. jejuni*, *Salmonella*, *Shigella*, *Yersinia*, *N. gonorrhoeae*, *C. trachomatis*, or *C. difficile* infections. Synovial fluid cultures are negative. Reactive arthritis is usually asymmetrical and is monoarticular/oligoarticular.

**Therapeutic Considerations:** See specific pathogen, below. Treat septic bursitis as septic arthritis.

### **Staphylococcus aureus (MSSA/MRSA)**

**Diagnostic Considerations:** Painful hot joint; unable to bear weight. Diagnosis by synovial fluid pleocytosis and positive culture for joint pathogen. Examine synovial fluid to rule out gout (doubly birefringent crystals) and pseudogout (calcium pyrophosphate crystals). May occur in setting of endocarditis with septic emboli to joints; other manifestations of endocarditis are usually evident.

**Pitfalls:** Rule out causes of noninfectious arthritis (sarcoidosis, Whipple's disease, Ehlers-Danlos, etc.), which are less severe, but may mimic septic arthritis. In reactive arthritis following urethritis (*C. trachomatis*, *Ureaplasma urealyticum*, *N. gonorrhoeae*) or diarrhea (*Shigella*, *Campylobacter*, *Yersinia*, *Salmonella*), synovial fluid culture is negative, and synovial fluid WBCs counts are usually  $< 10,000/\text{mm}^3$  with normal synovial fluid lactic acid and glucose. Do not overlook infective endocarditis in mono/polyarticular MSSA/MRSA septic arthritis without apparent source.

**Therapeutic Considerations:** For MRSA septic arthritis, vancomycin penetration into synovial fluid is poor; use linezolid instead. Immobilization of infected joint during therapy is helpful. Local installation of antibiotics into synovial fluid has no advantage over IV/PO antibiotics.

**Prognosis:** Treat as early as possible to minimize joint damage. Repeated aspiration/open drainage may be needed to preserve joint function.

### **Group A, B, C, G Streptococci**

**Diagnostic Considerations:** Usually monoarticular. Not usually due to septic emboli from endocarditis.

**Prognosis:** Related to extent of joint damage and rapidity of antibiotic treatment.

### **Enterobacteriaceae**

**Diagnostic Considerations:** Diagnosis by isolation of gram-negative bacilli from synovial fluid.

**Pitfalls:** Septic arthritis involving an unusual joint (e.g., sternoclavicular, sacral) should suggest IV drug abuse until proven otherwise.

**Therapeutic Considerations:** Joint aspiration is essential in suspected septic arthritis of the hip and may be needed for other joints; obtain orthopedic surgery consult. Local installation of antibiotics into joint fluid is of no proven value.

**Prognosis:** Related to extent of joint damage and rapidity of antibiotic treatment.

### **Pseudomonas aeruginosa**

**Diagnostic Considerations** *P. aeruginosa* septic arthritis/osteomyelitis may occur after water contaminated puncture wound (e.g., nail puncture of heel through shoes). Sternoclavicular/sacroiliac joint involvement is common in IV drug abusers (IVDAs).

**Pitfalls:** Suspect IVDA in *P. aeruginosa* septic arthritis without a history of trauma.

**Therapeutic Considerations:** If ciprofloxacin is used, treat with 750 mg (not 500 mg) dose for *P. aeruginosa* septic arthritis/osteomyelitis.

**Prognosis:** Related to extent of joint damage and rapidity of antibiotic treatment.

**Neisseria gonorrhoeae**

**Diagnostic Considerations:** Gonococcal arthritis may present as a monoarticular arthritis, or multiple joints may be affected as part of gonococcal arthritis-dermatitis syndrome (disseminated gonococcal infection). Bacteremia with positive blood cultures occurs early during rash stage while synovial fluid cultures are negative. Joint involvement follows with typical findings of septic arthritis and synovial fluid cultures positive for *N. gonorrhoeae*; blood cultures are negative at this stage. Acute tenosynovitis is often a clue to gonococcal septic arthritis.

**Pitfalls:** Spectinomycin is ineffective against pharyngeal gonorrhea.

**Therapeutic Considerations:** Gonococcal arthritis-dermatitis syndrome is caused by very susceptible strains of *N. gonorrhoeae*. Cephalosporins also eliminate incubating syphilis.

**Prognosis:** Excellent with arthritis-dermatitis syndrome; worse with only monoarticular arthritis.

**Salmonella sp.**

**Diagnostic Considerations:** Occurs in sickle cell disease and hemoglobinopathies. Diagnosis by blood/joint cultures.

**Pitfalls:** *S. aureus*, not *Salmonella*, is the most common cause of septic arthritis in sickle cell disease.

**Prognosis:** Related to severity of infection and underlying health of host.

**Septic Arthritis Secondary to Animal Bite Wound**

**Clinical Presentation:** Penetrating bite wound into joint space.

**Diagnostic Considerations:** Diagnosis by smear/culture of synovial fluid/blood cultures.

**Pitfalls:** May develop metastatic infection from bacteremia.

**Therapeutic Considerations:** Treat for at least 2 weeks of combined IV/PO therapy.

**Prognosis:** Related to severity of infection and underlying health of host.

**Chronic Septic Arthritis**

**Clinical Presentation:** Subacute/chronic joint pain with decreased range of motion and little or no fever. Able to bear weight on joint.

**Diagnostic Considerations:** Diagnosis by smear/culture of synovial fluid/synovial biopsy.

**Coccidioides immitis**

**Diagnostic Considerations:** Must grow organisms from synovium/synovial fluid for diagnosis.

**Pitfalls:** Synovial fluid the same as in TB (lymphocytic pleocytosis, low glucose, increased protein).

**Therapeutic Considerations:** Oral therapy is preferred; same cure rates as amphotericin regimens. HIV patients need life-long suppressive therapy.

**Prognosis:** Related to severity of infection and underlying health of host.

**Sporothrix schenckii**

**Diagnostic Considerations:** Usually a monoarticular infection secondary to direct inoculation/trauma.

**Pitfalls:** Polyarticular arthritis suggests disseminated infection.

**Therapeutic Considerations:** SSKI is useful for lymphocutaneous sporotrichosis, not bone/joint involvement.

**Prognosis:** Excellent for localized disease (e.g., lymphocutaneous sporotrichosis). In disseminated disease, prognosis is related to host factors.

### ***Mycobacterium tuberculosis* (TB)**

**Diagnostic Considerations:** Clue is subacute/chronic tenosynovitis over involved joint. Unlike other forms of septic arthritis, which are usually due to hematogenous spread, TB arthritis may complicate adjacent TB osteomyelitis. Synovial fluid findings include lymphocytic pleocytosis, low glucose, and increased protein.

**Pitfalls:** Send synovial biopsy for AFB smear/culture in unexplained chronic monoarticular arthritis.

**Therapeutic Considerations:** TB arthritis is usually treated for 6–9 months.

**Prognosis:** Related to severity of infection and underlying health of host.

## **Lyme Disease\*/Lyme Arthritis**

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	PO Therapy or IV-to-PO Switch
Lyme disease	<i>Borrelia burgdorferi</i>	Ceftriaxone 1 gm (IV) q24h × 2 weeks  <b>or</b> Doxycycline 200 mg (IV) q12h × 3 days, then 100 mg (IV) q12h × 4 weeks	Ceftizoxime 2 gm (IV) q8h × 2 weeks	Doxycycline 200 mg (PO) q12h × 3 days, then 100 mg (PO) q12h × 11 days <sup>†</sup>  <b>or</b> Amoxicillin 1 gm (PO) q8h × 2 weeks
Lyme arthritis		Ceftriaxone 1 gm (IV) q24h × 4 weeks  <b>or</b> Doxycycline 200 mg (IV) q12h × 3 days, then 100 mg (IV) q12h × 4 weeks	Ceftizoxime 2 gm (IV) q8h × 4 weeks	Amoxicillin 1 gm (PO) q8h × 4 weeks  <b>or</b> Doxycycline 200 mg (PO) q12h × 3 days, then 100 mg (PO) q12h × 4 weeks

Duration of therapy represents total time IV, PO, or IV + PO. Most patients on IV therapy able to take PO meds should be switched to PO therapy after clinical improvement.

\* See p. 21 for Lyme neuroborreliosis and p. 78 for Lyme myocarditis.

† Doxycycline therapy × 10 days as effective as 2 weeks.

‡ For adult patients intolerant of Amoxicillin, Doxycycline, and Cefuroxime axetil, Azithromycin (500 mg orally per day for 7–10 days), Clarithromycin (500 mg orally twice per day for 14–21 days, if the patient is not pregnant), or Erythromycin (500 mg orally 4 times per day for 14–21 days) may be given. The recommended dosages of these agents for children are as follows: Azithromycin, 10 mg/kg per day (maximum of 500 mg per day); Clarithromycin, 7.5 mg/kg twice per day (maximum of 500 mg per dose); and Erythromycin, 12.5 mg/kg 4 times per day (maximum of 500 mg per dose).

## **Lyme Disease**

**Clinical Presentation:** Presents acutely with local or disseminated disease following bite of tick infected with *Borrelia*. Erythema migrans (annular lesion with central clearing) occurs in ~ 75% within 2 weeks of tick bite and may be associated with fever, headache, arthralgias/myalgias, meningismus. Other possible acute manifestations include aseptic meningitis, Bell's palsy, peripheral neuropathy, mild hepatitis, or heart block. Chronic disease may present with arthritis, peripheral neuropathy, or acrodermatitis chronica atrophicans (usually > 10 years after infection).

**Diagnostic Considerations:** Diagnosis by clinical presentation plus elevated IgM Lyme titers. If the IgM ELISA Lyme titer is borderline or suspected to be a false-positive, obtain an IgM Western blot to confirm the diagnosis. False negative early IgM titers are common. IgM cross-reactivity common with EBV, CMV. After ruling out EBV/CMV, retest for Lyme disease IgM and IgG by ELISA/immunoblot 4 weeks later (false + IgM titers revert to negative). IgM titers may take 4–6 weeks to increase after tick bite.

**Pitfalls:** Rash is not always seen, and tick bite often goes unnoticed (tick often spontaneously falls off after 1–2 days of feeding). Do not overlook Lyme disease in patients with sudden unexplained heart block in areas where Ixodes ticks are endemic.

**Therapeutic Considerations:** *B. burgdorferi* is highly susceptible to all beta-lactams. For Bell's palsy or neuroborreliosis, minocycline (100 mg PO q12h x 2 weeks) may be preferred to doxycycline. Highest failure rates with macrolide therapy. Symptoms may persist for 1 year or more after adequate therapy. No rationale to re-treat persistent symptoms.

**Prognosis:** Excellent in normal hosts.

## **Lyme Arthritis**

**Clinical Presentation:** Acute Lyme arthritis presents with joint pain, decreased range of motion, ability to bear weight on joint, and little or no fever. Chronic Lyme arthritis resembles rheumatoid arthritis.

**Diagnostic Considerations:** Usually affects children and large weight-bearing joints (e.g., knee). Acute Lyme arthritis is diagnosed by clinical presentation plus elevated IgM Lyme titer. Chronic Lyme arthritis is suggested by rheumatoid arthritis-like presentation with negative ANA and rheumatoid factor, and positive IgG Lyme titer and synovial fluid PCR.

**Pitfalls:** Acute Lyme arthritis joint is red but not hot, in contrast to septic arthritis. In chronic Lyme arthritis, a negative IgG Lyme titer essentially rules out chronic Lyme arthritis, but an elevated IgG Lyme titer indicates only past exposure to *B. burgdorferi* and is not diagnostic of Lyme arthritis. Joint fluid in chronic Lyme disease is usually negative by culture, but positive by PCR; synovial fluid PCR, however, does not differentiate active from prior infection.

**Therapeutic Considerations:** IgG Lyme titers remain elevated for life, and do not decrease with treatment. Joint symptoms often persist for months/years after effective antibiotic therapy due to autoimmune joint inflammation; treat with anti-inflammatory drugs, not repeat antibiotic courses. Oral therapy as effective as IV therapy.

**Prognosis:** Good in normal hosts. Chronic/refractory arthritis may develop in genetically predisposed patients with DRW 2/4 HLA types.

## Infected Joint Prosthesis

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	IV-to-PO Switch
Staphylococcal  (Treat initially for MSSA; if later identified as MRSA/CoNS, treat accordingly)	S. epidermidis (CoNS)	Linezolid 600 mg (IV) q12h* <b>or</b> Vancomycin 1 gm (IV) q12h*	Cefotaxime 2 gm (IV) q6h** <b>or</b> Ceftizoxime 2 gm (IV) q8h**	Linezolid 600 mg (PO) q12h*
	S. aureus (MSSA)	Nafcillin 2 gm (IV) q4h* <b>or</b> Cefazolin 1 gm (IV) q8h* <b>or</b> Ceftriaxone 1 gm (IV) q24h†	Meropenem 1 gm (IV) q8h* <b>or</b> Clindamycin 600 mg (IV) q8h*	Clindamycin 300 mg (PO) q8h* <b>or</b> Linezolid 600 mg (PO) q12h* <b>or</b> Cephalexin 1 gm (PO) q6h*
	S. aureus (MRSA)	Linezolid 600 mg (IV) q12h* <b>or</b> Vancomycin 1 gm (IV) q12h* <b>or</b> Minocycline 100 mg (IV) q12h* <b>or</b> Quinupristin/dalfopristin 7.5 mg/kg (IV) q8h*		Linezolid 600 mg (PO) q12h* <b>or</b> Minocycline 100 mg (PO) q12h*

MRSA/MSSA = methicillin-resistant/sensitive *S. aureus*; MSSE/MRSE = methicillin-sensitive/resistant *S. epidermidis*. Duration of therapy represents total time IV or IV + PO. Most patients on IV therapy able to take PO meds should be switched to PO therapy after clinical improvement.

\* Treat for 1 week after joint prosthesis is replaced.

† Only if MSSE strain susceptible.

**Clinical Presentation:** Pain in area of prosthesis with joint loosening/instability ± low-grade fevers.

**Diagnostic Considerations:** Infected prosthesis is suggested by prosthetic loosening/lucent areas adjacent to prosthesis on plain films ± positive blood cultures. Diagnosis confirmed by bone scan. Use joint aspiration to identify organism.

**Pitfalls:** An elevated ESR with prosthetic loosening suggests prosthetic joint infection. Mechanical loosening without infection is common many years after joint replacement, but ESR is normal.

**Therapeutic Considerations:** Infected prosthetic joints usually must be removed for cure. Replacement prosthesis may be inserted anytime after infected prosthesis is removed. To prevent infection of new joint prosthesis, extensive debridement of old infected material is important. *If replacement of*

*infected joint prosthesis is not possible, chronic suppressive therapy* may be used with oral antibiotics e.g., minocycline 100 mg (PO) q12h. TMP-SMX 5 mg/kg (PO) q6h may be successful in long-term suppression/cure in total hip replacement (treat × 6 months) or total knee replacement (treat × 9 months) due to susceptible strains of MSSA/MSSE; adding rifampin 300 mg (PO) q12h may be helpful.

**Prognosis:** Related to adequate debridement of infected material when prosthetic joint is removed.

## Osteomyelitis

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	PO Therapy or IV-to-PO Switch
Acute  (Treat initially for MSSA; if later identified as MRSA or Enterobacteriaceae, treat accordingly)	S. aureus (MSSA)	Cefazolin 1 gm (IV) q8h × 4–6 weeks <b>or</b> Ceftriaxone 1 gm (IV) q24h × 4–6 weeks <b>or</b> Meropenem 1 gm (IV) q8h × 4–6 weeks	Cefotaxime 2 gm (IV) q6h × 4–6 weeks <b>or</b> Ceftizoxime 2 gm (IV) q8h × 4–6 weeks	Clindamycin 300 mg (PO) q8h × 4–6 weeks <b>or</b> Cephalexin 1 gm (PO) q6h × 4–6 weeks <b>or</b> Respiratory quinolone <sup>†</sup> (PO) q24h × 4–6 weeks
	S. aureus (MRSA)	Linezolid 600 mg (IV) q12h × 4–6 weeks <b>or</b> Minocycline 100 mg (IV) q12h × 4–6 weeks <b>or</b> Vancomycin 2 gm (IV) q12h × 4–6 weeks <b>or</b> Quinupristin/dalfopristin 7.5 mg/kg (IV) q8h × 4–6 weeks		Linezolid 600 mg (PO) q12h × 4–6 weeks <b>or</b> Minocycline 100 mg (PO) q12h × 4–6 weeks
	Enterobacteriaceae	Ceftriaxone 1 gm (IV) q24h × 4–6 wks <b>or</b> Quinolone <sup>†</sup> (IV) × 4–6 weeks <b>or</b> Tigecycline 100 mg (IV) × 1 dose, then 50 mg (IV) q12h × 4–6 weeks	Cefotaxime 2 gm (IV) q6h × 4–6 wks <b>or</b> Ceftizoxime 2 gm (IV) q8h × 4–6 wks	Quinolone <sup>†</sup> (PO) × 4–6 weeks



## Osteomyelitis (cont'd)

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	PO Therapy or IV-to-PO Switch
Chronic <i>Diabetes mellitus/Peripheral vascular disease</i>	Group A, B streptococci S. aureus (MSSA) E. coli P. mirabilis K. pneumoniae B. fragilis	Tigecycline 200 mg (IV) × 1 dose, then 100 mg (IV) q24h × 2–4 weeks <b>or</b> Meropenem 1 gm (IV) q8h* <b>or</b> Piperacillin/tazobactam 3.375 gm (IV) q6h* <b>or</b> Ertapenem 1 gm (IV) q24h*	Moxifloxacin 400 mg (IV) q24h* <b>or</b> Ceftizoxime 2 gm (IV) q8h* <b>or</b> Ampicillin/sulbactam 3 gm (IV) q6h* <b>or</b> <b>combination therapy with</b> Ceftriaxone 1 gm (IV) q24h* <b>plus</b> Metronidazole 1 gm (IV) q24h*	Clindamycin 300 mg (PO) q8h* <b>plus</b> Levofloxacin 500 mg (PO) q24h* <b>or monotherapy with</b> Moxifloxacin 400 mg (PO) q24h*
	Above pathogens/ S. aureus (MRSA)	Same as above <b>plus either</b> Minocycline 100 mg (IV) q12h × 2–4 weeks <b>or</b> Linezolid 600 mg (IV) q12h × 4–6 weeks <b>or</b> <b>monotherapy with</b> Tigecycline 200 mg (IV) × 1 dose, then 100 mg (IV) q24h × 2–4 weeks	Minocycline 100 mg (IV) q12h × 2–4 weeks <b>plus either</b> Levofloxacin 500 mg (IV) q24h <b>or</b> Ceftriaxone 1 gm (IV) q24h × 2–4 weeks	Minocycline 100 mg (PO) q12h × 2–4 weeks <b>plus</b> Quinolone <sup>†</sup> (PO) q24h × 2–4 weeks
	Above pathogens/ B. fragilis (foul discharge)	Tigecycline 200 mg (IV) × 1 dose, then 100 mg (IV) q24h × 2–4 weeks	Moxifloxacin 400 mg (IV) q24h × 2–4 weeks <b>or</b> Ertapenem 1 gm (IV) q24h × 2–4 weeks	Moxifloxacin 400 mg (PO) q24h × 2–4 weeks

## Osteomyelitis (cont'd)

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	PO Therapy or IV-to-PO Switch
Chronic (cont'd)	Brucella	Streptomycin 1 gm (IM) q24h until cured <b>plus</b> Doxycycline 200 mg (IV) q12h × 3 days, then 100 mg (IV) q12h until cured	Gentamicin 5 mg/kg (IV) q24h until cured <b>plus</b> Doxycycline 200 mg (IV) q12h × 3 days, then 100 mg (IV) q12h until cured	Doxycycline 200 mg (PO) q12h × 3 days, then 100 mg (PO) q12h until cured† <b>plus</b> Rifampin 600 mg (PO) q24h until cured
TB osteomyelitis	M. tuberculosis	Treat the same as pulmonary TB (p. 53), but extend treatment to 6–9 months		

MSSA/MRSA = methicillin-sensitive/resistant *S. aureus*. Duration of therapy represents total time IV, PO, or IV + PO. Most patients on IV therapy able to take PO meds should be switched to PO therapy soon after clinical improvement.

\* Treat for 1 week after adequate debridement or amputation.

† Moxifloxacin 400 mg or Levofloxacin 500 mg.

### Acute Osteomyelitis

**Clinical Presentation:** Tenderness over infected bone. Fever and positive blood cultures common.

**Diagnostic Considerations:** Diagnosis by elevated ESR with positive bone scan. Bone biopsy is not needed for diagnosis. Bone scan is positive for acute osteomyelitis in first 24 hours.

**Pitfalls:** Earliest sign on plain films is soft tissue swelling; bony changes evident after 2 weeks.

**Therapeutic Considerations:** Treat 4–6 weeks with antibiotics. Debridement is not necessary for cure.

**Prognosis:** Related to adequacy/promptness of treatment.

### Chronic Osteomyelitis

#### Diabetes Mellitus

**Clinical Presentation:** Afebrile or low-grade fever with normal WBC counts and deep penetrating ulcers ± draining sinus tracts.

**Diagnostic Considerations:** Diagnosis by elevated ESR and bone changes on plain films. Bone scan is not needed for diagnosis. Bone biopsy is preferred method of demonstrating organisms, since blood cultures are usually negative and cultures from ulcers/sinus tracts are unreliable.

**Pitfalls:** *P. aeruginosa* is a common colonizer and frequently cultured from deep ulcers/sinus tracts, but is not a pathogen in chronic osteomyelitis in diabetics.

**Therapeutic Considerations:** Surgical debridement is needed for cure; antibiotics alone are ineffective. Revascularization procedures usually do not help, since diabetes is a microvascular disease. Do not culture penetrating foot ulcers/draining sinus tracts; culture results reflect superficial flora. Bone biopsy during debridement is the best way to identify pathogen; if biopsy not possible, treat empirically.

**Prognosis:** Related to adequacy of blood supply/surgical debridement.

**Non-Diabetics with Peripheral Vascular Disease (PVD)**

**Clinical Presentation:** Absent or low-grade fever with normal WBC counts ± wet/dry digital gangrene.

**Diagnostic Considerations:** Diagnosis by clinical appearance of dusky/cold foot ± wet/dry gangrene. Chronic osteomyelitis secondary to PVD/open fracture is often polymicrobial.

**Pitfalls:** Wet gangrene usually requires surgical debridement/antibiotic therapy; dry gangrene may not.

**Therapeutic Considerations:** Surgical debridement needed for cure. Antibiotics alone are ineffective. Revascularization procedure may help treat infection by improving local blood supply.

**Prognosis:** Related to degree of vascular compromise.

**Brucella Osteomyelitis (*Brucella sp.*)**

**Diagnostic Considerations:** May be evidence of brucellosis elsewhere (epididymo-orchitis). Suspect in patients with a history of animal/raw milk/cheese exposure. Serologic diagnosis by serum SAT, ELISA, or PCR.

**Pitfalls:** *Brucella* has predilection for vertebra, sacroiliac joints. Erosions of anterior vertebra adjacent to disc space typical vs. TB with diffuse vertebral destruction.

**Therapeutic Considerations:** May require 6 weeks of antibiotic therapy.

**Prognosis:** Joint destruction usually permanent.

**TB Osteomyelitis (*Mycobacterium tuberculosis*)**

**Clinical Presentation:** Presents similarly to chronic bacterial osteomyelitis. Vertebral TB (Pott's disease) affects disk spaces early and presents with chronic back/neck pain ± inguinal/paraspinal mass.

**Diagnostic Considerations:** Diagnosis by AFB on biopsy/culture of infected bone. T-spot PPD-positive.

**Pitfalls:** Chest x-ray is normal in 50%. May be confused with cancer, brucella osteomyelitis or chronic bacterial osteomyelitis.

**Therapeutic Considerations:** Treated the same as TB arthritis.

**Prognosis:** Good for non-vertebral TB/vertebral (if treated before paraparesis/paraplegia).

## Empiric Therapy of Skin and Soft Tissue Infections

**Cellulitis, Erysipelas, Mastitis (uSSSIs)**

Subset	Usual Pathogens	Preferred IV Therapy	PO Therapy or IV-to-PO Switch
Above-the-waist	Group A, B streptococci	Ceftriaxone 1–2 gm (IV) q24h × 2 weeks  <b>or</b> Cefazolin 1 gm (IV) q8h × 2 weeks	Cephalexin 500 mg (PO) q6h × 2 weeks  <b>or</b> Clindamycin 300 mg (PO) q6h × 2 weeks  <b>or</b> Quinolone* (PO) q24h × 2 weeks
		Quinolone* (IV) of q24h × 2 weeks  <b>or</b> Clindamycin 600 mg (IV) q8h × 2 weeks	

**Cellulitis, Erysipelas, Mastitis (uSSSIs) (cont'd)**

Subset	Usual Pathogens	Preferred IV Therapy	PO Therapy or IV-to-PO Switch
Below-the-waist  (Treat initially for MSSA; if later identified as MRSA, treat accordingly)	Group A, B streptococci P. mirabilis K. pneumoniae E. coli	Cefazolin 1 gm (IV) q8h × 2 weeks  <b>or</b> Ceftriaxone 1 gm (IV) q24h × 2 weeks	Quinolone* (IV) q24h × 2 weeks  <b>or</b> Cephalexin 500 mg (PO) q6h × 2 weeks  Quinolone* (PO) q24h × 2 weeks

uSSSIs = uncomplicated skin skin structure infections.

MRSA/MSSA = methicillin-resistant/sensitive *S. aureus*. Duration of therapy represents total time IV, PO, or IV + PO. Most patients on IV therapy able to take PO meds should be switched to PO therapy soon after clinical improvement.

\* Moxifloxacin 400 mg or Levofloxacin 500 mg.

**Clinical Presentation:** Cellulitis presents as warm, painful, flat, non-pruritic skin erythema without discharge. Erysipelas resembles cellulitis but is raised and sharply demarcated. Mastitis presents as cellulitis of the breast cellulitis caused by groups A, B, C or G streptococci. Aspiration of streptococcal cellulitis is serous/serosanguinous fluid not purulent (as with MSSA/MRSA abscesses). MSSA/MRSA causes cutaneous abscesses, not cellulitis. MSSA/MRSA abscesses raised, warm, and tender ± bullae. No regional adenopathy or lymphangitis.

**Diagnostic Considerations:** Diagnosis by clinical appearance ± culture of pathogen from aspirated skin lesion(s). Group B streptococci are important pathogens in diabetics. Lower extremity cellulitis tends to recur. Chronic lymphedema or edema of an extremity predisposes to recurrent/persistent cellulitis.

**Pitfalls:** Streptococcal cellulitis often accompanied by lymphangitis and regional adenopathy; fever/chills common.

**Therapeutic Considerations:** Lower extremity cellulitis requires ~ 1 week of antibiotics to improve. Patients with peripheral vascular disease, chronic venous stasis, alcoholic cirrhosis, and diabetes take 1–2 weeks longer to improve and often require 3–4 weeks of treatment. Treat mastitis as cellulitis above-the-waist, and drain surgically if an abscess is present.

**Prognosis:** Related to degree of micro (DM) and or macrovascular (PVD) insufficiency.

## Complicated Skin/Skin Structure Infections (cSSSIs)

Subset	Usual Pathogens	Preferred IV Therapy	IV-to-PO Switch	
Mixed aerobic-anaerobic deep soft tissue infection	Enterobacteriaceae Group A streptococci <i>S. aureus</i> (MSSA) Anaerobic streptococci <i>Fusobacterium</i>	Piperacillin/tazobactam 3.375 gm (IV) q6h x 2 weeks <b>or</b> Meropenem 1 gm (IV) q8h x 2 weeks	Ertapenem 1 gm (IV) q24h x 2 weeks <b>or</b> Doripenem 1 gm (IV) q8h <b>or</b> Moxifloxacin 400 mg (IV) q24h x 2 weeks	Moxifloxacin 400 mg (PO) q24h x 2 weeks <b>or combination therapy with</b> Levofloxacin 500 mg (PO) q24h x 2 weeks <b>plus</b> Clindamycin 300 mg (PO) q8h x 2 weeks
Clostridial myonecrosis (gas gangrene)	<i>Clostridium</i> sp.	<u>Preferred IV</u> Penicillin G 10 mu (IV) q4h x 2 weeks <b>or</b> Clindamycin 600 mg (IV) q8h x 2 weeks	<u>Alternate IV</u> Piperacillin 4 gm (IV) q8h x 2 weeks <b>or</b> Ertapenem 1 gm (IV) q24h x 2 weeks	Not applicable
Necrotizing fasciitis/synergistic gangrene/Fournier's gangrene	Group A streptococci Enterobacteriaceae Anaerobic streptococci <i>S. aureus</i> (MSSA)	Piperacillin/tazobactam 3.375 gm (IV) q6h x 2 weeks <b>or</b> Ertapenem 1 gm (IV) q24h x 2 weeks	Clindamycin 600 mg (IV) q8h x 2 weeks <b>plus</b> Levofloxacin 500 mg (IV) q24h x 2 weeks	Clindamycin 300 mg (PO) q8h x 2 weeks <b>plus</b> Levofloxacin 500 mg (PO) q24h x 2 weeks
Phegmon, abscesses	MRSA/MSSA	Tigecycline 200 mg (IV) x 1 dose, then 100 mg (IV) q24h x 2 weeks <b>or</b> Daptomycin 6 mg/kg (IV) q24h x 2 weeks <b>or</b> Linezolid 600 mg (IV) q12h x 2 weeks <b>or</b> Tedizolid 200 mg (IV) q24h x 6 days, then 200 mg (PO) q24h x 6 days <b>or</b> Telavancin 10 mg/kg (IV) q24h x 2 weeks <b>or</b> Dalbavancin 1 gm (IV) x 1 dose, then 500 mg (IV) 7 days later	Linezolid 600 mg (PO) q12h x 2 weeks <b>or</b> Minocycline 100 mg (PO) q12h x 2 weeks	

cSSSIs = complicated skin skin structure infections.

CA-MRSA = community-acquired methicillin-resistant *S. aureus* (see p. 15), MSSA = methicillin-sensitive *S. aureus*. Duration of therapy represents total time IV or IV + PO. Most patients on IV therapy able to take PO meds should be switched to PO therapy after clinical improvement.

**Complicated Skin/Skin Structure Tissue Infections (cSSIs) (cont'd)**

Subset	Usual Pathogens	Preferred IV Therapy	IV-to-PO Switch
Pyomyositis/necrotizing abscesses	Community-acquired MRSA (CA-MRSA) <sup>§</sup>	Daptomycin 6 mg/kg (IV) q24h × 2 weeks <b>or</b> Linezolid 600 mg (IV) q12h × 2 weeks	Linezolid 600 mg (PO) q12h × 2 weeks <b>or</b> Minocycline 100 mg (PO) q12h × 2 weeks

CA-MRSA = community-acquired methicillin-resistant *S. aureus* (see p. 15), MSSA = methicillin-sensitive *S. aureus*.

§ CA-MRSA PVL-positive strains (see p. 15).

**Mixed Aerobic/Anaerobic Deep Soft Tissue Infection**

**Clinical Presentation:** Local pain/tenderness ± gross gas deep in soft tissues and usually high fevers. More common in diabetics.

**Diagnostic Considerations:** Diagnosed clinically. Bacteriologic diagnosis by gram stain/culture of aspirated fluid. Patients usually have high fevers. Wound discharge is foul when present.

**Pitfalls:** Gross creptance/prominent gas in soft tissues on x-ray suggests a mixed aerobic/anaerobic necrotizing infection, not gas gangrene. Gas gangrene extremely rare in diabetics.

**Therapeutic Considerations:** Prompt empiric therapy and surgical debridement may be lifesaving.

**Prognosis:** Related to severity of infection, adequacy of debridement, and underlying health of host.

**Gas Gangrene (Clostridial Myonecrosis)**

**Clinical Presentation:** Fulminant infection of muscle with little or n fever. Infected area is extremely painful, indurated, and discolored with or without bullae.

**Diagnostic Considerations:** Diagnosis is clinical. Aspiration of infected muscle shows few PMNs and gram-positive bacilli without spores (*C. perfringens* only). Gas gangrene is not accompanied by high fever. Patients are often apprehensive with relative bradycardia ± diarrhea. Wound discharge, if present, is sweetish and not foul. Rapidly progressive hemolytic anemia is characteristic.

**Pitfalls:** Gas gangrene (clostridial myonecrosis) has little visible gas on plain film x-rays; abundant gas should suggest a mixed aerobic/anaerobic infection, not gas gangrene. Gas gangrene extremely rare in diabetics.

**Therapeutic Considerations:** Surgical debridement is life saving and the only way to control infection.

**Prognosis:** Related to speed/extent of surgical debridement. Progression/death may occur in hours.

**Necrotizing Fasciitis/Synergistic Gangrene**

**Clinical Presentation:** Acutely ill patient with high fevers and extreme local pain without gas in tissues. If scrotum involved (± abdominal wall involvement), the diagnosis is Fournier's gangrene.

**Diagnostic Considerations:** Diagnosis by CT/MRI of involved extremity showing infection confined to one or more muscle compartments/fascial planes. Patients are febrile and ill. Gas is not present on exam or x-rays. May be polymicrobial or due to a single organism. Foul smelling exudate from infected soft tissues indicates anaerobes are present.

**Pitfalls:** Extreme pain in patients with deep soft tissue infections should suggest a compartment syndrome/necrotizing fasciitis. No hemolytic anemia, diarrhea, or bullae as with gas gangrene.

**Therapeutic Considerations:** Control/cure of infection requires surgical, debridement of dead tissue, and antimicrobial therapy. Clindamycin may be added for its anti-exotoxin effects if pathogen group A streptococci.

**Prognosis:** Related to rapidity/extent of surgical debridement.

### **Phegmon, Abscesses**

**Clinical Presentation:** MSSA/MRSA infection often accompanied by bullae. A phegmon is a pre-abscess before abscess wall formation. MSSA/MRSA abscesses are clinically indistinguishable from other abscesses, but remain the most common pathogens in cSSSI abscesses.

**Diagnostic Considerations:** Diagnosis of cellulitis and abscesses is clinical. Demonstration of fasciitis, phegmons, of abscesses are by imaging studies by CT/MRI.

**Pitfalls:** Bullae with cellulitis should suggest MSSA/MRSA (not group A streptococci). Bullae in DM/bullous diseases are not accompanied by fever/cellulitis.

**Therapeutic Considerations:** Cellulitis and phegmons may be treated with antibiotic therapy alone. Fasciitis and abscesses usually require debridement (fasciitis) or drainage (abscesses).

**Prognosis:** Good if treated appropriately/early.

### **Pyomyositis/Necrotizing Abscesses (CA-MRSA)**

**Clinical Presentation:** Abrupt onset of severe/deep muscle infection ± large abscesses should suggest community-acquired MRSA (CA-MRSA).

**Diagnostic Considerations:** The diagnosis of CA-MRSA is made on the basis of the distinctively fulminant/severe clinical presentation and by culturing MRSA from muscle/abscess. If available, test isolate for SCC *mec IV* ± Panton-Valentine leukocidin (PVL) gene.

**Pitfalls:** CA-MRSA susceptible to clindamycin, TMP-SMX, and doxycycline.

**Therapeutic Considerations:** Prompt/complete incision and drainage of abscesses and early use of anti-CA-MRSA drugs may be life-saving. Antibiotics effective against CA-MRSA (TMP-SMX, clindamycin, doxycycline) are ineffective against CO-MRSA/HA-MRSA; antibiotics effective against CO-MRSA/HA-MRSA are also effective against CA-MRSA (see p. 15). Use minocycline instead of doxycycline for CA-MRSA/CO-MRSA. Doxycycline resistance with CA-MRSA of concern. Minocycline more effective than doxycycline, TMP-SMX or clindamycin for CA-MRSA.

**Prognosis:** Related to presence of PVL gene CA-MRSA. CA-MRSA PVL negative infections are similar in severity to MSSA infections.

## Skin Ulcers

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	PO Therapy or IV-to-PO Switch
Decubitus ulcers (above the waist)	Group A streptococci E. coli, P. mirabilis K. pneumoniae S. aureus (MSSA)	Cefazolin 1 gm (IV) q8h* <b>or</b> Ceftriaxone 1 gm (IV) q24h*	Cefotaxime 2 gm (IV) q6h* <b>or</b> Ceftizoxime 2 gm (IV) q8h*	Cephalexin 500 mg (PO) q6h*
(sacral ulcers/ ulcers below the waist) stage III/ IV ulcers = <b>chronic osteomyelitis</b>	B. fragilis Group A streptococci E. coli P. mirabilis K. pneumoniae S. aureus (MSSA) S. aureus (MRSA) <sup>†</sup> (not <i>P. aeruginosa</i> ) <sup>§</sup>	Tigecycline 100 mg (IV) × 1 dose, then 50 mg (IV) q12h* <b>or</b> Ertapenem 1 gm (IV) q24h* <b>or</b> Piperacillin/tazobactam 3.375 gm (IV) q6h* <b>or</b> Meropenem 1 gm (IV) q8h*	Moxifloxacin 400 mg (IV) q24h* <b>or</b> <b>combination therapy with</b> Clindamycin 600 mg (IV) q6h* <b>plus</b> Levofloxacin 500 mg (IV) q24h*	Moxifloxacin 400 mg (PO) q24h* <b>or</b> <b>combination therapy with</b> Clindamycin 300 mg (PO) q8h* <b>plus</b> Levofloxacin 500 mg (PO) q24h*

Duration of therapy represents total time IV, PO, or IV + PO. Most patients on IV therapy able to take PO meds should be switched to PO therapy soon after clinical improvement.

- \* Treat Stages I/II (superficial) decubitus ulcers with local care. If no underlying chronic osteomyelitis, treat Stages III/IV (deep) decubitus ulcers with antibiotics and adequate bone debridement.
- † MRSA: Tigecycline 100 mg (IV) × 1 dose then 50 mg (IV) q12h\* alone or one of above non-MRSA antibiotics plus [Daptomycin 4 mg/kg (IV) q24h\* or Linezolid 600 mg (IV) q12h\*] or Vancomycin 1 gm (IV) q12h\* or Minocycline 100 mg (IV) q12h\*. PO therapy or IV-to-PO switch if MRSA: Linezolid 600 mg (PO) q12h\* or Minocycline 100 mg (PO) q12h\* plus one of above antibiotics.
- § Deep penetrating ulcers (stage III/IV) in diabetics represent underlying chronic osteomyelitis and are not due to *P. aeruginosa*. However, *P. aeruginosa* (a water associated organism) can be cultured from > 90% of deep diabetic foot ulcers and represents superficial colonization (from moist socks, moist dressings, whirlpool baths, etc.)



## Skin Ulcers (cont'd)

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	PO Therapy or IV-to-PO Switch
Diabetic foot ulcers (deep/complicated)  (Treat initially for MSSA; if later identified as Group A streptococci, MRSA, etc., treat accordingly)	S. aureus (MRSA)	Tigecycline 200 mg (IV) × 1 dose, then 100 mg (IV) q24h <sup>†</sup> <b>or</b> Vancomycin 1 gm (IV) q12h <sup>†</sup> <b>or</b> Daptomycin 4 mg/kg (IV) q24h <sup>†</sup> <b>or</b> Linezolid 600 mg (IV) q12h <sup>†</sup> <b>or</b> Minocycline 100 mg (IV) q12h <sup>†</sup>	Telavancin 10 mg/kg (IV) q24h <sup>†</sup> <b>or</b> Tedizolid 200 mg (IV) q24h × 6 days, then 200 mg (PO) × 6 days <b>or</b> Dalbavancin 1 gm (IV) × 1 dose, then 500 mg (IV) 7 days later	Linezolid 600 mg (PO) q12h <sup>†</sup> <b>or</b> Minocycline 100 mg (PO) q12h <sup>†</sup>
	S. aureus (MSSA) Group A, B streptococci E. coli P. mirabilis K. pneumoniae B. fragilis	Tigecycline 200 mg (IV) × 1 dose, then 100 mg (IV) q24h <sup>†</sup> <b>or</b> Ertapenem 1 gm (IV) q24h <sup>†</sup> <b>or</b> Piperacillin/tazobactam 3.375 gm (IV) q6h <sup>†</sup> <b>or</b> Moxifloxacin 400 mg (IV) q24h <sup>†</sup>	Cefoperazone 2 gm (IV) q12h <sup>†</sup> <b>or</b> Doripenem 1 gm (IV) q8h <sup>†</sup> <b>or</b> Ampicillin/sulbactam 3 gm (IV) q6h <sup>†</sup> <b>or combination therapy with</b> Ceftriaxone 1 gm (IV) q24h <sup>†</sup> <b>or</b> Levofloxacin 500 mg (IV) q24h <sup>†</sup> <b>plus either</b> Metronidazole 1 gm (IV) q24h <sup>†</sup> <b>or</b> Clindamycin 600 mg (IV) q8h <sup>†</sup>	Moxifloxacin 400 mg (PO) q24h <sup>†</sup> <b>or</b> <b>combination therapy with</b> Minocycline 100 mg (PO) q12h <sup>†</sup> <b>plus</b> Levofloxacin 500 mg (PO) q24h <sup>†</sup>

† Treat × 1 week after adequate debridement/amputation.

## Skin Ulcers (cont'd)

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	PO Therapy or IV-to-PO Switch
Ischemic foot ulcers	<i>S. aureus</i> (MRSA)	Treat the same as for deep/complicated diabetic foot ulcers, above		
(Treat initially for MSSA; if later identified as Gp. A strep, MRSA, etc., treat accordingly)	Group A streptococci <i>E. coli</i> <i>S. aureus</i> (MSSA)	Cefazolin 1 gm (IV) q8h × 2 weeks <b>or</b> Ceftriaxone 1 gm (IV) q24h × 2 weeks	Quinolone* (IV) q24h × 2 weeks	Quinolone* (PO) q24h × 2 weeks
	Any pathogen(s) above <b>plus</b> <i>S. aureus</i> (MRSA)	IV Therapy: MRSA drug <b>plus</b> a non-MRSA drug (see diabetic foot ulcers) PO Therapy or IV-to-PO Switch: MRSA drug <b>plus</b> a non-MRSA drug (see diabetic foot ulcers)		

MSSA = methicillin-sensitive *S. aureus*. Duration of therapy represents total time IV, PO, or IV + PO. Most patients on IV therapy able to take PO meds should be switched to PO therapy soon after clinical improvement.

\* Levofloxacin 750 mg or Moxifloxacin 400 mg.

### Decubitus Ulcers

**Clinical Presentation:** Painless ulcers with variable depth and infectious exudate ± fevers ≤ 102°F.

**Diagnostic Considerations:** Diagnosis by clinical appearance. Obtain ESR/bone scan to rule out underlying osteomyelitis with deep (Stage III/IV) decubitus ulcers.

**Pitfalls:** Superficial decubitus ulcers do not require systemic antibiotics.

**Therapeutic Considerations:** Deep decubitus ulcers require antibiotics and debridement, superficial ulcers do not. Coverage for *B. fragilis* is needed for deep perianal decubitus ulcers. Good nursing care is important in preventing/limiting extension of decubitus ulcers.

**Prognosis:** Related to fecal contamination of ulcer and bone involvement (e.g., osteomyelitis).

### Diabetic Foot Ulcers/Chronic Osteomyelitis

**Clinical Presentation:** Ulcers/sinus tracts on bottom of foot/between toes; usually painless. Fevers ≤ 102°F and a foul smelling exudate are common.

**Diagnostic Considerations:** In diabetics, deep, penetrating, chronic foot ulcers/draining sinus tracts are diagnostic of chronic osteomyelitis. ESR ≥ 100 mm/hr in a diabetic with a foot ulcer/sinus tract is diagnostic of chronic osteomyelitis. Foot films confirm chronic osteomyelitis. Bone scan is needed only in acute osteomyelitis.

**Pitfalls:** Do not rely on culture results of deep ulcers/sinus tracts to choose antibiotic coverage, since cultures reflect skin colonization, not bone pathogens. Treat empirically.

**Therapeutic Considerations:** *B. fragilis* coverage is required in deep penetrating diabetic foot ulcers/fetid foot infection. *P. aeruginosa* is often cultured from diabetic foot ulcers/sinus tracts, but represents colonization, not infection. *P. aeruginosa* is a “water” organism that colonizes feet from moist socks/dressings, irrigant solutions, or whirlpool baths. Surgical debridement is essential for cure of chronic osteomyelitis in diabetics. Treat superficial diabetic foot ulcers the same as cellulitis in non-diabetics (see p. 132).

**Prognosis:** Related to adequacy of debridement of infected bone.

### Ischemic Foot Ulcers

**Clinical Presentation:** Ulcers often clean/dry  $\pm$  digital gangrene. No fevers/exudate.

**Diagnostic Considerations:** Diagnosis by clinical appearance/location in a patient with peripheral vascular disease. Ischemic foot ulcers most commonly affect the toes, medial malleoli, dorsum of foot, or lower leg.

**Pitfalls:** In contrast to ulcers in diabetics, ischemic ulcers due to peripheral vascular disease usually do not involve the plantar surface of the foot.

**Therapeutic Considerations:** Dry gangrene should not be treated with antibiotics unless accompanied by signs of systemic infection. Wet gangrene should be treated as a mixed aerobic/anaerobic infection. Both dry/wet gangrene may require debridement for cure/control. Do not rely on ulcer cultures to guide treatment; treat empirically if necessary. Evaluate for revascularization.

**Prognosis:** Related to degree of vascular insufficiency.

### Skin Abscesses/Infected Cysts (Skin Pustules, Skin Boils, Furunculosis) (uSSSIs/cSSSIs)

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	PO Therapy or IV-to-PO Switch
<b>cSSSIs</b> (Deep/multiple skin abscesses)  (Treat initially for MSSA; if later identified as MRSA, treat accordingly)	S. aureus (MRSA)	Tigecycline 200 mg (IV) $\times$ 1 dose, then 100 mg (IV) q24h $\times$ 2 weeks <b>or</b> Linezolid 600 mg (IV) q12h $\times$ 2 weeks <b>or</b> Daptomycin 4 mg/kg (IV) q24h $\times$ 2 weeks <b>or</b> Minocycline 100 mg (IV) q12h $\times$ 2 weeks <b>or</b> Ceftaroline 600 mg (IV) of q12h $\times$ 2 weeks <b>or</b> Vancomycin 1 mg (IV) q12h $\times$ 2 weeks <b>or</b> Tedizolid 200 mg (IV) q24h $\times$ 6 days, then 200 mg (PO) $\times$ 6 days <b>or</b> Dalbavancin 1 gm (IV) $\times$ 1 dose, then 500 mg (IV) 7 days later <b>or</b> Telavancin 10 mg/kg (IV) q24h $\times$ 2 weeks <b>or</b> Oritavancin 1200 mg (IV) $\times$ 1 dose	Linezolid 600 mg (PO) q12h $\times$ 2 weeks <b>or</b> Minocycline 100 mg (PO) q12h $\times$ 2 weeks	
	S. aureus (MSSA)	Cefazolin 1 gm (IV) q8h $\times$ 2 weeks <b>or</b> Nafcillin 2 gm (IV) q4h $\times$ 2 weeks	Clindamycin 600 mg (IV) q8h $\times$ 2 weeks <b>or</b> Ceftriaxone 1 gm (IV) q24h $\times$ 2 weeks	Cephalexin 500 mg (PO) q6h $\times$ 2 weeks <b>or</b> Clindamycin 300 mg (PO) q8h $\times$ 2 weeks

uSSSIs = uncomplicated skin skin structure infections; cSSSIs = complicated skin skin structure infections.

MRSA/MSSA = methicillin-resistant/sensitive *S. aureus*.

## Skin Abscesses/Infected Cysts (Skin Pustules, Skin Boils, Furunculosis) (cSSSIs/uSSSIs) (cont'd)

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	PO Therapy or IV-to-PO Switch
<b>uSSSIs</b> <i>Infected pilonidal cysts</i>	Group A streptococci E. coli P. mirabilis K. pneumoniae S. aureus (MSSA)	Cefazolin 1 gm (IV) q8h × 2 weeks <b>or</b> Ceftriaxone 1 gm (IV) q24h × 2 weeks	Levofloxacin 500 mg (IV) q24h × 2 weeks <b>or</b> Moxifloxacin 400 mg (IV) q24h × 2 weeks <b>or</b> Ceftizoxime 2 gm (IV) q8h × 2 weeks	Levofloxacin 500 mg (PO) q24h × 2 weeks <b>or</b> Moxifloxacin 400 mg (PO) q24h × 2 weeks
<i>Hydradenitis suppurativa</i>	S. aureus (MSSA)	Not applicable	TMP-SMX 1 SS tablet (PO) q12h × 2–4 weeks <b>or</b> Clindamycin 300 mg (PO) q8h × 2–4 weeks <b>or</b> Minocycline 100 mg (PO) q12h × 2–4 weeks	
	S. aureus (MRSA)	Not applicable	Minocycline 100 mg (PO) q12h × 2–4 weeks	

uSSSIs = uncomplicated SSSIs.

cSSSIs = complicated SSSIs.

MRSA/MSSA = methicillin-resistant/sensitive *S. aureus*. Duration of therapy represents total time IV, PO, or IV + PO. Most patients on IV therapy able to take PO meds should be switched to PO therapy soon after clinical improvement.

### Skin Abscesses

**Clinical Presentation:** Warm painful nodules ± bullae, low-grade fever ± systemic symptoms, no lymphangitis. Skin boils/furunculosis present as acute, chronic, or recurrent skin pustules, and remain localized without lymphangitis.

**Diagnostic Considerations:** Specific pathogen diagnosed by gram stain of abscess. Recurring *S. aureus* abscesses are not uncommon and should be drained. Blood cultures are rarely positive. Suspect Job's syndrome if recurring abscesses with peripheral eosinophilia. Skin boils/furunculosis are diagnosed by clinical appearance (skin pustules).

**Pitfalls:** Recurring *S. aureus* skin infections may occur on immune basis but immunologic studies are usually negative.

**Therapeutic Considerations:** Repeated aspiration of abscesses may be necessary. Surgical drainage is required if antibiotics fail. Treat boils/furunculosis as in hydradenitis suppurativa.

**Prognosis:** Excellent if treated early.

### Infected Pilonidal Cysts

**Clinical Presentation:** Chronic drainage from pilonidal cysts.

**Diagnostic Considerations:** Diagnosis by clinical appearance. Deep/systemic infection is rare.

**Pitfalls:** Culture of exudate is usually unhelpful.

**Therapeutic Considerations:** Surgical debridement is often necessary.

**Prognosis:** Good with adequate excision.

### Hydradenitis Suppurativa

**Clinical Presentation:** Chronic, indurated, painful, raised axillary/groin lesions ± drainage/sinus tracts.

**Diagnostic Considerations:** Diagnosis by clinical appearance/location of lesions. Infections are often bilateral and tend to recur.

**Pitfalls:** Surgical debridement is usually not necessary unless deep/extensive infection.

**Therapeutic Considerations:** Most anti-*S. aureus* antibiotics have poor penetration and usually fail.

**Prognosis:** Good with recommended antibiotics. Surgery, if necessary, is curative.

## Skin Vesicles (Non-Genital)

Subset	Pathogen	Therapy
Herpes simplex	Herpes simplex virus (HSV)	Acyclovir 400 mg (PO) q8h × 10 days <b>or</b> Valacyclovir 1 gm (PO) q12h × 7–10 days <b>or</b> Famciclovir 250 mg (PO) q8h × 10 days
Chickenpox	Varicella zoster virus (VZV)	Acyclovir 800 mg (PO) q6h × 5 days <b>or</b> Valacyclovir 1 gm (PO) q8h × 5 days <b>or</b> Famciclovir 500 mg (PO) q8h × 5 days
Herpes zoster <i>Dermatomal zoster (shingles)</i>  <i>Disseminated zoster</i>	Varicella zoster virus (VZV)	Acyclovir 800 mg (PO) 5x/day × 7–10 days <b>or</b> Valacyclovir 1 gm (PO) q8h × 7–10 days <b>or</b> Famciclovir 500 mg (PO) q8h × 7–10 days  <u>IV therapy:</u> Acyclovir 10 mg/kg (IV) q8h × 7–10 days <u>PO therapy:</u> Valacyclovir 1 gm (PO) q8h × 7–10 days <b>or</b> Famciclovir 500 mg (PO) q8h × 7–10 days
Herpes whitlow	HSV-1	Acyclovir 400 mg (PO) q8h × 7–10 days <b>or</b> Valacyclovir 1 gm (PO) q12h × 7–10 days <b>or</b> Famciclovir 250 mg (PO) q8h × 7–10 days

### Herpes Simplex (HSV)

**Clinical Presentation:** Painful, sometimes pruritic vesicles that form pustules or painful erythematous ulcers. Associated with fever, myalgias.

**Diagnostic Considerations:** Diagnosis by clinical appearance and demonstration of HSV by culture of vesicle fluid/vesicle base. May be severe in HIV.

**Pitfalls:** Painful vesicular lesions surrounded by prominent induration distinguishes HSV from insect bites (pruritic) and cellulitis (no induration).

**Therapeutic Considerations:** Topical acyclovir ointment may be useful early when vesicles erupt, but is ineffective after vesicles stop erupting. For severe/refractory cases, use acyclovir 5 mg/kg (IV) q8h × 2–7 days, then if improvement, switch to acyclovir 400 mg (PO) q8h to complete 10-day course.

**Prognosis:** Related to extent of tissue involvement/degree of cellular immunity dysfunction.

### **Chickenpox (VZV)**

**Clinical Presentation:** Abrupt appearance of discrete/diffuse pruritic vesicles. Appear in successive crops over 3 days, then no more new lesions. Patients do not appear toxic.

**Diagnostic Considerations:** Chickenpox lesions are central and typically concentrated on the trunk, although vesicles may also occur in the mouth, GI or GU tract. Vesicles are seen at different stages of development, and are superficial with the classic “dew drop on a rose petal” appearance. Tzanck test is positive in chickenpox (negative in smallpox).

**Pitfalls:** Vesicles are not deep/umbilicated like smallpox. Smallpox patients are sick/toxic, and vesicles are at the same stage of development in each anatomical area. Vesicles begin and are concentrated on the face with smallpox.

**Therapeutic Considerations:** Begin therapy as early as possible before appearance of successive crops of vesicles appear. Treat VZV pneumonia early with acyclovir.

**Prognosis:** Children do better than adults. Worst prognosis in smokers/pregnancy (may develop chickenpox/VZV pneumonia).

### **Herpes Zoster (VZV)**

**Clinical Presentation:** Painful, vesicular eruption in dermatomal distribution. Pain may be difficult to diagnose. Risk of disseminated VZV with steroids or statins.

**Diagnostic Considerations:** Diagnosis by appearance/positive Tzanck test of vesicle base scrapings.

**Pitfalls:** Begin therapy within 2 days of vesicle eruption.

**Therapeutic Considerations:** Higher doses of acyclovir are required for VZV than HSV. See p. 343 for disseminated VZV, ophthalmic nerve/visceral involvement, or acyclovir-resistant strains.

**Prognosis:** Good if treated early. Some develop painful post-herpetic neuralgia of involved dermatomes.

### **Herpes Whitlow**

**Clinical Presentation:** Multiple vesicopustular lesions on fingers and hands. Lymphangitis, adenopathy, fever/chills are usually present, suggesting a bacterial infection.

**Diagnostic Considerations:** Common in healthcare workers giving patients oral care/suctioning.

**Pitfalls:** No need for antibiotics even though lesions appear infected with “pus.”

**Therapeutic Considerations:** Never incise/drain herpes whitlow; surgical incision will flare/prolong the infection.

**Prognosis:** Excellent, unless incision/drainage has been performed.

**Wound Infections** (for rabies, see pp. 362, 374–375)

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	PO Therapy or IV-to-PO Switch
Animal bite wounds <sup>†</sup>	Group A streptococci <i>P. multocida</i> Capnocytophaga canimorsus (DF2) <i>S. aureus</i> (MSSA)	Ampicillin/sulbactam 3 gm (IV) q6h × 2 weeks <b>or</b> Tigecycline 200 mg (IV) × 1 dose, then 100 mg (IV) q24h × 2 weeks	Ertapenem 1 gm (IV) q24h × 2 weeks <b>or</b> Piperacillin/tazobactam 3.375 gm (IV) q6h × 2 weeks	Amoxicillin/clavulanic acid 500/125 mg (PO) q8h or 875/125 mg (PO) q12h × 2 weeks <b>or</b> Doxycycline 200 mg (PO) q12h × 3 days, then 100 mg (PO) q12h × 11 days
Human bite wounds	Oral anaerobes Group A streptococci <i>E. corrodens</i> <i>S. aureus</i> (MSSA)	Same as for animal bite wounds, above	Same as for animal bite wounds, above	Same as for animal bite wounds, above
Cat scratch disease (CSD)	<i>Bartonella henselae</i> (invasive disease)	Doxycycline 200 mg (IV) q12h × 3 days, then 100 mg (IV) q12h × 4–8 weeks <b>or</b> Azithromycin 500 mg (IV) q24h × 4–8 weeks	Chloramphenicol 500 mg (IV) q6h × 4–8 weeks <b>or</b> Erythromycin 500 mg (IV) q6h × 4–8 weeks	Doxycycline 200 mg (PO) q12h × 3 days, then 100 mg (PO) q12h × 4–8 weeks* <b>or</b> Azithromycin 250 mg (PO) q24h × 4–8 weeks <b>or</b> Quinolone <sup>‡</sup> (PO) × 4–8 weeks
	<i>B. henselae</i> (lymphadenopathy only)	<u>PO therapy:</u> Azithromycin 500 mg (PO) × 1 dose, then 250 mg (PO) × 4 days		

MSSA = methicillin-sensitive *S. aureus*. Duration of therapy represents total time IV, PO, or IV + PO.

\* Loading dose is not needed PO if given IV with the same drug.

‡ Ciprofloxacin 400 mg (IV) or 500 mg (PO) q12h or Levofloxacin 500 mg (IV or PO) q24h or Moxifloxacin 400 mg (IV or PO) q24h.

† Rabies (see p. 362)

**Wound Infections (cont'd) (for rabies, see pp. 362, 374–375)**

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	PO Therapy or IV-to-PO Switch
Burn wounds (severe) <sup>†</sup>	Group A streptococci <i>S. aureus</i> (MSSA) <i>Enterobacter</i> <i>P. aeruginosa</i>	Doripenem 1 gm (IV) q8h × 2 weeks <b>or</b> Meropenem 1 gm (IV) q8h × 2 weeks	Cefepime 2 gm (IV) q8h × 2 weeks <b>or</b> Cefoperazone 2 gm (IV) q12h × 2 weeks	Not applicable
Freshwater-exposed wounds	<i>Aeromonas hydrophilia</i>	Quinolone <sup>‡</sup> (IV) × 2 weeks <b>or</b> TMP-SMX 2.5 mg/kg (IV) q6h × 2 weeks	Ceftriaxone 1 gm (IV) q24h × 2 weeks <b>or</b> Aztreonam 2 gm (IV) q8h × 2 weeks <b>or</b> Gentamicin 240 mg (IV) q24h × 2 weeks	Quinolone <sup>‡</sup> (PO) × 2 weeks <b>or</b> TMP-SMX 1 SS tablet (PO) q12h × 2 weeks
Saltwater-exposed wounds	<i>Vibrio vulnificus</i> <i>Vibrio</i> sp.	Doxycycline 200 mg (IV) q12h × 3 days, then 100 mg (IV) q12h × 11 days <b>or</b> Quinolone <sup>‡</sup> (IV) × 2 weeks	Ceftriaxone 2 gm (IV) q12h × 2 weeks <b>or</b> Chloramphenicol 500 mg (IV) q6h × 2 weeks	Doxycycline 200 mg (PO) q12h × 3 days, then 100 mg (PO) q12h × 11 days* <b>or</b> Quinolone <sup>‡</sup> (PO) × 2 weeks

MSSA = methicillin-sensitive *S. aureus*. Duration of therapy represents total time IV, PO, or IV + PO. Most patients on IV therapy able to take PO meds should be switched to PO therapy soon after clinical improvement.

† Treat only IV or IV-to-PO switch.

‡ Ciprofloxacin 400 mg (IV) or 500 mg (PO) q12h or Levofloxacin 500 mg (IV or PO) q24h or Moxifloxacin 400 mg (IV or PO) q24h.

**Animal Bite Wounds (for rabies, see p. 362)**

**Clinical Presentation:** Cellulitis surrounding bite wound.

**Diagnostic Considerations:** Diagnosis by culture of bite wound exudate. Deep bites may also cause tendinitis/osteomyelitis, and severe bites may result in systemic infection with bacteremia.

**Pitfalls:** Avoid erythromycin in penicillin-allergic patients (ineffective against *P. multocida*).

**Therapeutic Considerations:** For facial/hand bites, consult a plastic surgeon.

**Prognosis:** Related to adequacy of debridement and early antibiotic therapy.

**Human Bite Wounds**

**Clinical Presentation:** Cellulitis surrounding bite wound.



**Diagnostic Considerations:** Diagnosis by culture of bite wound exudate. Infection often extends to involve tendon/bone.

**Pitfalls:** Compared to animal bites, human bites are more likely to contain anaerobes, *S. aureus*, and Group A streptococci.

**Therapeutic Considerations:** Avoid primary closure of human bite wounds.

**Prognosis:** Related to adequacy of debridement and early antibiotic therapy.

### **Cat Scratch Disease (*Bartonella henselae*)**

**Clinical Presentation:** Subacute presentation of obscure febrile illness associated with cat bite/contact. Usually accompanied by adenopathy.

**Diagnostic Considerations:** Diagnosis by wound culture/serology for *Bartonella*. Cat scratch fever/disease may follow a cat scratch, but a lick from a kitten contaminating an inapparent micro-laceration is more common. May present as an FUO. Culture of exudate/node is unlikely to be positive, but silver stain of biopsy material shows organisms.

**Pitfalls:** Rule out lymphoma, which may present in similar fashion. Titers may cross react with *C. burnetii* (Q fever).

**Therapeutic Considerations:** For invasive disease, treat until symptoms/signs resolve. For lymphadenopathy, oral azithromycin decreases the size of nodes but may not reduce fever/systemic symptoms. *Bartonella* are sensitive in vitro to cephalosporins and TMP-SMX, but these antibiotics are ineffective in vivo.

**Prognosis:** Related to health of host.

### **Burn Wounds**

**Clinical Presentation:** Severe (3rd/4th degree) burns  $\pm$  drainage.

**Diagnostic Considerations:** Semi-quantitative bacterial counts help differentiate colonization (low counts) from infection (high counts). Burn wounds quickly become colonized.

**Pitfalls:** Treat only infected 3rd/4th degree burn wounds with systemic antibiotics.

**Therapeutic Considerations:** Meticulous local care/eschar removal/surgical debridement is key in preventing and controlling infection.

**Prognosis:** Related to severity of burns and adequacy of eschar debridement.

### **Freshwater-Exposed Wounds (*Aeromonas hydrophilia*)**

**Clinical Presentation:** Fulminant wound infection with fever and diarrhea.

**Diagnostic Considerations:** Diagnosis by stool/wound/blood culture.

**Pitfalls:** Suspect *A. hydrophilia* in wound infection with fresh water exposure followed by diarrhea.

**Therapeutic Considerations:** Surgical debridement of devitalized tissue may be necessary.

**Prognosis:** Related to severity of infection and health of host.

### **Saltwater-Exposed Wounds (*Vibrio vulnificus/Vibrio sp.*)**

**Clinical Presentation:** Fulminant wound infection with fever, painful hemorrhagic bullae, diarrhea.

**Diagnostic Considerations:** Diagnosis by stool/wound/blood culture. *Vibrio vulnificus* is a fulminant, life-threatening infection that may be accompanied by hypotension.

**Pitfalls:** Suspect *V. vulnificus* in acutely ill patients with fever, diarrhea, and bullous lesions after salt-water exposure.

**Therapeutic Considerations:** Surgical debridement of devitalized tissue may be necessary.

**Prognosis:** Related to extent of infection and health of host.

## Superficial Fungal Infections of Skin and Nails

Subset	Usual Pathogens	Topical Therapy	PO Therapy
Mucocutaneous (local/non-disseminated) candidiasis	<i>C. albicans</i>	Clotrimazole 1% cream twice daily × 2 weeks	Fluconazole 400 mg (PO) × 1 dose, then 200 mg (PO) q24h × 2 weeks
Tinea corporis (body ringworm)	<i>Trichophyton rubrum</i> <i>Epidermophyton floccosum</i> <i>Microsporum canis</i> <i>Trichophyton mentagrophytes</i>	Clotrimazole 1% cream twice daily × 4–8 weeks <b>or</b> Miconazole 2% cream twice daily × 2 weeks <b>or</b> Econazole 1% cream twice daily × 2 weeks	Terbinafine 250 mg (PO) q24h × 4 weeks <b>or</b> Ketoconazole 200 mg (PO) q24h × 4 weeks <b>or</b> Fluconazole 200 mg (PO) weekly × 4 weeks
Tinea capitis (scalp ringworm)	Same as Tinea corporis, above	Selenium sulfide shampoo daily × 2–4 weeks	Same as Tinea corporis (see above)
Tinea cruris (jock itch)	<i>T. cruris</i>	Same as Tinea corporis, above	Same as Tinea corporis (see above)
Tinea pedis (athlete's foot)	Same as Tinea corporis, above	Same as Tinea corporis, above	Terbinafine 250 mg (PO) q24h × 2 weeks <b>or</b> Ketoconazole 200 mg (PO) q24h × 4 weeks <b>or</b> Itraconazole 200 mg (PO) q24h × 4 weeks*
Tinea versicolor (pityriasis)	<i>Malassezia furfur</i> ( <i>Pityrosporum orbiculare</i> )	Clotrimazole cream (1%) or miconazole cream (2%) or ketoconazole cream (2%) daily × 7 days	Ketoconazole 200 mg (PO) q24h × 7 days <b>or</b> Itraconazole 200 mg (PO) q24h × 7 days* <b>or</b> Fluconazole 400 mg (PO) × 1 dose

\* Itraconazole solution provides more reliable absorption than Itraconazole capsules.

## Superficial Fungal Infections of Skin and Nails (cont'd)

Subset	Usual Pathogens	Topical Therapy	PO Therapy
Onychomycosis (nail infection)	Epidermophyton floccosum Trichophyton mentagrophytes Trichophyton rubrum C. albicans	Not applicable	Fluconazole 200 mg (PO) q24h <i>pulse dosed, i.e., 1 week per month</i> × 3 months (fingernail infection) or 6 months (toenail infection)  <b>or</b> Itraconazole 200 mg (PO) q24h* <i>pulse dosed, i.e., 1 week per month</i> × 2 months (fingernail infection) or 3 months (toenail infection)  <b>or</b> Terbinafine 250 mg (PO) q24h × 6 weeks (fingernail infection) or 12 weeks (toenail infection)

\* Itraconazole solution provides more reliable absorption than itraconazole capsules.

### **Mucocutaneous (Local/Non-disseminated) Candidiasis**

**Clinical Presentation:** Primary cutaneous findings include an erythematous rash with satellite lesions, which may be papular, pustular, or ulcerated. Lesions can be limited or widespread over parts of body. Chronic mucocutaneous candidiasis manifests as recurrent candidal infections of skin, nails, or mucous membranes.

**Diagnostic Considerations:** Diagnosis by demonstrating organism by stain/culture in tissue specimens. In HIV, *Candida* is very common on skin/mucous membranes.

**Pitfalls:** Do not confuse with the isolated, multinodular lesions of disseminated disease, which may resemble ecthyma gangrenosa or purpura fulminans.

**Therapeutic Considerations:** Diabetics and other compromised hosts may require prolonged therapy. In contrast to local disease, nodular cutaneous candidiasis represents disseminated disease (p. 290).

**Prognosis:** Related to extent of disease/host defense status.

### **Tinea Corporis (Body Ringworm)**

**Clinical Presentation:** Annular pruritic lesions on trunk/face with central clearing.

**Diagnostic Considerations:** Diagnosis by clinical appearance/skin scraping.

**Pitfalls:** Do not confuse with erythema migrans, which is not pruritic.

**Therapeutic Considerations:** If topical therapy fails, treat with oral antifungals.

**Prognosis:** Excellent.

### **Tinea Capitis (Scalp Ringworm)**

**Clinical Presentation:** Itchy, annular scalp lesions.

**Diagnostic Considerations:** Scalp lesions fluoresce with ultraviolet light. Culture hair shafts.

**Pitfalls:** *T. capitis* is associated with localized areas of alopecia.

**Therapeutic Considerations:** Selenium sulfide shampoo may be used first for 2–4 weeks. Treat shampoo failures with oral ketoconazole, terbinafine, or fluconazole.

**Prognosis:** Excellent.

### **Tinea Cruris (Jock Itch)**

**Clinical Presentation:** Groin, inguinal, perineal, or buttock lesions that are pruritic and serpiginous with scaling borders/central clearing.

**Diagnostic Considerations:** Diagnosis by clinical appearance/culture.

**Pitfalls:** Usually spares penis/scrotum, unlike Candida.

**Therapeutic Considerations:** In addition to therapy, it is important to keep area dry.

**Prognosis:** Excellent.

### **Tinea Pedis (Athlete's Foot)**

**Clinical Presentation:** Painful cracks/fissures between toes.

**Diagnostic Considerations:** Diagnosis by clinical appearance/skin scraping.

**Pitfalls:** Must keep feet dry or relapse/reinfection may occur.

**Therapeutic Considerations:** In addition to therapy, it is important to keep area dry.

**Prognosis:** Excellent.

### **Tinea Versicolor (Pityriasis)**

**Clinical Presentation:** Oval hypo- or hyperpigmented scaly lesions that coalesce into large confluent areas typically on upper trunk; chronic/relapsing.

**Diagnostic Considerations:** Diagnosis by clinical appearance and culture of affected skin lesions.

**Pitfalls:** *M. furfur* also causes seborrheic dermatitis, but seborrheic lesions are typically on the face/scalp.

**Therapeutic Considerations:** If topical therapy fails, treat with oral antifungals. Treat non-scalp seborrheic dermatitis with ketoconazole cream (2%) daily until cured.

**Prognosis:** Excellent.

### **Dermatophyte Nail Infections**

**Clinical Presentation:** Chronically thickened, discolored nails.

**Diagnostic Considerations:** Diagnosis by culture of nail clippings.

**Pitfalls:** Nail clipping cultures often contaminated by bacterial/fungal colonizers. Green nail discoloration suggests *P. aeruginosa*, not a fungal nail infection; treat with ciprofloxacin 500 mg (PO) q12h x 2–3 weeks.

**Therapeutic Considerations:** Lengthy therapy is required. However, terbinafine and itraconazole remain bound to nail tissue for months following dosing and thus therapy with these compounds is not continued until clearance of the nail bed.

**Prognosis:** Excellent if infection is totally eradicated from nail bed. New nail growth takes months.

## Skin Infestations

Subset	Usual Pathogens	Therapy
Scabies	<i>Sarcoptes scabiei</i>	Treat whole body with Permethrin cream 5% (Elimite); leave on for 8–10 hours <b>or</b> Ivermectin 18 mg (three 6-mg pills) (PO) × 1 dose
Head lice	<i>Pediculus humanus var. capitis</i>	Shampoo with Permethrin 5% (Elimite) or 1% (NIX) cream × 10 minutes
Body lice	<i>Pediculus humanus var. corporis</i>	Body lice removed by shower. Removed clothes should be washed in hot water or sealed in bags for 1 month, or treated with DDT powder 10% or malathion powder 1%
Pubic lice (crabs)	<i>Phthirus pubis</i>	Permethrin 5% (Elimite) or 1% (NIX) cream × 10 minutes to affected areas

### **Scabies (*Sarcoptes scabiei*)**

**Clinical Presentation:** Punctate, serpiginous, intensely pruritic black spots in webbed spaces of hands/feet and creases of elbows/knees.

**Diagnostic Considerations:** Diagnosis by visualization of skin tracts/burrows. Incubation period up to 6 weeks after contact. Spread by scratching from one part of body to another.

**Pitfalls:** Mites are not visible, only their skin tracks, but mites may be scraped out of tracts for diagnosis.

**Therapeutic Considerations:** Permethrin cream is usually effective. If itching persists after treatment, do not retreat (itching is secondary to hypersensitivity reaction of eggs in skin burrows). Treat close contacts. Vacuum bedding/furniture.

**Prognosis:** Norwegian scabies is very difficult to eradicate.

### **Head Lice (*Pediculus humanus var. capitis*)**

**Clinical Presentation:** White spots may be seen on hair shafts of head/neck, but not eyebrows.

**Diagnostic Considerations:** Nits on hair are unhatched lice eggs, seen as white dots attached to hair shaft. May survive away from body × 2 days.

**Pitfalls:** May need to retreat in 7 days to kill any lice that hatched from surviving nits.

**Therapeutic Considerations:** Shampoo with Permethrin 5% (Elimite) or 1% (NIX) cream kills lice/nits. Clothes and non-washables should be tied off in plastic bags × 2 weeks to kill lice. Alternately, wash and dry clothes/bed linens; heat from dryer/iron kills lice.

**Prognosis:** Related to thoroughness of therapy.

### **Body Lice (*Pediculus humanus var. corporis*)**

**Clinical Presentation:** Intense generalized pruritus.

**Diagnostic Considerations:** Smaller than head lice and more difficult to see. May survive away from body × 1 week.

**Pitfalls:** Body lice live in clothes; leave only for a blood meal, then return to clothing.

**Therapeutic Considerations:** Can survive in seams of clothing × 1 week.

**Prognosis:** Good if clothes are also treated.

**Pubic Lice (*Phthirus pubis*) Crabs****Clinical Presentation:** Genital pruritus.**Diagnostic Considerations:** Seen on groin, eyelashes, axilla. May survive away from body  $\times$  1 day.**Pitfalls:** Smaller than head lice, but easily seen.**Therapeutic Considerations:** Treat partners. Wash, dry, and iron clothes; heat from dryer/iron kills lice. Non-washables may be placed in a sealed bag  $\times$  7 days.**Prognosis:** Good if clothes are also treated.**Ischiorectal/Perirectal Abscess**

Subset	Pathogens	Preferred Therapy
Ischiorectal/ perirectal abscess	Enterobacteriaceae B. fragilis	Treat the same as mild/severe peritonitis (p. 100) $\pm$ surgical drainage depending on abscess size/severity

**Clinical Presentation:** Presents in normal hosts with perirectal pain, pain on defecation, leukocytosis, erythema/tenderness over abscess  $\pm$  fever/chills. In febrile neutropenia, there is only tenderness.**Diagnostic Considerations:** Diagnosis by erythema/tenderness over abscess or by CT/MRI.**Pitfalls:** Do not confuse with perirectal regional enteritis (Crohn's disease) in normal hosts, or with ecthyma gangrenosum in febrile neutropenics.**Therapeutic Considerations:** Antibiotic therapy may be adequate for mild cases. Large abscesses require drainage plus antibiotics  $\times$  1–2 weeks post-drainage. With febrile neutropenia, use an antibiotic that is active against both *P. aeruginosa* and *B. fragilis* e.g., meropenem.**Prognosis:** Good with early drainage/therapy.**Sepsis/Septic Shock****Sepsis/Septic Shock**

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	IV-to-PO Switch
Unknown source	Entero- bacteriaceae B. fragilis E. faecalis (VSE) <sup>†</sup>	Meropenem 1 gm (IV) q8h $\times$ 2 weeks <b>or</b> Piperacillin/ tazobactam 3.375 gm (IV) q6h $\times$ 2 weeks <b>or</b> Moxifloxacin 400 mg (IV) q24h $\times$ 2 weeks	Quinolone* (IV) $\times$ 2 weeks <b>plus either</b> Metronidazole 1 gm (IV) q24h $\times$ 2 weeks <b>or</b> Clindamycin 600 mg (IV) q8h $\times$ 2 weeks	Moxifloxacin 400 mg (PO) q24h $\times$ 2 weeks

VSE/VRE = *vancomycin-sensitive/resistant enterococci*. Duration of therapy represents total time IV or IV + PO.

\* Ciprofloxacin 400 mg (IV) q12h or Levofloxacin 500 mg (IV or PO) q24h.

<sup>†</sup> Treat initially for *E. faecalis* (VSE); if later identified as *E. faecium* (VRE), treat accordingly (urosepsis see pp. 154–155).

## Sepsis/Septic Shock (cont'd)

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	IV-to-PO Switch
Lung source <i>Community-acquired pneumonia</i> <sup>‡</sup>	S. pneumoniae H. influenzae K. pneumoniae**	Respiratory quinolone <sup>†</sup> (IV) q24h × 2 weeks <b>or</b> Ceftriaxone 1 gm (IV) q24h × 2 weeks	Any 2 <sup>nd</sup> generation cephalosporin (IV) × 2 weeks	Respiratory quinolone <sup>†</sup> (PO) q24h × 2 weeks <b>or</b> Doxycycline 200 mg (PO) q12h × 3 days, then 100 mg (PO) q12h × 11 days
	Influenza A (ILI <b>with</b> rapidly cavitating <72 hours multiple pulmonary infiltrates)	MSSA/MRSA	(see pp. 62–63)	(see p. 52)
<i>Nosocomial pneumonia</i>	P. aeruginosa K. pneumoniae E. coli S. marcescens (not MSSA/MRSA)	Same as ventilator-associated pneumonia (see p. 68)		
CVC source*** <i>Bacteremia</i>  (Treat initially for MSSA; if later identified as MRSA, etc., treat accordingly)	S. epidermidis (CoNS) S. aureus (MSSA) Klebsiella Enterobacter Serratia	Meropenem 1 gm (IV) q8h × 2 weeks <b>or</b> Cefepime 2 gm (IV) q12h × 2 wks	Ceftriaxone 1 gm (IV) q24h × 2 wks <b>or</b> Respiratory quinolone* (IV) q24h × 2 wks	Respiratory quinolone* (PO) q24h × 2 weeks <b>or</b> Cephalexin 500 mg (PO) q6h × 2 weeks
	S. aureus (MRSA)	Daptomycin 6 mg/kg (IV) q24h × 2 weeks <b>or</b> Linezolid 600 mg (IV) q12h × 2 weeks		Linezolid 600 mg (PO) q12h × 2 weeks

ILI = Influenza like illness.

MRSA/MSSA = methicillin resistant/sensitive *S. aureus*. Duration of therapy represents total time IV or IV + PO.

\* Moxifloxacin 400 mg or Levofloxacin 500 mg.

\*\* In alcoholics only

\*\*\* If clinically possible, remove CVC as soon as possible.

† Patients with ILI/influenza A (human/swine) presenting with *simultaneous* MSSA/MRSA CAP often present in shock.

‡ Levofloxacin 750 mg (IV) q24h or Moxifloxacin 400 mg (IV) q24h.

§ Uncomplicated by cardiopulmonary decompensation/failure, CAP does not present with hypotension/shock in normal hosts. Hyposplenia/asplenia should be suspected if CAP presents with hypotension/shock in patients with good cardiopulmonary function.

## Sepsis/Septic Shock (cont'd)

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	IV-to-PO Switch
		Quinupristin/dalfopristin 7.5 mg/kg (IV) q8h × 2 weeks  or Vancomycin 1 gm (IV) q12h × 2 weeks		Minocycline 100 mg (PO) q12h × 2 weeks
<p><i>Candidemia</i> (disseminated/invasive)</p> <p>Unless species is known, empiric therapy as for non-albicans/possibly fluconazole-resistant <i>Candida</i> is preferred if recent prior azole therapy, severe illness, neutropenia or high risk for infection with <i>C. glabrata</i> or <i>C. krusei</i></p>	<p><i>Candida albicans</i> (or other fluconazole-susceptible species)<sup>¶</sup></p>	<p>If less critically ill, not neutropenic, and no recent azole exposure: fluconazole is usual first choice alternates an echinocandin may be used.</p> <p>In critically ill, neutropenia or recent azole exposure: an echinocandin is preferred.</p> <p><b>or</b> Micafungin 100 mg (IV) q24h × 2 weeks<sup>†</sup> <b>or</b> Caspofungin 70 mg (IV) × 1 dose, then 50 mg (IV) q24h × 2 weeks<sup>†</sup> Fluconazole 800 mg (IV) × 1, then 400 mg (IV) q24h × 2 weeks<sup>†</sup> <b>or</b> Lipid-associated formulation of amphotericin B (p. 525) (IV) q24h × 2 weeks<sup>†</sup> <b>or</b> Amphotericin B deoxycholate 0.7 mg/kg (IV) q24h × 2 weeks<sup>†</sup> <b>or</b> Voriconazole (see "usual dose," p. 714) <b>or</b> Anidulafungin 200 mg (IV) × 1 dose, then 100 mg (IV) q24h × 2 weeks<sup>†</sup></p>		<p>Fluconazole 400 mg (PO) q24h × 2 weeks<sup>†</sup> <b>or</b> Voriconazole (see "usual dose," p. 714)</p>
	<p>Non-albicans <i>Candida</i><sup>¶</sup> (possibly fluconazole-resistant)</p>	<p>Choices are as for <i>C. albicans</i> (see above), but Fluconazole should <i>not</i> be used and an echinocandin is preferred. Micafungin 100 mg (IV) q24h × 2 weeks<sup>†</sup> <b>or</b> Caspofungin (see <i>C. albicans</i>, above) Use an Amphotericin or Voriconazole if additional mould coverage is desired <b>or</b> Lipid amphotericin B (p. 525) (IV) q24h<sup>†</sup> <b>or</b> Amphotericin B deoxycholate (see <i>C. albicans</i>, above) × 2 weeks<sup>†</sup> <b>or</b> Voriconazole (see "usual dose," p. 714) × 2 weeks<sup>†</sup> <b>or</b> Itraconazole (see <i>C. albicans</i>, above) <b>or</b> Anidulafungin 200 mg (IV) × 1 dose, then 100 mg (IV) q24h × 2 weeks<sup>†</sup></p>		<p>Voriconazole (see "usual dose," p. 714) × 2 weeks<sup>†</sup></p>

¶ Best agent depends on infecting species. Fluconazole-susceptibility varies predictably by species. (See Fluconazole Drug Summary) also, see Amphotericin B (deoxycholate and lipid-associated formulations) Drug Summaries.

† Treat candidemia for 2 weeks after negative blood cultures.



## Sepsis/Septic Shock (cont'd)

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	IV-to-PO Switch
Intra-abdominal/ pelvic source	Enterobacteriaceae <i>B. fragilis</i>	Ertapenem 1 gm (IV) q24h × 2 weeks <b>or</b> Tigecycline 200 mg (IV) × 1 dose, then 100 mg (IV) q24h × 2 weeks <b>or</b> Meropenem 1 gm (IV) q8h × 2 weeks <b>or</b> Piperacillin/tazobactam 3.375 gm (IV) q6h × 2 weeks	<b>Combination therapy with either</b> Ceftriaxone 1 gm (IV) q24h × 2 weeks <b>or</b> Levofloxacin 500 mg (IV) q24h × 2 weeks <b>plus</b> Metronidazole 1 gm (IV) q24h × 2 weeks	Moxifloxacin 400 mg (PO) q24h × 2 weeks <b>or</b> <b>combination therapy with</b> Levofloxacin 500 mg (PO) q24h × 2 weeks <b>plus</b> Clindamycin 300 mg (PO) q8h × 2 weeks
Urosepsis (community-acquired)	Enterobacteriaceae <i>E. faecalis</i> (VSE)	Levofloxacin 500 mg (IV) q24h × 1–2 weeks <b>or</b> Piperacillin/tazobactam 3.375 gm (IV) q6h × 1–2 weeks	Meropenem 1 gm (IV) q8h × 1–2 weeks <b>or</b> Levofloxacin 500 mg (IV) q24h	Levofloxacin 500 mg (PO) q24h × 1–2 weeks
	<i>E. faecium</i> (VRE)	Daptomycin 6 mg/kg (IV) q24h × 1–2 weeks <b>or</b> Linezolid 600 mg (IV) q12h × 1–2 weeks	Quinupristin/dalfopristin 7.5 mg/kg (IV) q8h × 1–2 weeks	Linezolid 600 mg (PO) q12h × 1–2 weeks <b>or</b> Minocycline 100 mg (PO) q12h × 1–2 weeks
(nosocomial)	<i>P. aeruginosa</i> Enterobacteriaceae	Meropenem 1 gm (IV) q8h × 1–2 weeks <b>or</b> Doripenem 1 gm (IV) q8h × 1–2 weeks <b>or</b> Levofloxacin 750 mg (IV) q24h × 1–2 weeks	Aztreonam 2 gm (IV) q8h × 1–2 weeks <b>or</b> Cefepime 2 gm (IV) q12h × 1–2 weeks	Levofloxacin 750 mg (PO) q24h × 1–2 weeks
	CRE	Ceftazidime/avibactam 2.5 gm (IV) q8h × 1–2 weeks	Colistin 5 mg/kg (IV) q8h × 1–2 weeks	

**Sepsis/Septic Shock (cont'd)**

<b>Subset</b>	<b>Usual Pathogens</b>	<b>Preferred IV Therapy</b>	<b>Alternate IV Therapy</b>	<b>IV-to-PO Switch</b>
Urosepsis (group D enterococci)	E. faecalis (VSE)	Ampicillin 2 gm (IV) q4h × 1–2 weeks <b>or</b> Piperacillin/tazobactam 3.375 gm (IV) q6h × 1–2 weeks <b>or combination therapy with</b> Vancomycin 1 gm (IV) q12h × 1–2 weeks <b>plus</b> Gentamicin 240 mg (IV) q24h × 1 week		Amoxicillin 1 gm (PO) q8h × 1–2 weeks <b>or</b> Levofloxacin 500 mg (PO) q24h × 1–2 weeks
	E. faecium (VRE)	Linezolid 600 mg (IV) q12h × 1–2 weeks <b>or</b> Quinupristin/dalfopristin 7.5 mg/kg (IV) q8h × 1–2 weeks		Linezolid 600 mg (PO) q12h × 1–2 weeks <b>or</b> Minocycline 100 mg (PO) q12h × 3 days, then 100 mg (PO) q12h × 4–11 days
Overwhelming sepsis with purpura (asplenia or hyposplenia)	S. pneumoniae H. influenzae N. meningitidis	Ceftriaxone 2 gm (IV) q24h × 2 weeks <b>or</b> Levofloxacin 500 mg (IV) q24h × 2 weeks	Cefepime 2 gm (IV) q12h × 2 weeks <b>or</b> Cefotaxime 2 gm (IV) q6h × 2 weeks	Levofloxacin 500 mg (PO) q24h × 2 weeks <b>or</b> Amoxicillin 1 gm (PO) q8h × 2 weeks
Steroids (high chronic dose)	Aspergillus	Treat as Aspergillus pneumonia (see p. 55)		
Miliary TB	M. tuberculosis	Treat as pulmonary TB (see p. 53) plus steroids × 1–2 wks		
Miliary BCG (disseminated BCG)	Bacille Calmette-Guérin (BCG)	Treat with INH plus RIF × 9 months; EMB, until susceptibilities known; may add steroids e.g., prednisolone 40 mg q24h × 1–2 weeks		
Septic shock	Gram-negative or gram-positive bacteria	Meropenem 1 gm (IV) q8h × 1–2 weeks*		

Duration of therapy represents total time IV or IV + PO.

\* Plus surgical decompression/drainage if needed.

### **Sepsis (Unknown Source)**

**Clinical Presentation:** Abrupt onset of high spiking fevers, rigors ± hypotension.

**Diagnostic Considerations:** Diagnosis suggested by high-grade bacteremia (2/4–4/4 positive blood cultures) with unexplained hypotension. Rule out pseudosepsis (GI bleed, myocardial infarction, pulmonary embolism, acute pancreatitis, adrenal insufficiency, etc.). Sepsis usually occurs from a GI, GU, or IV source, so coverage is directed against GI and GU pathogens if IV line infection is unlikely.

**Pitfalls:** Most cases of fever/hypotension are *not* due to sepsis. Before the label of “sepsis” is applied to febrile/hypotensive patients, first consider treatable/reversible mimics (see p. 151).

**Therapeutic Considerations:** Resuscitate shock patients initially with rapid adequate volume replacement, followed by pressors, if needed. Do not give pressors before volume replacement or hypotension may continue/worsen. Use normal saline, plasma expanders, or blood for volume replacement, not D<sub>5</sub>W. If patient is persistently hypotensive despite volume replacement, consider relative adrenal insufficiency: Obtain a serum cortisol level, then give cortisone 100 mg (IV) q6h × 24–72h; blood pressure will rise promptly if relative adrenal insufficiency is the cause of volume-unresponsive hypotension. Do not add/change antibiotics if patient is persistently hypotensive/febrile; look for GI bleed, myocardial infarction, pulmonary embolism, pancreatitis, undrained abscess, adrenal insufficiency, or IV line infection. Drain abscesses as soon as possible. Remove IV lines if the entry site is red or a central line has been in place for ≥ 7 days and there is no other explanation for fever/hypotension. Early antibiotic therapy/surgical drainage of abscesses, debridement of necrotic tissue eg. necrotizing fasciitis or relief of obstruction is critical.

**Prognosis:** Related to severity of septic process and underlying cardiopulmonary/immune status.

### **Sepsis (Lung Source)**

**Clinical Presentation:** Normal hosts with community-acquired pneumonia (CAP) do not present with sepsis. CAP with sepsis suggests the presence of impaired immunity/hyposplenic function (see “sepsis in hyposplenia/asplenia,” p. 157). Nosocomial pneumonia uncommonly presents as (or is complicated by) sepsis with otherwise unexplained hypotension.

**Diagnostic Considerations:** Impaired splenic function may be inferred by finding Howell-Jolly bodies (small, round, pinkish or bluish inclusion bodies in red blood cells) in the peripheral blood smear. The number of Howell-Jolly bodies is proportional to the degree of splenic dysfunction.

**Pitfalls:** CAP with hypotension/sepsis should suggest hyposplenic function, impaired immunity, or an alternate diagnosis that can mimic CAP/shock. Be sure to exclude acute MI, acute heart failure/COPD, PE/infarction, overzealous diuretic therapy, concomitant GI bleed, and acute pancreatitis.

**Therapeutic Considerations:** Patients with malignancies, myeloma, or SLE are predisposed to CAP, which is not usually severe or associated with shock. Be sure patients with CAP receiving steroids at less than “stress doses” do not have hypotension/shock from relative adrenal insufficiency. In patients with SLE, try to distinguish between lupus pneumonitis and CAP; SLE pneumonitis usually occurs as part of a SLE flare, CAP usually does not.

**Prognosis:** Related to underlying cardiopulmonary/immune status. Early treatment is important.

### **Sepsis (Central Venous Catheter [CVC] Source)**

**Clinical Presentation:** Temperature ≥ 102°F ± IV site erythema.

**Diagnostic Considerations:** Diagnosis by semi-quantitative catheter tip culture with ≥ 15 colonies plus blood cultures with same pathogen. If no other explanation for fever and line has been in place ≥ 7 days, remove CVC and obtain semi-quantitative catheter tip culture. Suppurative thrombophlebitis presents with hectic/septic fevers and pus at IV site ± palpable venous cord.

**Pitfalls:** Temperature  $\geq 102^{\circ}\text{F}$  with CVC infection, in contrast to phlebitis (temperature  $\leq 102^{\circ}\text{F}$ ). Acute bacterial endocarditis (ABE) may complicate intracardiac or CVC (not peripheral) infection.

**Therapeutic Considerations:** Line removal is usually curative, but antibiotic therapy is usually given for 1 week after CVC removal for gram-negative bacilli or 2 weeks after CVC removal for *S. aureus* (MSSA/MRSA). Antifungal therapy is also usually given for 2 weeks after CVC removal for candidemia. Dilated ophthalmoscopy by an ophthalmologist is important to exclude candidal endophthalmitis following candidemia.

**Prognosis:** Good if CVC is removed before endocarditis/metastatic spread.

### **Sepsis (Intra-abdominal/Pelvic Source)**

**Clinical Presentation:** Fever, peritonitis  $\pm$  hypotension. Usually a history of an intra-abdominal disorder that predisposes to sepsis (e.g., diverticulosis, gallbladder disease, recent intra-abdominal/pelvic surgery). Signs and symptoms are referable to the abdomen/pelvis.

**Diagnostic Considerations:** Clinical presentation plus imaging studies (e.g., abdominal/pelvic CT or MRI to demonstrate pathology) are diagnostic.

**Pitfalls:** Elderly patients may have little/no fever and may not have rebound tenderness. Be sure to exclude intra-abdominal mimics of sepsis (e.g., GI bleed, pancreatitis).

**Therapeutic Considerations:** Empiric coverage should be directed against aerobic gram-negative bacilli plus *B. fragilis*. Anti-enterococcal coverage is not essential. Antibiotic therapy is ineffective unless ruptured viscus is repaired, obstruction is relieved, abscesses are drained.

**Prognosis:** Related to rapidity/adequacy of abscess drainage and repair/lavage of ruptured organs. The preoperative health of the host is also important.

### **Sepsis (Urinary Source)**

**Clinical Presentation:** Fever/hypotension in a patient with diabetes mellitus, SLE, myeloma, pre-existing renal disease, stone disease, or partial/total urinary tract obstruction.

**Diagnostic Considerations:** Urine gram stain determines initial empiric coverage. Pyuria is also present. Diagnosis confirmed by culturing the same isolate from urine and blood.

**Pitfalls:** Pyuria without bacteriuria and bacteremia due to same pathogens is not diagnostic of urosepsis. Urosepsis does not occur in normal hosts; look for host defect (e.g., diabetes, renal disease).

**Therapeutic Considerations:** If stones/obstruction are not present, urosepsis resolves rapidly with appropriate therapy. Delayed/no response suggests infected/obstructed stent, stone, partial/total urinary tract obstruction, or renal abscess.

**Prognosis:** Good if stone/stent removed, obstruction relieved, abscess drained. Fatalities rare with urosepsis.

### **Sepsis (Hyposplenism/Asplenia)**

**Clinical Presentation:** Presents as overwhelming septicemia/shock with petechiae.

**Diagnostic Considerations:** Diagnosis by gram stain of buffy coat of blood or by blood cultures. Organism may be stained/cultured from aspirated petechiae. Howell-Jolly bodies in the peripheral smear are a clue to decreased splenic function. Conditions associated with hyposplenism include sickle cell trait/disease, cirrhosis, rheumatoid arthritis, SLE, systemic necrotizing vasculitis, amyloidosis, celiac disease, chronic active hepatitis, Fanconi's syndrome, IgA deficiency, intestinal lymphangiectasia, intravenous gammaglobulin therapy, myeloproliferative disorders, non-Hodgkin's lymphoma, regional enteritis, ulcerative colitis, Sezary syndrome, splenic infarcts/malignancies, steroid therapy, systemic mastocytosis, thyroiditis, infiltrative diseases of spleen, mechanical compression of splenic artery/spleen, Waldenström's macroglobulinemia, hyposplenism of old age, congenital absence of spleen.

**Pitfalls:** Suspect hyposplenism/asplenia in unexplained overwhelming infection.

**Therapeutic Considerations:** In spite of early aggressive antibiotic therapy and supportive care, patients often die within hours from overwhelming infection, especially due to *S. pneumoniae*.

**Prognosis:** Related to degree of splenic dysfunction.

### **Sepsis (Chronic High Dose Steroids)**

**Clinical Presentation:** Subacute onset of fever with disseminated infection (*Candida*, *Aspergillus*) in multiple organs.

**Diagnostic Considerations:** Diagnosis by positive blood cultures for fungi or demonstration of invasive fungal infection from tissue biopsy specimens. Sepsis is most commonly due to fungemia.

**Pitfalls:** Obtain blood cultures to diagnose fungemias and rule out bacteremias (uncommon).

**Therapeutic Considerations:** Empirical approach is the same as for invasive candidiasis (p. 153) or invasive *aspergillosis* (p. 159 and p. 164). Therapy focused solely on candidiasis (e.g., fluconazole alone) should be used only if *aspergillosis* seems unlikely following careful review of the epidemiologic and clinical presentation.

**Prognosis:** Related to degree of immunosuppression.

### **Miliary (Disseminated) TB (Mycobacterium tuberculosis)**

**Clinical Presentation:** Unexplained, prolonged fevers without localizing signs.

**Diagnostic Considerations:** Diagnosis by AFB on biopsy/culture of liver or bone marrow.

**Pitfalls:** Chest x-ray is negative early in 1/3. Subtle miliary (2 mm) infiltrates on chest x-ray 1–4 weeks.

**Therapeutic Considerations:** Treated the same as pulmonary TB ± steroids initially.

**Prognosis:** Death within weeks without treatment.

### **Miliary (Disseminated) BCG (Bacille Calmette-Guérin)**

**Clinical Presentation:** Fever, circulatory collapse, DIC days to weeks after intravesicular BCG.

**Diagnostic Considerations:** Usually occurs in compromised hosts (e.g., transplants, active TB, congenital/acquired immunodeficiencies [e.g., HIV], leukemias/lymphomas). Rare in normal hosts.

**Pitfalls:** Avoid intravesicular BCG immediately after traumatic catheterization, bladder biopsy, TURP.

**Therapeutic Considerations:** Treat with 4 anti-TB drugs plus steroids. Do not repeat BCG therapy.

**Prognosis:** Good with early treatment.

## Febrile Neutropenia

### Febrile Neutropenia

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	IV-to-PO Switch
Febrile leukopenia < 7 days	<i>P. aeruginosa</i> Enterobacteriaceae (not MSSA/MRSA)*	Meropenem 1 gm (IV) q8h*  or Levofloxacin 750 mg (IV) q24h*	Cefepime 2 gm (IV) q8h*  or Doripenem 1 gm (IV) q8h*	Levofloxacin 750 mg (PO) q24h*

**Febrile Neutropenia (cont'd)**

<b>&gt; 7 days</b>	C. albicans Non-albicans Candida Aspergillus	Micafungin 100 mg (IV) q24h* <b>or</b> Voriconazole (see "usual dose," p. 714)* <b>or</b> Caspofungin 70 mg (IV) × 1 dose, then 50 mg (IV) q24h* <b>or</b> Lipid-associated formulation of amphotericin B (IV) (p. 525) q24h*	Amphotericin B deoxycholate 1.5 mg/kg (IV) q24h until 1–2 gm given	Itraconazole 200 mg (PO) q12h* <b>or</b> Voriconazole (see "usual dose," p. 714)*
	MSSA (2° to CVC)†	Same as above (have good GNB and MSSA activity)	Same as above (have good GNB and MSSA activity)	Linezolid 600 mg (PO) q12h <b>or</b> Minocycline§ 100 mg (PO) q12h
	MRSA (2° to CVC)†	Meropenem 1 gm (IV) q8h* <b>or</b> Levofloxacin 750 mg (IV) q24h* <b>plus</b> Daptomycin§ 6 mg/kg (IV) q24h <b>or</b> Vancomycin§ 1 gm (IV) q12h	Cefepime 2 gm (IV) q8h* <b>or</b> Doripenem 1 gm (IV) q8h* <b>plus either</b> Linezolid 600 mg (IV) q12h <b>or</b> Quinupristin/dalfopristin§ 7.5 mg/kg (IV) q8h	Linezolid 600 mg (PO) q12h <b>or</b> Minocycline§ 100 mg (PO) q12h

MSSA = methicillin-sensitive *S. aureus*. Duration of therapy represents total time IV or IV + PO.

\* Treat until neutropenia resolves.

† If clinically possible, remove CVC.

§ Treat for 2 weeks post-CVC removal.

¶ Neutropenia, per se, does not predispose to MSSA/MRSA bacteremias. MSSA/MRSA bacteremias in febrile neutropenia are 2° to CVC associated infection, *not* neutropenia.

**Clinical Presentation:** Incidence of infection rises as PMN counts fall below 1000/mm<sup>3</sup>.

**Diagnostic Considerations:** Febrile neutropenia < 7 days ± positive blood cultures. After blood cultures are drawn, anti-*P. aeruginosa* coverage should be initiated. Do not overlook ischioirectal or perirectal abscess as sources of fever.

**Pitfalls:** Suspect fungemia if abrupt rise in temperature occurs after 7 days of appropriate anti-*P. aeruginosa* antibiotic therapy. Fungemias usually do not occur in first 7 days of neutropenia. Viridans streptococci (*S. mitis*) may present with bacteremia, shock, ARDs, and rash in febrile neutropenia.

**Therapeutic Considerations:** If a patient is neutropenic for > 2 weeks and develops RUQ/LUQ pain/increased alkaline phosphatase, suspect hepatosplenic candidiasis; confirm diagnosis with abdominal CT/MRI showing mass lesions in liver/spleen and treat as systemic/invasive candidiasis (p. 79). *S. aureus* is not a common pathogen in neutropenic compromised hosts without central IV lines, and *B. fragilis*/anaerobes are not usual pathogens in febrile neutropenia. If IV line infection/perirectal abscess are ruled out, consider tumor fever or drug fever before changing antibiotic therapy.

**Prognosis:** Related to degree and duration of neutropenia.

## Transplant Infections

### Transplant Infections

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	PO Therapy or IV-to-PO Switch
<b>BACTEREMIA OR CANDIDEMIA</b>				
<b>Bacteremia</b> Post-BMT (leukopenic pre-engraftment) < 7 days	<i>P. aeruginosa</i> Enterobacteriaceae <i>S. aureus</i> (MSSA) Viridans streptococci <i>E. faecalis</i> (VSE)	Meropenem 1 gm (IV) q8h*  <b>or</b> Piperacillin 4 gm (IV) q6h*	Quinolone <sup>†</sup> (IV) q24h*  <b>plus either</b> Aztreonam 2 gm (IV) q8h*  <b>or</b> Amikacin 1 gm (IV) q24h*	Quinolone <sup>†</sup> (PO) q24h*  <b>or</b> Ciprofloxacin 750 mg (PO) q12h*
	<i>S. aureus</i> (CO-MRSA/HA-MRSA)	Daptomycin 6 mg/kg (IV) q24h × 2 weeks  <b>or</b> Linezolid 600 mg (IV) q12h × 2 weeks  <b>or</b> Quinupristin/dalfopristin 7.5 mg/kg (IV) q8h × 2 weeks  <b>or</b> Vancomycin 1 gm (IV) q12h × 2 weeks		Linezolid 600 mg (PO) q12h × 2 weeks  <b>or</b> Minocycline 100 mg (PO) q12h × 2 weeks
	<i>E. faecium</i> (VRE)	Linezolid 600 mg (IV) q12h × 1–2 weeks	Daptomycin 6 mg/kg (IV) q24h × 1–2 weeks	Linezolid 600 mg (PO) q12h × 1–2 weeks  <b>or</b> Minocycline 100 mg (PO) q12h × 1–2 weeks

BMT/SOT = bone marrow/solid organ transplant, CO/HA-MRSA = community-onset/hospital-acquired methicillin-resistant *S. aureus* (see p. 14), MSSA = Methicillin-sensitive *S. aureus*.

<sup>†</sup> Levofloxacin 750 mg or Moxifloxacin 400 mg.

\* Treat until neutropenia resolves.

## Transplant Infections (cont'd)

Subset	Usual Pathogens	Preferred IV Therapy	Alternate IV Therapy	PO Therapy or IV-to-PO Switch
<b>Candidemia</b> Post- <b>BMT</b> (leukopenic pre-engraftment) > 7 days	<i>C. albicans</i> Non-albicans <i>Candida</i>	Micafungin 100 mg (IV) q24h Ambisome (L-Amb) 3–5 mg/kg (IV) q24h* <b>or</b> Caspofungin 70 mg (IV) × 1 dose, then 50 mg (IV) q24h* <b>or</b> Voriconazole (see “usual dose,” p. 714)* <b>or</b> Anidulafungin 200 mg (IV) × 1 dose, then 100 mg (IV) q24h	Itraconazole 200 mg (IV) q12h × 2 days, then 200 mg (IV) q24h** <b>or</b> Amphotericin B deoxycholate 0.6–1.5 mg/kg (IV) q24h until 1–2 gm given (significant toxicity likely—alternate therapy preferred)	Itraconazole 200 mg (PO) q12h** <b>or</b> Voriconazole (see “usual dose,” p. 714)** <b>or</b> Fluconazole 800 mg (PO) × 1, then 400 mg (PO) q24h**
<b>Bacteremia</b> Post- <b>SOT</b>  (Treat initially for MSSA; if later identified as MRSA, treat accordingly)	<i>S. aureus</i> (MSSA) Enterobacteriaceae	Meropenem 1 gm (IV) q8h × 2 weeks <b>or</b> Ceftriaxone 1 gm (IV) q24h × 2 weeks	Quinolone <sup>†</sup> (IV) q24h × 2 weeks <b>or</b> Cefepime 2 gm (IV) q12h × 2 weeks	Quinolone <sup>†</sup> (PO) q24h × 2 weeks <b>or</b> Cephalexin 500 mg (PO) q6h × 2 weeks
	<i>S. aureus</i> (CO-MRSA/HA-MRSA)	Linezolid 600 mg (IV) q12h × 2 weeks <b>or</b> Daptomycin 6 mg/kg (IV) q24h × 2 weeks <b>or</b> Vancomycin 1 gm (IV) q12h × 2 weeks <b>or</b> Quinupristin/dalfopristin 7.5 mg/kg (IV) q8h × 2 weeks		Linezolid 600 mg (PO) q12h × 2 weeks <b>or</b> Minocycline 100 mg (PO) q12h × 2 weeks

BMT/SOT = bone marrow/solid organ transplant, CO/HA-MRSA = community-onset/hospital-acquired methicillin-resistant *S. aureus* (see p. 14), MSSA = methicillin-sensitive *S. aureus*.

† Levofloxacin 750 mg or Moxifloxacin 400 mg.

\* Treat until neutropenia resolves.

‡ Loading dose is not needed PO if given IV with the same drug.

¶ Significant drug interactions are possible with usual immunosuppressive agents (e.g., sirolimus, cyclosporine, tacrolimus).

Review all concomitant medications for potential interactions. *Voriconazole contraindicated in patients with sirolimus.*



## Transplant Infections (cont'd)

<b>Viremia</b>		
CMV	<p>Ganciclovir 5 mg/kg (IV) q12h until clinical/ virologic response (not &lt; 2 weeks)</p> <p style="text-align: center;"><b>or</b></p> <p>Valganciclovir 900 mg (PO) q12h until clinical/virologic response (not &lt; 2 weeks)</p>	<p>Follow therapy with 1-3 months of CMV prophylaxis (see p. 368) or frequent clinical/virologic followup. Risk of CMV disease greatest 3-6 months post-organ transplant in D+/R- patients during the period of maximum immuno-suppression. Coinfection with HHV-6/7 may increase risk of CMV disease. Risk also high where ALG is used for induction therapy to treat organ rejection. Also, CMV disease risk higher in lung, pancreas and intestine transplants (kidney, liver have lower risk). Lowest risk of CMV disease in D-/R- transplants (should be given leukodepleted blood products and CMV negative blood). Consider IVIG<sup>†</sup> in patients with CMV pneumonia or consider CMVIG<sup>††</sup> in patients with CMV pneumonia or severe CMV infection.</p>
HHV-6	<p>Ganciclovir: 5 mg/ kg (IV) q12h until viremia/infection clears.</p> <p style="text-align: center;"><b>or</b></p> <p>Foscarnet 90 mg/ kg (IV) q12h until viremia/infection clears.</p> <p style="text-align: center;"><b>or</b></p> <p>Cidofovir 5 mg/kg (IV) q week × 2, then q other week* until viremia/infection clears.</p>	<p><i>Reduction of immunosuppression remains the mainstay of treatment.</i> HHV-6-A may be gancyclovir resistant. HHV-6-B variants susceptible to Ganciclovir and Foscarnet</p> <p>Treatment recommended for HHV-6 CNS disease with Ganciclovir, Foscarnet or a combination of both.</p>
BK virus	<p>Cidofovir 0.25 –1 mg/kg (IV) q2 weeks*</p> <p style="text-align: center;"><b>or</b></p>	<p><i>Reduction of immunosuppression remains the mainstay of treatment.</i> Treatment indicated for BKV viremia/BKV nephropathy. <i>No treatment necessary for asymptomatic BKV infection.</i> BV viremia predicts presence of BK nephropathy which precedes viruria. "Decoy cells" in urine indicate BK viruria. BK viruria is common and precedes viremia and invasive infection. ALG: antilymphocyte globulin (thymoglobulin).</p>

\* IVIG: 500 mg/kg (IV) q48h.

†† CMVIG:

**Renal Transplants:** 150 mg/kg (IV) within 72 hours post-transplant, then 100 mg/kg (IV) at weeks 2, 4, 6, 8, then 50 mg/kg (IV) at weeks 12 and 16.

**Non-Renal Transplants:** 150 mg/kg (IV) given within 72 hours post-transplant, then 150 mg/kg (IV) at weeks 2, 4, 6, 8, then 100 mg/kg (IV) at weeks 12 and 16 after transplant.

## Transplant Infections (cont'd)

Viremia (cont'd)		
	Leflunomide (given as mycophenolate replacement) <b>Loading Dose:</b> 100 mg (PO) q24h × 5 days, then <b>Maintenance Dose:</b> 40 mg (PO) q24h until viremia/infection clears.	Measure leflunomide trough levels to ensure therapeutic concentrations (50–100 mcg/ml).
EBV/PTLD	Rituximab <b>or</b> Chemotherapy/surgery	<i>Reduction or discontinuation of immunosuppression remains the mainstay of treatment of PTLD (spontaneous regression in ¼ – ½ of cases).</i>
RSV	Ribavirin 20 mg/mL (aerosol) for 12–18 hours q24h × 3–7 days.	<i>Reduction of immunosuppression remains the mainstay of treatment. Palivizumab may prevent progression from upper to lower RTI. IVIG may be added in severe cases.</i>
Adenovirus	Ribavirin <b>Loading Dose:</b> 30 mg/kg (IV), then <b>Maintenance Dose:</b> 16 mg/kg (IV) q6h × 4 days, then 8 mg/kg (IV) q8h until viremia/infection clears. <b>or</b> Cidofovir 5 mg/kg (IV) q 1–2 weeks (with saline hydration and probenecid) until viremia/infection clears. <sup>§</sup>	<i>Reduction of immunosuppression remains the mainstay of treatment.</i> Oral ribavirin may be effective, but outcomes are variable and may be adenovirus-serotype specific.

§ Probenecid 2 gm (PO) 3 hours prior to cidofovir, 1 gm at 2 hours and 8 hours after completion of infusion. Patient should receive 1 liter normal saline (IV) prior to each infusion of cidofovir; a second liter may be administered over 1–3 hours at the start of cidofovir infusion or immediately following infusion.

\* With hydration/probenecid (see Cidofovir Drug Summary).

## Transplant Infections (cont'd)

CNS INFECTIONS				
Encephalitis/ meningitis	CMV HHV-6	Ganciclovir 5 mg/kg (IV) q12h × 2–4 weeks or until resolution of symptoms and viremia IVIg or CMVIG may be beneficial in severe cases Valganciclovir 900 mg (PO) q24h × 3 months. After successful treatment, a course of 2° prophylaxis is recommended		
	CMV (ganciclovir-resistant CMV: MIC ≥ 3 mcg/mL) or UL 54 or UL 97 mutations	Cidofovir 5 mg/kg (IV) of 1–2 weeks with saline plus probenecid (see p. 163) <b>or</b> Foscarnet 60 mg/kg (IV) q8h × 2 weeks <b>or</b> Foscarnet plus Ganciclovir (see above) <b>or</b> Cidofovir (see above) may be used in severe cases		
	Listeria, HSV, C. neoformans, M. tuberculosis treated the same as in normal/other compromised hosts (see pp. 25–26)			
Brain abscess/ mass lesions	Aspergillus	Voriconazole (see “usual dose,” p. 714) until cured*	Ambisome (L-Amb) 5 mg/kg (IV) q24h until cured <b>or</b> Amphotericin B 1.5 mg/kg (IV) q24h until cured	Voriconazole (see “usual dose,” p. 714) until cured*
	Nocardia	TMP–SMX 5 mg/kg (IV) q6h <b>or</b> Minocycline 200 mg (IV) q12h until clinical improvement, then (PO) therapy for at least 9–12 months or until cured	Linezolid 600 mg (IV) q12h until clinical improvement, then (PO) therapy for at least 9–12 months or until cured <b>or</b> Meropenem 2 gm (IV) q8h for at least 9–12 months or until cured	TMP–SMX 2 DS (PO) q8h <b>or</b> Minocycline 200 mg (PO) q12h <b>or</b> Linezolid 600 mg (PO) q12h for at least 9–12 months or until cured
	T. gondii	<u>Preferred Therapy</u> Sulfadiazine 1–1.5 gm (PO) q6h + Pyrimethamine 200 mg (PO) × 1 dose then 50 mg (PO) q6h + folinic acid 10 mg (PO) q24h × 6–8 weeks until CT/MRI clinical response.		

\* Significant drug interactions are possible with usual immunosuppressive agents (e.g., sirolimus, cyclosporine tacrolimus). Review all concomitant medications for potential interactions. *Voriconazole contraindicated in patients on sirolimus.*

## Transplant Infections (cont'd)

CNS INFECTIONS (cont'd)		
		Follow with sulfadiazine 1 gm (PO) q12h + Pyrimethamine 50 mg (PO) q24h + folinic acid 10 mg (PO) q24h until cured <u>Alternate Therapy</u> Clindamycin 600 mg (IV or PO) q6h + Pyrimethamine 200 mg (PO) × 1 dose then 50 mg (PO) q6h + folinic acid 10 mg (PO) q24h × 6–8 weeks until CT/MRI clinical response. Follow with Sulfadiazine 1 gm (PO) q12h + Pyrimethamine 50 mg (PO) q24h + folinic acid 10 mg (PO) q24h until cured
	C. neoformans	Treat the same as in chronic meningitis (see p. 21)
PNEUMONIAS		
Focal/ segmental/ nodular infiltrates <i>Acute</i>	Legionella sp.	Treat the same as in normal hosts (see p. 56–57)
	L. micdadei	Treat the same as in normal hosts (see p. 56–57) <u>Alternate Therapy</u> TMP-SMX (see p. 55)
<i>Subacute</i>	Aspergillus	<u>Preferred Therapy</u> Voriconazole (see “usual dose,” p. 714) until cured** <b>or</b> Isavuconazole 200 mg (IV) q8h × 48 hours, then 200 mg (IV/PO) q24h until cured <b>or</b> Lipid amphotericin (p. 525) (IV) q24h until cured <u>Alternate Therapy</u> Itraconazole 200 mg (IV) q12h × 2 days, then 200 mg (IV) q24h × 1–2 weeks, then 200 mg (PO) solution q12h until cured*
	M. tuberculosis C. neoformans Nocardia	For TB (see p. 53). For C. neoformans (see p. 269). For Nocardia (see p. 268).
Diffuse infiltrates	S. stercoralis (hyperinfection syndrome)	<u>Preferred Therapy</u> Ivermectin 200 mcg/kg (PO) q24h until cured <u>Alternate Therapy</u> Thiabendazole 25–50 mg/kg (PO) q12h (max. 3 gm/day) until cured

Duration of therapy represents total time PO, IV, or IV + PO. Most patients on IV therapy able to take PO medications should be switched to PO therapy after clinical improvement.

\* If < 40 kg, use 100 mg (PO) maintenance dose.

† Significant drug interactions are possible with usual immunosuppressive agents (e.g., sirolimus, cyclosporine, tacrolimus). Review all concomitant medications for potential interactions. *Voriconazole contraindicated in patients on sirolimus.*

## Transplant Infections (cont'd)

PNEUMONIAS (cont'd)		
	PCP/RSV	Treat the same as in other compromised hosts (see p. 325)
	CMV, HHV-6	Treat the same as CMV (see p. 55), HHV-6 (see p. 164)
HEPATITIS		
Viral hepatitis	CMV	Treat the same as for CMV pneumonia (see p. 55)
	HBV, HCV	Treat the same as in normal hosts (see pp. 97–98)

### **Bacteremia/Candidemia (Bacteremia)**

**Clinical Presentation:** Fever and shaking chills  $\pm$  localizing signs. If localizing signs are present, the organ involved indicates the origin of the bacteremia (e.g., urinary tract findings suggest urosepsis).

**Diagnostic Considerations:** Diagnosis is clinical and is confirmed by positive blood cultures.

**Pitfalls:** 3/4 or 4/4 positive blood cultures indicates bacteremia. Even 1/4 positive blood cultures of an unusual pathogen may be clinically significant in BMT/SOT. The significance of 1/4 blood cultures with coagulase-negative staphylococci is less clear. *S. epidermidis* bacteremia is usually IV-line related, but in some compromised hosts, it may be pathogenic without an IV line focus.

**Therapeutic Considerations:** In SOT, coverage should be directed against *S. aureus* and Enterobacteriaceae. Anti-*P. aeruginosa* coverage is usually not needed since these patients are not neutropenic. If the source of infection is a central IV line, the line should be removed. In pre-engraftment BMT, coverage should be directed against *P. aeruginosa* until leukopenia resolves. Continued fever after 1 week of appropriate antibiotic therapy suggests the presence of candidemia, or another invasive fungal infection.

**Prognosis:** Good with early antibiotic therapy and, if appropriate, IV line removal.

### **Bacteremia/Candidemia (Candidemia)**

**Clinical Presentation:** Fever and shaking chills  $\pm$  localizing signs. If localizing signs are present, the organ involved indicates the origin of the fungemia (e.g., reddened central IV line site suggests IV line-related fungemia).

**Diagnostic Considerations:** *Candida* and *Fusarium* are the most common fungi associated with fungemia in BMT/SOT.

**Pitfalls:** Do not assume that all *Candida* are *C. albicans*. Non-*albicans* *Candida* are more common in SOT patients. Empiric therapy should be directed against non-*albicans* *Candida* pending speciation, which will also cover *C. albicans* (including fluconazole-resistant strains). Because mortality/morbidity associated with candidemia exceeds that of bacteremia, empiric therapy should be started as soon as candidemia is suspected. *Aspergillus*, common after BMT/SOT, rarely grows in blood cultures.

**Prognosis:** Related to underlying immune status and promptness of empiric antifungal therapy, and if appropriate, CVC removal.

### **CNS Infections (Encephalitis/Meningitis)**

**Clinical Presentation:** Typical encephalitis/meningitis presentation (fever, headache,  $\pm$  stiff neck, change in mental status).

**Diagnostic Considerations:** CSF usually reveals a lymphocytic predominance with a normal or normal CSF lactic acid and low glucose. The diagnosis of HSV, CMV, HHV-6 encephalitis can be made by CSF PCR.

**Pitfalls:** Patients with *Listeria* encephalitis often have a negative CSF Gram stain, but *Listeria* nearly always grows on CSF culture. HSV and *Listeria* encephalitis typically have RBCs in the CSF. Head CT/MRI rules out CNS mass lesions. Nuchal rigidity may be absent in meningitis.

**Therapeutic Considerations:** CMV encephalitis is rare but treatable, resulting in clinical/radiological improvement. However, neurological deficits usually remain. CMV retinitis, common in HIV (p. 327), is unusual in BMT/SOT.

**Prognosis:** Related to underlying immune status and promptness of therapy.

### **CNS Infections (Brain Abscess/Mass Lesions)**

**Clinical Presentation:** BMT/SOT patients with brain abscesses/mass lesions present with seizures/cranial nerve abnormalities. Mental status is clear, in contrast to patients with encephalitis, and nuchal rigidity is absent, in contrast to most patients with meningitis.

**Diagnostic Considerations:** Head CT/MRI is the preferred diagnostic modality, and brain biopsy is the definitive diagnostic method. CSF analysis is not usually helpful in mass lesions, with the exception of infection due to *M. tuberculosis* or *C. neoformans*. With *C. neoformans*, the CSF cryptococcal antigen test is positive, and the CSF India ink preparation may be positive. With *M. tuberculosis*, acid fast smear of the CSF is sometimes positive, but culture has a higher yield and PCR is the preferred diagnostic modality.

**Pitfalls:** Patients with brain abscesses/mass lesions should have a head CT/MRI before lumbar puncture. To avoid herniation during lumbar puncture when a mass lesion is present, lumbar puncture should be performed by an experienced operator, and a minimal amount of CSF should be withdrawn.

**Therapeutic Considerations:** *M. tuberculosis* and *C. neoformans* are readily treatable. Be sure to use antimicrobial therapy that penetrates into CSF/brain. If TMP, TMP-SMX, or minocycline cannot be used for CNS *Nocardia*, linezolid or meropenem (meningeal doses) may be useful.

**Prognosis:** Related to underlying immune status and promptness of therapy.

### **Pneumonias (Focal, Segmental, or Nodular Pulmonary Infiltrates)**

**Clinical Presentation:** Acute or subacute community-acquired pneumonia (CAP) with respiratory symptoms and fever.

**Diagnostic Considerations:** BMT/SOT patients with focal/segmental infiltrates are most commonly infected with the usual CAP pathogens affecting normal hosts (e.g., *S. pneumoniae*, *H. influenzae*, *Legionella*). The clinical presentation of CAP in organ transplants is indistinguishable from that in normal hosts. However, BMT/SOT patients presenting subacutely with focal/segmental infiltrates are usually infected with pulmonary pathogens with a slower clinical onset (e.g., *Nocardia*, *Aspergillus*). *L. micdadei* common in BMT/SOT, and may cavitate. Empiric therapy will not cover all possible pathogens; tissue biopsy is necessary for definitive diagnosis and specific therapy. Preferred diagnostic modalities include transbronchial lung biopsy, percutaneous thin needle biopsy, or open lung biopsy, not BAL which may be negative in tissue invasive infections, e.g., *Aspergillus*.

**Pitfalls:** Patients presenting with subacute onset of CAP have a different pathogen distribution than those presenting with acute CAP. PCP/CMV does not present with focal/segmental infiltrates. Unlike other *Legionella* sp., *L. micdadei* partially acid fast, and susceptible to TMP-SMX (as well as doxycycline and quinolones).

**Therapeutic Considerations:** BMT/SOT patients with acute onset of CAP are treated with the same antibiotics used to treat CAP in normal hosts. Empiric coverage is directed against both typical and atypical

bacterial pathogens. If no improvement in clinical status after 72 hours, proceed to transbronchial biopsy or lung biopsy to identify nonbacterial pathogens (e.g., *Nocardia*, *Aspergillus*). *Nocardia* reinfection may require 6–12 months of therapy.

**Prognosis:** Best with acute focal/segmental infiltrates. Not as good with subacute or chronic focal/segmental infiltrates.

### **Pneumonias (Diffuse Pulmonary Infiltrates)**

**Clinical Presentation:** Insidious onset of interstitial pneumonia usually accompanied by low-grade fevers. Focal/segmental infiltrates are absent.

**Diagnostic Considerations:** Bilateral diffuse infiltrates, which can be minimal or extensive, fall into two clinical categories: those with and without hypoxemia/ $\uparrow$  A-a gradient. Diffuse pulmonary infiltrates without hypoxemia suggest a noninfectious etiology (e.g., CHF, pulmonary drug reaction, pulmonary hemorrhage). The differential diagnosis of diffuse pulmonary infiltrates with hypoxemia includes PCP, CMV, HSV, RSV, HHV-6, VZV. Quantitative PCR testing of BAL fluid for CMV, VZV, RSV, HHV-6 may be diagnostically helpful. For interstitial infiltrates with hypoxemia, the chest x-ray may be only minimally abnormal, but chest CT scans are more revealing. CMV/PCP may require tissue biopsy for definitive diagnosis. A very highly elevated LDH suggests PCP. PCP =  $\beta$  1,3 D-glucan +, aspergillus galactomannan -. Transbronchial biopsy is preferable, but BAL may be used. The incidence of CMV pneumonia is highest in lung transplants and BMTs.

**Pitfalls:** Because infections in BMT/SOT are sequential, many patients with PCP pneumonia may have underlying CMV or HHV-6. In BMT, CMV found alone on lung biopsy suggests it is the primary pathogen. Serological tests are helpful for CMV ( $\uparrow$  CMV IgM);  $\uparrow$  CMV viral loads reflective of peripheral WBC reactivation, not pneumonia. CMV pneumonia diagnosed by demonstrations CMV CPEs (intracellular inclusions) on lung BAL/biopsy. *Candida* pneumonia does not exist as a separate entity but only rarely as part of disseminated/invasive candidiasis.

**Therapeutic Considerations:** Among the subacute diffuse pneumonias, PCP is readily treatable. Initiate treatment for CMV pneumonia with ganciclovir IV; after clinical improvement, complete therapy with valganciclovir (PO) until cured. If after treatment, there is an  $\uparrow$  in CMV antigen levels or quantitative PCR, treat pre-emptively to prevent CMV pneumonia with valganciclovir 900 mg (PO) q24h until CMV antigen levels return to previous levels or quantitative PCR becomes undetectable.

**Prognosis:** Related to underlying immune status, promptness of therapy, and general health of host.

### **Hepatitis (Viral)**

**Clinical Presentation:** Fever with  $\uparrow$  SGOT/SGPT  $\pm$  RUQ pain.

**Diagnostic Considerations:** Because CMV is of such critical importance in BMT/SOT, CMV testing should always be done in organ transplants with  $\uparrow$  SGOT/SGPT. Liver biopsy with immunohistochemical staining is diagnostic; quantitative PCR may be helpful. Prior to SOT, HIV, HBV, HCV testing should be performed on donor and recipient. Other viruses causing acute viral hepatitis post-transplant include EBV, HHV-6, VZV, adenovirus, and influenza. Viral hepatitis is usually accompanied by some degree of leukopenia. A few atypical lymphocytes may be present, and serum transaminases may be mildly or markedly elevated. CMV viremia is the commonest manifestation of CMV infection in BMT/SOT patients. CMV has a predilection for infecting the organ transplanted, and CMV hepatitis is particularly common in liver transplants. Anicteric hepatitis is more common than icteric hepatitis.

**Pitfalls:** The diagnosis of active CMV hepatitis in organ transplant patients is critical because it is an immunomodulating virus, adding to the net immunosuppressive effect of immunosuppressive therapy. Do not rely on CMV IgM/IgG titers for diagnosis.

**Therapeutic Considerations:** CMV should be treated aggressively to minimize its potentiating immunoregulatory defects, which may predispose to non-viral opportunistic pathogens and ↑ risk of chronic rejection in SOT. CMV antigen levels or quantitative PCR increase before CMV infection. Therefore, when CMV antigen levels or quantitative PCR increase, begin early pre-emptive therapy with valganciclovir 900 mg (PO) q12h until CMV antigen levels or quantitative PCR return to previous levels. There is no treatment for EBV.

**Prognosis:** Treated early, CMV responds well to therapy. CMV is associated with chronic allograft rejection in SOT recipient so that prevention or early diagnosis/therapy are critical.

## Toxin-Mediated Infectious Diseases

### Toxin-Mediated Infectious Diseases

Subset	Usual Pathogens	IV Therapy	PO Therapy or IV-to-PO Switch
Toxic shock syndrome (TSS)*  (Treat initially for MRSA; if later identified as MSSA, treat accordingly)	<i>S. aureus</i> (MRSA)	<u>Preferred IV Therapy</u> Vancomycin 1 gm (IV) q12h × 2 weeks <b>or</b> Linezolid 600 mg (IV) q12h × 2 weeks <u>Alternate IV Therapy</u> Minocycline 100 mg (IV) q12h × 2 weeks <b>or</b> Daptomycin 6 mg/kg (IV) q24h × 2 weeks	Linezolid 600 mg (PO) q12h × 2 weeks <b>or</b> Minocycline 100 mg (PO) q12h × 2 weeks
	<i>S. aureus</i> (MSSA)	<u>Preferred IV Therapy</u> Cefazolin 1 gm (IV) q8h × 2 weeks <u>Alternate IV Therapy</u> Nafcillin 2 gm (IV) q4h × 2 weeks <b>or</b> Clindamycin 600 mg (IV) q8h × 2 weeks	Cephalexin 500 mg (PO) q6h × 2 weeks <b>or</b> Clindamycin 300 mg (PO) q6h × 2 weeks
	<i>Clostridium sordelli</i>	<u>Preferred IV Therapy</u> Penicillin G 10 mu (IV) q4h × 2 weeks <b>or</b> Clindamycin 600 mg (IV) q8h × 2 weeks <b>or</b> Piperacillin 4 gm (IV) q8h × 2 weeks <u>Alternate IV Therapy</u> Meropenem 1 gm (IV) q8h × 2 weeks <b>or</b> Ertapenem 1 gm (IV) q24h × 2 weeks	Not applicable

MRSA/MSSA = methicillin-resistant/sensitive *S. aureus*. Duration of therapy represents total time IV, PO, or IV + PO. Most patients on IV therapy able to take PO meds should be switched to PO therapy after clinical improvement.

\* Treat only IV or IV-to-PO switch.



## Toxin-Mediated Infectious Diseases (cont'd)

Subset	Usual Pathogens	IV Therapy	PO Therapy or IV-to-PO Switch
Botulism (food, infant, wound)	Clostridium botulinum	<u>Preferred Therapy</u> 2 vials of type-specific trivalent (types A,B,E) or polyvalent (types A,B,C,D,E) antitoxin (IV)	<u>Alternate Therapy</u> Amoxicillin 1 gm (PO) q8h × 7 days (wound botulism only)
Tetanus	Clostridium tetani	<u>Preferred Therapy</u> Tetanus immune globulin (TIG) 3,000–6,000 units (IM) (50% into deltoid, 50% into wound site) <b>plus either</b> Penicillin G 4 mu (IV) q4h × 10 days <b>or</b> Doxycycline 200 mg (IV or PO) q12h × 3 days, then 100 mg (IV or PO) × 7 days	<u>Alternate Therapy</u> Tetanus antitoxin 1,500–3,000 units (IM/IV) <b>plus</b> Metronidazole 1 gm (IV) q12h × 10 days
Diphtheria (pharyngeal, nasal, wound, myocarditis)	Corynebacterium diphtheriae C. ulcerans	Diphtheria antitoxin (IV) over 1 hour (pharyngeal diphtheria = 40,000 units; nasopharyngeal diphtheria = 60,000 units; systemic diphtheria or diphtheria > 3 days duration = 100,000 units) <b>plus either</b> Penicillin G 1 mu (IV) q4h × 14 days <b>or</b> Erythromycin 500 mg (IV) q6h × 14 days	Diphtheria antitoxin (IV) over 1 hour (pharyngeal diphtheria = 40,000 units; nasopharyngeal diphtheria = 60,000 units; systemic diphtheria or diphtheria > 3 days duration = 100,000 units) <b>plus</b> Procaine penicillin 600,000 units (IM) q24h × 14 days

Duration of therapy represents total time IV, PO, or IV + PO. Most patients on IV therapy able to take PO meds should be switched to PO therapy after clinical improvement.

### Toxic Shock Syndrome (*S. aureus*)

**Clinical Presentation:** Scarlatiniform rash ± hypotension. Spectrum ranges from minimal infection to multiorgan system failure/shock. ↑ CPK common.

**Diagnostic Considerations:** Diagnosis by clinical presentation with mucous membrane, renal, liver, and skin involvement/culture of TSS-1 toxin-producing strain of *S. aureus* from mouth, nares, vagina, or wound.

**Pitfalls:** Toxic shock syndrome wound discharge is clear, not purulent.

**Therapeutic Considerations:** Remove source of toxin production if possible (e.g., remove tampon, drain collections). Clindamycin may be added for its anti-toxin effect.

**Prognosis:** Good in early/mild form. Poor in late/multisystem disease form.

### **Toxic Shock Syndrome (C. sordelli)**

**Clinical Presentation:** Resembles clostridial myonecrosis (gas gangrene) with soft tissue necrotizing infection are local edema. Hypotension with acute onset of nausea/vomiting and weakness characteristic. Associated with IVDA (black tar heroin), trauma, parturition, abortion, cadaver graft surgery. Hemoconcentration with leukocytosis typical; leukemoid reactions common with WBC counts  $> 50k/mm^3$ .

**Diagnostic Considerations:** Clinical diagnosis. Culture of *C. sordelli* from necrotic soft tissue.

**Pitfalls:** Gas gangrene like clinical presentation but with hemoconcentration not hemolytic anemia. Nausea/vomiting instead of diarrhea as with gas gangrene. Shock with no/low fever and  $\uparrow$  WBC should suggest the diagnosis. Muscle involvement (clostridial myonecrosis) and bullae typical of gas gangrene not a feature of *C. sordelli* TSS.

**Therapeutic Considerations:** Early/adequate debridement critical. Anti-anaerobic antibiotics and supportive measures important.

**Prognosis:** Like gas gangrene, prognosis related to early diagnosis and early/adequate surgical debridement.

### **Botulism (Clostridium botulinum)**

**Clinical Presentation:** Descending symmetrical paralysis beginning with cranial nerve involvement, induced by botulinum toxin. Onset begins with blurry vision, followed rapidly by ocular muscle paralysis, difficulty speaking, and inability to swallow. Respiratory paralysis may occur in severe cases. Mental status is unaffected. Usual incubation period is 10–12 hours. Incubation is shortest for Type E strain (hours), longest for Type A strain (up to 10 days), and is inversely proportional to the quantity of toxin consumed (food botulism). Wound botulism (Types A or B) may follow *C. botulinum* entry into IV drug abuser injection site, surgical or traumatic wounds. Infant (< 1 year) botulism (most commonly Type A or B) is acquired from *C. botulinum* containing honey. Patients with botulism are afebrile, and have profuse vomiting without diarrhea.

**Diagnostic Considerations:** Detection of botulinum toxin from stool, serum, or food (especially home canned foods with neutral or near neutral pH [ $\sim 7$ ] or smoked fish [Type E]) is diagnostic of food botulism. Wound botulism is diagnosed by culturing *C. botulinum* from the wound or by detecting botulinum toxin in the serum.

**Pitfalls:** Clinical diagnosis based on descending paralysis with cranial nerve involvement in an afebrile patient must be differentiated from Guillain-Barre (fever, ascending paralysis, sensory component) and polio (fever, pure ascending motor paralysis). Do not diagnose botulism in the absence of ocular/pharyngeal paralysis.

**Therapeutic Considerations:** Antitoxin neutralizes only unbound toxin, and does not reverse toxin-induced paralysis. Botulism is a toxin-mediated infection and antibiotic therapy (wound botulism) is adjunctive. Guanidine has been used with variable effect. Ventilator support is needed for respiratory paralysis. Bioterrorist botulism presents clinically and is treated the same as naturally-acquired botulism.

**Prognosis:** Good if treated early, before respiratory paralysis.

### **Tetanus (*Clostridium tetani*)**

**Clinical Presentation:** Begins with jaw stiffness/difficulty chewing induced by *C. tetani* toxin (tetanospasmin). Trismus rapidly follows with masseter muscle spasm, followed by spasm of the abdominal/back muscles. Rigidity and convulsions may occur. Patients are afebrile unless there is hypothalamic involvement (central fever), in which case fevers may exceed 106°F. Usual incubation period is 3–21 days.

**Diagnostic Considerations:** Diagnosis suggested by muscle spasms/rigidity in a patient with trismus.

**Pitfalls:** In rabies, muscle spasms are localized and usually involve the face/neck, rather than primary involvement of the extremities, as in tetanus.

**Therapeutic Considerations:** Tetanus is self-limited with intensive supportive care. Sedation is important, and avoidance of all stimuli is mandatory to reduce the risk of convulsions. Avoid unnecessary handling/movement of patient. Antitoxin is effective only in neutralizing unbound toxin. Tracheostomy/respiratory support can be lifesaving in severe cases.

**Prognosis:** Good if not complicated by spinal fractures, aspiration pneumonia, or CNS involvement (hyperpyrexia, hyper/hypotension).

### **Diphtheria (*Corynebacterium diphtheriae*/*Corynebacterium ulcerans*)**

**Clinical Presentation:** Within 1 week following insidious onset of sore throat without fever, pharyngeal patches coalesce to form a gray diphtheric membrane (surrounded by a red border), which is adherent/bleeds easily when removed. Membrane begins unilaterally; may extend to the soft palate, uvula and contralateral posterior pharynx; are accompanied by prominent bilateral anterior adenopathy; become necrotic (green/black); and have a foul odor (fetor oris). Submandibular edema ("bull neck") and hoarseness (laryngeal stridor) precede respiratory obstruction/death. Cutaneous diphtheria may follow *C. diphtheriae* contaminated wounds (traumatic, surgical) or insect/human bites, and is characterized by a leathery eschar (cutaneous membrane) covering a deep punched out ulcer. Serosanguineous discharge is typical of nasal diphtheria (membrane in nares). Diphtheric myocarditis may complicate any form of diphtheria (most commonly follows pharyngeal form), and usually occurs in the second week, but may occur up to 8 weeks after infection begins. Diphtheric polyneuritis is a common complication. Cardiac/neurologic complications are due to elaboration of a potent toxin.

**Diagnostic Considerations:** Diagnosis is suggested by unilateral membranous pharyngitis/palatal paralysis, absence of fever, and relative tachycardia. Diagnosis is confirmed by culture of *C. diphtheriae* from nares, membrane, or wound.

**Pitfalls:** Differentiated from *Arcanobacterium* (*Corynebacterium*) *haemolyticum* (which also forms a pharyngeal membrane) by culture and absence of scarlatiniform rash with *C. diphtheriae*. *C. ulcerans* has the same clinical features as *C. diphtheriae*.

**Therapeutic Considerations:** Antibiotic therapy treats the infection and stops additional toxin production. Antitoxin is effective against unbound toxin, but will not reverse toxin-mediated myocarditis/neuropathy. Serum sickness is common 2 weeks after antitoxin. Respiratory/cardiac support may be lifesaving. *C. ulcerans* is treated the same as *C. diphtheriae*.

**Prognosis:** Poor with airway obstruction or myocarditis. Myocarditis may occur despite early treatment.

## Bioterrorist Agents

### Bioterrorist Agents in Adults<sup>¶</sup>

Subset	Pathogen	IV/IM Therapy	IV-to-PO Switch
Anthrax <i>Inhalation, oropharyngeal, gastrointestinal</i>	Bacillus anthracis	Quinolone* (IV) × 2 weeks <b>or</b> Doxycycline 200 mg (IV) q12h × 3 days, then 100 mg (IV) q12h × 11 days <sup>‡</sup> <b>or</b> Penicillin G 4 MU (IV) q4h ± Clindamycin 600 mg (IV) q8h × 2 weeks	Quinolone* (PO) × 2 weeks <b>or</b> Doxycycline 200 mg (PO) q12h × 3 days, then 100 mg (PO) q12h × 11 days (loading dose not needed PO if given IV). Duration of IV + PO therapy = 60 days
		Treat severe cases with same (PO) antibiotics as for inhalation anthrax	
		If penicillin susceptible, treat with Penicillin G 4 MU (IV) q24h <b>or</b> Meropenem 2 gm (IV) q8h for at least 2 weeks or markedly improved and complete 60–100 days of therapy as described in the inhalation, oropharyngeal or gastrointestinal forms, in addition to quinolones or doxycycline. Consider adjunctive steroid therapy. Penicillin should NOT be used as a single agent. Not susceptible to cephalosporins or TMP–SMX.	
Tularemia pneumonia	Francisella tularensis	Streptomycin 1 gm (IM) q12h × 10 days <b>or</b> Gentamicin 5 mg/kg (IM or IV) q24h × 10 days <b>or</b> Doxycycline 200 mg (IV) q12h × 3 days, then 100 mg (IV) q12h × 11–18 days <sup>‡</sup> <b>or</b> Chloramphenicol 500 mg (IV) q6h × 14 days <b>or</b> Quinolone* (IV) × 10 days <u>If meningitis</u> ; add Chloramphenicol	Doxycycline 200 mg (PO) q12h × 3 days, then 100 mg (PO) q12h × 11–18 days <b>or</b> Quinolone* (PO) × 10 days  <u>If meningitis</u> ; add Chloramphenicol

Duration of therapy represents total treatment time.

‡ Patients who remain critically ill after Doxycycline 200 mg (IV) q12h × 3 days should continue receiving 200 mg (IV) q12h for the full course of therapy. For patients who have improved after 3 days, the dose may be decreased to 100 mg (IV or PO) q12h to complete the course of therapy. Total duration of IV + PO therapy = 60 days.

\* Ciprofloxacin 400 mg (IV) q12h or 500 mg (PO) q12h or Levofloxacin 500 mg (IV or PO) q24h.

¶ Additional information can be obtained at [www.bt.cdc.gov](http://www.bt.cdc.gov). For post-exposure prophylaxis, see p. 358.

**Bioterrorist Agents in Adults<sup>¶</sup>** (cont'd)

Subset	Pathogen	IV/IM Therapy	IV-to-PO Switch
Pneumonic plague	<i>Yersinia pestis</i>	Treat the same as tularemic pneumonia (see p. 176)	
Botulism	<i>Clostridium botulinum</i>	Contrary to the package insert, administer 50 mg/kg up to 1 vial of type-specific trivalent (types A,B,E) or polyvalent (types A,B,C,D,E) antitoxin (IV) after skin testing. Antitoxin administration is not repeated (circulating antitoxin's half-life = 5–8 days). Treatment with 1 vial resulted in adverse effects in < 1%; treatment with 2–4 times present dose resulted in hypersensitivity reactions in 9%. Antibiotics do not neutralize toxin	
Smallpox	Variola virus	Smallpox vaccine ≤ 4 days after exposure. Cidofovir 5 mg/kg (IV) × 1 dose may be protective for up to 6 days post-exposure.	
Ebola	Ebola virus	No specific therapy. Supportive therapy can be life saving	

Duration of therapy represents total treatment time.

\* Ciprofloxacin 400 mg (IV) q12h or 500 mg (PO) q12h or Levofloxacin 500 mg (IV or PO) q24h.

¶ Additional information can be obtained at [www.bt.cdc.gov](http://www.bt.cdc.gov). For post-exposure prophylaxis, see p. 358.

‡ Patients who remain critically ill after Doxycycline 200 mg (IV) q12h × 3 days should continue receiving 200 mg (IV) q12h for the full course of therapy. For patients who have improved after 3 days, the dose may be decreased to 100 mg (IV or PO) q12h to complete the course of therapy. Total duration of IV + PO therapy = 60 days.

**Anthrax (*B. anthracis*)**

**Clinical Presentation:** Bioterrorist anthrax usually presents as cutaneous or inhalational anthrax. Cutaneous anthrax has the same clinical presentation as naturally-acquired anthrax: Lesions begin as painless, sometimes mildly pruritic papules, usually on the upper extremities, neck, or face, and evolve into a vesicular lesion which may be surrounded by satellite lesions. A “gelatinous halo” surrounds the vesicle as it evolves into an ulcer, and a black eschar eventually develops over the ulcer. Inhalational anthrax is a biphasic illness. Initially, there is a viral illness-like prodrome with fever, chills, and myalgias with chest discomfort 3–5 days after inhaling anthrax spores. Bacteremia is common. Patients often improve somewhat over the next 1–2 days, only to rapidly deteriorate and become critically ill with high fevers, dyspnea, cyanosis, crushing substernal chest pain, and shock. Oropharyngeal anthrax presents with fever, soft tissue edema, painful cervical adenopathy. Lesions in oropharynx ulcerate in ~ 2 weeks. GI anthrax presents with fever, malaise ± syncope, followed in 24 hours by mild nausea/vomiting, severe abdominal pain, and then ascites, ↑ abdominal pain, flushed face, and shock.

**Diagnostic Considerations:** Cutaneous anthrax is a clinical diagnosis suggested by the lack of pain relative to the size of the lesion. A presumptive microbiologic diagnosis is made by finding gram-positive bacilli in the fluid from the gelatinous halo surrounding the ulcer or from under

the eschar. Blood cultures may reveal *B. anthracis*. Definitive diagnosis depends on identifying *B. anthracis* from culture of the skin lesions or blood cultures. Inhalation anthrax is suspected in patients with fevers, chest pain, and mediastinal widening accompanied by bilateral pleural effusions on chest x-ray. If chest x-ray findings are equivocal, then a chest CT/MRI is recommended to demonstrate mediastinal lymph node enlargement. Inhalational anthrax presents as a hemorrhagic mediastinitis, not community-acquired pneumonia. The diagnosis is clinical but supported by Gram stain of hemorrhagic pleural fluid demonstrating gram-positive bacilli. Patients with inhalational anthrax often have positive blood cultures and may have associated anthrax meningitis. If meningitis is present, the CSF is hemorrhagic and CSF Gram stain shows gram-positive bacilli, which, when cultured, is *B. anthracis*.

**Pitfalls:** Cutaneous anthrax is most often initially confused with ringworm or a brown recluse spider bite. Subacute/chronic lesions may initially resemble ringworm, but the skin lesion in ringworm has an annular configuration, is painless, and is accompanied by prominent pruritus, particularly at the edges of the lesion. Patients with ringworm have no fever or systemic symptoms. Brown recluse spider bites produce extremely painful lesions with irregular edges, which eventually develop a necrotic center followed by eschar formation. The lesions of the brown recluse spider bite are irregular, not accompanied by fever, and intensely painful. In contrast, cutaneous anthrax lesions are painless, round, and are not primarily pruritic in nature. Be alert to the possibility of smallpox following outbreaks of other bioterrorist agents such as anthrax, as the genome of smallpox is easily modified and can be incorporated into bacteria.

**Therapeutic Considerations:** *B. anthracis* is highly susceptible to nearly all antibiotics; in the U.S. bioterrorist experience, no strains were resistant to antibiotics. Traditionally, penicillin has been used to treat natural anthrax, but because of concern for resistant bioterrorist strains, doxycycline or quinolones are preferred. Because meningitis is frequently associated with inhalational anthrax, penicillin in (IV) meningeal doses may be added as a second or third antibiotic to quinolones or doxycycline. For meningeal anthrax use penicillin G or meropenem in meningeal doses. Clindamycin is active against *B. anthracis* and has been used in combination therapy because of its potential anti-exotoxin activity. Some patients seemed to respond somewhat better when clindamycin 600 mg (IV) q8h or 300 mg (PO) q8h plus rifampin 300 mg (PO) q12h is added to either a quinolone or doxycycline. Corticosteroids should be considered for severe mediastinal edema or meningitis. Depending upon antimicrobial susceptibility testing, rifampin, vancomycin, penicillin, ampicillin, chloramphenicol, imipenem, clindamycin or clarithromycin may be added if the need arises. Prolonged therapy of 100 days with or without anthrax vaccine has been recommended by some authors. Three doses of anthrax vaccine (BioThraxT, formerly AVA - anthrax vaccine absorbed) have been recommended by the ACIP and the John Hopkins Working Group on Civilian Bio-Defense with antimicrobials for prophylaxis after aerosolized exposure, but as it is not licensed, it must be administered under an IND application. Some *B. anthracis* strains produce cephalosporinase and inducible beta-lactamase that make penicillins drugs less suitable for initial therapy. In general, the organism is resistant to trimethoprim-sulfamethoxazole.

**Prognosis:** Prognosis of cutaneous anthrax is uniformly good. With inhalational anthrax, prognosis is related to the inhaled dose of the organism, underlying host status, and rapidity of initiating antimicrobial therapy. Inhalational anthrax remains a highly lethal infectious disease, but with early intervention/supportive care, some patients survive. Patients with associated anthrax meningitis have a poor prognosis.

### **Tularemia Pneumonia (*F. tularensis*)**

**Clinical Presentation:** Fever, chills, myalgias, headache, dyspnea and a nonproductive cough may occur, but encephalopathy is absent. Chest x-ray resembles other causes of community-acquired pneumonia, but tularemia pneumonia is usually accompanied by hilar adenopathy and pleural effusion, which is serosanguineous or frankly bloody. Cavitation sometimes occurs. Relative bradycardia is not present and serum transaminases are not elevated.

**Diagnostic Considerations:** Tularemia pneumonia can resemble other atypical pneumonias, but in a patient presenting with community-acquired pneumonia, the presence of hilar adenopathy with pleural effusions should suggest the diagnosis. *F. tularensis* may be seen in the Gram stain of the sputum or bloody pleural effusion fluid as a small, bipolar staining, gram-negative bacillus. Diagnosis is confirmed serologically or by culture of the organism from respiratory fluid/blood.

**Pitfalls:** Gram-negative bacilli in the sputum may resemble *Y. pestis* but are not bipolar staining. Chest x-ray may resemble inhalational anthrax (hilar adenopathy/mediastinal widening). Both tularemia pneumonia and inhalational anthrax may be accompanied by bloody pleural effusions. In contrast to inhalational anthrax (which may be accompanied by anthrax meningitis), CNS involvement is not a feature of tularemia pneumonia.

**Therapeutic Considerations:** Streptomycin is the antibiotic traditionally used to treat tularemia. Gentamicin may be substituted for streptomycin if it is not available. Doxycycline, chloramphenicol, or a quinolone are also effective.

**Prognosis:** Depends on inoculum size and health of host. Mortality rates for severe untreated infection can be as high as 30%, although early treatment is associated with mortality rates < 1%.

### **Pneumonic Plague (*Y. pestis*)**

**Clinical Presentation:** Bioterrorist plague presents as pneumonic plague and has the potential for person-to-person spread. After an incubation period of 1–4 days, the patient presents with acute onset of fever, chills, headache, myalgias and dizziness, followed by pulmonary manifestations including cough, chest pain, dyspnea. Hemoptysis may occur, and increasing respiratory distress and circulatory collapse are common. Compared to community-acquired pneumonia, patients presenting with plague pneumonia are critically ill. Sputum is pink and frothy and contains abundant bipolar staining gram-negative bacilli. Chest x-ray is not diagnostic.

**Diagnostic Considerations:** *Yersinia pestis* may be demonstrated in sputum Gram stain (bipolar staining gram-negative bacilli) and may be recovered from blood cultures. Laboratory confirmation requires isolation of *Y. pestis* from body fluid or tissue culture. Consider the diagnosis in any critically ill patient with pneumonia and bipolar staining gram-negative bacilli in the sputum.

**Pitfalls:** Plague pneumonia can resemble tularemia pneumonia, but there are several distinguishing features. Unlike plague, tularemia pneumonia is usually associated with hilar enlargement and pleural effusion. Although gram-negative bacilli may be present in the sputum of patients with tularemia, the organisms are not bipolar staining.

**Therapeutic Considerations:** Streptomycin is the preferred drug for pneumonic plague. Doxycycline or a quinolone is also effective.

**Prognosis:** Depends on inoculum size, health of the host, and the rapidity of treatment. Left untreated, mortality rates exceed 50%. ARDS, DIC, and other manifestations of gram-negative sepsis are more common when treatment is delayed.

**Botulism (*C. botulinum*)** (see p. 171)**Smallpox**

**Clinical Presentation:** After an incubation period of 1–12 days, typical smallpox is heralded by high fever, headache, and gastrointestinal complaints (vomiting, colicky pain). No rash is present at this time. After 1–2 days, the fever decreases to near normal level, and macules begin to appear on the head, usually at the hairline. Macules progress to papules, then vesicles, then finally pustules. The rash begins on the face/head and rapidly spreads to the extremities with relative sparing of the trunk. The mucous membranes of the oropharynx and upper/lower airways are also affected early. Lesions initially are umbilicated, then later lose their umbilication. The fully formed smallpox pustule is located deep in the dermis. The appearance of the pustules is accompanied by recrudescence of fever. Hemorrhagic smallpox is a fulminant form of smallpox that begins with petechial lesions in a “swimming trunk” distribution and results in widespread hemorrhage into the skin and mucous membranes. Patients look toxemic and have high fevers with no other signs of smallpox; death from toxemia often occurs before the typical rash appears.

**Diagnostic Considerations:** Smallpox is most likely to be confused with chickenpox or drug eruptions. Patients with chickenpox are less toxemic and the lesion distribution is different from smallpox. Chickenpox lesions occur in crops for the first 72 hours, then stop. The lesions of chickenpox are superficial, not deep in the dermis like smallpox, and chickenpox vesicles are predominantly centripetal rather than centrifugal. The chickenpox vesicle has been described as a “dewdrop on a rose petal” because of its fragility and superficial location on the skin. If there is any doubt, a Tzanck test should be performed by unroofing the vesicle, scraping cells from the base of the vesicle, and staining the cells. A positive Tzanck test indicates chickenpox, not smallpox. Alternatively, a monoclonal VZV test can be performed on vesicle base cells. Drug eruptions are not accompanied by toxemia and are usually accompanied by relative bradycardia if fever is present.

**Pitfalls:** Smallpox is easily missed before the rash and is difficult to diagnose. Look for the combination of high fever/headache with gastrointestinal symptoms (e.g., abdominal pain) that precedes the rash. GI complaints may be confused with appendicitis. A petechial rash in a swimming trunk distribution does not occur with any other infectious disease and should immediately suggest smallpox. Recently human monkeypox has occurred in the Western Hemisphere after transmission via imported African rodent pets. After an incubation period of 7–19 days, patients develop fever, headache, and malaise. Skin lesions appear on head, trunk, and extremities (including palms/soles). Rash begins like smallpox as macules, then papules, and finally umbilicated vesicles. Some exudative pharyngitis/tonsillitis with cervical adenopathy may be present. Encephalitis is very rare. Laboratory results are nonspecific. Unlike smallpox, human monkeypox patients are not toxic, have pharyngitis/tonsillitis with cervical adenopathy, and have focal hemorrhage into some lesions (in hemorrhagic smallpox, hemorrhages are extensive/widespread). Patients immunized against smallpox are unlikely to acquire human monkeypox.

**Therapeutic Considerations:** Smallpox vaccination should be initiated as soon as the diagnosis is suspected. Smallpox vaccine may be given at full strength or in a 1:5 dilution, which is also protective. Cidofovir may prove useful but dose for smallpox is not established.

**Prognosis:** Variable in typical smallpox, with deep, permanent scarring, especially on the face. Hemorrhagic smallpox is highly lethal.



## **Ebola/Lassa Fever**

**Clinical Presentation:** After an incubation period of 3–9 days, abrupt onset of high fevers, severe headache/myalgias followed by diarrhea, extreme malaise. Hemorrhagic phenomenon—GI, renal, vaginal, conjunctival bleeding—occur at 5–7 days. Patients rapidly become critically ill. Fever is biphasic. Patients usually have leukopenia, thrombocytopenia, and hepatic/renal dysfunction. Conjunctival suffusion is also an early finding in half the cases. If a patient is not a traveler from an endemic area (e.g., Africa), suspect bioterrorist Ebola/Lassa fever. Lassa fever differs from Ebola in having prominent head/neck edema. CNS finding (oculogyric crisis, seizures, deafness) are characteristic of Lassa fever.

**Diagnostic Considerations:** Ebola is a hemorrhagic fever clinically indistinguishable from Yellow fever and other African hemorrhagic fevers (e.g., Lassa fever, Marburg virus disease). Presumptive diagnosis is clinical; definitive diagnosis is confirmed by specific virologic/serologic studies.

**Pitfalls:** Patients with Ebola may complain initially of a sore throat and dry cough, with or without chest pain. Diarrhea/abdominal pain is not uncommon. The rash is maculopapular before it becomes hemorrhagic. Failure to consider the diagnosis may occur early when sore throat/GI symptoms are prominent (i.e., before hemorrhagic manifestations appear).

**Therapeutic Considerations:** There is no effective therapy available for Ebola infection. Supportive therapy can be life saving.

**Prognosis:** Varies with severity of infection and health of the host.

---

## REFERENCES AND SUGGESTED READINGS

---

- Azad AF. Pathogenic rickettsiae as bioterrorism agents. *Clin Infect Dis* 45 Suppl 1:S52–S55, 2007.
- Alvarez-Lerma F, Grau S, Alvarez-Beltran. Levofloxacin in the treatment of ventilator-associated pneumonia. *Clin Microbiol Infect* 12:81–92, 2006.
- Baddour LM, Bettmann MA, Bolger AF, et al. Nonvascular cardiovascular device-related infections. *Circulation* 108:2015–2031, 2003.
- Bartlett JG, Inglesby TV, JR, Bono L. Management of anthrax. *Clin Infect Dis* 35:851–858, 2002.
- Bouza E, Burillo A, Munoz P. Antimicrobial Therapy of Clostridium difficile-associated diarrhea. *Med Clin N Am* 90:1141–63, 2006.
- Boyce JM. Methicillin-resistant Staphylococcus aureus. *Lancet Infect Dis* 5:653–663, 2005.
- Bradley SF. Infections in long-term-care residents. *Infect Med* 22:168–172, 2005.
- Breman JG, Henderson DA. Diagnosis and management of smallpox. *N Engl J Med* 346:1300–1308, 2002.
- Carratala J, Martin-Herrero JE, Mykietiuk A, et al. Clinical experience in the management of community-acquired pneumonia: lessons from the use of fluoroquinolones. *Clin Microbiol Infect* 12:2–11, 2006.
- Castro P, Soriano A, Escrich C, et al. Linezolid treatment of ventriculoperitoneal shunt infection without implant removal. *Eur J Clin Microbiol Infect Dis* 24:603–06, 2005.
- Celebi G, Baruonu F, Ayoglu F, et al. Tularemia, a reemerging disease in northwest Turkey: epidemiological investigation and evaluation of treatment responses. *Jpn J Infect Dis* 59:229–234, 2006.
- Chalmers JD, Rother C, Salih W, et al. Healthcare-associated pneumonia does not accurately identify potentially resistant pathogens: A systemic review and meta-analysis. *Clin Infect Dis* 58:330–339, 2014.
- Chastre J, Wolff M, Fagon JY, et al. Comparison of 8 vs 15 days of antibiotic therapy for ventilator-associated pneumonia in adults. *JAMA* 290:2588–2598, 2003.
- Chen XM, Leithly JS, Paya CV, et al. Cryptosporidiosis. *N Engl J Med* 346:1723–1731, 2002.
- Chiou CC. Does penicillin remain the drug of choice for pneumococcal pneumonia in view of emerging in vitro resistance? *Clin Infect Dis* 42:234–7, 2006.
- Cunha BA. Antibiotic selection in the penicillin-allergic patient. *Med Clin North Am* 90:1257–64, 2006.
- Cunha BA. Antimicrobial therapy of multidrug-resistant Streptococcus pneumoniae, vancomycin-resistant enterococci, and methicillin-resistant Staphylococcus aureus. *Med Clin North Am* 90:1165–82, 2006.

- Cunha BA. The atypical pneumonias: clinical diagnosis and importance. *Clin Microbiol Infect* 12:12–4, 2006.
- Cunha BA. Oral antibiotic therapy of serious systemic infections. *Med Clin North Am* 90:1197–222, 2006.
- Cunha BA. Ventilator-associated pneumonia: monotherapy is optimal if chosen wisely. *Crit Care* 10:141, 2006.
- Cunha BA. Herpes Simplex-1 (HSV-1) pneumonia. *Infect Dis Practice* 29:375–78, 2005.
- Cunha BA. Malaria vs. Typhoid fever: a diagnostic dilemma? *Am J of Med* 118:1442–43, 2005.
- Cunha BA. Methicillin-resistant *Staphylococcus aureus*: clinical manifestations and antimicrobial therapy. *Clin Microbiol Infect* 11:33–42, 2005.
- Cunha BA. *Pseudomonas aeruginosa*: resistance and therapy. *Semin Respir Infect* 17:231–9, 2002.
- Cunha BA. Clinical relevance of penicillin-resistant *Streptococcus pneumoniae*. *Semin Respir Infect* 17: 204–14, 2002.
- Cunha BA. Smallpox: An Oslerian primer. *Infectious Disease Practice* 26:141–148, 2002.
- Cunha BA. Osteomyelitis in the elderly. *Clin Infect Dis* 35:287–273, 2002.
- Cunha BA. Bioterrorism in the emergency room: Anthrax, tularemia, plague, ebola and smallpox. *Clinical Microbiology & Infection* 8:489–503, 2002.
- Cunha BA. Central nervous system infections in the compromised host. A diagnostic approach. *Infect Dis Clin* 15:67–590, 2001.
- Cunha BA. Effective antibiotic resistance and control strategies. *Lancet* 357:1307–1308, 2001.
- Cunha BA. Nosocomial pneumonia: Diagnostic and therapeutic considerations. *Medical Clinics of North America* 85:79–114, 2001.
- Cunha BA. Community acquired pneumonia: diagnostic and therapeutic considerations. *Medical Clinics of North America* 85:43–77, 2001.
- Cunha BA. Pneumonias in the compromised host. *Infect Dis Clin* 15:591–612, 2001.
- Cunha BA. Community-acquired pneumonias re-visited. *Am J Med* 108:436–437, 2000.
- Daver NG, Shelburne SA, Atmar RL, et al. Oral step-down therapy is comparable to intravenous therapy for *Staphylococcus aureus* osteomyelitis. *Journal of Infection* 54:539–544, 2007.
- Dangerfield B, Chung A, Webb B, et al. Predictive value of methicillin-resistant *Staphylococcus aureus* (MRSA) nasal swab PCR assay for MRSA pneumonia. *Antimicrob Agents Chemother* 58:589–864, 2014.
- Dunning J, Baillie JK, Cao B, et al. Antiviral combinations for severe influenza. *Lancet* 14:1259–1270, 2014.
- Eckburg PB, Schneider JJ, Renault CA. Avian influenza in humans: A practical review for clinicians. *Infect Med* 22:535–542, 2005.
- Esposito S, Bosis S, Semino M, et al. Infections and systemic lupus erythematosus. *Eur J Clin Microbiol & Infect Dis* 33:1467–1475, 2014.
- Fihn SD. Acute uncomplicated urinary tract infection in women. *N Engl J Med* 349:259–66, 2003.
- File TM Jr. Community-acquired pneumonia. *Lancet* 362:1991–2001, 2003.
- Fishbain J, Peleg AY. Treatment of *Acinetobacter* Infections. *Clin Infect. Dis.* 51:79–84, 2010.
- Fishman, JA. Infection in solid-organ transplant recipients. *N Engl J Med* 357:2601–2614, 2007.
- Furin J, Nardell EA. Multidrug-resistant tuberculosis: An update on the best regimens. *J Respir Dis* 27:172–82, 2006.
- Gagliotti C, Nobilio L, Milandri M, et al. Macrolide prescriptions and erythromycin resistance for *Streptococcus pyogenes*. *Clin Infect Dis* 42:1153–6, 2006.
- Gomes CC, Vormittag E, Santos CR, Levin AS. Nosocomial infection with cephalosporin-resistant *Klebsiella pneumoniae* is not associated with increased mortality. *Infect Control and Hosp Epidemiol* 27:907–12, 2006.
- Gould IM. The clinical significance of methicillin-resistant *Staphylococcus aureus*. *J Hosp Infect* 61:277–282, 2005.
- Gupta R, Warren T, Wald A. Genital herpes. *Lancet* 370: 2127–2137, 2007.
- Haque NZ, Zervos MJ. Vancomycin-Resistant Enterococcal Infections: clinical manifestations and Management. *Infections in Medicine* 23:14–19, 2006.
- Hayakawa K, Nakagawa K. Treatment of infected total knee arthroplasty. *Infect Diseases in Clin Practice* 14:211–215, 2006.
- Heldman AW, Harter TV, Ray SC, et al. Oral antibiotic treatment of right-sided staphylococcal endocarditis in injection drug users: prospective randomized comparison with parenteral therapy. *Am J Med* 101: 68–76, 1996.
- Johnson DH, Cunha BA. Infections in alcoholic cirrhosis. *Infectious Disease Clinics* 16:363–372, 2001.
- Johnson LB, Saravolatz LD. Community-acquired MRSA: current epidemiology and management issues. *Infections in Medicine* 23:6–10, 2006.
- Joseph SM, Peiris MD, D. Phil et al. The severe acute respiratory syndrome. *N Engl J Med* 349:2431–41, 2003.
- Kallen AJ, Mu Y, Bulens S, et al. Health care-associated invasive *MRSA* infections, 2005–2008. *JAMA* 304: 641–648, 2010.

- Kanafani ZA, Khalifeh N, Kanj SS, Araj GF, Khalifeh M, Sharara AI. Antibiotic use in acute cholecystitis: practice patterns in the absence of evidence-based guidelines. *J Infect* 51:128–134, 2005.
- Kassakian SZ, Mermel LA. Changing epidemiology of infections due to extended spectrum betalactamase producing bacteria. *Antimicrob Resist Infect Control* 25:9, 2014.
- Kauffman CA. Managing fungal pneumonias: a review of the new therapies. *J Crit Illness* 20:30–35, 2005.
- Kauffman CA. Endemic mycoses in patients with hematologic malignancies. *Semin Respir Infect* 17:106–12, 2002.
- Kawai N, Idematsu H, Iwaki N, et al. Comparisons of the effectiveness of zanamivir and oseltamivir against influenza A/H<sub>1</sub>N<sub>1</sub>, A/H3N2, and B. *Clin Infect Dis* 48:996–99, 2009.
- Khadori N, Kanchanapoom T. Overview of biological terrorism: Potential agents and preparedness. *Clin Microbiol News* 27:1, 2005.
- Kimberlin DW, Rouse DJ. Genital herpes. *N Engl J Med* 350:1970–7, 2004.
- Kmeid JG, Youssef MM, Kanafani ZA, et al. Combination therapy for Gram-negative bacteria: what is the evidence? *Expert Rev Anti Infect Ther* 11:1355–1362, 2013.
- Kohli A, Shaffer A, Sherman A, et al. Treatment of hepatitis C. A systematic review. *J Am Med Assoc.* 312:631–640, 2014.
- Lamas C. Diagnostic strategy for blood culture-negative endocarditis. *Clin Infect. Dis.* 51:141–142, 2010.
- Leather HL, Wingard JR. Infections following hematopoietic stem cell transplantation. *Infect Dis Clin North Am* 15:483–520, 2001.
- Linden PK. Treatment options for vancomycin-resistant enterococcal infections. *Drugs* 62: 425–41, 2002.
- Lipsky BA, Byren I, Hoey CT. Treatment of bacterial prostatitis. *Clin Infect Dis.* 50:1641–1652, 2010.
- Loddenkemper R, Sagebiel D, Brendel A. Strategies against multidrug-resistant tuberculosis. *Eur Respir J Suppl* 36:66s–77s, 2002.
- Maartens G, Willinson RJ. Tuberculosis. *Lancet* 370:2030–2043, 2007.
- Martin-Pena A, Aguilar-Guisado M, Espigado I, Cisneros JM. Antifungal combination therapy for invasive aspergillosis. *Clin Infect Dis* 59:1437–1445, 2014.
- McLaughlin SP, Carson CC. Urinary tract infections in women. *Med Clin North Am* 88:417–29, 2004.
- Mead PA, Safdieh JE, Nizza P, et al. Ommaya reservoir infections: A 16 year retrospective analysis. *Journal of Infection* 68:225–230, 2014.
- Meddings J, Rogers MAM, Macy M, et al. Systematic review and meta-analysis: reminder systems to reduced catheter-associated urinary tract infections and urinary catheter use in hospitalized patients. *Clin Infect Dis.* 51:550–560, 2010.
- Mermel LA. What is the Predominant Source of Intravascular Catheter Infections? *Clin Infect Dis* 52:211–212, 2011.
- Metlay JP, Fishman NO, Joffe MM, et al. Macrolide resistance in adults with bacteremic pneumococcal pneumonia. *Emerging Infect Dis* 12:1223–30, 2006.
- Michalopoulos A, Falagas ME. Treatment of Acinetobacter infections. *Expert Opin Pharmacotherapy.* 11:779–788, 2010.
- Minnaganti V, Cunha BA. Infections associated with uremia and dialysis. *Infect Dis Clin* 16:385–406, 2001.
- Musher DM, Musher BJ. Contagious acute gastrointestinal infections. *N Engl J Med* 351:2417–27, 2004.
- Musher DM, Aslam S, Logan N, et al. Relatively poor outcome after treatment of Clostridium difficile colitis with metronidazole. *Clin Infect Dis* 40:1586–1590, 2005.
- Mylonakis E, Calderwood SB. Infective endocarditis in adults. *N Engl J Med* 345:1318, 2001.
- Nicholson KG, Wood JM, Zambon M. Influenza. *Lancet* 362:1733–45, 2003.
- Nucci M, Anaissie E, Betts RF, et al. Early removal of central venous catheter in patients with candidemia does not improve outcome: analysis of 842 patients from 2 randomized clinical trials. *Clin Infect. Dis.* 51:295–303, 2010.
- Parry CM, Hien TT, Dougan G, et al. Typhoid fever. *N Engl J Med* 347:1770–82, 2002.
- Patterson KC, Streck M. Diagnosis and treatment of pulmonary aspergillosis syndromes. *Chest* 146:1358–1368, 2014.
- Peipert JF. Genital Chlamydial Infections. *N Engl J Med* 349:2424–30, 2003.
- Perea S, Patterson TF. Invasive Aspergillus infections in hematologic malignancy patients. *Semin Respir Infect* 17:99–105, 2002.
- Perencevich EN, Diekema DJ. Decline in invasive MRSA infection. where to go from here? *JAMA* 304: 687–689, 2010.
- Peterson LR. Penicillin for treatment of pneumococcal pneumonia: does In vitro resistance really matter? *Clin Infect Dis* 42:224–33, 2006.
- Pontoizeau C, Dangers L, Jarlier V, et al. Ruling out False Positive Urinary Legionella pneumophila serogroup 1 and Streptococcus pneumonia antigen test results by heating urine. *J Clin Micro* 52:4347–4349, 2014.
- Razonable RR. Management of cytomegalovirus infection after renal transplantation. *Infect Dis in Clinical Practice* 17:220–230, 2009.

- Rex JH. Approach to the treatment of systemic fungal infections. In: Kelley WN (ed), *Kelley's Textbook of Internal Medicine*, 4<sup>th</sup> ed., 2279–2281, 2000.
- Rodriguez-Bano J. Selection of empiric therapy in patients with catheter-related infections. *Clin Microbiol Infect* 8:275–81, 2002.
- Ross AGP, Bartley PB, Sleight AC, et al. Schistosomiasis. *N Engl J Med* 346:1212–1220, 2002.
- Rovers MM, Schilder AGM, Zielhuis GA, et al. Otitis media. *Lancet* 363:465–73, 2004.
- Ruhe JJ, Monson T, Bradsher RW, Menon A. Use of long-acting tetracyclines for methicillin-resistant *Staphylococcus aureus* infections: case series and review of the literature. *Clin Infect Dis* 40:1429–1434, 2005.
- Safdar A, Bryan CS, Stinfor S, et al. Prosthetic valve endocarditis due to vancomycin-resistant *Enterococcus faecium*: treatment with chloramphenicol plus minocycline. *Clin Infect Dis* 34:61–3, 2002.
- Sagnelli E, Santantonio T, Coppola N, et al. Acute hepatitis C: clinical and laboratory diagnosis, course of the disease, treatment. *Infection* 42:601–610, 2014.
- Schoen RT. Better laboratory testing for Lyme disease: No more western blot. *Clin Infect Dis* 57:341–343, 2013.
- Slenczka W, Klenk HD. Fort years of Marburg virus. *J Infect Dis* 196:S131–S135, 2007.
- Sobel JD. Botulism. *Clin Infect Dis* 41:1167–1173, 2005.
- Sobel JD, Wiesenfeld HC, Martens M, et al. Maintenance fluconazole therapy for recurrent vulvovaginal candidiasis. *N Engl J Med* 351:876–83, 2004.
- Spellberg BJ, Filler SG, Edwards JE. Current treatment strategies for disseminated candidiasis. *Clin Infect Dis* 42:244–251, 2006.
- Steffen R, Hill DR, DuPont HL. Traveler's diarrhea: A clinical review. *JAMA* 313:71–80, 2015.
- Szvalb AD, Raad II, Weinberg JS, et al. Ommaya reservoir-related infections: The clinical manifestations and treatment outcomes. *Infection* 68:216–224, 2014.
- Tegnell A, Wahren B, Elgh G. Smallpox - eradicated, but a growing terror threat. *Clin Microbiol Infect* 8:504–509, 2002.
- Teshae EH, Hu DJ, Holmberg SD. The two faces of hepatitis E virus. *Clin Infect Dis* 51:328–334, 2010.
- Thielman NM, Guerrant RL. Acute infectious diarrhea. *N Engl J Med* 350:38–47, 2004.
- Tolkoff-Rubin NE, Rubin RH. Recent advances in the diagnosis and management of infection in the organ transplant recipient. *Semin Nephrol* 20:148–163, 2000.
- Tunkel AR, Hartman BJ, Kaplan SL, et al. Practical guidelines for the management of bacterial meningitis. *Clin Infect Dis* 39:1267–84, 2004.
- van de Beek D, de Gans J, Spanjaard L, et al. Clinical features and prognostic factors in adults with bacterial meningitis. *N Engl J Med* 351:1849–59, 2004.
- van de Beek D, de Gans J, Tunkel AR, Wijdicks EFM. Community-acquired bacterial meningitis in adults. *N Engl J Med* 354:44–53, 2006.
- Viale P, Stefani S. Vascular catheter-associated infections: a microbiological and therapeutic update. *Journal of Chemotherapy* 18:235–49, 2006.
- Walsh TJ, Rex JH. All catheter-related candidemia is not the same: assessment of the balance between the risks and benefits of removal of vascular catheters. *Clin Infect Dis* 34:600–602, 2002.
- Wikie MD, Hanson MF, Statham PF, Brennan PM. Infections of cerebrospinal fluid diversion devices in adults: The role of intraventricular antimicrobial therapy. *J Infection* 63:239–246, 2013.
- Wormser, GP. Discovery of new infectious diseases *Bartonella* species. *N Engl J Med* 356:2346–2347, 2007.
- Wormser, GP. Early Lyme Disease. *N Engl J Med* 354:2794–801, 2006.
- Wright AJ, Fishman JA. Central nervous system syndromes in solid organ transplant recipients. *Clin Infect Dis* 59:1001–1011, 2014.

## GUIDELINES

- Bisno AL, Gerber MA, Gwaltney JM Jr, Kaplan EL, Schwartz RH. Practice guidelines for the diagnosis and management of group A streptococcal pharyngitis. *Infectious Diseases Society of America. Clin Infect Dis* 35:113–25, 2002.
- CDC MMWR. Sexually transmitted diseases treatment guidelines, 2010. (RR-12) 59:1–110, 2010.
- Chapman SW, Bradsher RW Jr, Campbell CD Jr, et al. Practice guidelines for the management of patients with blastomycosis. *Infectious Diseases Society of America. Clin Infect Dis* 30:679–83, 2000.
- Cordonnier C, Pautas C, Maury S, Vekhoff A, et al. Empirical versus preemptive antifungal therapy for high-risk, febrile, neutropenic patients: A randomized, controlled trial. *Clin Infect Dis* 48:1042–1051, 2009.
- Dworkin RH, Johnson RW, Breuer J. Recommendations for the management of Herpes zoster. *Clin Infect Dis* 44:51–26, 2007.
- Dykevicz CA. Summary of the guidelines for preventing opportunistic infectious among hematopoietic stem cell transplant recipients. *Clin Infect Dis* 33:139–44, 2001.
- File TM. Recommendations for treatment of hospital-acquired and ventilator-associated pneumonia: review of recent international guidelines. *Clin Infect Dis* 51:S42–S47, 2010.

- Freifeld A, Bow E, Sepkowitz K, et al. Clinical Practice Guideline for the Use of Antimicrobial Agents in Neutropenic Patients with Cancer: 2010 Update by the Infectious Diseases Society of America. *Clin Infect Dis* 52:e56–e93, 2011.
- Galgiani JN, Ampel NM, Blair JE, et al. Coccidioidomycosis. *Clin Infect Dis* 41:1217–23, 2005.
- Goodman EL. Practice guidelines for evaluating new fever in critically ill adult patients. *Clin Infect Dis* 30:234, 2000.
- Guerrant RL, Van Gilder T, Sterner TS, et al. Practice guidelines for the management of infectious diarrhea. *Clin Infect Dis* 15:321–4, 2001.
- Gupta K, Hooton T, Naber K, et al. International Clinical Practice Guidelines for the Treatment of Acute Uncomplicated Cystitis and Pyelonephritis in Women: A 2010 Update by the Infectious Diseases Society of America and the European Society for Microbiology and Infectious Diseases. *Clinical Infectious Diseases*. 52:e103–e120, 2011.
- Horsburgh CR Jr, Feldman S, Ridzon R. Practice guidelines for the treatment of tuberculosis. *Clin Infect Dis* 31:633–9, 2000.
- Hughes WT, Armstrong D, Bodey GP, et al. Guidelines for the use of antimicrobial agents in neutropenic patients with cancer. *Clin Infect Dis* 34:730–51, 2002.
- Kauffman CA, Hajjeh R, Chapman SW. Practice guidelines for the management of patients with sporotrichosis. For the mycoses study group. *Infectious Diseases Society of America. Clin Infect Dis* 30:684–7, 2000.
- Krol V, Hamid NS, Cunha BA. Neurosurgically related nosocomial *Acinetobacter baumannii* meningitis: report of two cases and literature review. *J Hospital Infection*. 71:176–180, 2009.
- Lipsky BA, Berendt AR, Deery HG, et al. Diagnosis and treatment of diabetic foot infections. *Infect Dis* 39:885–910, 2004.
- Liu C, Bayer A, Cosgrove S, et al. Clinical Practice Guidelines by the Infectious Diseases Society of America for the Treatment of Methicillin-Resistant *Staphylococcus Aureus* Infections in Adults and Children. *Clinical Infectious Diseases*. 52:1–38, 2011.
- Mandell LA, Wunderink RG, Anzueto A, et al. Infectious diseases society of america/american thoracic society consensus guidelines on the management of community-acquired pneumonia in adults. *Clin Infect Dis* 44(S2):S27–S72, 2007.
- Marr K, Boeckh M. Practice guidelines for fungal infections: a risk-guided approach. *Clin Infect Dis* 32:331–51, 2001.
- Mermel LA, Allon M, Bouza E, Craven DE, et al. Clinical practice guidelines for the diagnosis and management of intravascular catheter-related infection: 2009 update by the Infectious Diseases Society of America. *Clin Infect Dis* 49:1–45, 2009.
- Nicolle LE, Bradley S, Colgan R, et al. Infectious Diseases Society of America guidelines for the diagnosis and treatment of asymptomatic bacteriuria in adults. *Clin Infect Dis* 40:643–54, 2005.
- O'Grady NP, Barie PS, Bartlett JG, et al. Practice guidelines for evaluating new fever in critically ill adult patients. *Clin Infect Dis* 26:1042–1059, 1998.
- Pappas PG, Rex JH, Sobel JD, et al. Guidelines for treatment of candidiasis. *Clin Infect Dis* 38:161–89, 2004.
- Pappas PG, Kauffman CA, Andes D, et al. Clinical practice guidelines for the management of candidiasis: 2009 update by the Infectious Diseases Society of America. *Clin Infect Dis* 48:503–535, 2009.
- Rolston KV. The Infectious Diseases Society of America 2002 guidelines for the use of antimicrobial agents in patients with cancer and neutropenia: salient features and comments. *Clin Infect Dis* 39 Suppl 1:S44–8, 2004.
- Rubin LG, Levin MJ, Ljungman P, et al. 2013 IDSA Clinical Practice Guideline for Vaccination of the Immunocompromised Host. *Clin Infect Dis* 58:309–318, 2014.
- Saag MS, Graybill RJ, Larsen RA, Pappas PG, et al. Practice guidelines for the management of cryptococcal disease. *Infectious Diseases Society of America. Clin Infect Dis* 30:710–8, 2000.
- Sobel JD. Practice guidelines for the treatment of fungal infections. For the mycoses study group. *Infectious diseases society of america. Clin Infect Dis* 30:652, 2000.
- Solomkin JS, Mazuski JE, Bradley JS, et al. Diagnosis and management of complicated intra-abdominal infection in adults and children: guidelines by the surgical infection society and the infectious diseases society of america. *Clin Infect Dis* 50:133–164, 2010.
- Solomon JS, Mazuski JE, Baron EJ, et al. Guidelines for the selection of anti-infective agents for complicated intra-abdominal infections. *Clin Infect Dis* 37:997–1005, 2003.
- Stevens DA, Dan VL, Judson MA, Morrison VA, et al. Practice guidelines for diseases caused by *Aspergillus*. *Infectious diseases society of america. Clin Infect Dis* 30:696–709, 2000.
- Stevens DL, Bisno AL, Chambers HF, et al. Practice guidelines for the diagnosis and management of skin and soft tissue infections. *Clin Infect Dis* 41:1373–1406, 2005.
- Stevens DL, Bisno AL, Chambers HF, et al. Practice guidelines for the diagnosis and management of skin and soft tissue infections. *Clin Infect Dis* 59:147–159, 2014.
- Talmor M, Li P, Barie PS. Acute paranasal sinusitis in critically ill patients: guidelines for prevention, diagnosis, and treatment. *Clin Infect Dis* 25:1441–6, 1997.

- Tice AD, Rehm SJ, Dalovisio JR, et al. Practice guidelines for outpatient parenteral antimicrobial therapy. IDSA guidelines. *Clin Infect Dis* 38:1651–72, 2004.
- Tunkel AR, Hartman BJ, Kaplan SL, et al. Practice guidelines for the management of bacterial meningitis. *Clin Infect Dis* 39:1267–1284, 2004.
- Wheat LJ, Freifeld AG, Kleiman MB, et al. Clinical practice guidelines for the management of patients with histoplasmosis: 2007 update by the Infectious Diseases Society of America. *Clin Infect Dis* 45:807–25, 2007.
- Wheat LJ, Musial CE, Jenny-Avital E. Diagnosis and management of central nervous system histoplasmosis. *Clin Infect Dis* 40:844–852, 2005.
- Workowski KA. Sexually transmitted diseases treatment guidelines. *Clin Infect Dis* 44:573–174, 2007.
- Wormser GP, Nadelman RB, Batwyler RJ, et al. Practice guidelines for the treatment of Lyme disease. Infectious Disease Society of America. *Clin Infect Dis* 31 (Suppl 1): 1–14, 2000.
- Yu VL, Ramirez J, Roig J, et al. Legionnaires disease and the updated IDSA guidelines for community-acquired pneumonia. *Clin Infect Dis* 39:1734–7, 2004.
- Cimolai N (ed). *Laboratory Diagnosis of Bacterial Infections*. New York, Marcel Dekker, 2001.
- Cohen J, Powderly WG, Opal S (Eds). *Infectious Diseases* (4th Ed). Elsevier, Philadelphia 2015.
- Cunha BA (ed). *Infectious Disease in Critical Care Medicine* (3rd Ed), Informa Healthcare, New York, 2010.
- Cunha BA (ed). *Tick-Borne Infectious Diseases*. Marcel Dekker, New York, 2005.
- Cunha BA (ed). *Infectious Disease in the Elderly*. John Wright & Co., London, 1988.
- Despommier DD, Gwadz RW Hotez PJ, Knirsh CA (eds). *Parasitic Diseases* (5th Ed), Apple Tree Productions, LLC, New York, New York, 2005.
- Faro S, Soper DE (eds). *Infectious Diseases in Women*. WB Saunders Company, Philadelphia, 2001.
- Farrar J (eds). *Manson's Tropical Diseases* (23rd Ed). Elsevier Saunders, London, 2014.
- Glauser MP, Pizzo PA (eds). *Management of Infection in Immunocompromised Patients*. W.B. Saunders, London, 2000.
- Gorbach SL, Bartlett JG, Blacklow NR (eds). *Infectious Diseases* (3rd Ed), Philadelphia, Lippincott, Williams & Wilkins, 2004.
- Grayson ML (ed). *Kucers' The Use of Antibiotics* (6th Ed), ASM Press, Washington, DC, 2010.
- Guerrant RL, Walker DH, Weller PF (eds). *Tropical Infectious Disease: Principles, Pathogens & Practice* (3rd Ed), Elsevier, Philadelphia, 2011.
- Halperin JJ. *Encephalitis: Diagnosis and Treatment*. Informa Healthcare, New York, 2008.
- Hauser AR, Rello J (eds). *Severe Infections Caused by Pseudomonas Aeruginosa*. Kluwer Academic Publishers, Boston, Massachusetts, 2003.
- Koff RS *Hepatitis Essentials*, Jones & Bartlett, Sudbury, MA, 2011.
- Madkour MM (ed). *Tuberculosis*. Springer-Verlag, Berlin Germany, 2004.
- Maertens JA, Marr KA. *Diagnosis of Fungal Infections*. Informa Healthcare, New York, 2007.
- McMillan A, Young H, Ogilvie MM, Scott GR (eds). *Clinical practice in Sexually Transmissible Infections*, Saunders, London, England, 2002.
- Pilch RF, Ziliinskas RA (eds). *Encyclopedia of Bioterrorism Defense*. Wiley-Liss, Hoboken, New Jersey, 2005.
- Raoult D, Parola P. *Rickettsial Diseases*. Informa Healthcare, New York, 2007.
- Rom WN, Garay SM (eds). *Tuberculosis*. Lippincott Williams & Wilkins, Philadelphia, Pennsylvania, 2004.

## TEXTBOOKS

- Arikan S Rex JH (eds). *Antifungal Drugs in Manual of Clinical Microbiology* (8th Ed), 2003.
- Baddour L, Gorbach SL (eds). *Therapy of Infectious Diseases*. Saunders, Philadelphia, Pennsylvania, 2003.
- Bennett JE, Dolin R, Blaser MJ (eds). *Mandell, Douglas, and Bennett's Principles and Practice of infectious Diseases* (8th Ed), Elsevier Churchill Livingstone, 2015.
- Bodey GP, Feinstein V (eds). *Candidiasis*. New York, Raven Press, 1985.
- Bowden RA, Ljungman P, Snyderman DR (eds). *Transplant Infections* (3rd Ed), Lippincott Williams & Wilkins, Philadelphia, PA, 2010.
- Brandstetter R, Cunha BA, Karetsky M (eds). *The Pneumonias*. Mosby, Philadelphia, 1999.
- Brook I (ed). *Sinusitis*. Taylor & Francis Group, New York, New York, 2006.
- Brusch JL *Endocarditis Essentials*. Jones & Bartlett, Sudbury, MA, 2010.
- Brusch, JL. *Infective Endocarditis: Management in the Era of Intravascular Devices*. Informa Healthcare, New York 2007.
- Bryskier A (ed). *Antimicrobial Agents*. ASM Press, Washington, D.C., 2005.
- Calderone RA (ed). *Candida and Candidiasis*. ASM Press, Washington DC, 2002.

- Schedl W, Whitley R, Marra C (eds). *Infections of the Central Nervous System* (3<sup>rd</sup> Ed), Lippincott Williams & Wilkins, Philadelphia, 2005.
- Schlossberg D (ed). *Tuberculosis & Nontuberculous Mycobacterial Infections* (6<sup>th</sup> Ed), New York, McGraw-Hill, 2011.
- Schlossberg D (ed). *Clinical Infections Diseases* (2<sup>nd</sup> Ed), Cambridge University Press, London, 2015.
- Schlossberg D (ed). *Current Therapy of Infectious Disease* (3<sup>rd</sup> Ed), Mosby-Yearbook, St. Louis, 2008.
- Schlossberg D (ed). *Medical Interventions for Bioterrorism and Emerging Infections. Handbooks in Healthcare Co.*, Newtown, Pennsylvania, 2004.
- Singh N, Aguado JM (eds). *Infectious Complications in Transplant Patients*. Kluwer Academic Publishers, Boston, 2000.
- Studahl M, Cinque P, Bergstrom T (eds). *Herpes Simplex Viruses*. Taylor & Francis Group, New York, New York, 2006.
- Wingard JR, Anaissie EL (eds). *Fungal Infections in the Immunocompromised Patient*. Taylor & Francis Group, Boca Raton, Florida, 2005.
- Woods JB (ed). *USAMRIID's Medical management of biological casualties handbook* (6<sup>th</sup> Ed), US Army Medical Research Institute of Infectious Disease, Fort Detrick, Frederick, Maryland, 2005.
- Yoshikawa TT, Rajagopalan S (eds). *Antibiotic Therapy for Geriatric Patients*. Taylor & Francis Group, New York, New York, 2006.
- Yu, V, Edwards G, McKinnon PS, Peloquin C, Morse G (eds). *Antimicrobial Therapy and Vaccines, Volume II: Antimicrobial Agents* (2<sup>nd</sup> Ed), ESun Technologies, Pittsburgh, Pennsylvania, 2005.

## Chapter 3

# Antibiotic Susceptibility Profiles and Initial Therapy of Isolates Pending Susceptibility Results

**Burke A. Cunha, MD, Paul E. Schoch, PhD, Edward J. Bottone, PhD  
John H. Rex, MD, Cheston B. Cunha, MD**

## Antibiotic Susceptibility Profiles (Tables 3.1–3.3).

Table 3.1. <b>Antibiotic Susceptibility Profiles</b> (Penicillins, Macrolides, Tetracyclines, Clindamycin, Metronidazole, Rifampin, TMP-SMX, and Chloramphenicol) .....	186
Table 3.2. <b>Antibiotic Susceptibility Profiles</b> (Cephalosporins) .....	191
Table 3.3. <b>Antibiotic Susceptibility Profiles</b> (Aminoglycosides, Fluoroquinolones, Colistin, Polymyxin B, Aztreonam, Tigecycline, Vancomycin, Quinupristin/Dalfopristin, Linezolid, Tedizolid, Daptomycin, Telavancin, Dalbavancin, Oritavancin, Nitrofurantoin, Fosfomycin, and Carbapenems) ...	198

## Gram Stain Characteristics of Isolates (by Morphology, Arrangement,

Oxygen Requirements) .....	203-206
Table 3.4. <b>Key Factors in Antibiotic Selection</b> (Isolate Known) .....	207
Table 3.5. <b>Antibiotic Selection Based on Resistance Potential</b> .....	207

## To determine the Clinical Significance of an Isolate in Selecting Appropriate Initial Antibiotic Therapy Pending Susceptibility Testing (Tables 3.6–3.9)

Table 3.6. <b>Clinical Significance of AEROBIC Isolates</b> Pending Susceptibility Testing .....	208
Table 3.7. <b>Clinical Significance of CAPNOPHILIC Isolates</b> Pending Susceptibility Testing .....	235
Table 3.8. <b>Clinical Significance of ANAEROBIC Isolates</b> Pending Susceptibility Testing .....	236
Table 3.9. <b>Clinical Significance of YEAST/FUNGI</b> Pending Susceptibility Testing .....	242
Table 3.10. Technique for Gram Stain and Giemsa Stain .....	246
Table 3.11. Clinical Use of CSF Gram Stain, WBC Type, Glucose .....	246
Table 3.12. Clinical Use of the Sputum Gram Stain .....	248
Table 3.13. Clinical Use of the Urine Gram Stain .....	248
Table 3.14. Clinical Use of the Fecal Gram Stain .....	249
References and Suggested Readings .....	249

If isolate is a pathogen (P), in the appropriate body site, pathogens should be treated. Pathogens isolated from body sites not causing infections *at the site cultured* should ordinarily not be treated, e.g., *S. pneumoniae* from urine or a wound. Non-pathogens (NP), colonizers (C), or skin contaminants (C\*) ordinarily should not be "covered" or treated.



Table 3.1. Antibiotic Susceptibility Profiles (Penicillins, Macrolides, Tetracyclines, and Others)

ORGANISMS	Penicillins						Anti-Pseudomonal Penicillins				Macrolides				Tetra-Cyclines			Miscellaneous					
	Penicillin G (IV)	Penicillin V (PO)	Ampicillin (IV/PO)	Ampicillin/Subactam (IV)	Amoxicillin (PO)	Amoxicillin/Clavulanate (PO)	Nafcillin (IV)	Ticarcillin (IV)	Ticarcillin/Clavulanate (IV)	Piperacillin (IV)	Piperacillin/ tazobactam (IV)	Erythromycin (IV/PO)	Clarithromycin (PO)	Azithromycin (IV/PO)	Telithromycin (PO)	Tetracycline (IV/PO)	Doxycycline (IV/PO)	Minocycline (IV/PO)	Cindamycin (IV/PO)	Metronidazole (IV/PO)	Rifampin (PO)	TMP-SMX (IV/PO)	Chloramphenicol (IV/PO)
<b>Aerobic Gram Positive Cocci (Clusters)</b>																							
<i>Staphylococcus aureus</i> (MSSA)	0	0	1	2	0	2	3	0	0	3	3	0	0	0	0	2	1	1	1	0	2	1	2
<i>Staphylococcus aureus</i> (HA/CO-MRSA)	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2 <sup>†</sup>	1	2 <sup>†</sup>	2 <sup>†</sup>	0	2	3 <sup>†</sup>	0
<i>S. aureus</i> (CA-MRSA)	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2	1	2	2	0	0	2	0
<i>S. epidermidis</i> (CoNS)	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2	2	2	2	0	2	2	0
<b>Aerobic Gram Positive Cocci (Chains)</b>																							
<i>Enterococcus faecalis</i> (VSE)	0	0	1	2	1	1	0	2	2	2	3	0	0	0	0	0	0	0	0	0	0	3	3
<i>Enterococcus faecium</i> (VRE) <sup>††</sup>	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	3	2	0	0	0	0	0	3
Streptococci (Groups A, B, C, F, G)	1	1	2	2	1	2	0	2	2	2	3	3	3	0	0	0	0	1	0	0	0	0	3

Streptococcus (bovis) galloyticus	1	1	2	2	1	2	0	2	2	2	2	3	0	0	0	0	0	0	0	0			
<i>Viridans streptococci</i> ( <i>S. mitior, milleri, mitis, mutans, oralis, sanguis, parasanguis, salivarius</i> )	1	1	2	2	1	2	0	2	2	2	3	3	3	3	0	0	0	2	0	0	3		
<b>Aerobic Gram Positive Cocci (Pairs)</b>																							
<i>Streptococcus pneumoniae</i> (PSSP)	1	1	3*	2	1	2	0	2	2	2	2	2	3*	3*	2	3*	1	2	1	0	3*	3	
<i>Streptococcus pneumoniae</i> (PRSP)	1	0	0*	2	1	2	0	3	3	3	3	0	0	0	2	0	1	1	2	0	2	0	3
<i>Streptococcus pneumoniae</i> (MDRSP)	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2	0	2	3	3	0	3	0	3
<b>Aerobic Gram Negative Cocci (Pairs)</b>																							
<i>Neisseria gonorrhoeae</i>	0	0	0	2	0	2	0	0	2	0	2	3	0	2	0	3*	2	2	0	0	2	0	3
<i>Neisseria meningitidis</i>	1	2	2	2	2	2	0	2	0	2	0	0	0	0	0	0	3	3	0	0	2	0	2
<b>Aerobic Gram Positive Bacilli</b>																							
<i>Bacillus anthracis</i>	2	3	2	3	2	3	0	0	0	0	0	3	3	3	0	2	1	0	2	0	2	0	2
<i>Corynebacterium diphtheriae</i>	2	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0

1 = Preferred antibiotic—High degree of activity against the isolate and likely to be clinically effective, most strains susceptible, has a favorable PK/PD characteristics, has a good safety profile, and has a “low resistance” potential (see Chapter 11 Drug Summaries for antibiotic dosing details).  
 2 = Alternate choice—With many attributes of a preferred antibiotic. 3 = Acceptable, but preferably select an alternate antibiotic. 0 = No activity or no data or limited experience. \* = May be effective initially but as a “high resistance” antibiotic potential, resistance may develop during/after therapy.  
 † = Preferably use another CA-MRSA antibiotic, e.g., minocycline. †† = Same for vancomycin resistant *E. faecalis*.



<i>Escherichia coli</i>	0	0	3*	2	2	2	2	2	2	0	0	0	0	2	3	0	0	0	3	2
<i>Francisella tularensis</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2	0	0	0	3	3
<i>Haemophilus sp.</i>	0	0	3*	2	2	0	3	2	3	2	0	3	3	2	3*	1	3	0	3	2
<i>Klebsiella sp.</i>	0	0	0	2	0	3	0	3	3	3	0	0	0	0	0	0	0	0	0	3
<i>Moraxella catarrhalis</i>	0	0	0	3	0	3	0	3	2	3	2	0	0	3	2	3	1	3	0	3*
<i>Morganella sp.</i>	0	0	0	3	0	0	3	3	3	3	0	0	0	0	0	0	0	0	0	3*
<i>P. aeruginosa</i>	0	0	0	0	0	0	3	3	3	3	0	0	0	0	0	0	0	0	0	0
<i>Proteus sp.</i>	0	0	0	3	0	0	3	3	3	3	0	0	0	0	0	0	0	0	0	3*
<i>Providencia sp.</i>	0	0	0	3	0	0	3	3	3	3	0	0	0	0	0	0	0	0	0	3*
<i>Salmonella sp.</i>	0	0	0	3	3	3	0	3	3	3	0	0	3	0	3*	2	0	0	0	3
<i>Serratia marcescens</i>	0	0	0	0	0	0	0	3	0	3	0	0	0	0	0	0	0	0	0	3
<i>Shigella sp.</i>	0	0	0	3	3	3	0	3	3	3	0	0	3	0	3	0	0	0	0	3*
<i>Stenotrophomonas (Pseudomonas) maltophilia</i>	0	0	0	0	0	0	0	3	0	3	0	0	0	0	0	2	0	0	0	2
<i>Vibrio vulnificus</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0	3	1	3	0	0	3
<i>Yersinia enterocolitica</i>	0	0	0	2	0	3	0	3	2	2	0	0	0	0	0	0	0	0	0	1
<b>Anaerobic Gram Positive Cocci (chains)</b>																				
<i>Peptostreptococcus</i>	1	1	2	2	2	2	0	2	2	2	2	3	3	3	0	3	2	2	2	0

1 = Preferred antibiotic—High degree of activity against the isolate and likely to be clinically effective, most strains susceptible, has a favorable PK/PD characteristics, has a good safety profile, and has a “low resistance” potential (see Chapter 11 Drug Summaries for antibiotic dosing details). 2 = Alternate choice—With many attributes of a preferred antibiotic. 3 = Acceptable, but preferably select an alternate antibiotic. 0 = No activity or no data or limited experience. \* = May be effective initially but as a “high resistance” antibiotic potential, resistance may develop during/after therapy.

§ = Always use in combination with another antibiotic.

Table 3.1. Antibiotic Susceptibility Profiles (Penicillins, Macrolides, Tetracyclines, and Others) (cont'd)

ORGANISMS	Penicillins						Anti-Pseudomonal Penicillins				Macrolides				Tetra-Cyclines				Miscellaneous				
	Penicillin G (IV)	Penicillin V (PO)	Ampicillin (IV/PO)	Ampicillin/Sulbactam (IV)	Amoxicillin (PO)	Amoxicillin/Clavulanate (PO)	Nafcillin (IV)	Ticarcillin (IV)	Ticarcillin/Clavulanate (IV)	Piperacillin (IV)	Piperacillin/Tazobactam (IV)	Erythromycin (IV/PO)	Clarithromycin (PO)	Azithromycin (IV/PO)	Telithromycin (PO)	Tetracycline (IV/PO)	Doxycycline (IV/PO)	Minocycline (IV/PO)	Clindamycin (IV/PO)	Metronidazole (IV/PO)	Rifampin (PO) <sup>§</sup>	TMP-SMX (IV/PO)	Chloramphenicol (IV/PO)
<b>Anaerobic Gram Positive Bacilli</b>																							
<i>Actinomyces</i> sp.	1	1	2	2	1	2	0	0	0	0	2	2	2	2	2	2	2	2	0	0	0	0	0
<b>Anaerobic Gram Negative Bacilli</b>																							
<i>Bacteroides fragilis</i> group ( <i>B. distasonis</i> , <i>ovatus</i> , <i>thetaiotaomicron</i> , <i>vulgatus</i> )	0	0	2	2	0	2	0	2	2	2	3	3	3	0	3	2	2	1	1	0	0	0	2
<i>Prevotella</i> sp.	1	1	1	1	1	1	0	2	2	2	2	2	2	2	2	2	2	1	1	0	0	0	2

1 = Preferred antibiotic—High degree of activity against the isolate and likely to be clinically effective, most strains susceptible, has a favorable PK/PD characteristics, has a good safety profile, and has a “low resistance” potential (see Chapter 11 Drug Summaries for antibiotic dosing details).  
 2 = Alternate choice—With many attributes of a preferred antibiotic. 3 = Acceptable, but preferably select an alternate antibiotic. 0 = No activity or no data or limited experience. § = Always use in combination with another antibiotic.

Table 3.2. Antibiotic Susceptibility Profiles (Cephalosporins)

		CEPHALOSPORINS																	Anti-MRSA (IV)				
ORGANISMS	1 <sup>st</sup> GC (IV)	1 <sup>st</sup> GCs (PO)		2 <sup>nd</sup> GCs (IV)			2 <sup>nd</sup> GCs (PO)			3 <sup>rd</sup> GCs (IV)			3 <sup>rd</sup> GCs (PO)				Anti-Pseudomonal Cephalosporins (IV)						
		Cefadroxil (PO)	Cephalexin (PO)	Cefoxitin (IV)	Cefuroxime (IV)	Cefotetan (IV)	Cefaclor (PO)	Loracarbef (PO)	Cefprozil (PO)	Cefuroxime axetil (PO)	Cefotaxime (IV)	Ceftizoxime (IV)	Ceftriaxone (IV)	Cefixime (PO)	Ceftibuten (PO)	Cefpodoxime (PO)	Cefdinir (PO)	Cefditoren (PO)	Cefoperazone (IV)	Ceftazidime/ Avibactam (IV)	Ceftolozane/ Tazobactam (IV)	Cefepime (IV)	
Aerobic Gram Positive Cocci (Clusters)																							
<i>Staphylococcus aureus</i> (MSSA)	1	2	1	2	2	3	3	3	2	2	2	2	0	2	2	2	0	2	3	3	3	2	2
<i>Staphylococcus aureus</i> (HA/CO-MRSA)	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1
<i>S. aureus</i> (CA-MRSA)	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2
<i>S. epidermidis</i> (CoNS)	3	0	0	3	3	3	3	0	0	3	3	3	0	0	0	0	0	3	0	3	3	3	0

1 = Preferred antibiotic—High degree of activity against the isolate and likely to be clinically effective, most strains susceptible, has a favorable PK/PD characteristics, has a good safety profile, and has a "low resistance potential" (see Chapter 11 Drug Summaries for antibiotic dosing details); 2 = Alternate choice—With many attributes of a preferred antibiotic; 3 = Acceptable, but preferably select an alternate antibiotic; 0 = No activity or no data or limited experience.

Table 3.2. Antibiotic Susceptibility Profiles (Cephalosporins) (cont'd)

CEPHALOSPORINS																								
ORGANISMS	1 <sup>st</sup> GC (IV)	1 <sup>st</sup> GCs (PO)		2 <sup>nd</sup> GCs (IV)			2 <sup>nd</sup> GCs (PO)			3 <sup>rd</sup> GCs (IV)			3 <sup>rd</sup> GCs (PO)					Anti-Pseudomonal Cephalosporins (IV)					Anti-MRSA (IV)	
		Cefadroxil (PO)	Cephalexin (PO)	Cefoxitin (IV)	Cefuroxime (IV)	Cefotetan (IV)	Cefaclor (PO)	Cefprozil (PO)	Cefuroxime axetil (PO)	Cefotaxime (IV)	Ceftizoxime (IV)	Ceftriaxone (IV)	Cephixime (PO)	Ceftibuten (PO)	Cefpodoxime (PO)	Cefdinir (PO)	Cefditoren (PO)	Cefoperazone (IV)	Ceftazidime (IV)	Ceftazidime/ Avibactam (IV)	Ceftolozame/ Tazobactam (IV)	Cefepime (IV)	Ceftaroline fosamil (IV)	
Cefazolin (IV)	1																							
<i>Enterococcus faecalis</i> (VSE)	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Enterococcus faecium</i> (VRE)	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Streptococci (groups A, B, C, E, G)	1	1	2	2	2	2	2	1	2	1	1	1	1	2	2	2	2	1	3	3	3	2	2	2 <sup>†</sup>
<i>Streptococcus (bovis) galloyticus</i>	1	1	2	2	2	2	2	2	2	1	1	1	2	3	2	2	2	1	3	3	3	2	2	0

<i>Viridans streptococci</i> ( <i>S. mitis</i> , <i>milleri</i> , <i>mitis</i> , <i>mutans</i> , <i>oralis</i> , <i>sanguis</i> , <i>parvulus</i> , <i>salivarius</i> )	1	1	2	2	2	3	2	2	1	1	1	1	2	3	2	2	2	1	2	3	3	2	0
	<b>Aerobic Gram Positive Cocci (Pairs)</b>																						
<i>Streptococcus pneumoniae</i> (PSSP)	1	1	1	2	2	2	2	1	2	2	2	1	2	2	2	3	3	2	3	3	3	3	2
	<i>Streptococcus pneumoniae</i> (PRSP)																						
<i>Streptococcus pneumoniae</i> (PRSP)	1	1	1	2	2	2	2	2	1	2	2	1	2	2	2	0	2	2	3	2	3	3	2
	<i>Streptococcus pneumoniae</i> (MDRSP)																						
<b>Aerobic Gram Negative Cocci (Pairs)</b>	1	1	1	2	2	2	2	2	1	2	2	1	2	2	0	2	2	3	2	3	3	3	2
	<i>Neisseria gonorrhoeae</i>																						
<i>Neisseria meningitidis</i>	2	0	0	2	2	3	3	3	3	3	2	2	1	1	3	2	2	2	2	3	3	2	0
	3	0	0	3	2	3	3	3	3	3	2	3	1	3	3	0	0	0	2	3	3	3	0

1 = Preferred antibiotic—High degree of activity against the isolate and likely to be clinically effective, most strains susceptible, has a favorable PK/PD characteristics, has a good safety profile, and has a "low resistance potential" (see Chapter 11 Drug Summaries for antibiotic dosing details).  
 2 = Alternate choice—With many attributes of a preferred antibiotic. 3 = Acceptable, but preferably select an alternate antibiotic. 0 = No activity or no data or limited experience.  
 † Group A & B only.



Table 3.2. Antibiotic Susceptibility Profiles (Cephalosporins) (cont'd)

CEPHALOSPORINS																								
ORGANISMS	1 <sup>st</sup> GC (IV)	1 <sup>st</sup> GCs (PO)		2 <sup>nd</sup> GCs (IV)		2 <sup>nd</sup> GCs (PO)			3 <sup>rd</sup> GCs (IV)			3 <sup>rd</sup> GCs (PO)				Anti-Pseudomonal Cephalosporins (IV)					Anti-MRSA (IV)			
	Cefazolin (IV)	Cefadroxil (PO)	Cephalexin (PO)	Cefoxitin (IV)	Cefuroxime (IV)	Cefotetan (IV)	Cefaclor (PO)	Loracarbef (PO)	Cefprozil (PO)	Cefuroxime axetil (PO)	Cefotaxime (IV)	Ceftizoxime (IV)	Ceftriaxone (IV)	Cephixime (PO)	Ceftibuten (PO)	Cefpodoxime (PO)	Cedirir (PO)	Cefditoren (PO)	Cefoperazone (IV)	Ceftazidime (IV)	Ceftazidime/Avibactam (IV)	Ceftolozane/Tazobactam (IV)	Cefepime (IV)	Ceftaroline fosamil (IV)
<b>Aerobic Gram Positive Bacilli</b>																								
<i>Bacillus anthracis</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Corynebacterium diphtheriae</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Corynebacterium jeikeium</i> (JK)	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Listeria monocytogenes</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Nocardia</i> sp.	0	0	0	0	3	0	0	0	0	0	3	0	3	0	0	0	0	0	0	0	0	0	0	0
<b>Aerobic Gram-Negative Bacilli</b>																								
<i>Acinetobacter</i> sp.	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2	2	2	2	2	0

<i>Aeromonas hydrophila</i>	0	0	0	3	3	0	0	0	3	2	2	3	0	0	0	0	0	0	2	3	3	3	2	0
<i>Bordetella pertussis</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Brucella</i> sp.	3	0	0	2	2	2	0	3	2	2	2	2	0	0	0	0	0	0	0	3	3	3	3	0
<i>Burkholderia (Pseudomonas) cepacia</i>	2	0	0	0	0	0	0	0	0	3	3	3	0	3	0	0	0	0	3	2	2	2	3	0
<i>Campylobacter</i> sp.	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Citrobacter</i> sp.	0	0	0	2	3	3	0	0	3	2	2	2	0	0	0	0	0	0	0	2	2	2	2	0
<i>Enterobacter</i> sp.	0	0	0	3	3	3	0	0	0	2	2	2	0	0	0	0	0	0	2	3	1	1	1	0
<i>Escherichia coli</i>	2	2	2	2	2	2	2	2	2	1	1	2	2	2	2	2	2	2	1	2	1	1	1	3
<i>Francisella tularensis</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Haemophilus</i> sp.	3	1	1	2	2	2	3	2	2	1	1	1	1	2	2	2	2	2	1	1	1	1	1	2 <sup>†</sup>
<i>Klebsiella</i> sp.	2	2	2	2	2	2	2	2	2	2	1	1	2	2	2	2	2	2	2	3	2	2	1	0
<i>Moraxella catarrhalis</i>	3	2	2	3	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	0
<i>Morganella</i> sp.	3	0	0	3	3	3	0	0	0	3	2	2	3	0	0	0	0	0	2	3	2	2	2	0
<i>P. aeruginosa</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	3	1	1	1	0
<i>Providencia</i> sp.	3	0	0	3	3	3	0	0	0	3	2	2	3	0	0	0	0	0	2	3	2	2	2	0
<i>Proteus</i> sp.	3	0	0	3	3	3	0	0	0	3	2	2	3	0	0	0	0	0	2	3	2	2	2	0
<i>Salmonella</i> sp.	0	0	0	3	3	3	0	0	0	0	1	1	0	0	0	0	0	0	3	3	3	3	2	0

1 = Preferred antibiotic—High degree of activity against the isolate and likely to be clinically effective, most strains susceptible, has a favorable PK/PD characteristics, has a good safety profile, and has a "low resistance potential" (see Chapter 11 Drug Summaries for antibiotic dosing details). 2 = Alternate choice—With many attributes of a preferred antibiotic. 3 = Acceptable, but preferably select an alternate antibiotic. 0 = No activity or no data or limited experience. <sup>†</sup>H. influenzae only.

Table 3.2. Antibiotic Susceptibility Profiles (Cephalosporins) (cont'd)

ORGANISMS	CEPHALOSPORINS														Anti-MRSA (IV)									
	1 <sup>st</sup> GC (IV)	1 <sup>st</sup> GCs (PO)		2 <sup>nd</sup> GCs (IV)		2 <sup>nd</sup> GCs (PO)		3 <sup>rd</sup> GCs (IV)			3 <sup>rd</sup> GCs (PO)					Anti-Pseudomonal Cephalosporins (IV)								
	Cefazolin (IV)	Cefadroxil (PO)	Cephalexin (PO)	Cefoxitin (IV)	Cefuroxime (IV)	Cefotetan (IV)	Cefaclor (PO)	Loracarbef (PO)	Cefprozil (PO)	Cefuroxime axetil (PO)	Cefotaxime (IV)	Ceftizoxime (IV)	Ceftriaxone (IV)	Cefixime (PO)	Ceftibuten (PO)	Cefpodoxime (PO)	Cefdinir (PO)	Ceftiofen (PO)	Cefoperazone (IV)	Ceftazidime (IV)	Ceftazidime/Avibactam (IV)	Ceftiozane/Tazobactam (IV)	Cefepime (IV)	Ceftriaxone fosamil (IV)
<i>Serratia marcescens</i>	0	0	0	3	0	2	0	0	0	0	1	1	1	0	0	3	0	3	1	3	2	2	1	0
<i>Shigella</i> sp.	0	0	0	3	0	3	0	0	0	0	2	2	2	0	0	0	0	0	2	3	3	3	2	0
<i>Stenotrophomonas (Pseudomonas) maltophilia</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	3	0	0	0	0	2	2	0	0
<i>Vibrio vulnificus</i>	0	0	0	0	0	0	0	0	0	0	2	0	0	0	0	0	0	0	0	2	0	0	0	0
<i>Yersinia enterocolitica</i>	0	0	0	3	3	3	0	0	0	0	2	2	2	2	2	0	2	0	2	3	0	0	2	0



Table 3.3. Antibiotic Susceptibility Profiles (Aminoglycosides, Fluoroquinolones, Carbapenems, and Others)

ORGANISMS	Amino-Glycosides			Fluoro-Quinolones			Miscellaneous										Carbapenems								
	Gentamicin (IV)	Tobramycin (IV)	Amikacin (IV)	Ciprofloxacin (IV/PO)	Levofloxacin (IV/PO)	Moxifloxacin (IV/PO)	Colistin (IV)	Polymyxin B (IV)	Aztreonam (IV)	Tigecycline (IV)	Vancomycin (IV)	Q/D* (IV)	Linezolid (IV/PO)	Tedizolid (IV/PO)	Daptomycin (IV)	Ortavancin (IV)	Telavancin (IV)	Dalbavancin (IV)	Nitrofurantoin † (PO)	Fosfomycin † (PO)	Imipenem (IV)	Meropenem (IV)	Ertapenem (IV)	Doripenem (IV)	
<b>Aerobic Gram Positive Cocci (Clusters)</b>																									
<i>Staphylococcus aureus</i> (MSSA)	2	3	3	3*	2	2	0	0	1	3	1	1	1	1	1	1	1	1	0	0	2	2	2	2	
<i>Staphylococcus aureus</i> (HA/CO-MRSA)	0	0	0	0	0	0	0	0	1	2	1	1	1	1	1	1	1	1	0	0	0	0	0	0	
<i>S. aureus</i> (CA-MRSA)	0	0	0	0	0	0	0	0	1	1	1	1	1	1	1	1	1	1	0	0	0	0	0	0	
<i>S. epidermidis</i> (CoNS)	0	0	0	0	3	3	0	0	1	1	1	1	1	1	1	1	1	1	0	0	3	3	3	3	
<b>Aerobic Gram Positive Cocci (Chains)</b>																									
<i>Enterococcus faecalis</i> (VSE)	0	0	0	0	3	2	0	0	1	2	0	1	1	2	3	3	3	1	2	2	2	2	0	3	
<i>Enterococcus faecium</i> (VRE)	0	0	0	0	0	0	0	0	1	0	1	1	1	2	3	3	3	1	3	0	0	0	0	3	
Streptococci (groups A, B, C, E, G)	0	0	0	0	3	2	0	0	2	2	3	2	2	2	2	2	2	2	0	2	2	2	2	2	
<i>Streptococcus (bovis) galloyticus</i>	0	0	0	0	3	2	0	0	2	2	2	3	3	3	3	3	3	0	0	2	2	2	0	2	

\* Q/D: Quinupristin/Dalfopristin

<i>Viridans streptococci</i> ( <i>S. mitis</i> , <i>milleri</i> , <i>mitis</i> , <i>mutans</i> , <i>oralis</i> , <i>sanguis</i> , <i>parosanguis</i> , <i>salivarius</i> )	0	0	0	0	3	2	0	0	2	2	2	2	2	2	2	2	2	0	0	2	2	0	2
<b>Aerobic Gram Positive Cocci (Pairs)</b>																							
<i>Streptococcus pneumoniae</i> (PSSP)	0	0	0	3*	1	1	0	0	1	2	2	2	2	3	2	2	2	2	0	0	2	2	2
<i>Streptococcus pneumoniae</i> (PRSP)	0	0	0	3*	1	1	0	0	1	2	2	2	2	3	2	2	2	2	0	0	2	2	2
<i>Streptococcus pneumoniae</i> (MDRSP)	0	0	0	3*	1	1	0	0	1	2	2	2	2	3	2	2	2	2	0	0	2	2	2
<b>Aerobic Gram Negative Cocci (Pairs)</b>																							
<i>Neisseria gonorrhoeae</i>	0	0	0	1	1	2	0	2	0	0	0	0	0	0	0	0	0	0	0	0	3	3	0
<i>Neisseria meningitidis</i>	0	0	0	2	2	2	0	2	0	0	0	0	0	0	0	0	0	0	0	0	2	2	0
<b>Aerobic Gram Positive Bacilli</b>																							
<i>Bacillus anthracis</i>	0	0	0	1	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Corynebacterium diphtheriae</i>	0	0	0	0	0	0	0	0	3	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Corynebacterium jeikeium</i> (JK)	0	0	0	0	0	0	0	2	1	3	2	2	0	0	0	0	0	0	0	0	0	0	0

1 = Preferred antibiotic—High degree of activity against the isolate and likely to be clinically effective, most strains susceptible, has a favorable PK/PD characteristics, has a good safety profile, and has a "low resistance" potential (see Chapter 11 Drug Summaries for antibiotic dosing details). 2 = Alternate choice—With many attributes of a preferred antibiotic. 3 = Acceptable, but preferably select an alternate antibiotic. 0 = No activity or no data or limited experience. \* = May be effective initially but as a "high resistance" antibiotic potential, resistance may develop during/after therapy.

Table 3.3. Antibiotic Susceptibility Profiles (Aminoglycosides, Fluoroquinolones, Carbanemns, and Others) (cont'd)

ORGANISMS	Amino-Glycosides			Fluoro-Quinolones			Miscellaneous										Carbanemns							
	Gentamicin (IV)	Tobramycin (IV)	Amikacin (IV)	Ciprofloxacin (IV/PO)	Levofloxacin (IV/PO)	Moxifloxacin (IV/PO)	Colistin (IV)	Polymyxin B (IV)	Aztreonam (IV)	Tigecycline (IV)	Vancomycin (IV)	Q/D* (IV)	Linezolid (IV/PO)	Tedizolid (IV/PO)	Daptomycin (IV)	Oritavancin (IV)	Telavancin (IV)	Dalbavancin	Nitrofurantoin † (PO)	Fostomycin † (PO)	Imipenem (IV)	Meropenem (IV)	Ertapenem (IV)	Doripenem (IV)
<i>Listeria monocytogenes</i>	3	3	3	0	0	0	0	0	3	0	3	3	3	3	0	0	0	0	0	0	3	2	0	0
<i>Nocardia</i>	0	0	2	0	0	0	0	0	0	0	0	3	3	0	0	0	0	0	0	3	2	0	0	0
<b>Aerobic Gram-Negative Bacilli</b>																								
<i>Acinetobacter</i> sp.	1	2	2	3*	2	3	2	1	2*	0	0	0	0	0	0	0	0	0	0	3	2*	1	0	1
<i>Aeromonas hydrophila</i>	2	2	2	1	1	1	0	3	2	0	0	0	0	0	0	0	0	0	0	0	2	2	3	0
<i>Bordetella</i> sp.	0	0	0	3	3	3	0	0	0	0	0	0	0	0	0	0	0	0	0	0	3	3	3	0
<i>Brucella</i> sp.	2	0	0	3	3	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Burkholderia (Pseudomonas) cepacia</i>	0	0	0	3*	3	3	0	0	0	0	0	0	0	0	0	0	0	0	0	0	3	2	3	0
<i>Campylobacter</i> sp.	3	0	0	1	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	3	3	3	3
<i>Citrobacter</i> sp.	1	1	2	3	2	3	3	0	2	0	0	0	0	0	0	0	0	0	2	0	1	1	1	1
<i>Enterobacter</i> sp.	1	1	1	2*	2	2	3	2	2	0	0	0	0	0	0	0	0	0	3	3	1	1	1	1
<i>Escherichia coli</i>	2	2	1	2	2	2	3	2	1	0	0	0	0	0	0	0	0	0	2	2	2	2	2	2
<i>Francisella tularensis</i>	1	3	0	2	2	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0

\* Q/D: Quinupristin/Dalfopristin

<i>Haemophilus sp.</i>	3	3	3	2	2	2	3	3	2	0	0	0	0	0	0	0	0	0	0	0	2	2	2	2
<i>Klebsiella sp.</i>	2	2	1	2	1	1	1	1	1	0	0	0	0	0	0	0	0	0	0	0	2	1	1	1
<i>Moraxella catarrhalis</i>	3	3	3	2	2	2	0	2	2	0	0	0	0	0	0	0	0	0	0	0	2	2	2	2
<i>Morganella sp.</i>	2	2	1	3	2	2	0	2	0	0	0	0	0	0	0	0	0	0	0	0	3	2	2	2
<i>P. aeruginosa</i>	3	3	1	1*	1	0	1	1*	0	0	0	0	0	0	0	0	0	0	0	0	1	2*	1	0
<i>Proteus sp.</i>	2	2	1	3	2	2	0	2	0	0	0	0	0	0	0	0	0	0	0	0	2	2	2	2
<i>Providencia sp.</i>	2	2	1	3	2	2	0	2	0	0	0	0	0	0	0	0	0	0	0	0	2	2	2	2
<i>Salmonella sp.</i>	0	0	0	1	1	1	3	2	2	0	0	0	0	0	0	0	0	0	0	0	2	2	2	2
<i>Serratia marcescens</i>	3	2	3	1*	1	2	0	2	2	0	0	0	0	0	0	0	0	0	0	0	3	2	2	2
<i>Shigella sp.</i>	3	3	3	1	1	1	3	2	2	0	0	0	0	0	0	0	0	0	0	0	2	2	2	2
<i>Stenotrophomonas (Pseudomonas) maltophilia</i>	0	0	0	3	2	2	2	0	3	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Vibrio vulnificus</i>	0	0	0	3	3	3	3	0	3	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Yersinia enterocolitica</i>	2	2	2	2	2	2	0	2	3	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<b>Anaerobic** Gram Positive Cocci (chains)</b>																								
<i>Peptostreptococcus</i>	0	0	0	0	0	3	0	0	2	3	3	2	2	0	0	0	0	0	0	0	2	2	2	2
<b>Anaerobic Gram Positive Bacilli</b>																								
<i>Actinomyces sp.</i>	0	0	0	0	0	3	0	0	3	0	0	3	3	0	0	0	0	0	0	0	0	2	2	0

1 = Preferred antibiotic—High degree of activity against the isolate and likely to be clinically effective, most strains susceptible, has a favorable PK/PD characteristics, has a good safety profile, and has a "low resistance" potential (see Chapter 11 Drug Summaries for antibiotic dosing details). 2 = Alternate choice—With many attributes of a preferred antibiotic. 3 = Acceptable, but preferably select an alternate antibiotic. 0 = No activity or no data or limited experience. \* = May be effective initially but as a "high resistance" antibiotic potential, resistance may develop during/after therapy.



Table 3.3. Antibiotic Susceptibility Profiles (Aminoglycosides, Fluoroquinolones, Carbapenems, and Others) (cont'd)

ORGANISMS	Amino-Glycosides			Fluoro-Quinolones			Miscellaneous										Carbapenems								
	Gentamicin (IV)	Tobramycin (IV)	Amikacin (IV)	Ciprofloxacin (IV/PO)	Levofloxacin (IV/PO)	Moxifloxacin (IV/PO)	Colistin (IV)	Polymyxin B (IV)	Aztreonam (IV)	Tigecycline (IV)	Vancomycin (IV)	Q/D* (IV)	Linezolid (IV/PO)	Tedizolid (IV/PO)	Daptomycin (IV)	Oritavancin (IV)	Telavancin (IV)	Dalbavancin (IV)	Nitrofurantoin † (PO)	Fosfomycin † (PO)	Imipenem (IV)	Meropenem (IV)	Ertapenem (IV)	Doripenem (IV)	
<b>Anaerobic Gram Negative Bacilli</b> <i>Bacteroides fragilis</i> group ( <i>B. distasonis</i> , <i>ovatus</i> , <i>thetaiotaomicron</i> , <i>vulgatus</i> )	0	0	0	0	0	2	0	0	1	0	3	0	0	0	0	0	0	0	0	0	1	1	1	1	1
<i>Prevotella</i> sp.	0	0	0	2	2	2	0	0	2	0	0	0	0	0	0	0	0	0	0	2	2	2	2	2	

1 = Preferred antibiotic—High degree of activity against the isolate and likely to be clinically effective; most strains susceptible, has a favorable PK/PD characteristics, has a good safety profile, and has a "low resistance" potential (see Chapter 11 Drug Summaries for antibiotic dosing details).

2 = Alternate choice—With many attributes of a preferred antibiotic.

3 = Acceptable, but preferably select an alternate antibiotic.

0 = No activity or no data or limited experience.

\* Q/D: Quinupristin/Dalfopristin

† = For lower tract infections (UTIs)/catheter associated bacteriuria (CAB) only.

## Gram Stain Characteristics of Isolates (by Morphology, Arrangement, Oxygen Requirements)

### AEROBIC ISOLATES

<b>GRAM-POSITIVE COCCI (CLUSTERS)</b>	
Staphylococcus aureus (MSSA/MRSA/ VISA/VRSA) <sup>†‡</sup> .....	208
Staphylococcus (coagulase negative) epidermidis (CoNS) <sup>†‡</sup> .....	209
Staphylococcus saprophyticus <sup>†‡</sup> .....	210
<b>GRAM-POSITIVE COCCI (CHAINS)</b>	
Enterococcus faecalis (VSE) <sup>α‡</sup> .....	210
Enterococcus faecium (VRE) <sup>α‡</sup> .....	211
Group A streptococci <sup>β</sup> .....	211
Group B streptococci (S. agalactiae) <sup>β</sup> .....	211
Group C, F, G streptococci <sup>β</sup> .....	212
Streptococcus anginosus (S. milleri) group (S. intermedius, S. anginosus, S. constellatus).....	213
Streptococcus (bovis) gallolyticus <sup>α‡</sup> .....	213
Viridans streptococci (S. mitior, mitis, mutans, oralis, sanguis, parasanguis, salivarius) <sup>α‡</sup> .....	213
<b>GRAM-POSITIVE COCCI B(PAIRS)</b>	
Group B streptococci (S. agalactiae) <sup>β</sup> .....	211
Leuconostoc <sup>α</sup> .....	213
Streptococcus pneumoniae <sup>α</sup> .....	213
<b>GRAM-NEGATIVE COCCI (PAIRS)</b>	
Neisseria gonorrhoeae <sup>†</sup> .....	214
Neisseria meningitidis <sup>†</sup> .....	214
<b>GRAM-POSITIVE BACILLI</b>	
Arcanobacterium (Corynebacterium) haemolyticum <sup>β</sup> .....	215
Bacillus anthracis <sup>†‡</sup> .....	215
Bacillus cereus, subtilis, megaterium <sup>†‡mv</sup> .....	215
Corynebacterium diphtheriae <sup>†‡</sup> .....	216
Corynebacterium jeikeium (JK) <sup>†‡</sup> .....	216
Erysipelothrix rhusiopathiae <sup>α</sup> .....	216
Listeria monocytogenes <sup>†‡mv</sup> .....	217
Nocardia asteroides, braziliensis <sup>†‡</sup> .....	217
Rhodococcus equi <sup>†‡</sup> .....	217
<b>GRAM-NEGATIVE BACILLI</b>	
Acinetobacter baumannii, Iwoffii, calcoaceticus, haemolyticus <sup>*†v</sup> .....	218
Aeromonas hydrophila <sup>†m</sup> .....	218
Aggregatibacter (Actinobacillus) actinomycetemcomitans <sup>*</sup> .....	218
Alcaligenes (Achromobacter) xylooxidans <sup>†m</sup> .....	219
Bartonella henselae, quintana, bacilliformis <sup>*</sup> .....	219
Bordetella pertussis, parapertussis <sup>†</sup> .....	219
Brucella abortus, canis, suis, melitensis <sup>†</sup> .....	219
Burkholderia (Pseudomonas) cepacia <sup>†m</sup> .....	220
Burkholderia (Pseudomonas) pseudomallei <sup>†m</sup> .....	220
Campylobacter fetus <sup>†m</sup> .....	220
Campylobacter jejuni <sup>†m</sup> .....	221
Cardiobacterium hominis.....	221
Chromobacterium violaceum <sup>†m</sup> .....	221
Chryseobacterium (Flavobacterium) meningosepticum <sup>†</sup> .....	221
Citrobacter diversus, freundii, koseri <sup>†*m</sup> .....	222
Edwardsiella tarda <sup>*†m</sup> .....	222
Enterobacter agglomerans, aerogenes, cloacae <sup>*†m</sup> .....	223
Escherichia coli <sup>*†m</sup> .....	223
Francisella tularensis <sup>*†</sup> .....	223
Hafnia alvei <sup>*†m</sup> .....	224

Helicobacter (Campylobacter)	Pseudomonas (Chryseomonas)
pylori <sup>†m</sup> ..... 224	luteola (Ve-1) <sup>*†sm</sup> ..... 229
Hemophilus influenzae, parainfluenzae,	Pseudomonas (Flavimonas)
aphrophilus, paraphrophilus*..... 225	oryzihabitans (Ve-2) <sup>*†sm</sup> ..... 230
Kingella (Moraxella) kingae..... 225	Salmonella typhi, non-typhi <sup>*†m</sup> ..... 230
Klebsiella pneumoniae, oxytoca <sup>*†</sup> ..... 225	Serratia marcescens <sup>*†sm</sup> ..... 230
Klebsiella ozaenae,	Shigella boydii, sonnei, flexneri,
rhinoscleromatis <sup>*†</sup> ..... 226	dysenteriae <sup>*†s</sup> ..... 231
Legionella sp <sup>†m</sup> ..... 226	Stenotrophomonas (Pseudomonas,
Leptospira interrogans <sup>m</sup> ..... 226	Xanthomonas) maltophilia <sup>*†sm</sup> ..... 231
Moraxella (Branhamella)	Streptobacillus moniliformis*..... 231
catarrhalis <sup>†s</sup> ..... 227	Vibrio cholerae <sup>†sm</sup> ..... 232
Morganella morganii <sup>*†sm</sup> ..... 227	Vibrio parahaemolyticus <sup>†sm</sup> ..... 232
Ochrobactrum anthropi (Vd) <sup>†sm</sup> ..... 227	Vibrio vulnificus, alginoliticus <sup>†sm</sup> ..... 232
Pasteurella multocida <sup>†s</sup> ..... 227	Yersinia enterocolitica <sup>*†s</sup> ..... 232
Plesiomonas shigelloides <sup>†sm</sup> ..... 228	Yersinia pestis <sup>*†s</sup> ..... 233
Proteus mirabilis, penneri,	
vulgaris <sup>*†sm</sup> ..... 228	<b>SPIROCHETES</b>
Providencia alcalifaciens, rettgeri,	Borrelia burgdorferi <sup>m</sup> ..... 233
stuartii <sup>*†sm</sup> ..... 229	Borrelia recurrentis <sup>m</sup> ..... 234
Pseudomonas aeruginosa <sup>†sm</sup> ..... 229	Spirillum minus <sup>m</sup> ..... 234

- \* Oxidase negative.   <sup>(α)</sup> = α (alpha) hemolysis on BAP.   <sup>(m)</sup> = motile.  
† Catalase positive.   <sup>(β)</sup> = β (beta) hemolysis on BAP.   <sup>(v)</sup> = Gram variable bacilli.  
<sup>(s)</sup> Non-lactose fermenter.   <sup>(γ)</sup> = γ (gamma) hemolysis on BAP.

### CAPNOPHILIC ISOLATES\*

<b>GRAM-NEGATIVE BACILLI</b>	Capnocytophaga ochraceus
Capnocytophaga canimorsus/	(DF-1) <sup>m</sup> ..... 235
cynodegni (DF-2 like) <sup>m</sup> ..... 235	Eikenella corrodens..... 235
<b>+ Capnophilic organisms grow best under increased CO<sub>2</sub> tension.</b>	<sup>(m)</sup> = motile.

### ANAEROBIC ISOLATES

<b>GRAM-POSITIVE COCCI (CHAINS)</b>	Bifidobacterium sp..... 237
Peptococcus..... 236	Clostridium botulinum <sup>m</sup> ..... 237
Peptostreptococcus..... 236	Clostridium difficile <sup>m</sup> ..... 238
	Clostridium perfringens, septicum,
	novyi <sup>mv</sup> ..... 238
<b>GRAM-POSITIVE BACILLI</b>	Clostridium tetani <sup>m</sup> ..... 238
Actinomyces israelii, odontolyticus <sup>++</sup> ... 236	Eubacterium sp..... 239
Arachnia propionica <sup>++</sup> ..... 237	

Lactobacillus sp.**	239	Fusobacterium nucleatum	240
Propionibacterium acnes**†	239	Prevotella (Bacteroides) bivia	240
		Prevotella (Bacteroides) melaninogenicus, intermedius	241
<b>GRAM-NEGATIVE BACILLI</b>			
Bacteroides fragilis group (B. distasonis, ovatus, thetaiotaomicron, vulgatus) . . .	240		

**++ Microaerophilic organisms. Grow best under decreased O<sub>2</sub> concentration.**

**\* Oxidase negative.** <sup>(m)</sup> = motile.

**† Catalase positive.** <sup>(v)</sup> = Gram variable bacilli.

### YEASTS/FUNGI

Aspergillus fumigatus, flavus, niger†	242	Candida albicans	242
Candida (non-albicans): C. krusei, lusitaniae, tropicalis, pseudotropicalis, glabrata, guilliermondii, dublinensis, lipolytica	243	Cryptococcus neoformans	243
		Histoplasma capsulatum	244
		Malassezia furfur	244
		Penicillium marneffei	245

### Alphabetical Index of Isolates

Acinetobacter baumannii, Iwoffii, calcoaceticus, haemolyticus	218	Borrelia recurrentis	234
Actinomyces israelii, odontolyticus	236	Brucella abortus, canis, suis, melitensis	219
Aeromonas hydrophila	218	Burkholderia (Pseudomonas) cepacia	220
Aggregatibacter (Actinobacillus) actinomycetemcomitans	218	Burkholderia (Pseudomonas) pseudomallei	220
Alcaligenes (Achromobacter) xylosoxidans	219	Campylobacter fetus	220
Arachnia propionica	237	Campylobacter jejuni	221
Arcanobacterium (Corynebacterium) haemolyticum	215	Candida albicans	242
Aspergillus fumigatus, flavus, niger	242	Candida (non-albicans): C. krusei, lusitaniae, tropicalis, pseudotropicalis, glabrata, guilliermondii, dublinensis, lipolytica	243
Bacillus anthracis	215	Capnocytophaga canimorsus/ cynodegni (DF-2 like)	235
Bacillus cereus, subtilis, megaterium	215	Capnocytophaga ochraceus (DF-1)	235
Bacteroides fragilis group (B. distasonis, ovatus, thetaiotaomicron, vulgatus)	240	Cardiobacterium hominis	221
Bartonella henselae, quintana, bacilliformis	219	Chromobacterium violaceum	221
Bifidobacterium sp.	237	Chryseobacterium (Flavobacterium) meningosepticum	221
Bordetella pertussis, parapertussis	219	Citrobacter diversus, freundii, koseri	222
Borrelia burgdorferi	233	Clostridium botulinum	237

<i>Clostridium difficile</i> .....	238	<i>Pasteurella multocida</i> .....	227
<i>Clostridium perfringens</i> , septicum, novyi .....	238	<i>Penicillium marneffei</i> .....	245
<i>Clostridium tetani</i> .....	238	<i>Peptococcus</i> .....	236
<i>Corynebacterium diphtheriae</i> .....	216	<i>Peptostreptococcus</i> .....	236
<i>Corynebacterium jeikeium</i> .....	216	<i>Plesiomonas shigelloides</i> .....	228
<i>Cryptococcus neoformans</i> .....	243	<i>Prevotella (Bacteroides) bivia</i> .....	240
<i>Edwardsiella tarda</i> .....	222	<i>Prevotella (Bacteroides)</i>	
<i>Eikenella corrodens</i> .....	235	<i>melaninogenicus, intermedius</i> .....	241
<i>Enterobacter agglomerans</i> , <i>aerogenes</i> ,		<i>Propionibacterium acnes</i> .....	239
<i>cloacae</i> .....	223	<i>Proteus mirabilis, vulgaris</i> .....	228
<i>Enterococcus faecalis</i> (VSE) .....	210	<i>Providencia alcalifaciens, rettgeri,</i>	
<i>Enterococcus faecium</i> (VRE) .....	211	<i>stuartii</i> .....	229
<i>Erysipelothrix rhusiopathiae</i> .....	216	<i>Pseudomonas aeruginosa</i> .....	229
<i>Escherichia coli</i> .....	223	<i>Pseudomonas (Chryseomonas)</i>	
<i>Eubacterium sp</i> .....	239	<i>luteola (Ve-1)</i> .....	229
<i>Francisella tularensis</i> .....	223	<i>Pseudomonas (Flavimonas)</i>	
<i>Fusobacterium nucleatum</i> .....	240	<i>oryzihabitans (Ve-2)</i> .....	230
Group A streptococci .....	211	<i>Rhodococcus equi</i> .....	217
Group B streptococci ( <i>S. agalactiae</i> ) .....	211	<i>Salmonella typhi</i> , non-typhi .....	230
Group C, F, G streptococci .....	212	<i>Serratia marcescens</i> .....	230
<i>Hafnia alvei</i> .....	224	<i>Shigella boydii, sonnei, flexneri,</i>	
<i>Helicobacter (Campylobacter) pylori</i> .....	224	<i>dysenteriae</i> .....	231
<i>Hemophilus influenzae</i> , <i>parainfluenzae</i> ,		<i>Spirillum minus</i> .....	234
<i>aphrophilus, paraphrophilus</i> .....	225	<i>Staphylococcus aureus</i> (MSSA/MRSA/ VISA/VRSA) .....	208
<i>Histoplasma capsulatum</i> .....	244	<i>Staphylococcus (coagulase negative)</i>	
<i>Kingella kingae</i> .....	225	<i>epidermidis (CoNS)</i> .....	209
<i>Klebsiella ozaenae, rhinoscleromatis</i> .....	226	<i>Staphylococcus saprophyticus</i> .....	210
<i>Klebsiella pneumoniae, oxytoca</i> .....	225	<i>Stenotrophomonas (Pseudomonas,</i>	
<i>Lactobacillus sp.</i> .....	239	<i>Xanthomonas) maltophilia</i> .....	231
<i>Legionella sp.</i> .....	226	<i>Streptobacillus moniliformis</i> .....	231
<i>Leptospira interrogans</i> .....	226	<i>Streptococcus (bovis) gallolyticus</i> .....	213
<i>Leuconostoc</i> .....	213	<i>Streptococcus pneumoniae</i> .....	213
<i>Listeria monocytogenes</i> .....	217	<i>Streptococcus anginosus (S. milleri) group</i>	
<i>Malassezia furfur</i> .....	244	( <i>S. intermedius, S. anginosus,</i>	
<i>Moraxella (Branhamella)</i>		<i>S. constellatus</i> ) .....	213
<i>catarrhalis</i> .....	227	<i>Viridans streptococci (S. mitior, mitis,</i>	
<i>Morganella morganii</i> .....	227	<i>mutans, oralis, sanguis,</i>	
<i>Neisseria gonorrhoeae</i> .....	214	<i>parasanguis, salivarius)</i> .....	213
<i>Neisseria meningitidis</i> .....	214	<i>Vibrio cholerae</i> .....	232
<i>Nocardia asteroides, brasiliensis</i> .....	217	<i>Vibrio parahaemolyticus</i> .....	232
<i>Ochrobactrum anthropi (Vd)</i> .....	227	<i>Vibrio vulnificus, alginolyticus</i> .....	232
		<i>Yersinia enterocolitica</i> .....	232
		<i>Yersinia pestis</i> .....	233

**Table 3.4. Key Factors in Antibiotic Selection (Isolate Known)**

- 
- **Select an antibiotic with a high degree of activity against the known pathogen** (not colonizer).
  - **Dose appropriate for the target tissue to assure therapeutic/effective concentrations at site of infection.** If necessary, adjust dose ↑ for tissue targets that require higher doses, e.g., bacterial meningitis, endocarditis, etc., or ↓ dose for sites with antibiotic concentrations that are above serum concentrations, e.g., skin, urine, etc.
  - **Select an empiric antibiotic with a “low resistance” potential** avoid, if possible, antibiotics with a “high resistance” potential (also with a high degree of activity against the presumed pathogen). Select a “low resistance” potential antibiotic for the same/different class with a high degree of activity.
  - **Select antibiotic with a good safety profile** and minimal potential for drug-drug interactions.
  - **Select antibiotic that is relatively cost effective** (first take into account the above principles).
- 

**Table 3.5. Antibiotic Selection Based on Resistance Potential**

- 
- Antibiotic resistance may be classified as natural (*P. aeruginosa* is naturally resistant to chloramphenicol, i.e., not in its spectrum). Acquired resistance may be relative or high level resistance (gentamicin resistant *P. aeruginosa* with extremely high MIC cannot be overcome by increased dosing). In contrast, relative resistance, e.g., meropenem relative resistance to *P. aeruginosa* may be overcome, if antibiotic concentrations can be achieved with normal/high dosing that exceed the MIC of the organism at the site of infection.
  - There are many mechanisms of antibiotic resistance, but mechanisms do not explain differences in resistance potential within antibiotic classes. Mechanism of resistance does not explain why ciprofloxacin is responsible for nearly all fluoroquinolone resistance to *S. pneumoniae* and *P. aeruginosa*. Similarly, among the five 3rd GC, only ceftazidime has been associated with *P. aeruginosa* resistance problems.
  - Clinically, antibiotics may be considered in terms of resistance potential, i.e., high, moderate, or low. “High resistance potential” antibiotics cause resistance even with minimal use, but with widespread use can cause (non-clonal) resistance problems when used in high volume or over time. While an occasional mutation may result sporadic resistance with any antibiotic, resistance to “low resistance potential” antibiotics is not related to antibiotic class, volume or duration of use. After decades of extensive worldwide use with “low resistance potential” antibiotics, there are no widespread resistance problems with “low resistance potential” antibiotics, e.g., nitrofurantoin, amikacin, ceftriaxone, doxycycline.
  - If possible, always try to preferentially select a “low resistance potential” antibiotic with the appropriate spectrum and PK/PD characteristics for the pathogen/body site being treated. For each resistant problem pathogen in each antibiotic class, there are usually “low resistance potential” antibiotics alternatives within the class. In addition, other “low resistance potential” antibiotics may be found in different antibiotic classes. Try to use “low resistance potential” antibiotics in place of “high resistance potential” antibiotics, e.g., TMP-SMX (*S. pneumoniae* and MSSA resistance), tetracycline (*S. pneumoniae* and MSSA resistance), gentamicin (*P. aeruginosa* resistance), ceftazidime (*S. pneumoniae* and *P. aeruginosa* resistance), imipenem (*P. aeruginosa* resistance).
-

Table 3.6. Clinical Significance of AEROBIC Isolates Pending Susceptibility Testing

GRAM-POSITIVE COCCI (CLUSTERS)			
Isolate	Isolate Significance	Therapy	Comments
Staphylococcus aureus (MSSA/MRSA)	<ul style="list-style-type: none"> <li>CSF = C*, P (CNS shunts)</li> <li>Blood = C*, P (from soft tissue/bone infection, abscess, IV line infection, ABE, PVE)</li> <li>Sputum = C, P (S. aureus pneumonia is rare; usually only after viral influenza)</li> <li>Urine = C, P (S. aureus in urine is usually due to skin contamination or rarely overwhelming S. aureus bacteremia)</li> <li>Stool = C, P (enterocolitis)</li> <li>Wound = C, P (cellulitis, abscess)</li> </ul>	<p>MSSA: Nafcillin (IV), Cefazolin (IV), Clindamycin (IV/PO), a "respiratory quinolone" (IV/PO), Minocycline (IV/PO), Daptomycin (IV), any carbapenem (IV), Linezolid (IV/PO), Tigecycline (IV), Telavancin (IV)</p> <p>Hospital-acquired MRSA (HA-MRSA)/Community-onset MRSA (CO-MRSA): Daptomycin (IV), Linezolid (IV/PO), Tigecycline (IV), Vancomycin (IV), Minocycline (IV/PO), Quinupristin/dalfopristin (IV), Telavancin (IV)</p> <p>Community-acquired MRSA (CA-MRSA): Doxycycline*, TMP-SMX, Clindamycin</p> <p>MSSA/MRSA: Linezolid (IV/PO), Daptomycin (IV), Telavancin (IV)</p>	<p>MSSA: For oral treatment, 1<sup>st</sup> generation cephalosporins are better than oral anti-staphylococcal penicillins (e.g., dicloxacillin)</p> <p>MRSA: <i>in-vitro</i> susceptibility testing is unreliable; treat MRSA infections empirically. Most effective drugs for MRSA are vancomycin, linezolid, ceftaroline fosamil (ABSSSI only), quinupristin/ceftaroline, minocycline, daptomycin, tigecycline.</p> <p><b>Preferentially use minocycline instead of doxycycline for MSSA/MRSA.</b></p> <p>Community-acquired MRSA (CA-MRSA) SCC mec IV, V CA-MRSA has different susceptibilities than HA-MRSA/CO-MRSA (see p. 14). CA-MRSA strains with Panton-Valentin Leukocidin PVL+ gene cause two distinct clinical syndromes: severe necrotizing pneumonia (with viral influenza) and severe necrotizing fasciitis/pyomyositis. <b>CA-MRSA is usually susceptible to doxycycline in vitro, but minocycline more effective in vivo than doxycycline</b>, TMP-SMX, clindamycin. <i>Drugs effective against HA-MRSA/CO-MRSA are also effective against CA-MRSA. However, drugs effective against CA-MRSA are may not be effective against HA-MRSA/CO-MRSA</i></p> <p>MSSA/MRSA: Noncontinuous low-grade blood culture positivity indicates skin contamination during venipuncture. Continuous high-grade blood culture positivity (3/4 or 4/4) indicates intravascular infection or abscess</p>

<p>VISA/VRSA; MICs for vancomycin sensitive (VSSA), heteroresistant vancomycin intermediate (hVISA), intermediate (VISA), and resistant (VRSA) <i>S. aureus</i> are &lt; 4 mcg/mL, &lt; 4 mcg/mL (with subpopulations &gt; 4 mcg/mL), 8–16 mcg/mL, and ≥ 32 mcg/mL, respectively.</p>			<p>Staphylococcus epidermidis (MSSE/MRSE) or coagulase negative staphylococci (CoNS) Staphylococcus lugdunensis</p>
<p>Usually non-pathogenic in absence of prosthetic/implant materials. Common cause of PVE; rare cause of native valve SBE. Treat foreign body-related infection until foreign body is removed.</p> <p><i>S. lugdunensis</i> is a CoNS but is often misidentified as <i>S. aureus</i> since it produces “clumping factor” which gives a <b>1</b> rapid short tube coagulase test (long tube test →) although a CoNS resembles <i>S. aureus</i> in terms of invasiveness/virulence.</p> <p>Unlike <i>S. aureus</i>, <i>S. lugdunensis</i> is pan-sensitive to antibiotics which is another clue the isolate is not <i>S. aureus</i>. <i>S. lugdunensis</i> bacteremia associated with community acquired (not nosocomial) SBE.</p>	<p>MSSE: Linezolid (IV/PO), Daptomycin (IV), Vancomycin (IV), Telavancin (IV), any carbapenem (IV), a “respiratory quinolone” (IV/PO), Minocycline (IV/PO)</p> <p>MRSE: Linezolid (IV/PO), Daptomycin (IV), Vancomycin (IV), Quinupristin/dalfopristin (IV), Minocycline (IV/PO)</p>	<ul style="list-style-type: none"> <li>• CSF = C*, P (CNS shunts)</li> <li>• Blood = C*, P (from IV lines, infected implants, prosthetic valve)</li> <li>• endocarditis [PVE], rarely from native valve</li> <li>• subacute bacterial endocarditis [SBE]</li> <li>• Sputum = C</li> <li>• Urine = C (may be reported as <i>S. saprophyticus</i>; request novobiocin sensitivity to differentiate)</li> <li>• <i>S. epidermidis</i> from other coagulase-negative staphylococci)</li> </ul>	<ul style="list-style-type: none"> <li>• Stool = NP</li> <li>• Wound = C, P (infected foreign body drainage)</li> </ul>

C = colonizer; C\* = skin contaminant; NP = non-pathogen at site; P = pathogen at site; (IV/PO) = IV or PO. See p. xi for all other abbreviations.

\* Minocycline (IV/PO) preferred.



Table 3.6. Usual Clinical Significance of AEROBIC Isolates Pending Susceptibility Testing (cont'd)

GRAM-POSITIVE COCCI (CLUSTERS)				
Isolate	Isolate Significance	Preferred Therapy	Alternate Therapy	
			Comments	
Staphylococcus saprophyticus (coagulase-negative staphylococci)	<ul style="list-style-type: none"> <li>• CSF = NP</li> <li>• Blood = NP</li> <li>• Sputum = NP</li> <li>• Urine = P (cystitis, pyelo)</li> <li>• Stool = NP</li> <li>• Wound = NP</li> </ul>	<p>Preferred therapy</p> <p>Amoxicillin (PO)</p> <p>TMP-SMX (PO)</p> <p>Nitrofurantoin (PO)</p> <p>Alternate therapy</p> <p>Any quinolone (PO)</p> <p>Any 1<sup>st</sup> generation cephalosporin (PO)</p>	<p>S. saprophyticus UTI is associated with a urinary "fishy odor", alkaline urine pH, and microscopic hematuria. Novobiocin sensitivity differentiates coagulase-negative staphylococci (sensitive) from S. saprophyticus (resistant).</p>	
GRAM-POSITIVE COCCI (CHAINS)				
Enterococcus faecalis (VSE)	<ul style="list-style-type: none"> <li>• CSF = NP (except from S. stercoralis hyperinfection or V-P shunt infection)</li> <li>• Blood = C*, P (from GI/GU source, SBE)</li> <li>• Sputum = NP</li> <li>• Urine = C, P (cystitis, pyelonephritis)</li> <li>• Stool = NP</li> <li>• Wound = C, P (cellulitis)</li> </ul>	<p>Non-SBE</p> <p>Ampicillin (IV)</p> <p>Amoxicillin (PO)</p> <p>Meropenem (IV)</p> <p>Piperacillin (IV)</p> <p>Linezolid (IV/PO)</p> <p>Tigecycline (IV)</p> <p>Daptomycin (IV)</p> <p>SBE</p> <p>Gentamicin + ampicillin (IV) or vancomycin (IV)</p> <p>Meropenem (IV)</p> <p>Piperacillin (IV)</p> <p>Linezolid (IV/PO)</p>	<p>Non-SBE</p> <p>Cefoperazone (IV)</p> <p>Chloramphenicol (IV)</p> <p>Levofloxacin (IV/PO)</p> <p>Moxifloxacin (IV/PO)</p> <p>Nitrofurantoin (PO) (UTIs only)</p> <p>SBE</p> <p>Any quinolone (IV/PO)</p> <p>Cefoperazone (IV)</p>	<p>Sensitive to ampicillin, not penicillin. Cause of intermediate (severity between ABE and SBE) endocarditis; hepatobiliary infections, and UTIs. Enterococci (E. faecalis, E. faecium) are the only cause of SBE (below the waist) from GI/GU sources. Permissive pathogen (i.e., usually does not cause infection alone) in the abdomen/pelvis (except in gallbladder or urinary bladder/kidneys). Cefoperazone is the only cephalosporin with anti-E. faecalis (VSE) activity (MIC ~ 32 mcg/mL). Quinupristin/dalfopristin is not active against E. faecalis (VSE).</p> <p>Treat vancomycin resistant E. faecalis as VRE (see E. faecium VRE p. 211).</p>

Enterococcus faecium (VRE)	<ul style="list-style-type: none"> <li>CSF = NP (except from <i>S. stercoalis</i> hyperinfection or V-P shunt infection)</li> <li>Blood = C* P (from GI/GU source, SBE)</li> <li>Sputum = C</li> <li>Urine = C, P (cystitis, pyelo)</li> <li>Stool = NP</li> <li>Wound = C, P (cellulitis)</li> </ul>	Non-SBE Linezolid (IV/PO), quinupristin/dalfopristin (IV), doxycycline (IV/PO), tigecycline (IV), chloramphenicol (IV/PO), nitrofurantoin (PO) (UTIs only)  SBE Linezolid (IV/PO) Quinupristin/dalfopristin (IV)	Same spectrum of infection as <i>E. faecalis</i> . Colonization common; infection uncommon. Fecal carriage is intermittent but prolonged. In vitro sensitivity = in vivo efficacy. Increased prevalence of <i>E. faecalis</i> (VRE) related to vancomycin IV (not PO) use. Nitrofurantoin preferred for VRE lower UTIs/catheter-associated bacteriuria.
Group A streptococci	<ul style="list-style-type: none"> <li>CSF = C*, P (rare cause of ABM)</li> <li>Blood = P (from skin/soft tissue infection)</li> <li>Sputum = P (rare cause of CAP)</li> <li>Urine = NP</li> <li>Stool = NP</li> <li>Wound = C, P (cellulitis)</li> <li>Throat = C, P (pharynx is colonized with Group A streptococci in ~30% of patients with EBV mono)</li> </ul>	Amoxicillin (PO) Any $\beta$ -lactam (IV/PO)	For Group A streptococcal pharyngitis, amoxicillin is preferred over penicillin. Clindamycin is best for elimination of carrier states, and for penicillin-allergic patients with streptococcal pharyngitis. Any $\beta$ -lactam is equally effective against Group A streptococci. Nafcillin is the most active anti-staphylococcal penicillin against Group A streptococci. Erythromycin is no longer reliable against Group A streptococci due to increasing resistance. Doxycycline has little/no activity against Group A streptococci.
Group B streptococci ( <i>S. agalactiae</i> )	<ul style="list-style-type: none"> <li>CSF = P</li> <li>Blood = P (from IV line/urine source, SBE)</li> </ul>	Non-SBE, non-CNS Clindamycin (IV/PO)	Cause of UTIs and IV line infections in diabetics and the elderly. Cause of neonatal meningitis. Infection is uncommon in the general population. Rarely a cause of SBE in

C = colonizer; C\* = skin contaminant; NP = non-pathogen at site; P = pathogen at site; (IV/PO) = IV or PO. See p. xi for all other abbreviations.

Table 3.6. Usual Clinical Significance of AEROBIC Isolates Pending Susceptibility Testing (cont'd)

GRAM-POSITIVE COCCI (CHAINS)				
Isolate	Isolate Significance	Preferred Therapy	Alternate Therapy	Comments
	<ul style="list-style-type: none"> <li>Sputum = NP</li> <li>Urine = P (CAB, UTIs, especially in diabetics, elderly)</li> <li>Stool = NP</li> <li>Wound = C, P (diabetic foot infections)</li> </ul>	Any 1 <sup>st</sup> , 2 <sup>nd</sup> , 3 <sup>rd</sup> generation cephalosporin (IV/PO)  <u>SBE</u> Ceftriaxone (IV) Penicillin (IV) Vancomycin (IV)  <u>CNS</u> Ceftriaxone (IV) Penicillin (IV)	SBE Meropenem (IV) Ertapenem (IV) Linezolid (IV/PO)  <u>CNS</u> Chloramphenicol (IV) Linezolid (IV/PO)	Non-pregnant adults. On gram stain, GBS appear larger/rounder than <i>S. pneumoniae</i> . Colonies of GBS on BAP have a "sheen" vs. <i>S. pneumoniae</i> . Aminoglycosides and tetracyclines are ineffective.
Group C, F, G streptococci	<ul style="list-style-type: none"> <li>CSF = P (meningitis)</li> <li>Blood = P (from skin/soft tissue infection, SBE)</li> <li>Sputum = P (rare cause of CAP)</li> <li>Throat = C (especially with viral pharyngitis), P (pharyngitis in medical personnel)</li> <li>Urine = NP</li> <li>Stool = NP</li> <li>Wound = P (cellulitis)</li> </ul>	Ceftriaxone (IV) Penicillin (IV) Ampicillin (IV) Clindamycin (IV/PO)	Vancomycin (IV) Amoxicillin (PO) Any 1 <sup>st</sup> , 2 <sup>nd</sup> , 3 <sup>rd</sup> generation cephalosporin (IV) Meropenem (IV) Ertapenem (IV)	Group C, G streptococci may cause pharyngitis, wound infections, and rarely SBE. Group G streptococci associated with malignancies. Common pharyngeal colonizers in medical personnel.

GRAM-POSITIVE COCCI (CHAINS)				
Streptococcus (bovis) galloyticus	<ul style="list-style-type: none"> <li>Blood = P (SBE from GI source)</li> <li>Urine = NP</li> </ul>	Ceftriaxone (IV) Ampicillin (IV) Clindamycin (IV/PO)	Vancomycin (IV) Amoxicillin (PO) Any cephalosporin (IV)	Associated with GI malignancies. Non-enterococcal Group D streptococci (e.g., <i>S. bovis</i> ) are sensitive to penicillin.
<i>S. anginosus</i> ( <i>S. milleri</i> ) group ( <i>S. intermedius</i> , <i>S. anginosus</i> , <i>S. constellatus</i> )	<ul style="list-style-type: none"> <li>CNS = P</li> <li>Blood = P</li> <li>Wound = P (head/neck abscesses)</li> </ul>	Ceftriaxone (IV) Vancomycin (IV)	Amoxicillin (PO) Any cephalosporin (PO)	<i>S. anginosus</i> ( <i>S. milleri</i> ) group ( <i>S. intermedius</i> , <i>S. anginosus</i> , <i>S. constellatus</i> ) prone to invasive disease, bacteremia and abscess formation.
<i>S. viridans</i> streptococci ( <i>S. mitis</i> , <i>S. mitis</i> , <i>S. mutans</i> , <i>S. oralis</i> , <i>S. sanguis</i> , <i>S. parasanguis</i> , <i>S. salivarius</i> )	<ul style="list-style-type: none"> <li>CSF = NP (aseptic meningitis with SBE)</li> <li>Blood = C*, P (1° bacteremia, SBE)</li> <li>Sputum = NP</li> <li>Urine = NP</li> <li>Stool = NP</li> <li>Wound = NP</li> </ul>	Ceftriaxone (IV) Penicillin (IV)	Amoxicillin (PO) Any 1 <sup>st</sup> , 2 <sup>nd</sup> , 3 <sup>rd</sup> generation cephalosporin (IV/PO) Meropenem (IV) Ertapenem (IV) Vancomycin (IV)	Low-grade blood culture positivity (1/4) indicates contamination during venipuncture. Continuous/high-grade blood culture positivity (3/4 or 4/4) indicates SBE until proven otherwise.
GRAM-POSITIVE COCCI (PAIRS)				
Leuconostoc	<ul style="list-style-type: none"> <li>CSF = NP</li> <li>Blood = P (PVE)</li> <li>Sputum = NP</li> <li>Urine = P (UTIs)</li> <li>Stool = NP</li> <li>Wound = NP</li> </ul>	Penicillin (IV) Ampicillin (IV) Clindamycin (IV/PO)	Amoxicillin (PO) Erythromycin (IV) Minocycline (IV/PO) Clarithromycin XL (PO)	Cocci bacillary forms resemble streptococci/enterococci. Cause of infection in compromised hosts. Rare cause of IV line infection. Usually vancomycin-resistant.
Streptococcus pneumoniae	<ul style="list-style-type: none"> <li>CSF = P (ABM)</li> <li>Blood = P (from respiratory tract source)</li> <li>Sputum = C, P</li> <li>Urine = NP</li> <li>Stool = NP</li> </ul>	Multidrug Resistant <i>S. pneumoniae</i> (MDRSP) A* respiratory quinolone* (IV/PO); telithromycin (PO); ertapenem (IV); meropenem (IV); ceftipime (IV); linezolid (IV/PO); vancomycin (IV)		Penicillin-resistant <i>S. pneumoniae</i> (PRSP) are still sensitive to full-dose/high-dose $\beta$ -lactams. If possible, avoid macrolides, as > 30% of <i>S. pneumoniae</i> are macrolide resistant (MRSP). (~ 20–25% are naturally resistant and 10–15% acquire macrolide resistance).

C = colonizer; C\* = skin contaminant; NP = non-pathogen at site; P = pathogen at site; (IV/PO) = IV or PO. See p. xi for all other abbreviations.

Table 3.6. Usual Clinical Significance of AEROBIC Isolates Pending Susceptibility Testing (cont'd)

GRAM-POSITIVE COCCI (PARIS)				
Isolate	Isolate Significance	Preferred Therapy	Alternate Therapy	
	<ul style="list-style-type: none"> <li>Wound = P (cellulitis only in SLE)</li> </ul>	<p><u>Sensitive or relatively PCN-resistant</u> Doxycycline (IV/PO); any cephalosporin (IV/PO); clindamycin (IV/PO); amoxicillin/clavulanic acid (PO)</p>		
GRAM-NEGATIVE COCCI (PARIS)				
Neisseria gonorrhoeae (GC)	<ul style="list-style-type: none"> <li>CSF = NP</li> <li>Blood = P (from pharyngitis, proctitis, ABE)</li> <li>Sputum = NP</li> <li>Urine = P (urethritis)</li> <li>Stool = NP</li> <li>Wound = NP</li> <li>Rectal discharge = P (GC proctitis)</li> </ul>	<p><u>Penicillin-sensitive</u> <u>N. gonorrhoeae</u> (PSNG) Ceftriaxone (IV/IM) Any quinolone (IV/PO)</p> <p><u>PRNG</u> Ceftriaxone (IV/IM)</p>	<p><u>Penicillin-sensitive</u> <u>N. gonorrhoeae</u> (PSNG) Penicillin (IV/IM) Amoxicillin (PO) Doxycycline (IV/PO)</p> <p><u>PPNG</u> Spectinomycin (IM) Any quinolone (PO) Any 1<sup>st</sup>, 2<sup>nd</sup>, 3<sup>rd</sup> gen. cephalosporin (IV/IM)</p>	<p>Cause of "culture negative" right-sided ABE. May be cultured from synovial fluid/blood in disseminated GC infection (arthritis-dermatitis syndrome). Spectinomycin is ineffective against pharyngeal GC/Incubating syphilis. PRNG are tetracycline-resistant (TRNG). GC strains from Hawaii/California have increased quinolone resistance; use cefixime or ceftriaxone for such strains. Treat possible Chlamydia trachomatis co-infection and sexual partners.</p>
Neisseria meningitidis	<ul style="list-style-type: none"> <li>CSF = P (ABM)</li> <li>Blood = P (acute/chronic meningococemia)</li> <li>Sputum = C, P (only in closed populations; e.g., military recruits)</li> </ul>	<p>Penicillin (IV) Ampicillin (IV) Any 3<sup>rd</sup> generation cephalosporin (IV)</p>	<p>Chloramphenicol (IV) Cefepime (IV) Meropenem (IV)</p>	<p>In ABM, do not decrease meningeal dose of <math>\beta</math>-lactam antibiotics as patient improves, since CSF penetration/concentration decreases as meningeal inflammation decreases. Chloramphenicol is an excellent choice for penicillin-allergic patients. Preferred meningococcal prophylaxis is an oral quinolone (single dose).</p>

<b>GRAM-POSITIVE BACILLI</b>			
	<ul style="list-style-type: none"> <li>Urine C, P (urethritis rarely)</li> <li>Stool = NP</li> <li>Wound = NP</li> </ul>		
Arcanobacterium (Corynebacterium) haemolyticum	<ul style="list-style-type: none"> <li>CSF = NP</li> <li>Blood = NP</li> <li>Sputum = P (oropharyngeal secretions)</li> <li>Urine = NP</li> <li>Stool/Wound = NP</li> </ul>	Doxycycline (PO)	Erythromycin (PO) Azithromycin (PO) Any 1 <sup>st</sup> , 2 <sup>nd</sup> , 3 <sup>rd</sup> generation cephalosporin (PO) Clarithromycin XL (PO)
Bacillus anthracis (naturally acquired) (For potential bioterrorist anthrax, see p. 173)	<ul style="list-style-type: none"> <li>CSF = P (ABM)</li> <li>Blood = P (septicemia; isolation required; dangerous)</li> <li>Sputum = P (mediastinitis; anthrax pneumonia rare)</li> <li>Urine = NP</li> <li>Stool = NP</li> <li>Wound = P (ulcer; isolation required; dangerous)</li> </ul>	Penicillin (IV) Doxycycline (IV/PO) Any quinolone (IV/PO)	Amoxicillin (PO) Ampicillin (IV)
Bacillus cereus, subtilis, megaterium	<ul style="list-style-type: none"> <li>CSF = NP</li> <li>Blood = C*</li> <li>P (leukopenic compromised hosts)</li> </ul>	Vancomycin (IV) Clindamycin (IV/PO)	Meropenem (IV) Any quinolone (IV/PO)

C = colonizer; C\* = skin contaminant; NP = non-pathogen at site; P = pathogen at site; (IV/PO) = IV or PO. See p. xi for all other abbreviations.

Causes membranous pharyngitis with scarlet fever-like rash. Differentiate from *C. diphtheriae* by culture. Penicillin and ampicillin are less effective than erythromycin or doxycycline.

Doxycycline may be used for therapy/outbreak prophylaxis. Streptobacillary configuration in blood. Causes hemorrhagic meningitis, wound infections, and bacteremia. Quinolones are effective. Alert microbiology laboratory of potentially biohazardous specimens.

Soil organisms not commonly pathogenic for humans. Suspect pseudoinfection if isolated from clinical specimens. Look for soil/dust contamination of blood culture tube top/

Table 3.6. Usual Clinical Significance of AEROBIC Isolates Pending Susceptibility Testing (cont'd)

GRAM-POSITIVE BACILLI				
Isolate	Isolate Significance	Preferred Therapy	Alternate Therapy	Comments
Corynebacterium diphtheriae	<ul style="list-style-type: none"> <li>• Sputum = NP</li> <li>• Urine = NP</li> <li>• Stool = NP</li> <li>• Wound = NP</li> </ul>	Penicillin (IV) Erythromycin (IV) Clindamycin (IV/PO)	Doxycycline (IV/PO) Clarithromycin XL (PO) Rifampin (PO)	<p>apparatus. Rare pathogen in leukopenic compromised hosts.</p> <p>Administer diphtheria antitoxin as soon as possible (p. 170). Antibiotic therapy is adjunctive, since diphtheria is a toxin-mediated disease. Patients may die unexpectedly from toxin-induced myocarditis during recovery.</p>
Corynebacterium jeikeium (JK)	<ul style="list-style-type: none"> <li>• CSF = C*, P (CSF shunts)</li> <li>• Blood = C*, P (from IV lines)</li> <li>• Sputum = NP</li> <li>• Urine/Stool = NP</li> <li>• Wound = C</li> </ul>	Vancomycin (IV) Linezolid (IV/PO)	Quinupristin/dalfopristin (IV)	Cause of IV line/foreign body infections. In-vitro testing is not always reliable. Highly resistant to most anti-gram positive antibiotics.
Erysipelothrix rhusiopathiae	<ul style="list-style-type: none"> <li>• CSF = NP</li> <li>• Blood = P (from SBE)</li> <li>• Sputum = NP</li> <li>• Urine = NP</li> <li>• Stool = NP</li> <li>• Wound = P (Chronic erysipelas-like skin lesions)</li> </ul>	Penicillin (IV) Ampicillin (IV)	Any 3 <sup>rd</sup> generation cephalosporin (IV) Any quinolone (IV/PO)	Cause of "culture-negative" SBE. Susceptible to clindamycin but resistant to vancomycin.

Listeria monocytogenes	<ul style="list-style-type: none"> <li>• CSF = P (ABM)</li> <li>• Blood = P (1° bacteremia, SBE)</li> <li>• Sputum = NP</li> <li>• Urine = NP</li> <li>• Stool = NP</li> <li>• Wound = NP</li> </ul>	<p>Ampicillin (IV) Amoxicillin (PO) Chloramphenicol (IV)</p> <p>CNS Ampicillin (IV) TMP-SMX (IV/PO) Chloramphenicol (IV/PO)</p> <p>SBE Ampicillin (IV)</p>	<p>Doxycycline (IV/PO) Erythromycin (IV) Meropenem (IV)</p> <p>CNS Meropenem (IV) (Meningeal dosed) Linezolid (IV/PO)</p> <p>SBE Meropenem (IV) Linezolid (IV/PO)</p>	<p>Listeria ABM is common in T-cell deficiencies (e.g., lymphoma, steroids, HIV). Causes SBE in normal hosts, and is the commonest cause of bacteremia in non-neutropenic cancer patients. 3<sup>rd</sup> generation cephalosporins are ineffective against Listeria.</p>
Nocardia asteroides, brasiliensis	<ul style="list-style-type: none"> <li>• CSF = P (brain abscess)</li> <li>• Blood = P (from lung/soft tissue source)</li> <li>• Sputum = P (pneumonia, lung abscess)</li> <li>• Urine/Stool = NP</li> <li>• Wound = P (skin lesions from direct inoculation or dissemination)</li> </ul>	<p>TMP-SMX (IV/PO) Minocycline (IV/PO)</p>	<p>Imipenem (IV) plus either amikacin (IV) or any 3<sup>rd</sup> generation cephalosporin (IV)</p>	<p>Branched, filamentous, beady hyphae are typical, but coccobacillary and bacillary forms are also common. Nocardia are gram-positive, aerobic, and acid fast. Linezolid is active against Nocardia and may be effective if other agents cannot be used. Quinolones and macrolides are usually ineffective.</p>
Rhodococcus equi	<ul style="list-style-type: none"> <li>• CSF = NP</li> <li>• Blood = P (from pneumonia, lung abscess)</li> <li>• Sputum = P (pneumonia with abscess/cavitation)</li> </ul>	<p>Any quinolone (IV/PO) Vancomycin (IV)</p>	<p>Erythromycin (IV) Imipenem (IV) Meropenem (IV) Doxycycline (IV/PO) TMP-SMX (IV/PO)</p>	<p>Causes TB-like community-acquired pneumonia in AIDS patients. Filamentous bacteria break into bacilli/cocci. Aminoglycosides and <math>\beta</math>-lactams are relatively ineffective.</p>

C = colonizer; C<sup>+</sup> = skin contaminant; NP = non-pathogen at site; P = pathogen at site; (IV/PO) = IV or PO. See p. xi for all other abbreviations.



Table 3.6. Usual Clinical Significance of AEROBIC Isolates Pending Susceptibility Testing (cont'd)

Isolate	Isolate Significance	GRAM-POSITIVE BACILLI		Comments
		Preferred Therapy	Alternate Therapy	
<b>GRAM-NEGATIVE BACILLI</b>				
Acinetobacter baumannii, lwoffii, calcoaceticus, haemolyticus	<ul style="list-style-type: none"> <li>CSF = C*, P (ABM)</li> <li>Blood = P (from IV line, lung, or urine source)</li> <li>Sputum = C, P (VAP)</li> <li>Urine = C, P (CAB)</li> <li>Stool = NP</li> <li>Wound = C (common), P (rare)</li> </ul>	Any carbapenem (IV) Ampicillin/sulbactam (IV) Colistin Polymyxin B Minocycline (IV/PO)	Any 3 <sup>rd</sup> generation cephalosporin (IV) (except ceftazidime) Cefepime (IV) Fosfomycin (PO)	Colonization common; infection uncommon. If possible, avoid treating Acinetobacter colonization in respiratory secretions or urine (CAB). Occurs in outbreaks of ventilator-associated pneumonia. Test susceptibility to each carbapenem (may be susceptible to one but not others). Use meropenem for MDR susceptible isolates. For meropenem resistant MDR isolates, use colistin, polymyxin B, tigacycline, minocycline, or doripenem.
Aeromonas hydrophila	<ul style="list-style-type: none"> <li>CSF = NP</li> <li>Blood = P (from wound, urine, or GI source)</li> <li>Sputum = NP</li> <li>Urine = C, P (CAB)</li> <li>Stool = P (diarrhea)</li> <li>Wound = P (cellulitis)</li> </ul>	Gentamicin (IV) TMP-SMX (IV/PO) Any quinolone (IV/PO)	Doxycycline (IV/PO) Any 3 <sup>rd</sup> generation cephalosporin (IV/PO) Any carbapenem (IV) Aztreonam (IV)	Cause of wound infection, septic arthritis, diarrhea, and necrotizing soft tissue infection resembling gas gangrene.
Aggregatibacter (Actinobacillus) actinomycetem-comitans	<ul style="list-style-type: none"> <li>CSF = NP</li> <li>Blood = P (from abscess, SBE)</li> <li>Sputum = NP</li> <li>Urine/Stool = NP</li> </ul>	Any quinolone (IV/PO) Any 3 <sup>rd</sup> generation cephalosporin (IV/PO)	Penicillin (IV) + gentamicin (IV) TMP-SMX (IV/PO)	Cause of "culture-negative" SBE. One of the HACEK organisms. Found with Actinomyces in abscesses. Resistant to erythromycin and clindamycin.

Alcaligenes (Achromobacter) xylosoxidans	<ul style="list-style-type: none"> <li>Wound = P (from abscess, draining fistulous tract)</li> <li>CSF = P (rarely ABM)</li> <li>Blood = P (from urine)</li> <li>Sputum = NP</li> <li>Urine = P (CAB)</li> <li>Stool = NP</li> <li>Wound = P (cellulitis rare)</li> </ul>	<ul style="list-style-type: none"> <li>Imipenem (IV)</li> <li>Meropenem (IV)</li> <li>Any 3<sup>rd</sup> generation cephalosporin (IV/PO)</li> </ul>	<ul style="list-style-type: none"> <li>Any quinolone (IV/PO)</li> <li>Cefepime (IV)</li> <li>Aztreonam (IV)</li> </ul>	Water-borne pathogen resembling Acinetobacter microbiologically. Resistant to aminoglycosides and 1 <sup>st</sup> , 2 <sup>nd</sup> generation cephalosporins.
Bartonella henselae, quintana, bacilliformis	<ul style="list-style-type: none"> <li>CSF = NP</li> <li>Blood = P (from skin source, SBE)</li> <li>Sputum = NP</li> <li>Urine = NP</li> <li>Stool = NP</li> <li>Wound = P (skin lesions)</li> </ul>	<ul style="list-style-type: none"> <li>Doxycycline (IV/PO)</li> <li>Azithromycin (PO)</li> </ul>	<ul style="list-style-type: none"> <li>Clarithromycin XL (PO)</li> <li>Any quinolone (IV/PO)</li> <li>Any aminoglycoside (IV)</li> </ul>	B. henselae (bacteremia, endocarditis, peliosis hepatis, bacillary angiomatosis); B. quintana (relapsing, trench fever, bacillary angiomatosis); B. bacilliformis (Oroyo fever, Carrion's disease). May present as FUO. Titters may cross react with C. burnetii (Q fever). TMP-SMX and cephalosporins are ineffective.
Bordetella pertussis, parapertussis	<ul style="list-style-type: none"> <li>CSF = NP</li> <li>Blood = P (from respiratory tract source)</li> <li>Sputum = C, P (pertussis)</li> <li>Urine = NP</li> <li>Stool = NP</li> <li>Wound = NP</li> </ul>	<ul style="list-style-type: none"> <li>Erythromycin (IV)</li> <li>Clarithromycin XL (PO)</li> <li>Azithromycin (IV/PO)</li> </ul>	<ul style="list-style-type: none"> <li>Any quinolone (IV/PO)</li> <li>TMP-SMX (IV/PO)</li> <li>Doxycycline (IV/PO)</li> </ul>	Causes pertussis in children and incompletely/non-immunized adults. Macrolides remain the preferred therapy. Resistant to penicillins, cephalosporins, and aminoglycosides.
Brucella abortus, canis, suis, melitensis	<ul style="list-style-type: none"> <li>CSF = P (meningitis)</li> <li>Blood = P (from abscess, SBE)</li> </ul>	<ul style="list-style-type: none"> <li>Doxycycline (IV/PO) + gentamicin (IV)</li> </ul>	<ul style="list-style-type: none"> <li>TMP-SMX (IV/PO) + gentamicin (IV)</li> </ul>	Causes prolonged relapsing infection. Zoonotic cause of brucellosis/Malta fever. Resistant to penicillins.

C = colonizer; C\* = skin contaminant; NP = non-pathogen at site; P = pathogen at site; (IV/PO) = IV or PO. See p. xiv for all other abbreviations.

Table 3.6. Usual Clinical Significance of AEROBIC Isolates Pending Susceptibility Testing (cont'd)

GRAM-NEGATIVE BACILLI				
Isolate	Isolate Significance	Preferred Therapy	Alternate Therapy	Comments
	<ul style="list-style-type: none"> <li>Sputum = NP</li> <li>Urine = P (pyelonephritis)</li> <li>Stool/Wound = NP</li> </ul>	Doxycycline + streptomycin (IM)	Doxycycline (IV/PO) + rifampin (PO) Any quinolone (IV/PO) + rifampin (PO)	
Burkholderia (Pseudomonas) cepacia	<ul style="list-style-type: none"> <li>CSF = NP</li> <li>Blood = P (usually from IV line/urinary tract infection)</li> <li>Sputum = C (not a cause of VAP)</li> <li>Urine = C</li> <li>Stool = NP</li> <li>Wound = NP</li> </ul>	TMP-SMX (IV/PO) Minocycline (IV/PO)	A "respiratory quinolone" (IV/PO) Chloramphenicol (IV/PO)	Rare cause of urosepsis following urologic instrumentation. Common water-borne colonizer in intensive care units. Opportunistic pathogen in cystic fibrosis/bronchiectasis. Resistant to aminoglycosides, colistin, and polymyxin B.
Burkholderia (Pseudomonas) pseudomallei	<ul style="list-style-type: none"> <li>CSF = NP</li> <li>Blood = P (from septicemic melioidosis)</li> <li>Sputum = P (chronic cavitary pneumonia)</li> <li>Urine = NP</li> <li>Stool = NP</li> <li>Wound = NP</li> </ul>	Imipenem (IV) Meropenem (IV) Ceftazidime (IV) Doxycycline (IV/PO)	Chloramphenicol (IV) TMP-SMX (IV/PO) Amoxicillin/clavulanic acid (PO)	Acute (septicemia)/chronic (cavitary CAP/abscesses) melioidosis endemic in S.E. Asia. Chronic melioidosis resembles reactivation TB, but has lower lobe distribution. Prolonged latency until reactivation years later. Slow response to effective therapy (1-2 weeks). Prolonged therapy needed to prevent relapse (≥ 3 months). Oxidase positive. Resistant to penicillin, aminoglycosides, colistin.
Campylobacter fetus	<ul style="list-style-type: none"> <li>CSF = P (ABM)</li> <li>Blood = P (from vascular source)</li> </ul>	Gentamicin (IV) Imipenem (IV) Meropenem (IV)	Chloramphenicol (IV) Ampicillin (IV)	Causes invasive infection with spread to CNS. CNS infection may be treated with meningeal doses of chloramphenicol, ampicillin,

	<ul style="list-style-type: none"> <li>• Sputum = NP</li> <li>• Urine = NP</li> <li>• Stool = NP</li> <li>• Wound = NP</li> </ul>		Any 3 <sup>rd</sup> generation cephalosporin (IV)	or a 3 <sup>rd</sup> generation cephalosporin. Resistant to erythromycin.
Campylobacter jejuni	<ul style="list-style-type: none"> <li>• CSF = NP</li> <li>• Blood = P (from GI source)</li> <li>• Sputum = NP</li> <li>• Urine = NP</li> <li>• Stool = P (diarrhea)</li> <li>• Wound = NP</li> </ul>	Any quinolone (IV/PO) Erythromycin (PO) Doxycycline (IV/PO)	Azithromycin (PO) Clarithromycin XL (PO)	Commonest cause of acute bacterial diarrhea. Resistant to TMP-SMX.
Cardiobacterium hominis	<ul style="list-style-type: none"> <li>• CSF = NP</li> <li>• Blood = P (from SBE)</li> <li>• Sputum = NP</li> <li>• Urine = NP</li> <li>• Stool = NP</li> <li>• Wound = NP</li> </ul>	Penicillin (IV) + gentamicin (IV) Ampicillin (IV) + gentamicin (IV)	Any 3 <sup>rd</sup> generation cephalosporin (IV) + gentamicin (IV)	Pleomorphic bacillus with bulbous ends. Often appears in clusters resembling rosettes. Indole positive. Cause of "culture-negative" SBE (one of the HACEK organisms). Rare cause of abdominal abscess. Grows best with CO <sub>2</sub> enhancement. Resistant to macrolides and clindamycin.
Chromobacterium violaceum	<ul style="list-style-type: none"> <li>• CSF = NP</li> <li>• Blood = P (from wound infection)</li> <li>• Sputum = NP</li> <li>• Urine = NP</li> <li>• Stool = NP</li> <li>• Wound = P (drainage from deep soft tissue infection)</li> </ul>	Gentamicin (IV) Doxycycline (IV/PO)	Chloramphenicol (IV)	Cause of cutaneous lesions primarily in tropical/subtropical climates. Often mistaken for Vibrio or Alcaligenes. Resistant to $\beta$ -lactams.
Chryseobacterium (Flavobacterium) meningosepticum	<ul style="list-style-type: none"> <li>• CSF = P (ABM)</li> <li>• Blood = P (from IV line infection, PVE)</li> <li>• Sputum = NP</li> </ul>	CNS TMP-SMX (IV/PO)	CNS Chloramphenicol (IV)	Rare cause of ABM in newborns and PVE in adults. Only unencapsulated Flavobacterium species. Clindamycin, clarithromycin, and vancomycin are useful only in non-CNS

C = colonizer; C\* = skin contaminant; NP = non-pathogen at site; P = pathogen at site; (IV/PO) = IV or PO. See p. xi for all other abbreviations.

Table 3.6. Usual Clinical Significance of AEROBIC Isolates Pending Susceptibility Testing (cont'd)

GRAM-NEGATIVE BACILLI				
Isolate	Isolate Significance	Preferred Therapy	Alternate Therapy	Comments
	<ul style="list-style-type: none"> <li>Urine = C, P (from urologic instrumentation)</li> <li>Stool = NP</li> <li>Wound = C, P (cellulitis)</li> </ul>	<p><u>Non-CNS</u> Vancomycin (IV) + rifampin (PO) Any quinolone (IV/PO)</p>	<p><u>Non-CNS</u> Clarithromycin XL (PO) + rifampin (PO) Clindamycin (IV/PO)</p>	infections. Resistant to aztreonam and carbapenems.
Citrobacter diversus; freundii, koseri	<ul style="list-style-type: none"> <li>CSF = C*, P (from NS procedure)</li> <li>Blood = C*, P (from IV line/urinary tract infection)</li> <li>Sputum = C (not pneumonia)</li> <li>Urine = C, P (from urologic instrumentation)</li> <li>Stool = NP</li> <li>Wound = C, P (rarely in compromised hosts)</li> </ul>	<p>Any carbapenem (IV) Cefepime (IV) Any quinolone (IV/PO)</p>	<p>Aztreonam (IV) Piperacillin (IV) Any 3<sup>rd</sup> generation cephalosporin (IV)</p>	Common wound/urine colonizer. Rare pathogen in normal hosts. Often aminoglycoside resistant. (C. freundii is usually more resistant than C. koseri).
Edwardsiella tarda	<ul style="list-style-type: none"> <li>CSF = NP</li> <li>Blood = P (from liver abscess)</li> <li>Sputum/Urine = NP</li> <li>Stool = P</li> <li>Wound C, P (wound infection)</li> </ul>	<p>Ampicillin (IV) Amoxicillin (PO) Any quinolone (IV/PO)</p>	<p>Doxycycline (IV/PO) Any 3<sup>rd</sup> generation cephalosporin (IV/PO)</p>	Cause of bacteremia, usually from liver abscess or wound source.

<p>Enterobacter agglomerans, aerogenes, cloacae</p>	<ul style="list-style-type: none"> <li>• CSF = C*, P (from NS procedure)</li> <li>• Blood = C* P (from IV line/urinary tract infection)</li> <li>• Sputum = C (not a cause of pneumonia)</li> <li>• Urine = C, P (post-urologic instrumentation)</li> <li>• Stool = NP</li> <li>• Wound = C, P (rarely in compromised hosts)</li> </ul>	<p>Any carbapenem (IV)</p>	<p>Any quinolone (IV/PO)</p> <p>Aztreonam (IV)</p> <p>Piperacillin (IV)</p> <p>Cefepime (IV)</p>	<p>Not a cause of community-acquired or nosocomial pneumonia. Common colonizer of respiratory secretions and wound/urine specimens. Antibiotic resistance to E. cloacae &gt; E. aerogenes &gt; E. agglomerans. Treatment of Enterobacter colonizers with ceftazidime or ciprofloxacin may result in MDR/ESBL Enterobacter sp.</p> <p>CRE usually susceptible only to tigacycline, colistin, polymyxin B, ceftazidime/avibactam, fosfomycin.</p>
<p>Escherichia coli</p>	<ul style="list-style-type: none"> <li>• CSF = P (ABM)</li> <li>• Blood = P (from GI/GU source)</li> <li>• Sputum = P (rarely CAP from urinary source, VAP)</li> <li>• Urine = C, P (CAB, cystitis, pyelonephritis)</li> <li>• Stool = C, P (diarrhea)</li> <li>• Wound = P (cellulitis)</li> </ul>	<p>Any 1<sup>st</sup>, 2<sup>nd</sup>, 3<sup>rd</sup> generation cephalosporin (IV/PO)</p> <p>Amoxicillin (PO)</p> <p>Any quinolone (IV/PO)</p> <p>Ceftriaxone (IV)</p> <p>Nitrofurantoin (PO) (UTIs only)</p>	<p>Aztreonam (IV)</p> <p>Gentamicin (IV)</p> <p>TMP-SMX (IV/PO)</p>	<p>Common pathogen, usually from GI/GU source. Many strains are resistant to ampicillin and some to 1<sup>st</sup> generation cephalosporins. ESBL-producing E. coli may be treated with a carbapenem.</p> <p>CRE usually susceptible only to tigacycline, colistin, polymyxin B, ceftazidime/avibactam, fosfomycin.</p>
<p>Francisella tularensis</p>	<ul style="list-style-type: none"> <li>• CSF = NP</li> <li>• Blood = P (isolation dangerous)</li> <li>• Sputum = P (tularemic pneumonia; isolation dangerous)</li> </ul>	<p>Doxycycline (IV/PO)</p> <p>Gentamicin (IV/IM)</p> <p>Streptomycin (IM)</p>	<p>Chloramphenicol (IV/PO)</p> <p>Any quinolone (IV/PO)</p>	<p>Six clinical tularemia syndromes. Alert microbiology laboratory of potentially biohazardous specimens. Do not culture. Resistant to penicillins and cephalosporins. Bioterrorist tularemia is treated the same as naturally-acquired tularemia.</p>

C = colonizer; C\* = skin contaminant; NP = non-pathogen at site; P = pathogen at site; (IV/PO) = IV or PO. See p. xi for all other abbreviations.

Table 3.6. Usual Clinical Significance of AEROBIC Isolates Pending Susceptibility Testing (cont'd)

Isolate	GRAM-NEGATIVE BACILLI			Comments
	Isolate Significance	Preferred Therapy	Alternate Therapy	
Hafnia alvei	<ul style="list-style-type: none"> <li>Urine/Stool = NP</li> <li>Wound = P (isolation dangerous)</li> <li>CSF = C, P (from NS procedure)</li> <li>Blood = C*, P (from IV line/urinary tract infection)</li> <li>Sputum = C (not pneumonia)</li> <li>Urine = C, P (post-urolologic instrumentation)</li> <li>Stool = NP</li> <li>Wound = C, P (rarely in compromised hosts)</li> </ul>	Cefepime (IV) Any quinolone (IV/PO) Aztreonam (IV)	Piperacillin (IV) Imipenem (IV) Meropenem (IV)	Formerly Enterobacter hafniae. Uncommon nosocomial pathogen. Rarely pathogenic in normal hosts. Cause of UTIs in compromised hosts.
Helicobacter (Campylobacter) pylori	<ul style="list-style-type: none"> <li>CSF = NP</li> <li>Blood = NP</li> <li>Sputum = NP</li> <li>Urine = NP</li> <li>Stool = P (from upper GI tract biopsy specimens, not stool)</li> <li>Wound = NP</li> </ul>	Omeprazole (PO) + clarithromycin XL (PO) Omeprazole (PO) + amoxicillin (PO) Metronidazole (PO) + amoxicillin (PO) + bismuth subsalicylate (PO)	Doxycycline (PO) + metronidazole (PO) + bismuth subsalicylate (PO)	Optimal therapy awaits definition. Treat until cured. Some strains of resistant H. pylori may respond to treatment with a quinolone. TMP-SMX is ineffective.

Haemophilus influenzae, parainfluenzae, aphrophilus, paraphrophilus	<ul style="list-style-type: none"> <li>• CSF = P (ABM)</li> <li>• Blood = P (from respiratory tract or cardiac source)</li> <li>• Sputum = C, P (CAP)</li> <li>• Urine = NP</li> <li>• Stool = NP</li> <li>• Wound = P</li> </ul>	<p><u>For all Haemophilus species</u></p> <p>Any 2<sup>nd</sup>, 3<sup>rd</sup> generation cephalosporin (IV/PO)</p> <p>Any quinolone (IV/PO)</p> <p>Doxycycline (IV/PO)</p>	<p><u>For all Haemophilus species</u></p> <p>Chloramphenicol (IV)</p> <p>TMP-SMX (IV/PO)</p> <p>Azithromycin (PO)</p> <p>Aztreonam (IV)</p> <p><u>Ampicillin-resistant H. influenzae</u></p> <p>Meropenem (IV)</p> <p>Imipenem (IV)</p> <p>Ertapenem (IV)</p> <p>Cefepime (IV)</p> <p>Aztreonam (IV)</p>	<p>1<sup>st</sup> generation cephalosporins, erythromycin, and clarithromycin have limited anti-H. influenzae activity; doxycycline and azithromycin are better. Hemophilus species are common colonizers of the respiratory tract. Rarely a cause of "culture-negative" SBE (H. parainfluenzae/aphrophilus are HACEK organisms). H. influenzae and H. parainfluenzae (pathogens) may be differentiated from the throat commensals H. hemolyticus and H. parahemolyticus (non-pathogens) by hemolysis on sheep agar.</p>
Kingella (Moraxella) kingae	<ul style="list-style-type: none"> <li>• CSF = NP</li> <li>• Blood = P (from skeletal or cardiac source)</li> <li>• Sputum = C</li> <li>• Urine = NP</li> <li>• Stool/wound = NP</li> </ul>	<p>Ampicillin (IV) + any aminoglycoside (IV)</p>	<p>Any 3<sup>rd</sup> generation cephalosporin + any aminoglycoside (IV)</p> <p>Imipenem (IV)</p> <p>Meropenem (IV)</p> <p>Any quinolone (IV/PO)</p>	<p>Common colonizer of respiratory tract, but rarely a respiratory pathogen. Causes septic arthritis/osteomyelitis in children and endocarditis in adults (one of HACEK organisms). Oxidase positive. Growth enhanced with CO<sub>2</sub>.</p>
Klebsiella pneumoniae, oxytoca	<ul style="list-style-type: none"> <li>• CSF = P (ABM)</li> <li>• Blood = P (from respiratory, GI, GU source)</li> </ul>	<p>Tigecycline (IV)</p> <p>Any carbapenem (IV)</p>	<p>Any 3<sup>rd</sup> generation cephalosporin (IV, PO) except ceftazidime</p> <p>Any quinolone (IV/PO)</p>	<p>TMP-SMX may be ineffective in systemic infection. Anti-pseudomonal penicillins have limited anti-Klebsiella activity. Klebsiella usually susceptible to carbapenems. CRE usually susceptible only to tigecycline, colistin, polymyxin B, ceftazidime/avibactam, fosfomycin.</p>

C = colonizer; C\* = skin contaminant; NP = non-pathogen at site; P = pathogen at site; (IV/PO) = IV or PO. See p. xi for all other abbreviations.



**Table 3.6. Usual Clinical Significance of AEROBIC Isolates Pending Susceptibility Testing (cont'd)**

GRAM-NEGATIVE BACILLI					
Isolate	Isolate Significance	Preferred Therapy	Alternate Therapy		Comments
			Therapy	Therapy	
Klebsiella ozaenae, rhinoscleromatis	<ul style="list-style-type: none"> <li>Sputum = C, P (CAP/VAP)</li> <li>Urine = C (CAB), P</li> <li>Stool = NP</li> <li>Wound = C, P</li> <li>CSF = NP</li> <li>Blood = NP</li> <li>Sputum = NP</li> <li>Urine = NP</li> <li>Stool = NP</li> <li>Wound = P (rhinoscleromatis lesions)</li> </ul>	Any quinolone (PO)	Aztreonam (IV) Cefepime (IV)	NDM-1 metallo $\beta$ -lactamase strains are carbapenem resistant and usually susceptible only to colistin, tigacycline.	
Legionella sp.	<ul style="list-style-type: none"> <li>CSF = NP</li> <li>Blood = NP</li> <li>Sputum = P (CAP or VAP)</li> <li>Urine = NP</li> <li>Stool = NP</li> <li>Wound = NP</li> </ul>	Any quinolone (IV/PO) Doxycycline (IV/PO) Tigacycline (IV) Azithromycin (IV/PO)	Clarithromycin XL (PO) Erythromycin (IV) Telithromycin (PO)	TMP-SMX (PO) + rifampin (PO)	Skin infection usually requires prolonged treatment for cure (weeks-to-months).  Anti-Legionella activity: quinolones > doxycycline > erythromycin. Erythromycin failures are not uncommon. Rarely a cause of culture-negative SBE/PVE.
Leptospira interrogans	<ul style="list-style-type: none"> <li>CSF = P (ABM)</li> <li>Blood = P (1° bacteremia)</li> <li>Sputum = NP</li> <li>Urine = P (excreted in urine)</li> <li>Stool = NP</li> <li>Wound = NP</li> </ul>	Doxycycline (IV/PO) Penicillin G (IV) Any 3 <sup>rd</sup> generation cephalosporin (IV/PO)	Amoxicillin (PO)		Blood/urine cultures may be positive during initial/bacteremic phase, but are negative during immune phase. Relapse is common. Resistant to chloramphenicol.

Moraxella (Branhamella) catarrhalis	<ul style="list-style-type: none"> <li>• CSF = NP</li> <li>• Blood = P (rarely from CAP)</li> <li>• Sputum = C, P (CAP)</li> <li>• Urine = NP</li> <li>• Stool = NP</li> <li>• Wound = NP</li> </ul>	<p>Any 2<sup>nd</sup>, 3<sup>rd</sup> generation cephalosporin (IV/PO)</p> <p>Any quinolone (IV/PO)</p> <p>Telithromycin (PO)</p> <p>Doxycycline (IV/PO)</p>	<p>Azithromycin (PO)</p> <p>Clarithromycin XL (PO)</p> <p>TMP-SMX (IV/PO)</p> <p>Amoxicillin/clavulanic acid (PO)</p>	<p>Almost all strains are <math>\beta</math>-lactamase positive and resistant to penicillin/ampicillin. <math>\beta</math>-lactamase-resistant <math>\beta</math>-lactams are effective. Resembles Acinetobacter on sputum gram stain.</p>
Morganella morganii	<ul style="list-style-type: none"> <li>• CSF = NP</li> <li>• Blood = P (from GU source)</li> <li>• Sputum = NP</li> <li>• Urine = P (CAB, cystitis, pyelonephritis)</li> <li>• Stool = NP</li> <li>• Wound = P (cellulitis rare)</li> </ul>	<p>Any quinolone (IV/PO)</p> <p>Any 3<sup>rd</sup> generation cephalosporin (IV)</p> <p>Any carbapenem (IV)</p>	<p>Any aminoglycoside (IV)</p> <p>Aztreonam (IV)</p> <p>Cefepime (IV)</p>	<p>Common uropathogen. Causes bacteremia with urosepsis. Rare cause of wound infections.</p>
Chrobactrum anthropi (CDC group Vd)	<ul style="list-style-type: none"> <li>• CSF = NP</li> <li>• Blood = P (from IV line infections)</li> <li>• Sputum = C</li> <li>• Urine = C</li> <li>• Stool/Wound = C</li> </ul>	<p>Any quinolone (IV/PO)</p> <p>TMP-SMX (IV/PO)</p>	<p>Any aminoglycoside (IV)</p> <p>Imipenem (IV)</p> <p>Meropenem (IV)</p>	<p>Pathogen in compromised hosts. Oxidase and catalase positive. Resistant to <math>\beta</math>-lactams.</p>
Pasteurella multocida	<ul style="list-style-type: none"> <li>• CSF = P (ABM)</li> <li>• Blood = P (from respiratory source, bite wound/abscess)</li> <li>• Sputum = C, P (CAP, bronchiectasis)</li> </ul>	<p>Amoxicillin (PO)</p> <p>Doxycycline (IV/PO)</p> <p>Penicillin G (IV)</p>	<p>Ampicillin/sulbactam (IV)</p> <p>Piperacillin (IV)</p> <p>Any quinolone (IV/PO)</p>	<p>Common cause of infection following dog/cat bites. Many antibiotics are effective, but erythromycin is ineffective. Resembles Hemophilus sp. on sputum gram stain.</p>

C = colonizer; C\* = skin contaminant; NP = non-pathogen at site; P = pathogen at site; (IV/PO) = IV or PO. See p. xi for all other abbreviations.

Table 3.6. Usual Clinical Significance of AEROBIC Isolates Pending Susceptibility Testing (cont'd)

GRAM-NEGATIVE BACILLI				
Isolate	Isolate Significance	Preferred Therapy	Alternate Therapy	Comments
<i>Plesiomonas shigelloides</i>	<ul style="list-style-type: none"> <li>• Urine = C, P (pyelonephritis)</li> <li>• Stool = NP</li> <li>• Wound = P (human/animal bites)</li> <li>• CSF = NP</li> <li>• Blood = P (from GU source)</li> <li>• Sputum = NP</li> <li>• Urine = NP</li> <li>• Stool = P (diarrhea)</li> <li>• Wound = NP</li> </ul>	Any quinolone (PO) TMP-SMX (PO)	Doxycycline (PO) Aztreonam (PO)	Infrequent cause of diarrhea, less commonly dysentery. Oxidase positive. $\beta$ -lactamase strains are increasing. Resistant to penicillins.
<i>Proteus mirabilis, vulgaris</i>	<ul style="list-style-type: none"> <li>• CSF = NP</li> <li>• Blood = P (from urinary source)</li> <li>• Sputum = C</li> <li>• Urine = C, P (from urologic instrumentation)</li> <li>• Stool = NP</li> <li>• Wound = C, P (wound infection)</li> </ul>	<p><u><i>P. mirabilis</i>, indole (-)</u></p> <p>Ampicillin (IV)</p> <p>Any 1<sup>st</sup>, 2<sup>nd</sup>, 3<sup>rd</sup> gen. cephalosporin (IV/PO)</p> <p><u><i>P. vulgaris</i>, indole (+)</u></p> <p>Any 3<sup>rd</sup> generation cephalosporin (IV/PO)</p> <p>Cefepime (IV)</p> <p>Any quinolone (IV/PO)</p>	<p><u><i>P. mirabilis</i>, indole (-)</u></p> <p>TMP-SMX (IV/PO)</p> <p>Amoxicillin (PO)</p> <p><u><i>P. vulgaris</i>, indole (+)</u></p> <p>Aztreonam (IV) Any carbapenem (IV)</p> <p>Any aminoglycoside (IV)</p>	Usually a uropathogen. Most antibiotics are effective against <i>P. mirabilis</i> (indole-negative); <i>P. penneri</i> (indole-negative <i>P. vulgaris</i> ) is resistant to ceftriaxone. Indole-positive <i>Proteus</i> sp. require potent antibiotics to treat non-UTIs. <i>P. penneri</i> (indole-negative <i>P. vulgaris</i> ) resistant to 3 <sup>rd</sup> gen. cephalosporins; use cefepime, carbapenem, or quinolone.

Providencia alcalifaciens, rettgeri, stuartii	<ul style="list-style-type: none"> <li>• CSF = NP</li> <li>• Blood = C*, P (from GU source)</li> <li>• Sputum/Stool = NP</li> <li>• Urine = C, P</li> <li>• Wound = C, P (rare)</li> </ul>	<p>Any quinolone (IV/PO)</p> <p>Any 3<sup>rd</sup> generation cephalosporin (IV/PO)</p> <p>Cefepime (IV)</p> <p>Meropenem (IV)</p> <p>Ertapenem (IV)</p>	<p>Any aminoglycoside (IV)</p> <p>Aztreonam (IV)</p> <p>Piperacillin (IV)</p> <p>Imipenem (IV)</p>	<p>Almost always a uropathogen. Formerly classified as indole-positive Proteus.</p>
Pseudomonas aeruginosa	<ul style="list-style-type: none"> <li>• CSF = NP</li> <li>• Blood = P (from respiratory, GU source)</li> <li>• Sputum = C (usually), P (rarely indicates VAP)</li> <li>• Urine = C, P (from urologic instrumentation)</li> <li>• Stool = NP</li> <li>• Wound = C (almost always)</li> </ul>	<p><u>Mono</u>therapy</p> <p>Meropenem (IV)</p> <p>Cefepime (IV)</p> <p><u>Combination</u> therapy</p> <p>either meropenem (IV) or cefepime (IV) <i>plus</i> amikacin</p>	<p>Doripenem (IV)</p> <p>Colistin (IV)</p> <p>Polymyxin B (IV)</p> <p>Amikacin (IV)</p> <p>Aztreonam (IV)</p>	<p>Colonization common; infection uncommon. If possible, avoid treating P. aeruginosa in respiratory secretions in ventilated patients (unless tracheobronchitis) or urine cultures (CAB). For serious systemic P. aeruginosa infection, double-drug therapy preferred. All double anti-P. aeruginosa regimens are equally effective. Individual differences in activity (MICs) are unimportant if combination therapy is used. If MDR P. aeruginosa meropenem susceptible, treat with meropenem. If MDR P. aeruginosa meropenem resistant, treat with colistin, polymyxin B, or doripenem.</p>
Pseudomonas (Chryseomonas) luteola (CDC group Ve-1)	<ul style="list-style-type: none"> <li>• CSF = NP</li> <li>• Blood = P (from IV line infection)</li> <li>• Sputum = NP</li> <li>• Urine = NP</li> <li>• Stool = NP</li> <li>• Wound = NP</li> </ul>	<p>Imipenem (IV)</p> <p>Meropenem (IV)</p> <p>Cefepime (IV)</p>	<p>Piperacillin/ tazobactam (IV)</p> <p>Aztreonam (IV)</p>	<p>Opportunistic pathogen primarily in compromised hosts.</p>

C = colonizer; C\* = skin contaminant; NP = non-pathogen at site; P = pathogen at site; (IV/PO) = IV or PO. See p. xi for all other abbreviations.

Table 3.6. Usual Clinical Significance of AEROBIC Isolates Pending Susceptibility Testing (cont'd)

GRAM-NEGATIVE BACILLI				
Isolate	Isolate Significance	Preferred Therapy	Alternate Therapy	Comments
<i>Pseudomonas</i> (Flavimonas) oryzihabitans (CDC group Ve-2)	<ul style="list-style-type: none"> <li>• CSF = P (NS procedures)</li> <li>• Blood = P (from IV line infection)</li> <li>• Sputum = NP</li> <li>• Urine = NP</li> <li>• Stool = NP</li> <li>• Wound = P (rare)</li> </ul>	Imipenem (IV) Meropenem (IV) Cefepime (IV)	Any 3 <sup>rd</sup> generation cephalosporin (IV) Piperacillin (IV) Aztreonam (IV)	Rare cause of central IV line infections in compromised hosts (usually in febrile neutropenics). Rare cause of peritonitis in CAPD patients. Oxidase negative, unlike other <i>Pseudomonas</i> species.
<i>Salmonella typhi</i> , non-typhi	<ul style="list-style-type: none"> <li>• CSF = NP</li> <li>• Blood = P (from GI source)</li> <li>• Sputum = NP</li> <li>• Urine = P (only with enteric fever)</li> <li>• Stool = C (carrier), P (gastroenteritis, enteric fever)</li> <li>• Wound = NP</li> </ul>	Any quinolone (IV/PO) Any 3 <sup>rd</sup> generation cephalosporin (IV)	Chloramphenicol (IV) TMP-SMX (IV/PO) Doxycycline (IV/PO)	Carrier state is best eliminated by a quinolone or TMP-SMX. If drug therapy fails to eliminate carrier state, look for hepatic/bladder calculi for persistent focus. Many strains are resistant to ampicillin/amoxicillin.
<i>Serratia marcescens</i>	<ul style="list-style-type: none"> <li>• CSF = P (from NS procedures)</li> <li>• Blood = P (from IV line or urinary source)</li> <li>• Sputum = C, P (rarely in VAP)</li> <li>• Urine = C, P (post-urologic instrumentation)</li> <li>• Stool = NP</li> <li>• Wound = C, P (rare)</li> </ul>	Any 3 <sup>rd</sup> generation cephalosporin (IV/PO) (except ceftazidime) Any quinolone (IV/PO) Cefepime (IV)	Any carbapenem (IV) Gentamicin (IV) Aztreonam (IV) Piperacillin (IV)	Enterobacteriaceae. Associated with water sources. Common colonizer of respiratory secretions/urine in ICU. <i>Serratia nosocomial</i> pneumonia and PVE are rare. Cause of septic arthritis, osteomyelitis, and SBE (IV drug abusers). Among the aminoglycosides, gentamicin has the greatest anti- <i>Serratia</i> activity.

Shigella boydii, sonnei, flexneri, dysenteriae	<ul style="list-style-type: none"> <li>• CSF = NP</li> <li>• Blood = P (from GI source)</li> <li>• Sputum = NP</li> <li>• Urine = NP</li> <li>• Stool = P (Shigella dysentery)</li> <li>• Wound = NP</li> </ul>	Any quinolone (IV/PO)	TMP-SMX (IV/PO) Azithromycin (IV/PO)	No carrier state. Severity of dysentery varies with the species: S. dysenteriae (most severe) > S. flexneri > S. sonnei/boydii (least severe).
Stenotrophomonas (Pseudomonas, Xanthomonas) maltophilia	<ul style="list-style-type: none"> <li>• CSF = C, P (from NS procedures)</li> <li>• Blood = C*, P (from IV line infection, GU source)</li> <li>• Sputum = C (not VAP)</li> <li>• Urine = C, P (from urologic instrumentation)</li> <li>• Stool = NP</li> <li>• Wound = C, P (rarely in compromised hosts)</li> </ul>	TMP-SMX (IV/PO) Minocycline (IV/PO)	Cefepime (IV) Any respiratory quinolone (IV/PO)	Potential pulmonary pathogen only in bronchiectasis/cystic fibrosis. Resistant to aminoglycosides and carbapenems. Although usually carbapenem resistant, ~60% of strains demonstrate synergy with meropenem + levofloxacin. Susceptible to chloramphenicol, rifampin, colistin, polymyxin B.
Streptobacillus moniliformis	<ul style="list-style-type: none"> <li>• CSF = P (brain abscess)</li> <li>• Blood = P (from wound)</li> <li>• Sputum = P (lung abscess)</li> <li>• Urine = NP</li> <li>• Stool = NP</li> <li>• Wound = P (from rat bite)</li> </ul>	Penicillin (IV) Ampicillin (IV) Amoxicillin (PO)	Oxyacycline (IV/PO) Erythromycin (IV) Clindamycin (IV/PO)	Cause of Haverhill fever and rat-bite fever, with abrupt onset of severe headache/arthritis after bite wound has healed. No regional adenopathy. Morbilliform/petechial rash. Arthritis in 50%. May cause SBE.

C = colonizer; C\* = skin contaminant; NP = non-pathogen at site; P = pathogen at site; (IV/PO) = IV or PO. See p. xi for all other abbreviations.

Table 3.6. Usual Clinical Significance of AEROBIC Isolates Pending Susceptibility Testing (cont'd)

GRAM-NEGATIVE BACILLI				
Isolate	Isolate Significance	Preferred Therapy	Alternate Therapy	Comments
<i>Vibrio cholerae</i>	<ul style="list-style-type: none"> <li>• CSF = NP</li> <li>• Blood = P (from GI source)</li> <li>• Sputum = NP</li> <li>• Urine = NP</li> <li>• Stool = P (cholera)</li> <li>• Wound = NP</li> </ul>	Doxycycline (IV/PO) Any quinolone (IV/PO)	TMP-SMX (IV/PO)	No carrier state. Treat for 3 days. Single-dose therapy is often effective. Resistant to ampicillin.
<i>Vibrio parahaemolyticus</i>	<ul style="list-style-type: none"> <li>• CSF = NP</li> <li>• Blood = P (from GI source)</li> <li>• Sputum = NP</li> <li>• Urine = NP</li> <li>• Stool = P (diarrhea)</li> <li>• Wound = P</li> </ul>	Doxycycline (IV/PO)	Any quinolone (IV/PO)	Most cases of gastroenteritis caused by <i>V. parahaemolyticus</i> are self-limited and require no treatment.
<i>Vibrio vulnificus</i> , <i>alginolyticus</i>	<ul style="list-style-type: none"> <li>• CSF = NP</li> <li>• Blood = P (from GI/wound source)</li> <li>• Sputum = NP</li> <li>• Urine = NP</li> <li>• Stool = P (diarrhea)</li> <li>• Wound = P (water-contaminated wound raw oysters, other shell fish ingestion)</li> </ul>	Doxycycline (IV/PO) Any quinolone (IV/PO)	Piperacillin (IV) Ampicillin/ sulbactam (IV)	Causes necrotizing soft tissue infection resembling gas gangrene. Patients are critically ill with fever, bullous lesions, diarrhea, and hypotension. Treat wound infection, bacteremia. Aminoglycoside susceptibilities are unpredictable.
<i>Yersinia enterocolitica</i>	<ul style="list-style-type: none"> <li>• CSF = NP</li> <li>• Blood = P (from GI source)</li> </ul>	Any quinolone (IV/PO)	TMP-SMX (IV/PO) Any 3 <sup>rd</sup> generation cephalosporin (IV/PO)	Cause of diarrhea with abdominal pain. If pain in is right lower quadrant, may be mistaken for acute appendicitis.

Yersinia pestis	<ul style="list-style-type: none"> <li>• Sputum = NP</li> <li>• Urine = NP</li> <li>• Stool = P (diarrhea)</li> <li>• Wound = NP</li> <li>• CSF = NP</li> <li>• Blood = P (septicemic plague; isolation required; dangerous)</li> <li>• Sputum = P (pneumonic plague; isolation required; dangerous)</li> <li>• Urine = NP</li> <li>• Stool = NP</li> <li>• Wound = P (lymph nodes, lymph node drainage; bubonic plague; isolation required; dangerous)</li> </ul>	Gentamicin (IV) Doxycycline (IV/PO)	Chloramphenicol (IV/PO)	Cause of bubonic, septicemic, and pneumonic plague. Doxycycline or any quinolone may be used for prophylaxis. Alert microbiology laboratory of potentially biohazardous specimens. Do not culture. Bioterrorist plague is treated the same as naturally-acquired plague.
<b>SPIROCHETES</b>				
Borrelia burgdorferi	<ul style="list-style-type: none"> <li>• CSF = P (neuroborreliosis)</li> <li>• Blood = P (rarely isolated; requires special media)</li> <li>• Sputum = NP</li> <li>• Urine = NP</li> <li>• Stool = NP</li> <li>• Wound = P (rarely isolated from erythema migrans lesions)</li> </ul>	Doxycycline (PO) Amoxicillin (PO)	Any cephalosporin (PO) Azithromycin (PO) Erythromycin (PO)	Cause of Lyme disease. $\beta$ -lactams and doxycycline are effective. Erythromycin least effective for erythema migrans. Minocycline may be preferred to doxycycline for neuroborreliosis.

C = colonizer; C\* = skin contaminant; NP = non-pathogen at site; P = pathogen at site; (IV/PO) = IV or PO. See p. xi for all other abbreviations.



Table 3.6. Usual Clinical Significance of AEROBIC Isolates Pending Susceptibility Testing (cont'd)

SPIROCHETES				
Isolate	Isolate Significance	Preferred Therapy	Alternate Therapy	Comments
Borrelia recurrentis	<ul style="list-style-type: none"> <li>• CSF = P (ABM)</li> <li>• Blood = P (1° bacteremia)</li> <li>• Sputum = NP</li> <li>• Urine = NP</li> <li>• Stool = NP</li> <li>• Wound = NP</li> </ul>	<p>Doxycycline (IV/PO)</p> <p>Azithromycin (IV/PO)</p>	<p>Erythromycin (IV)</p> <p>Penicillin (IV)</p> <p>Ampicillin (IV)</p> <p>Any 1<sup>st</sup>, 2<sup>nd</sup>, 3<sup>rd</sup> generation cephalosporin (IV/PO)</p>	Cause of relapsing fever. May be recovered from septic metastatic foci. Septic emboli may cause sacroiliitis, SBE, myositis, orchitis, or osteomyelitis.
Spirillum minus	<ul style="list-style-type: none"> <li>• CSF = NP</li> <li>• Blood = P (from wound source, SBE)</li> <li>• Sputum = NP</li> <li>• Urine = NP</li> <li>• Stool = NP</li> <li>• Wound = P (from rat bite)</li> </ul>	<p>Penicillin (IV)</p> <p>Amoxicillin (PO)</p>	<p>Doxycycline (IV/PO)</p> <p>Any quinolone (IV/PO)</p>	Cause of rat-bite fever. Bite wound heals promptly, but 1–4 weeks later becomes painful, purple and swollen, and progresses to ulceration and eschar formation. Painful regional adenopathy. Central maculopapular rash is common (rarely urticarial). Arthralgias/arthritis is rare compared to rat-bite fever from Streptobacillus moniliformis. Rarely causes SBE.

C = colonizer; C\* = skin contaminant; NP = non-pathogen at site; P = pathogen at site; (IV/PO) = IV or PO. See p. xi for all other abbreviations.

Table 3.7. Clinical Significance of CAPNOPHILIC Isolates Pending Susceptibility Testing

GRAM-NEGATIVE BACILLI				
Isolate	Isolate Significance	Preferred Therapy	Alternate Therapy	Comments
Capnocytophaga canimorsus/ cynodegmi (DF-2 like)	<ul style="list-style-type: none"> <li>• CSF = NP</li> <li>• Blood = P (from GI source, bite wound)</li> <li>• Sputum = NP</li> <li>• Urine = NP</li> <li>• Stool = NP</li> <li>• Wound = P (from dog/cat bite)</li> </ul>	Ampicillin/ sulbactam (IV) Piperacillin/ tazobactam (IV) Imipenem (IV) Meropenem (IV) Ertapenem (IV)	Clindamycin (IV/PO) Any quinolone (IV/PO) Doxycycline (IV/PO)	Associated with animal bites or cancer. May cause fatal septicemia in cirrhotics/asplenic. Resistant to aminoglycosides, metronidazole, TMP-SMX, and aztreonam.
Capnocytophaga ochraceus (DF-1)	<ul style="list-style-type: none"> <li>• CSF = NP</li> <li>• Blood = P (from GI, wound, abscess source)</li> <li>• Sputum = NP</li> <li>• Urine = NP</li> <li>• Stool = NP</li> <li>• Wound = P</li> </ul>	Ampicillin/ sulbactam (IV) Piperacillin/ tazobactam (IV) Imipenem (IV) Meropenem (IV) Ertapenem (IV)	Clindamycin (IV/PO) Any quinolone (IV/PO) Doxycycline (IV/PO)	Thin, spindle-shaped bacilli resemble Fusobacteria morphologically; "Gliding motility" seen in hanging drop preparations. Cause of septicemia, abscesses, and wound infections. Resistant to aminoglycosides, metronidazole, TMP-SMX, and aztreonam.
Eikenella corrodens	<ul style="list-style-type: none"> <li>• CSF = NP</li> <li>• Blood = P (SBE in IV drug abusers)</li> <li>• Sputum = NP</li> <li>• Urine = NP</li> <li>• Stool = NP</li> <li>• Wound = P (IV drug abusers)</li> </ul>	Penicillin (IV) Ampicillin (IV) Imipenem (IV) Meropenem (IV) Ertapenem (IV)	Piperacillin (IV) Ampicillin/ sulbactam (IV) Doxycycline (IV/PO) Amoxicillin (PO)	Cause of "culture-negative" SBE (one of the HACEK organisms). Resistant to clindamycin and metronidazole.

C = colonizer; C\* = skin contaminant; NP = non-pathogen at site; P = pathogen at site; (IV/PO) = IV or PO. See p. xi for all other abbreviations.

Table 3.8. Usual Clinical Significance of ANAEROBIC Isolates Pending Susceptibility Testing

GRAM-POSITIVE BACILLI				
Isolate	Isolate Significance	Preferred Therapy	Alternate Therapy	
Peptococcus	<ul style="list-style-type: none"> <li>• CSF = P (brain abscess)</li> <li>• Blood = P (from GI/pelvic source)</li> <li>• Sputum = C, P (aspiration pneumonia, lung abscess)</li> <li>• Urine/Stool = NP</li> <li>• Wound = P (rarely a sole pathogen)</li> </ul>	Penicillin (IV) Ampicillin (IV) Amoxicillin (PO) Clindamycin (IV/PO)	Chloramphenicol (IV) Erythromycin (IV) Any carbapenem (IV) Moxifloxacin (IV/PO)	Normal flora of mouth, GI tract, and pelvis. Associated with mixed aerobic/anaerobic dental, abdominal, and pelvic infections, especially abscesses.
Peptostreptococcus	<ul style="list-style-type: none"> <li>• CSF = P (brain abscess)</li> <li>• Blood = P (GI/pelvic source)</li> <li>• Sputum = C, P (aspiration pneumonia, lung abscess)</li> <li>• Urine/Stool = NP</li> <li>• Wound = P (rarely a sole pathogen)</li> </ul>	Penicillin (IV) Ampicillin (IV) Amoxicillin (PO) Clindamycin (IV/PO)	Chloramphenicol (IV) Erythromycin (IV) Any carbapenem (IV) Moxifloxacin (IV/PO)	Normal flora of mouth, GI tract, and pelvis. Associated with mixed aerobic/anaerobic dental, abdominal, and pelvic infections, especially abscesses.
GRAM-POSITIVE BACILLI				
Actinomyces israelii, odontolyticus	<ul style="list-style-type: none"> <li>• CSF = P (brain abscess)</li> <li>• Blood = NP</li> </ul>	Amoxicillin (PO) Doxycycline (PO)	Erythromycin (PO) Clindamycin (PO)	Anaerobic and non-acid fast. Usually presents as cervical, facial, thoracic, or abdominal masses/fistulas. Prolonged

<p><i>Arachnia propionica</i></p>	<ul style="list-style-type: none"> <li>• Sputum = C, P (lung abscess)</li> <li>• Urine = NP</li> <li>• Stool = NP</li> <li>• Wound = P (fistulas/underlying abscess)</li> </ul>	<p>Clindamycin (IV/PO) Ampicillin (IV) + gentamicin (IV)</p>	<p>Erythromycin (IV)</p>	<p>(6–12 month) treatment is needed for cure. Unlike <i>Nocardia</i>, <i>Actinomyces</i> rarely causes CNS infections. May be cultured from polymicrobial brain abscess of pulmonary origin. Quinolones, aminoglycosides, metronidazole, and TMP-SMX have little activity.</p> <p>Polymicrobial pathogen in dental, lung, and brain abscesses.</p>
<p><i>Bifidobacterium</i> sp.</p>	<ul style="list-style-type: none"> <li>• CSF = P (brain abscess)</li> <li>• Blood = NP</li> <li>• Sputum = C, P (lung abscess)</li> <li>• Urine = NP</li> <li>• Stool = NP</li> <li>• Wound = NP</li> </ul>	<p>Clindamycin (IV/PO) Ampicillin (IV) + gentamicin (IV)</p>	<p>Erythromycin (IV)</p>	<p>Usually part of polymicrobial infection.</p>
<p><i>Clostridium botulinum</i></p>	<ul style="list-style-type: none"> <li>• CSF = NP</li> <li>• Blood = NP</li> <li>• Sputum = NP</li> <li>• Urine/Stool = NP</li> <li>• Wound = P (wound botulism)</li> </ul>	<p>Penicillin (IV)</p>	<p>Clindamycin (IV/PO) Imipenem (IV) Meropenem (IV)</p>	<p>Give trivalent equine antitoxin (p. 170) as soon as possible. Antibiotic therapy is adjunctive.</p>

C = colonizer; C\* = skin contaminant; NP = non-pathogen at site; P = pathogen at site; (IV/PO) = IV or PO. See p. xi for all other abbreviations.

Table 3.8. Usual Clinical Significance of ANAEROBIC Isolates Pending Susceptibility Testing (Cont'd)

GRAM-POSITIVE BACILLI				
Isolate	Isolate Significance	Preferred Therapy	Alternate Therapy	Comments
<i>Clostridium difficile</i>	<ul style="list-style-type: none"> <li>• CSF = NP</li> <li>• Blood = P (rarely from GI source)</li> <li>• Sputum = NP</li> <li>• Urine = NP</li> <li>• Stool = C (normal fecal flora), P (antibiotic-associated diarrhea/colitis)</li> <li>• Wound = NP</li> </ul>	<p><i>C. difficile</i> diarrhea</p> <p>Vancomycin (PO)</p> <p>Nitazoxanide (PO)</p> <p><i>C. difficile</i> colitis</p> <p>Metronidazole (IV/PO)</p> <p>Nitazoxanide (PO)</p>	<p><i>C. difficile</i> diarrhea</p> <p>Metronidazole (PO)</p> <p><i>C. difficile</i> colitis</p> <p>Tigacycline (IV)</p>	<p><b>C. difficile diarrhea</b> PO vancomycin preferred. <i>PO vancomycin more reliably effective than PO metronidazole.</i></p> <p>Nitazoxanide also highly effective. PO metronidazole, not PO vancomycin, increases prevalence of VRE. <b>C. difficile colitis</b>, use IV or PO metronidazole (IV vancomycin ineffective). Nitazoxanide (PO) or tigacycline (IV) also highly effective. Diagnose <i>C. difficile</i> diarrhea by stool <i>C. difficile</i> toxin assay/PCR. Diagnose <i>C. difficile</i> colitis by <i>C. difficile</i> + toxin assay/PCR plus colitis on abdominal CT scan/colonoscopy.</p>
<i>Clostridium perfringens</i> , <i>septicum</i> , <i>novyi</i>	<ul style="list-style-type: none"> <li>• CSF = NP</li> <li>• Blood = P (from GI source/malignancy)</li> <li>• Sputum = NP</li> <li>• Urine = NP</li> <li>• Stool = NP</li> <li>• Wound = P (gas gangrene)</li> </ul>	<p>Penicillin (IV)</p> <p>Piperacillin/tazobactam (IV)</p> <p>Meropenem (IV)</p> <p>Ertapenem (IV)</p>	<p>Clindamycin (IV)</p> <p>Chloramphenicol (IV)</p> <p>Imipenem (IV)</p>	<p>Usual cause of myonecrosis (gas gangrene). Surgical debridement is crucial; antibiotic therapy is adjunctive. Also causes emphysematous cholecystitis/cystitis. Does not form spores in blood cultures as does <i>C. sordelli</i>.</p>
<i>Clostridium tetani</i>	<ul style="list-style-type: none"> <li>• CSF = NP</li> <li>• Blood = NP</li> </ul>	<p>Penicillin (IV)</p> <p>Clindamycin (IV)</p>	<p>Imipenem (IV)</p> <p>Meropenem (IV)</p>	<p>Prompt administration of tetanus immune globulin is crucial (p. 170). Antibiotic therapy is adjunctive.</p>

Eubacterium sp.	<ul style="list-style-type: none"> <li>• Sputum = NP</li> <li>• Urine/Stool = NP</li> <li>• Wound = P (wound tetanus)</li> <li>• CSF = P (brain abscess)</li> <li>• Blood = P (from dental, GI, GU, lung source)</li> <li>• Sputum = P (lung abscess)</li> <li>• Urine/Stool = NP</li> <li>• Wound = NP</li> </ul>	Clindamycin (IV/PO) Ampicillin (IV) + gentamicin (IV)	Erythromycin (IV)	Pathogen in lung/pelvic/brain abscesses, and chronic periodontal disease. Eubacterium bacteremias are associated with malignancies.
Lactobacillus sp.	<ul style="list-style-type: none"> <li>• CSF = P (ABM)</li> <li>• Blood = P (1° bacteremia, SBE, or from endometritis)</li> <li>• Sputum = NP</li> <li>• Urine = P (rare)</li> <li>• Stool = NP</li> <li>• Wound = NP</li> </ul>	Ampicillin (IV) + gentamicin (IV) Clindamycin (IV/PO)	Erythromycin (IV)	Uncommon pathogen in normal/compromised hosts. Rare cause of SBE. Variably resistant to cephalosporins and quinolones. Some clindamycin-resistant strains. Resistant to metronidazole and vancomycin.
Propionibacterium acnes	<ul style="list-style-type: none"> <li>• CSF = C*, P (meningitis from NS shunts)</li> <li>• Blood = C*, P (from IV line infection, SBE)</li> <li>• Sputum = NP</li> <li>• Urine = NP</li> <li>• Stool = NP</li> <li>• Wound = C</li> </ul>	Penicillin (IV) Clindamycin (IV/PO)	Doxycycline (IV/PO)	Common skin colonizer/blood culture contaminant. Rarely causes prosthetic joint infection, endocarditis, or CNS shunt infection. Resistant to metronidazole.

C = colonizer; C\* = skin contaminant; NP = non-pathogen at site; P = pathogen at site; (IV/PO) = IV or PO. See p. xi for all other abbreviations.

Table 3.8. Usual Clinical Significance of ANAEROBIC Isolates Pending Susceptibility Testing (cont'd)

GRAM-NEGATIVE BACILLI				
Isolate	Isolate Significance	Preferred Therapy	Alternate Therapy	Comments
Bacteroides fragilis group (B. distasonis, ovatus, thetaiotaomicron, vulgatus)	<ul style="list-style-type: none"> <li>• CSF = P (meningitis from Strongyloides hyperinfection)</li> <li>• Blood = P (from GI/ pelvic source)</li> <li>• Sputum = NP</li> <li>• Urine = NP, P (only from colonic fistula)</li> <li>• Stool = NP</li> <li>• Wound = NP</li> </ul>	Tigecycline (IV) Piperacillin/ tazobactam (IV) Any carbapenem (IV)	Moxifloxacin (IV/PO) Ampicillin/sulbactam (IV) Clindamycin (IV/PO) <i>or combination</i> <i>of Metronidazole</i> (IV/PO) <i>plus either</i> <i>ceftriaxone (IV) or</i> <i>levofloxacin (IV/PO)</i>	Major anaerobe below the diaphragm. Usually part of polymicrobial lower intra-abdominal and pelvic infections. Cefotetan is less effective against B. fragilis DOT strains (B. distasonis, B. ovatus, B. thetaiotaomicron). Resistant to penicillin.
Fusobacterium nucleatum	<ul style="list-style-type: none"> <li>• CSF = P (brain abscess)</li> <li>• Blood = P (from lung, GI source)</li> <li>• Sputum = P (aspiration pneumonia, lung abscess)</li> <li>• Urine = NP</li> <li>• Stool = NP</li> <li>• Wound = P (rarely)</li> </ul>	Clindamycin (IV/PO) Piperacillin/ tazobactam (IV) Ampicillin/ sulbactam (IV)	Chloramphenicol (IV) Metronidazole (IV/PO)	Mouth flora associated with dental infections and anaerobic lung infections. F. nucleatum is associated with jugular vein septic phlebitis and GI cancer. Resembles Capnophagia sp. on sputum gram stain.
Prevotella (Bacteroides) bivia	<ul style="list-style-type: none"> <li>• CSF = NP</li> <li>• Blood = P (from dental, lung, pelvic source)</li> <li>• Sputum = P (lung abscess)</li> </ul>	Penicillin (IV/PO) Any $\beta$ -lactam (IV/PO)	Any quinolone (IV/PO) Doxycycline (IV/PO) Clindamycin (IV/PO)	Cause of dental, oropharyngeal, and female genital tract infections.

<ul style="list-style-type: none"> <li>• Urine = NP</li> <li>• Stool = NP</li> <li>• Wound = NP</li> </ul>	<ul style="list-style-type: none"> <li>• CSF = P (brain abscess)</li> <li>• Blood = P (from oral/pulmonary source)</li> <li>• Sputum = P (from aspiration pneumonia, lung abscess)</li> <li>• Urine = NP</li> <li>• Stool = NP</li> <li>• Wound = NP</li> </ul>	<p>Aspiration pneumonia/lung abscess</p> <p>Any <math>\beta</math>-lactam (IV/PO)</p> <p>Any quinolone (IV/PO)</p> <p>Brain abscess</p> <p>Penicillin (IV)</p>	<p>Aspiration pneumonia/lung abscess</p> <p>Doxycycline (IV/PO)</p> <p>Brain abscess</p> <p>Chloramphenicol (IV)</p>	<p>Predominant anaerobic flora of mouth. Known as "oral pigmented" Bacteroides (e.g., B. melanogenicus). Antibiotics used to treat community-acquired pneumonia are effective against oral anaerobes (e.g., Prevotella) in aspiration pneumonia; does not require anti-B. fragilis coverage with clindamycin, metronidazole, or moxifloxacin.</p>
<p>Prevotella (Bacteroides) melaninogenicus, intermedium</p>				

C = colonizer; C\* = skin contaminant; NP = non-pathogen at site; P = pathogen at site; (IV/PO) = IV or PO. See p. xi for all other abbreviations.



Table 3.9. Clinical Significance of YEAST/FUNGI Pending Susceptibility Testing

YEAST/FUNGI				
Isolate	**Usual Isolate Significance	Preferred Therapy	Alternate Therapy	Comments
Aspergillus species	<ul style="list-style-type: none"> <li>CSF = P (only from disseminated infection)</li> <li>Blood = C, P (1° fungemia or from pulmonary source)</li> <li>Sputum = C, P (pneumonia)</li> <li>Urine = NP</li> <li>Stool = NP</li> <li>Wound = NP, P (rarely, but possible with extensive wounds, e.g., burns)</li> </ul>	<p>See p. 55:</p> <p>Voriconazole (IV/PO)</p> <p>Amphotericin B lipid formulation (IV)</p>	<p>See p. 55:</p> <p>Posaconazole (PO);</p> <p>Amphotericin B deoxycholate (IV)</p>	<p>A. fumigatus is the usual cause of invasive aspergillosis. Growth of Aspergillus sp. from a specimen can represent airborne contamination. Aspergillus pneumonia and disseminated aspergillosis are not uncommon in patients on chronic steroids or immunosuppressives (esp. organ transplants). Recovery of Aspergillus from sputum or BAL is not diagnostic of Aspergillus pneumonia. Definitive Dx is by lung biopsy demonstrating vessel/tissue invasion. <math>\beta</math> 1,3 D-glucan (BG)+, aspergillus galactomannan (AG)+.</p>
Candida albicans	<ul style="list-style-type: none"> <li>CSF = P (only from disseminated infection)</li> <li>Blood = P (1° candidemia or from IV line infection)</li> <li>Sputum = C, P (only from disseminated infection)</li> </ul>	<p>Fluconazole (IV/PO)</p> <p>Micafungin (IV)</p> <p>Caspofungin (IV)</p> <p>Anidulafungin (IV)</p> <p>Posaconazole (PO)</p>	<p>Amphotericin B deoxycholate (IV/PO)</p> <p>Amphotericin B lipid formulation (IV)</p> <p>Itraconazole (IV/PO)</p> <p>Voriconazole (IV/PO)</p>	<p>Common colonizer of GI/GU tracts. Colonization common in diabetics, alcoholics, patients receiving steroids/antibiotics. Commonest cause of fungemia in hospitalized patients. Candidemia secondary to central IV lines should always be treated as possible disseminated disease even though this is not invariably the case. Repeated blood cultures and careful follow-up (including</p>

	<ul style="list-style-type: none"> <li>• Urine = C, P (from cystitis, pyelonephritis)</li> <li>• Stool = C (source of candiduria)</li> <li>• Wound = NP</li> </ul>			ophthalmoscopy) should be undertaken to exclude possible occult dissemination following even a single positive blood culture. Primary candidal pneumonia is rare.
Candida non-albicans	<ul style="list-style-type: none"> <li>• CSF = P (only from disseminated infection)</li> <li>• Blood = P (1° candidemia or from IV line infection)</li> <li>• Sputum = NP</li> <li>• Urine = C (indwelling catheters), P (from cystitis, pyelonephritis)</li> <li>• Stool = C (source of candiduria)</li> <li>• Wound = NP</li> </ul>	<p>Micafungin (IV)          Caspofungin (IV)          Anidulafungin (IV)          Posaconazole (PO)          Voriconazole (IV/PO)</p>	<p>Fluconazole (IV/PO)          Amphotericin B deoxycholate (IV/PO)          Amphotericin B lipid formulation (IV)          Itraconazole (IV/PO)</p>	<p>Non-albicans Candida cause the same spectrum of invasive disease as C. albicans. Fluconazole-susceptibility varies predictably by species. C. glabrata (usually) and C. krusei (almost always) are resistant to fluconazole. C. lusitanae is often resistant to amphotericin B (deoxycholate and lipid-associated formulations). Other species are generally susceptible to all agents.</p>
Cryptococcus neoformans	<ul style="list-style-type: none"> <li>• CSF = P (meningitis, brain abscess)</li> <li>• Blood = P (from pulmonary source)</li> <li>• Sputum = P (pneumonia)</li> <li>• Urine = NP</li> <li>• Stool = NP</li> <li>• Wound = NP*</li> </ul>	<p>CNS          Amphotericin B deoxycholate (IV) ± flucytosine (PO)          Non-CNS          Amphotericin B deoxycholate (IV)</p>	<p>CNS          Fluconazole (IV/PO)          Non-CNS          Itraconazole (IV/PO)          Fluconazole (IV/PO)</p>	<p>C. neoformans meningitis may occur with or without dissemination. Cryptococcal pneumonia frequently disseminates to CNS. C. neoformans in blood cultures occurs in compromised hosts (eg, HIV) and indicates disseminated infection.</p>

C = colonizer; C\* = skin contaminant; NP = non-pathogen at site; P = pathogen at site; (IV/PO) = IV or PO. See p. xi for all other abbreviations.  
 \* Cutaneous cryptococcus represents disseminated infection. \*\* Fungi can produce disseminated infections that involve essentially any organ. Isolation of a fungus from any normal sterile site should be cause for a careful review of the patient's epidemiology, risk factors, and clinical presentation.

Table 3.9. Usual Clinical Significance of YEAST/FUNGI Pending Susceptibility Testing (cont'd)

YEAST/FUNGI				
Isolate	**Usual Isolate Significance	Preferred Therapy	Alternate Therapy	Comments
		Amphotericin B lipid formulation (IV)		
Histoplasma capsulatum	<ul style="list-style-type: none"> <li>• CSF = P (from disseminated infection, pneumonia)</li> <li>• Blood = P (1° fungemia, rarely SBE)</li> <li>• Sputum = P (pneumonia, mediastinitis)</li> <li>• Urine = P</li> <li>• Stool = P</li> <li>• Wound = P</li> </ul>	Itraconazole (IV/PO) Amphotericin B deoxycholate (IV)	Fluconazole (IV/PO) Amphotericin B lipid formulation (IV)	Histoplasma recovered from CSF/blood cultures indicates dissemination. Disseminated (reactivated latent) histoplasmosis is most common in compromised hosts (e.g., HIV). Itraconazole is ineffective for meningeal histoplasmosis, but is preferred for chronic suppressive therapy.
Malassezia furfur	<ul style="list-style-type: none"> <li>• CSF = NP</li> <li>• Blood = P (from IV line infection)</li> <li>• Sputum = NP</li> <li>• Urine = NP</li> <li>• Stool = NP</li> <li>• Wound = P (eosinophilic folliculitis)</li> </ul>	Itraconazole (IV/PO) ketoconazole (PO)	Fluconazole (IV/PO)	M. furfur IV line infections are associated with IV lipid hyperalimentation emulsions. Fungemia usually resolves with IV line removal. Morphology in blood is blunt buds on a broad base yeast. M. furfur requires long chain fatty acids for growth (overlay agar with thin layer of olive oil, Tween 80, or oleic acid).

<p>Penicillium marnieffei</p>	<ul style="list-style-type: none"> <li>• CSF = NP</li> <li>• Blood = P (usually from dissemination)</li> <li>• Sputum = P (pneumonia)</li> <li>• Urine = NP</li> <li>• Stool = NP</li> <li>• Wound = NP*</li> </ul>	<p>Amphotericin B deoxycholate (IV) Itraconazole (IV/PO)</p>	<p>Amphotericin B lipid formulation (IV)</p>	<p>Histoplasma-like yeast forms seen in lymph nodes, liver, skin, bone marrow, blood. Characteristic red pigment diffuses into agar. Closely resembles histoplasmosis yeast forms (<i>H. capsulatum</i> has narrow based budding yeast forms, but <i>P. marnieffei</i> has transverse septa). Skin lesions indicate dissemination. May resemble molluscum contagiosum. Hepatosplenomegaly is common.</p>
-------------------------------	---	--	--	--

C = colonizer; C\* = skin contaminant; NP = non-pathogen at site; P = pathogen at site; (IV/PO) = IV or PO. See p. xi for all other abbreviations.

\* Cutaneous lesions represents disseminated infection.

\*\* Fungi can produce disseminated infections that involve essentially any organ. Isolation of a fungus from any normal sterile site should be cause for a careful review of the patient's epidemiology, risk factors, and clinical presentation.

**Table 3.10. Technique for Gram Stain and Giemsa Stain**

GRAM STAIN
<p><b>Clinical applications:</b> CSF, sputum, urine</p> <p><b>Technique:</b></p> <ol style="list-style-type: none"> <li>1. Place specimen on slide.</li> <li>2. Heat fix smear on slide by passing it quickly over a flame.</li> <li>3. Place crystal violet solution on slide for 20 seconds.</li> <li>4. Wash gently with water.</li> <li>5. Apply Gram iodine solution to slide for 20 seconds.</li> <li>6. Decolorize the slide quickly in solution of acetone/ethanol.</li> <li>7. Wash slide gently with water.</li> <li>8. Counterstain slide with safranin for 10 seconds.</li> <li>9. Wash gently with water; air dry or blot dry with bibulous paper.</li> </ol> <p><b>Interpretation:</b> Gram-negative organisms stain red; gram-positive organisms stain blue. <i>B. fragilis</i> stains weakly pink. Fungi stain deep blue. For interpretation of Gram stain findings in CSF, urine, sputum, and feces see Tables 3.6–3.9.</p>
GIEMSA STAIN
<p><b>Clinical applications:</b> Blood buffy coat, bone marrow</p> <p><b>Technique:</b></p> <ol style="list-style-type: none"> <li>1. Place specimen on slide.</li> <li>2. Fix smear by placing slide in 100% methanol for 1 minute.</li> <li>3. Drain methanol off slide.</li> <li>4. Flood slide with Giemsa stain (freshly diluted 1:10 with distilled water) for 5 minutes.</li> <li>5. Wash slide gently with water; air dry.</li> </ol> <p><b>Interpretation:</b> Fungi/parasites stain light/dark blue.</p>

**Table 3.11. Clinical Use of CSF Gram Stain, WBC Type, Glucose (see Color Atlas of CSF Gram stains)**

Gram Stain	Organism/Condition
Gram-positive bacilli	Pseudomoniasis (Bacillus, Listeria Corynebacteria)
Gram-negative bacilli	<i>H. influenzae</i> (small, encapsulated, Non-enteric/enteric aerobic pleomorphic) bacilli (larger, unencapsulated)
Gram-positive cocci	Gp A, B, D, streptococci (pairs/chains) <i>S. aureus</i> (pairs/clusters) <i>S. pneumoniae</i> (pairs) <i>S. epidermidis</i> (pairs/clusters)
Gram-negative diplococci	<i>Neisseria meningitidis</i>

**Table 3.11. Clinical Use of CSF Gram Stain, WBC Type, Glucose (cont'd)**

<b>Gram Stain</b>	<b>Organism/Condition</b>	
Mixed organisms (polymicrobial)	Pseudomonas meningitis Anaerobic organisms (brain abscess with meningeal leak)	Neonatal meningitis Meningitis 2° to penetrating head trauma
<b>WBC Type/Glucose</b>	<b>Organism/Condition</b>	
Purulent CSF, no organisms	Neisseria meningitidis	Listeria
Clear CSF, no organisms	Viral meningitis Viral encephalitis TB/fungal meningitis Sarcoidosis meningitis Meningeal carcinomatosis Brain abscess Parameningeal infection Septic emboli 2° to SBE SLE cerebritis Lyme's disease	Lymphocytic choriomeningitis (LCM) Drug induced aseptic meningitis Listeria HIV Syphilis Leptospirosis Bacterial meningitis (very early/partially-treated) Meningitis (leukopenic host) Rocky Mountain Spotted Fever
Cloudy CSF, no WBCs	S. pneumoniae	
Predominantly PMNs, decreased glucose	Bacterial meningitis (partially-treated) Listeria HSV-1/2 encephalitis TB (early/beginning therapy) Sarcoidosis	Parameningeal infection Septic emboli 2° to SBE Amebic meningoencephalitis Syphilis (early) Posterior-fossa syndrome
Predominantly lymphocytes, normal glucose	Bacterial meningitis (partially-treated) Sarcoidosis Lyme's disease HIV Leptospirosis Rocky Mountain Spotted Fever	Viral meningitis Viral encephalitis Parameningeal infection TB/fungal meningitis Parasitic meningitis Meningeal carcinomatosis
Predominantly lymphocytes, decreased glucose	Bacterial meningitis (partially-treated) TB/fungal meningitis Sarcoidosis Lymphocytic choriomeningitis (LCM) Mumps	Enteroviral meningitis Listeria Leptospirosis Syphilis Meningeal carcinomatosis
Red blood cells	Traumatic tap CNS bleed/tumor Listeria Leptospirosis Herpes (HSV-1) encephalitis	TB meningitis Amebic (Naegleria) meningoencephalitis Anthrax

**Table 3.12. Clinical Use of the Sputum Gram Stain (see Color Atlas of Sputum Gram stains)**

Gram Stain	Organism	Comments
Gram-positive diplococci	<i>S. pneumoniae</i>	Lancet-shaped encapsulated diplococci (not streptococci)
Gram-positive cocci (grape-like clusters)	<i>S. aureus</i>	Clusters predominant. Short chains or pairs may also be present
Gram-positive cocci (short chains or pairs)	Group A streptococci	Virulence inversely proportional to length of streptococci. Clusters not present
Gram-positive beaded/filamentous branching organisms	<i>Nocardia</i>	Coccobacillary forms common
Gram-negative coccobacillary organisms	<i>H. influenzae</i>	Pleomorphic may be encapsulated. Gram negative cocci/bacilli. Lightly stained
Gram-negative bacilli	<i>Klebsiella</i> <i>P. aeruginosa</i>	Plump and encapsulated Thin and often arranged in end-to-end pairs
Gram-negative diplococci	<i>Moraxella</i> (Branhamella) <i>catarrhalis</i> <i>Neisseria meningitidis</i>	Kidney bean-shaped diplococci

**Table 3.13. Clinical Use of the Urine Gram Stain (see Color Atlas of Urine Gram stains)**

Gram Stain	Organism	Comments
Gram-positive cocci (clusters)*	<i>S. aureus</i> <i>S. epidermidis</i> <i>S. saprophyticus</i>	Skin flora contaminant Skin flora contaminant Uropathogen
Gram-positive cocci (chains)	Group B streptococci Group D streptococci <i>E. faecalis</i> <i>E. faecium</i>	Uropathogen Uropathogen May represent colonization or infection May represent colonization or infection
Gram-negative bacilli*	Coliform bacilli	Uropathogen; may represent colonization or infection
Gram-negative diplococci*	<i>N. gonorrhoeae</i> <i>N. meningitidis</i>	Gonococcal urethritis Rare cause of urethritis

\* *Staphylococci* (except *S. saprophyticus*), *S. pneumoniae*, and *B. fragilis* are *not* uropathogens.

**Table 3.14. Clinical Use of the Fecal Gram Stain**

Gram Stain	Possible Organisms	
Fecal leukocytes present	Enteropathogenic E. coli (EPEC) Enteroinvasive E. coli (EIEC) Shigella Yersinia Campylobacter Salmonella Vibrio parahaemolyticus Vibrio vulnificus	Aeromonas hydrophila Chlamydia trachomatis Plesiomonas shigelloides Neisseria gonorrhoeae (proctitis) Herpes simplex virus (HSV-1) <i>Noninfectious:</i> Ulcerative colitis
No fecal leukocytes	Enterovirus Rotavirus Coronavirus Enterotoxigenic E. coli (ETEC) S. aureus Clostridium perfringens Adenovirus	Norwalk virus Vibrio cholerae Bacillus cereus (food poisoning) Giardia lamblia Isospora belli Cryptosporidia Strongyloides stercoralis
Fecal leukocytes variable	Clostridium difficile Enterohemorrhagic E. coli (EHEC)	Cytomegalovirus (CMV) Herpes simplex virus (HSV-1)
Red blood cells present	Shigella Salmonella Campylobacter EPEC EHEC Enteroinvasive E. coli (EIEC) Enterohemorrhagic E. coli (EHEC)	Clostridium difficile Cytomegalovirus (CMV) Yersinia Plesiomonas shigelloides <i>Noninfectious:</i> Ulcerative colitis

## REFERENCES AND SUGGESTED READINGS

- Aridan S, Paetznick V, Rex JH. Comparative evaluation of disk diffusion with microdilution assay in susceptibility of caspofungin against *Aspergillus* and *Fusarium* isolates. *Antimicrob Agents Chemother* 46:3084–7, 2002.
- Baldis MM, Keidich SD, Mukhejee PK, et al. Mechanisms of fungal resistance: an overview. *Drugs* 62:1025–40, 2002.
- Bartlett JG. Antibiotic-associated diarrhea. *N Engl J Med* 346:334–9, 2002.
- Bingen E, Leclercq R, Fitoussi F, et al. Emergence of group A streptococcus strains with different mechanisms of macrolide resistance. *Antimicrob Agents Chemother* 46:1199–203, 2002.
- Bouza E, Cercenado E. Klebsiella and enterobacter: antibiotic resistance and treatment implications. *Semin Respir Infect* 17:215–30, 2002.
- Boyce JM. Methicillin-resistant *Staphylococcus aureus*. *Lancet Infect Dis* 5:653–63, 2005.



- Brueggemann AB, Coffmen SL, Rhomberg P, et al. Fluoroquinolone resistance in *Streptococcus pneumoniae* in United States. *Antimicrob Agents Chemother* 46:680–8, 2002.
- Courvalin P. Vancomycin resistance in gram-positive cocci. *Clin Infect Dis* 42:525–534, 2006.
- Cunha BA. Clinical relevance of penicillin-resistant *Streptococcus pneumoniae*. *Semin Respir Infect* 17:204–14, 2002.
- Cunha BA. *Pseudomonas aeruginosa*: resistance and therapy. *Semin Respir Infect* 17:231–9, 2002.
- Cunha BA. MRSA & VRE: In vitro susceptibility versus in vivo efficacy. *Antibiotics for Clinicians* 4:31–32, 2000.
- Cunha BA. The significance of antibiotic false sensitivity testing with in vitro testing. *J Chemother* 9:25–35, 1997.
- Daneman N, McGeer, Green K, Low DE. Macrolide Resistance in Bacteremic Pneumococcal Disease: Implications for Patient Management. *Clin Infect Dis* 43:432–8, 2006.
- Doern CD. When does 2 plus 2 equal 5? Review of Antimicrobial Synergy testing. *J Clin Microb* 52:4124–28, 2014.
- Doern GV. Macrolide and Ketolide Resistance with *Streptococcus pneumoniae*. *Med Clin N Am* 90:1109–24, 2006.
- Espinell-Ingroff A, Chaturvedi V, Fothergill A, et al. Optimal testing conditions for determining MICs and minimum fungicidal concentrations of new and established antifungal agents for uncommon molds: NCCLS collaborative study. *J Clin Microbiol* 40:3776–81, 2002.
- Falagas ME, Kasiakou SK. Colistin: the revival of polymyxins for the management of multidrug-resistant gram-negative bacterial infections. *Clin Infect Dis* 40:1333–41, 2005.
- Fuchs PC, Barry AL, Brown SD. In vitro activity of telithromycin against *Streptococcus pneumoniae* resistant to other antibiotics, including cefotaxime. *J Antimicrob Chemother* 49:399–401, 2002.
- Gould IM. The clinical significance of methicillin-resistant *Staphylococcus aureus*. *J Hosp Infect* 61:277–282, 2005.
- Halstead DC, Abid J, Dowzicky MJ. Antimicrobial susceptibility among *Acinetobacter calcoaceticus-baumannii* complex and enterobacteriaceae collected as part of the tigecycline evaluation and surveillance trial. *Journal of Infection* 55:49–57, 2007.
- Holmes A, Ganner M, McGuane S, et al. *Staphylococcus aureus* isolates carrying Panton-Valentine leucocidin genes in England and Wales: frequency, characterization, and association with clinical disease. *J Clin Microbiol* 43:2384–90, 2005.
- Jain R, Danziger LH. Multidrug-resistant *Acinetobacter* infections: an emerging challenge to clinicians. *Ann Pharmacother* 38:1449–59, 2004.
- Jevitt LA, Thorne GM, Traczewski MM, et al. Multicenter evaluation of the Etest and disk diffusion methods for differentiating Daptomycin-susceptible from non-Daptomycin-susceptible *Staphylococcus aureus* isolates. *J Clin Microbiol* 44:3098–3104, 2006.
- Jiang X, Zhang Z, Li M, et al. Detection of Extended-Spectrum B-Lactamases in Clinical Isolates of *Pseudomonas aeruginosa*. *Antimicrobial Agents and Chemotherapy* 50:2990–95, 2006.
- Kenneth S, Thomson. Lowering of Third Generation Cephalosporin Breakpoints. *Clin Infect Dis* 57:1663–64, 2013.
- King A, Bathgate T, Phillips I. Erythromycin susceptibility of viridans streptococci from the normal throat flora of patients treated with azithromycin or clarithromycin. *Clin Microbiol Infect* 8:85–92, 2002.
- Kocazeybek B, Arabaci U, Erenturk S. Investigation of various antibiotic combinations using the E-Test method in multiresistant *Pseudomonas aeruginosa* strains. *Chemotherapy* 48:31–5, 2002.
- Kontoyiannis DP, Lewis RE. Antifungal drug resistance of pathogenic fungi. *Lancet* 359:1135–44, 2002.
- Kralovic SM, Danko LH, Roselle GA. Laboratory reporting of *Staphylococcus aureus* with reduced susceptibility to vancomycin in United States Department of Veterans Affairs facilities. *Emerg Infect Dis* 8:402–7, 2002.
- Kremery V, Barnes AJ. Non-albicans *Candida* spp. causing fungaemia: pathogenicity and antifungal resistance. *J Hosp Infect* 50:243–60, 2002.
- Laverdiere M, Hoban D, Restieri C, et al. In vitro activity of three new triazoles and one echinocandin against *Candida* bloodstream isolates from cancer patients. *J Antimicrob Chemother* 50:119–23, 2002.
- Linden PK. Treatment options for vancomycin-resistant enterococcal infections. *Drugs* 62:425–41, 2002.
- Linder JA, Stafford RS. Erythromycin-resistant group A streptococci. *N Engl J Med* 347:614–5, 2002.
- Lonks JR, Garau J, Gomez L, et al. Failure of macrolide antibiotic treatment in patients with bacteremia due to erythromycin-resistant *Streptococcus pneumoniae*. *Clin Infect Dis* 35:556–64, 2002.
- Marchaim D, Pogue JM, Tzuman O, et al. Major Variation in MICs of Tigecycline in Gram-Negative Bacilli as a Function of Testing Method. *J Clin Micro* 52:1617–1621, 2014.
- Metlay JP, Fishman NO, Joffe MM, et al. Macrolide Resistance in Adults with Bacteremic Pneumococcal Pneumonia. *Emerging Infect Dis* 12, 1223–30, 2006.
- Midolo PD, Matthews D, Fernandez CD, et al. Detection of extended spectrum beta-lactamases in the routine clinical microbiology laboratory. *Pathology* 34:362–4, 2002.

- Møllering RC Jr. Problems with antimicrobial resistance in gram-positive cocci. *Clin Infect Dis* 26:1177–8, 1998.
- Naas T, Fortneau N, Spicq C, et al. Three-year survey of community-acquired methicillin-resistant *Staphylococcus aureus* producing Panton-Valentine leukocidin in a French university hospital. *J Hosp Infect* 61:321–9, 2005.
- Paterson DL, Bonomo RA. Extended-spectrum  $\beta$ -lactamases: a clinical update. *Clin Microbiol Rev* 18:657–86, 2005.
- Pelaez T, Alcalá L, Alonso R, et al. Reassessment of *Clostridium difficile* susceptibility to metronidazole and vancomycin. *Antimicrob Agents Chemother* 46:1647–50, 2002.
- Perri MB, Hershberger E, Ionescu M, et al. In vitro susceptibility of vancomycin-resistant enterococci (VRE) to fosfomycin. *Diagn Microbiol Infect Dis* 42:269–71, 2002.
- Poole K. Aminoglycoside resistance in *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother* 49:479–87, 2005.
- Rex JH, Pfaller MA, Walsh TJ, et al. Antifungal susceptibility testing: practical aspects and current challenges. *Clin Microbiol Rev* 14:643–658, 2002.
- Rex JH, Rinaldi MG, Pfaller MA. Resistance of *Candida* species to fluconazole. *Antimicrob Agents Chemother* 39:1–8, 1995.
- Rice LB. Rapid Diagnostics and Appropriate Antibiotic Use. *Clin Infect Dis* 52:S357–S360, 2011.
- Robinson DA, Sutcliffe JA, Tweedros W, et al. Evolution and global dissemination of macrolide-resistant group A streptococci. *Antimicrobial Agents and Chemotherapy* 50:2903–11, 2006.
- Ruhe JJ, Monson T, Bradsher RW, et al. Use of long-acting tetracyclines for methicillin-resistant *Staphylococcus aureus* infections: case series and review of the literature. *Clin Infect Dis* 40:142–934, 2005.
- Stratton CW. In vitro susceptibility testing versus in vivo effectiveness. *Med Clin N Am* 90:1077–88, 2006.
- Suzuki S, Yamazaki T, Narita M, et al. Clinical evaluation of macrolide-resistant *Mycoplasma pneumoniae*. *Antimicrobial Agents and Chemotherapy* 50:709–12, 2006.
- Swenson JM, Killgore GE, Tenover FC. Antimicrobial susceptibility testing of *Acinetobacter* spp. by NCCLS broth microdilution and disk diffusion methods. *J Clin Microbiol* 42:5102–8, 2004.
- Syndman DR, Jacobut NV, McDermott LA, et al. National survey on the susceptibility of *Bacteroides fragilis*: report and analysis of trends for 1997–2000. *Clin Infect Dis* 35(Suppl 1):S126–34, 2002.
- Taccone FS, Rodriguez-Villalobos H, DeBacker D, et al. Successful treatment of septic shock due to pan-resistant *Acinetobacter baumannii* using combined antimicrobial therapy including tigecycline. *Eur J Clin Microbiol Infect Dis* 25:257–60, 2006.
- Tenover FC, Moellering RC. The rationale for revising the clinical and laboratory standards institute vancomycin minimal inhibitory concentration interpretive criteria for *Staphylococcus aureus*. *Clin Infect Dis* 44:1208–1215, 2007.
- Tomioka H, Sato K, Shimizu T, et al. Anti-*Mycobacterium tuberculosis* activities of new fluoroquinolones in combination with other antituberculous drugs. *J Infect* 44:160–5, 2002.
- Turner PJ. Trends in antimicrobial susceptibilities among bacterial pathogens isolated from patients hospitalized in European medical centers: 6-year report of the MYSTIC Surveillance Study (1997–2002). *Diagn Microbiol Infect Dis* 51:281–9, 2005.
- Wagenvoort JHT, Ciraci VE, Penders RJR, et al. Superior discriminative performance of ceftizoxime disk diffusion test for detecting methicillin-resistant *Staphylococcus aureus*. *Eur J Clin Microbiol Infect Dis* 25:405–6, 2006.
- Walsh FM, Amyes SG. Microbiology and drug resistance mechanisms of fully resistant pathogens. *Curr Opin Microbiol* 7:439–44, 2004.
- Yong D, Lee K, Yum JH, et al. Imipenem-EDTA disk method for differentiation of metallo- $\beta$ -lactamase-producing clinical isolates of *Pseudomonas* spp. and *Acinetobacter* spp. *J Clin Microbiol* 40:3798–801, 2002.
- Yuan S, Astion ML, Shapiro J, et al. Clinical impact associated with corrected results in clinical microbiology testing. *J Clin Microbiol* 43:2188–93, 2005.

### TEXTBOOKS

- Anaissie EJ, McGinnis MR, Pfaller MA (eds). *Clinical Mycology*. Churchill Livingstone, New York, 2003.
- Bennett JE, Dolin R, Blaser MJ (eds). *Mandell, Douglas, and Bennett's Principles and Practice of Infectious Diseases (8th Ed)*, Elsevier Churchill Livingstone, New York, London, 2015.
- Bottone EJ (ed). *An Atlas of the Clinical Microbiology of Infectious Diseases, Volume 1. Bacterial Agents*. The Parthenon Publishing Group, Boca Raton, 2004.
- Bottone EJ (ed). *An Atlas of the Clinical Microbiology of Infectious Diseases, Volume 2. Viral, fungal, and parasitic agents*. Taylor and Francis, 2006.
- Bowden RA, Ljungman P, Snyderman DR (eds). *Transplant Infections (3rd Ed)*, Lippincott Williams & Wilkins, Philadelphia, PA, 2010.

- Bryskier A (ed). Antimicrobial Agents. ASM Press, Washington, D.C., 2005.
- Cohen J, Powderly WG, Opal S (Eds). Infectious Diseases (4th Ed). Elsevier, Philadelphia 2015.
- de la Maza LM, Pezzlo MT, Shigei JT, et al. (eds). Color Atlas of Medical Bacteriology. ASM Press, Washington, D.C., 2004.
- Forbes BA, Sahm DF, Weissfeld AS, et al. (eds). Bailey & Scott's Diagnostic Microbiology (12<sup>th</sup> Ed), St. Louis, Mosby, 2007.
- Gorbach SL, Bartlett JG, Blacklow NR (eds). Infectious Diseases (3<sup>rd</sup> Ed), Philadelphia, Lippincott, Williams & Wilkins, 2004.
- Grayson ML (ed). Kucers' The Use of Antibiotics (6<sup>th</sup> Ed), ASM Press, Washington, DC, 2010.
- Janda JM, Abbott SL (eds). The Enterobacteria (2<sup>nd</sup> Ed), ASM Press, Washington, D.C., 2006.
- Koneman EW, Allen SD, Janda WM, et al. (eds). Color Atlas and Textbook of Diagnostic Microbiology (5<sup>th</sup> Ed), Lippincott-Raven Publishers, Philadelphia, 1997.
- Lorian V (ed). Antibiotics in Laboratory Medicine (5<sup>th</sup> Ed), Lippincott Williams & Wilkins, Philadelphia, 2005.
- Madigan MT, Martinko JM, Parker J (eds). Brock Biology of Microorganisms (10<sup>th</sup> Ed), Prentice Hall, Upper Saddle River, NJ, 2003.
- Scholar EM Pratt WB (eds). The Antimicrobial Drugs (2<sup>nd</sup> Ed), Oxford University Press, New York, 2000.
- Versalovic J, Carroll KC, Funke G, et al. (eds). Manual of Clinical Microbiology (10<sup>th</sup> Ed), Washington, DC, ASM Press, 2011.

## Chapter 4

**Parasites, Fungi, Unusual Organisms\*****Kenneth F. Wagner, DO, James H. McGuire, MD****Burke A. Cunha, MD, Jean E. Hage, MD****John H. Rex, MD, Edward J. Bottone, PhD**

Parasites, Fungi, Unusual Organisms in Blood .....	254
Parasites, Fungi, Unusual Organisms in CSF/Brain .....	261
Parasites, Fungi, Unusual Organisms in Lungs. ....	265
Parasites, Fungi, Unusual Organisms in Heart .....	273
Parasites, Fungi, Unusual Organisms in the Liver .....	273
Parasites, Fungi, Unusual Organisms in Stool/Intestines .....	276
Parasites, Fungi, Unusual Organisms in Skin/Muscle. ....	283
References and Suggested Readings .....	298
Treatment of Malaria in Adults in the US .....	721

## Parasites, Fungi, Unusual Organisms in Blood

### Microfilaria in Blood

Subset	Pathogen	Preferred Therapy	Alternate Therapy
Filariasis	<i>Brugia malayi</i>	Doxycycline 100 mg (PO) of q12h × 6 weeks <b>plus</b> Diethylcarbamazine: day 1: 50 mg (PO) day 2: 50 mg (PO) q8h day 3: 100 mg (PO) q8h days 4–14: 2 mg/kg (PO) q8h	Ivermectin 200 mcg/kg (PO) × 1 dose ± albendazole 400 mg (PO) × 1 dose
	<i>Wuchereria bancrofti</i>	Doxycycline 100 mg (PO) q12h × 6 weeks <b>plus</b> Albendazole 400 mg (PO) × 1 dose <b>plus</b> Diethylcarbamazine: day 1: 50 mg (PO) day 2: 50 mg (PO) q8h day 3: 100 mg (PO) q8h days 4–14: 2 mg/kg (PO) q8h	Ivermectin 400 mcg/kg (PO) × 1 dose.

#### **Brugia malayi**

**Clinical Presentation:** May present as an obscure febrile illness, chronic lymphedema, lymphangitis, or cutaneous abscess. “Filarial fevers” usually last 1 week and spontaneously remit.

**Diagnostic Considerations:** Diagnosis by demonstrating microfilaria on Giemsa’s stained thick blood smear or by using the concentration method; yield is increased by passing blood through a Millipore filter before staining. Several smears should be taken over 24 hours. Adult worms may be detected in scrotal lymphatics by ultrasound. Common infection in Southeast Asia (primarily China, Korea, India, Indonesia, Malaysia, Philippines, Sri Lanka). Most species have nocturnal periodicity (microfilaria in blood at night). Eosinophilia is most common during periods of acute inflammation.

**Pitfalls:** Genital manifestations—scrotal edema, epididymitis, orchitis, hydrocele—are frequent with *W. bancrofti*, but rare with *B. malayi*.

**Prognosis:** Related to state of health and extent of lymphatic obstruction. No satisfactory treatment is available. Single-dose ivermectin is effective treatment for microfilaremia, but does not kill the adult worm (although diethylcarbamazine kills some). If no microfilaria in blood, full-dose diethylcarbamazine (2 mg/kg q8h) can be started on day one. Antihistamines or corticosteroids may decrease allergic reactions from disintegration of microfilaria.

#### **Wuchereria bancrofti**

**Clinical Presentation:** May present as an obscure febrile illness, chronic lymphedema, lymphangitis, or cutaneous abscess. Genital (scrotal) lymphatic edema, groin lesions, epididymitis, orchitis, hydroceles are characteristic. Chyluria may occur. “Filarial fevers” usually last 1 week and spontaneously remit. Lymphedema worsened by cellulitis associated with *Tinea pedis* infections.

**Diagnostic Considerations:** Diagnosis by demonstrating microfilaria on Giemsa’s stained thick blood smear or by using the concentration method; yield is increased by passing blood through a Millipore filter before staining. Several smears should be taken over 24 hours. *W. bancrofti* is the most common

human filarial infection, particularly in Asia (China, India, Indonesia, Japan, Malaysia, Philippines), South-east Asia, Sri Lanka, Tropical Africa, Central/South America, and Pacific Islands. Most species have nocturnal periodicity (microfilaria in blood at night). Eosinophilia is common.

**Pitfalls:** Differentiate from “hanging groins” of Loa Loa, which usually do not involve the scrotum.

**Prognosis:** Related to state of health and extent of lymphatic obstruction. No satisfactory treatment is available. Single-dose ivermectin is effective treatment for microfilaremia, but does not kill the adult worm (although diethylcarbamazine kills some). If no microfilaria in blood, full-dose diethylcarbamazine (2 mg/kg q8h) can be started on day one. Antihistamines or corticosteroids decrease allergic reactions from disintegration of microfilaria. Wolbachia bacteria are endosymbionts in *W. bancrofti* filariasis. Treatment with doxycycline effective against Wolbachia which are important in microfilarial reproduction.

## Trypanosomes in Blood

Subset	Pathogen	Preferred Therapy	Alternate Therapy
Chagas' disease (American trypanosomiasis)	<i>Trypanosoma brucei cruzi</i>	Nifurtimox 2–3 mg/kg/day (PO) q6h × 30–90 days	Benznidazole 2.5–3.5 mg/kg (PO) q12h × 60 days
Loa Loa (Loiasis)	<i>L. loa</i>	Diethylcarbamazine 2 mg/kg (PO) q8h × 3 weeks	Albendazole 200 mg (PO) q12h × 3 weeks
Sleeping sickness <i>West African (Gambian) trypanosomiasis</i>	<i>Trypanosoma brucei gambiense</i>	<u>Early disease</u> Pentamidine 4 mg/kg (IM) q24h × 7 days	
		<u>Late disease</u> Melarsoprol 2.2 mg/kg (IV) q24h × 10 days <b>plus</b> Nifurtimox 15 mg/kg (PO) q8h × 10 days <b>or</b> Eflornithine 100 mg/kg (PO) q6h × 10 days	
<i>East African (Rhodesian) trypanosomiasis</i>	<i>Trypanosoma brucei rhodesiense</i>	<u>Early disease</u> Suramin test dose of 4–5 mg/kg (IV) day 1, then five injections of 20 mg/kg (IV) q 7 days (max. dose 1 gm/injection day 3, 10, 17, 24, 31) max. dose 1 gm/injection	
		<u>Late disease</u> Melarsoprol 3 series of 1.8, 2.16, 2.52 mg/kg (IV) q24h; 3 series of 2.52, 2.88, 3.25 mg/kg (IV) q24h; 3 series of 3.6, 3.6, 3.6 mg/kg (IV) q24h; the series given at intervals of 7 days	

### **Chagas' Disease (*Trypanosoma brucei cruzi*) American Trypanosomiasis**

**Clinical Presentation:** Presents acutely after bite of infected reduviid bug with unilateral painless edema of the palpebrae/periorcular tissues (Romaña's sign), or as an indurated area of erythema and swelling with local lymph node involvement (chagoma). Fever, malaise, and edema of the face and lower extremities may follow. Generalized lymphadenopathy and hepatosplenomegaly occur. Patients with chronic disease may develop cardiac involvement (cardiomyopathy with arrhythmias, heart block, heart failure, thromboembolism), GI involvement (megaesophagus, megaduodenum, megacolon) or CNS involvement in HIV/immunosuppressed.

**Diagnostic Considerations:** Common in Central and South America. Acquired from infected reduviid bug, which infests mud/clay parts of primitive dwellings. Transmitted by blood transfusion (~10%), organ transplants, and congenitally. Diagnosis in acute disease by detecting trypanosomes in wet prep of anticoagulated blood or stained buffy coat smears. Amastigote forms present intracellularly in monocytes/histiocytes in Giemsa-stained smears, bone marrow or lymph node aspirates, or by xenodiagnosis. Screening test ELISA IFA; confirmatory test RIPA (radioimmuno precipitation assay).

**Pitfalls:** Do not overlook the diagnosis in patients from endemic areas with unexplained heart block ± apical ventricular aneurysms. May be transmitted by blood transfusion/organ transplantation.

**Prognosis:** Related to extent of cardiac GI, or CNS involvement.

### **Sleeping Sickness (*T. brucei gambiense/rhodesiense*) West African (Chronic)/East African (Acute) Trypanosomiasis**

**Clinical Presentation:** Sleeping sickness from *T. brucei gambiense* is milder than sleeping sickness from *T. brucei rhodesiense*, which is usually a fulminant infection. A few days to weeks after bite of tsetse fly, patients progress through several clinical stages:

- *Chancre stage:* Trypanosomal chancre occurs at bite site and lasts several weeks.
- *Blood/lymphatic stage:* Blood parasitemia is associated with intermittent high fevers, headaches and insomnia, followed by generalized adenopathy. Posterior cervical lymph node enlargement (Winterbottom's sign) is particularly prominent with *T. brucei gambiense*. Hepatosplenomegaly and transient edema/pruritus/irregular circinate rash are common. Myocarditis (tachycardia unrelated to fevers) occurs early (before CNS involvement) and is responsible for acute deaths from *T. brucei rhodesiense*.
- *CNS stage:* Occurs acutely with East African trypanosomiasis or chronically with West African trypanosomiasis after non-specific symptoms, and is characterized by increasing lethargy, somnolence (sleeping sickness), and many subtle CNS findings. Coma and death ensue without treatment. With melarsoprol, use prednisolone 1 mg/kg (PO) q24h (start steroid 1 day prior to first dose and continue to last dose).

**Diagnostic Considerations:** Diagnosis by demonstrating trypanosomes in blood, chancre, or lymph nodes aspirates by Giemsa-stained thin and thick preparations, light microscopy, or buffy coat concentrates with acridine orange. CSF determines early vs. late stage disease (> 20 WBCs/mm<sup>3</sup>).

**Pitfalls:** Do not miss other causes of prominent bilateral posterior cervical lymph node enlargement, e.g., lymphoma, EBV. Serum arginase a biomarker for effective therapy.

**Prognosis:** Related to extent of cardiac/CNS involvement. Relapse may occur.

### **Loa Loa (Loiasis)**

**Clinical Presentation:** Cutaneous swellings (Calabar swellings) with pruritus. Adults may be visible when migrations under the conjunctiva or under the skin. Disappear in 3 days. Calabar swellings are painless and appear on the extremities. Eosinophilia prominent.

**Diagnostic Considerations:** Demonstrates of microfilariae in blood (at noon) or by demonstration of *L. loa* in skin/eye. Immunodiagnosis unhelpful.

**Pitfalls:** Calabar swellings occur one at time and may last for hours/days.

**Prognosis:** Poorest with CNS involvement.

## Spirochetes in Blood

Subset	Pathogen	Preferred Therapy	Alternate Therapy
Relapsing fever <i>Louse-borne (LBRF)</i> <i>Tick-borne (TBRF)</i>	<i>Borrelia recurrentis</i> > 15 <i>Borrelia</i> species (U.S. <i>B. hermsi</i> ; Africa: <i>B. duttonii</i> ; Africa/Middle East: <i>B. crocidurae</i> )	<b>LBRF</b> Erythromycin 500 mg (IV or PO) q6h × 7 days  <b>TBRF</b> Doxycycline 200 mg (PO) × 3 days, then 100 mg (PO) q12h × 7 days	<b>TBRF with CNS involvement</b> Penicillin G 2 mu (IV) q4h × 2 weeks <b>or</b> Ceftriaxone 1 gm (IV) q12h × 2 weeks <b>or</b> Cefotaxime 3 gm (IV) q6h × 2 weeks
Rat bite fever	<i>Spirillum minus</i>	Penicillin G 4 mu (IV) q4h × 2 weeks <b>or</b> Ceftriaxone 1 g (IV) q24h × 2 weeks <b>or</b> Doxycycline 200 mg (IV or PO) q12h × 3 days, then 100 mg (IV or PO) q12h × 11 days	Amoxicillin 1 gm (PO) q8h × 2 weeks <b>or</b> Erythromycin 500 mg (IV or PO) q6h × 2 weeks <b>or</b> Chloramphenicol 500 mg (IV) q6h × 2 weeks

### **Relapsing Fever, Louse-Borne (LBRF) / Tick-Borne (TBRF)**

**Clinical Presentation:** Abrupt onset of “flu-like” illness with high fever, rigors, headache, myalgias, arthralgias, tachycardia, dry cough, abdominal pain after exposure to infected louse or tick. Truncal petechial rash and conjunctival suffusion are common. Hepatosplenomegaly/DIC may occur. Bleeding or rash at bite site. Complications more common in LBRF. Fevers last ~ 1 week, remit for a week, and usually relapse only once in LBRF, but several times in TBRF. Relapses usually last 2–3 days. Fevers are often higher in TBRF.

**Diagnostic Considerations:** *Borreliae* seen in Wright/Giemsa-stained blood smears. LBRF is endemic in South American Andes, Central and East Africa, and is associated with crowded, unhygienic conditions. Soft ticks (*Ornithodoros*) TBRF main vector. Bite at night, patients do not recall tick bite. TBRF is seen throughout the world, and is endemic in Western U.S., British Columbia, Mexico, Central/South America, Mediterranean, Central Asia, and Africa. With TBRF, meningismus ± facial nerve palsy common with *B. duttoni* rare with *B. hermsi*.

**Pitfalls:** Spirochetes are most likely to be seen during febrile periods. Blood smears may be negative if not obtained during fever.

**Prognosis:** Good if treated early. Usually no permanent sequelae.



**Rat Bite Fever (*Spirillum minus*)**

**Clinical Presentation:** Infection develops 1–4 weeks following bite of a rat. Healed rat bite becomes red, painful, swollen and ulcerated, with regional lymphangitis/adenopathy. Recurrent fevers occurs in 2–4 day fever cycles. Fevers are usually accompanied by chills, headache, photophobia, nausea, vomiting. Rash on palms/soles develops in > 50%. Arthritis, myalgias, and SBE are rare.

**Diagnostic Considerations:** Spirochetes are seen in Wright/Giemsa-stained blood smears. Differential diagnosis includes *Borrelia*, malaria, and lymphoma. VDRL is positive.

**Pitfalls:** May be confused with syphilis, due to rash on palms/soles and false-positive syphilis serology in 50%. SBE occurs with *Streptobacillus moniliformis*, not *S. minus* (unless there is preexisting valvular disease). Bite wound ulcerates in *S. minus*, not *Streptobacillus moniliformis*.

**Prognosis:** Patients with arthritis have a protracted course.

**Intracellular Inclusion Bodies in Blood**

Subset	Pathogen	Preferred Therapy	Alternate Therapy
Babesiosis	<i>Babesia microti</i>	Azithromycin 500 mg (PO) × 1, then 250 mg (PO) q24h × 7 days <b>plus</b> Atovaquone (suspension) 750 mg (PO) q12h × 7 days	Clindamycin 600 mg (PO) q8h × 7 days <b>plus</b> Quinine 650 mg (PO) q8h × 7 days
Ehrlichiosis/ anaplasmosis <i>Human monocytic ehrlichiosis (HME)</i>  <i>Human granulocytic anaplasmosis (HGA)</i>	<i>Ehrlichia chaffeensis</i> , ewubgum  <i>Anaplasma (Ehrlichia) phagocytophilum</i>	Doxycycline 200 mg (IV or PO) q12h × 3 days, then 100 mg (IV or PO) q12h × 1–2 weeks total	Any once-daily quinolone (IV or PO) × 1–2 weeks <b>or</b> Rifampin 300 mg (PO) q12h × 1–2 weeks* <b>or</b> Chloramphenicol 500 mg (IV or PO) q6h × 1–2 weeks*
<b>Severe Malaria</b> (Usually <i>P. falciparum</i> )		<b>Quinidine gluconate plus either doxycycline or clindamycin</b> 6.25 mg base/kg (= 10 mg salt/kg) loading dose (IV) over 1–2 h, then 0.0125 mg base/kg/min (= 0.02 mg salt/kg/min) continuous infusion for at least 24h. An alternative regimen is 15 mg base/kg (= 24 mg salt/kg) loading dose infused over 4 hours, followed by 7.5 mg base/kg (= 12 mg salt/kg) infused over 4 hours every 8 hours, starting 8 hours after the loading dose. Once parasitemia <1%; and patient can take oral medications, complete therapy with oral quinine or an oral regimen. (see below chloroquine-sensitive or resistant). <b>Quinidine/quinine course = 7 days in Southeast Asia; = 3 days in Africa or south America.</b>	

\* May be used in pregnancy.

## Intracellular Inclusion Bodies in Blood (cont'd)

Subset	Therapy
	<p style="text-align: center;"><b>or</b></p> <p><b>Quinine</b> 20 mg (salt)/kg (IV) over 4 hours (in D<sub>2</sub>W), then 10 mg (salt)/kg (IV) over 2 hours q8h until able to take oral meds; complete 7 days total therapy with <b>doxycycline</b> or <b>oral regimen</b> (see below chloroquine-sensitive or resistant).</p> <p style="text-align: center;"><b>or</b></p> <p><b>Artemether-lumefantrine</b> 3.2 mg/kg (IM) × 1 dose, then 1.6 mg/kg (IM) q24h until able to take oral meds; complete 7 days total therapy with <b>doxycycline</b> or <b>oral regimen</b> (see below chloroquine-sensitive or resistant).</p> <p style="text-align: center;"><b>or</b></p> <p><b>Artesunate</b> 2.4 mg/kg (IV) initially and at 12, 24, and 48 hours, then q24h until able to take oral meds; complete 7 days total therapy with <b>doxycycline</b> or <b>oral regimen</b> (see below chloroquine-sensitive or resistant).</p>
<p><b>Uncomplicated Malaria</b> (<i>P. falciparum</i>, <i>P. malariae</i>, <i>P. knowlesi</i> or unidentified species)</p> <p><i>P. vivax</i> and <i>P. ovale</i></p>	<p><b>Chloroquine-sensitive</b></p> <p><b>Chloroquine phosphate</b> 600 mg base (= 1000 mg salt) (PO) immediately, followed by 300 mg base (= 500 mg salt) (PO) at 6, 24, and 48 hours. Total dose = 1500 mg base (= 2500 mg salt).</p> <p style="text-align: center;"><b>or</b></p> <p><b>Hydroxychloroquine</b> 620 mg base (= 800 mg salt) (PO) loading dose, then 310 mg base (= 400 mg salt) (PO) at 6, 24, and 48 hours.</p> <p><b>Plus primaquine phosphate</b> 30 mg base (PO) q24h × 2 weeks.</p> <p><b>Chloroquine-resistant</b></p> <p><b>Quinine sulfate</b> 625 mg base (= 625 mg salt) (PO) q8h × 7 days <i>plus</i> <b>doxycycline</b> 200 mg (PO) q12h × 3 days, then 100 mg (PO) q12h × 4 days.</p> <p style="text-align: center;"><b>or</b></p> <p><b>Atovaquone/proguanil</b> (250/100 gm PO tab) 4 tablets as single dose or 4 tablets (PO) q12h × 3 days.</p> <p style="text-align: center;"><b>or</b></p> <p><b>Artesunate</b> 4 mg/kg (PO) × 3 days <i>plus</i> <b>Mefloquine</b> 684 mg base (= 750 mg salt) (PO) as initial dose followed by 456 mg base (= 500 mg salt) (PO) given 6–12 hours after initial dose. Total dose = 1250 mg salt.</p> <p style="text-align: center;"><b>or</b></p> <p><b>Artemether/lumefantrine</b> 4 tabs = 1 dose (20/120 mg tablets) give initial dose, followed by second dose 8h later, then 1 dose (PO) q12h for the following 2 days.</p> <p style="text-align: center;"><b>or</b></p> <p><b>Dihydroartemisinin</b> 40 mg <i>plus</i> piperazine 320 mg (PO) q24h × 3 days.</p>

For CDC guidelines for malaria in US (see p. 721).

**Babesiosis (*Babesia microti*)**

**Clinical Presentation:** “Malarial-like illness” with malaise, fever, relative bradycardia, shaking chills, myalgias, arthralgias, headache, abdominal pain, and splenomegaly. Laboratory abnormalities include anemia, atypical lymphocytes, relative lymphopenia, thrombocytopenia, mildly elevated LFTs, highly elevated  $\uparrow$  ESR,  $\uparrow$  ferritin, and  $\uparrow$  LDH. Transmitted by infected Ixodes ticks.

**Diagnostic and Considerations:** Characteristic four merozoites (often pear shaped) arranged in “Maltese cross” formation (tetrads). Serology diagnostic of acute infection. Hyposplenic patients may have profound hemolytic anemia and life-threatening infection.

**Pitfalls:** Co-infection with Lyme disease may occur. No serological cross-reactivity between *Babesia* and *Borrelia* (Lyme disease). Merozoites only may be confused with *P. falciparum* malaria. Extra-RBC forms and vacuolated RBCs distinguish babesiosis from malaria. Travel history is important. Doxycycline ineffective.

**Prognosis:** May be more severe with Lyme disease co-infection. Severe/fatal if  $\downarrow$ /absent splenic function. Exchange transfusions may be life saving.

**Ehrlichiosis (HME)/Anaplasmosis (HGA)**

**Clinical Presentation:** Acute febrile illness with chills, headache, malaise, myalgias, leukopenia, relative lymphopenia, atypical lymphocytes, thrombocytopenia,  $\uparrow$  LFTs,  $\downarrow$  ESR,  $\uparrow$  ferritin. No vasculitis. Resembles Rocky Mountain spotted fever (RMSF), but without rash.

**Diagnostic Considerations:** Characteristic “morula” (spherical, basophilic, mulberry-shaped, cytoplasmic inclusion bodies) may be seen in peripheral blood neutrophils in HGA. PCR from blood is 86% sensitive and highly specific for early diagnosis. Obtain acute and convalescent IFA serology. HGA vector is Ixodes ticks clinical co-infection with *B. burgdorferi* (Lyme Disease) is rare, but may occur. Main HME vector *Amblyomma americanum* (lone star tick).

**Pitfalls:** No morula with HME. Rash uncommon in HME and rare in HGA. *E. chaffeensis* (HME) titers will not be elevated with *A. phagocytophilium* (HGA) and vice versa. PCR/blood smears positive early. Seropositivity increases over time.

**Prognosis:** Good if treated early. Delayed response/more severe with Lyme co-infection disease.

**Malaria (*Plasmodium ovale/vivax/falciparum/malariae/knowlesi*)**

**Clinical Presentation:** Presents acutely with fever/chills, severe headaches, cough, nausea/vomiting, diarrhea, abdominal/back pain. Typical “malarial paroxysm” consists of chills, fever and profuse sweating, followed by extreme prostration. There are a paucity of physical findings, but most have tender hepatomegaly/splenomegaly and relative bradycardia.  $\uparrow$  T. bilirubin, thrombocytopenia, atypical lymphocytes, and  $\uparrow$  LFTs are common.

**Diagnostic Considerations:** Diagnosis by visualizing *Plasmodium* on thick/thin Giemsa or Wright-stained smears.

**Pitfalls:** Be wary of diagnosing malaria without headache. Dengue most closely resembles malaria. On abdominal US, dengue patients have gallbladder wall thickening/splenomegaly but not hepatomegaly vs. malaria patients which have a normal gallbladder, splenomegaly/heptomegaly. If no atypical lymphocytes on smear (auto cell counters are insensitive to atypical lymphocytes), question the diagnosis of malaria. *P. knowlesi* (monkey malaria emerging cause of human malaria. Resembles *P. malariae* (microscopically) but may be severe resembling *P. falciparum* (clinically). Treat *P. knowlesi* as chloroquine sensitive (*P. vivax*, *P. ovale*, *P. malariae*) malaria.

**Prognosis:** Related to species degree of parasitemia *P. falciparum* with high-grade parasitemia is most severe, and may be complicated by coma, hypoglycemia, renal failure, or non-cardiogenic pulmonary edema. If parasitemia exceeds 15%, consider exchange transfusions.

**Fungi/Mycobacterium in Blood**

See histoplasmosis (p. 341), Mycobacterium tuberculosis (treat as pulmonary TB, p. 326), Mycobacterium avium-intracellulare (p. 321, 342).

**Parasites, Fungi, Unusual Organisms in CSF/Brain****Cysts/Mass Lesions in CSF/Brain**

Subset	Pathogens	Preferred Therapy	Alternate Therapy
Cerebral nocardiosis	Nocardia sp.	<u>Preferred IV Therapy:</u> TMP-SMX (TMP 5 mg/kg, SMX 15 mg/kg) (IV) q6h until clinical improvement, then (PO) therapy  <u>Alternate IV PO Therapy:</u> Minocycline or Doxycycline 100 mg (IV) q12h until clinical improvement, then (PO) therapy	<u>Preferred PO Therapy</u> TMP-SMX 1 DS tablet (PO) q12h × 6 months  <u>Alternate PO Therapy</u> Minocycline 100 mg (PO) q12h × 6 months  <b>or</b> Doxycycline 100 mg (PO) q12h × 6 months
Cryptococcal meningitis/cryptococcomas	Cryptococcus neoformans	See p. 22	
Cerebral amebiasis	Entamoeba histolytica	Metronidazole 750 mg (PO) q8h × 10 days  <b>or</b> Tinidazole 800 mg (PO) q8h × 5 days	
Primary amebic meningo-encephalitis	Naegleria fowleri	See p. 21	
Granulomatous amebic encephalitis	Acanthamoeba	See p. 22	
Cerebral echinococcosis (hydatid cyst disease)	Echinococcus granulosus or multilocularis	Surgical resection plus Albendazole 400 mg* (PO) q12h until cured	Surgical resection plus Mebendazole 50 mg/kg (PO) q24h until cured
Cerebral gnathostomiasis	Gnathostoma spinigerum	Surgical resection	Albendazole 400 mg (PO) q12h × 3 weeks <b>or</b> Ivermectin 200 mcg/kg/d × 2 days
Cerebral coenurosis	Taenia multiceps	Surgical resection	

**Cysts/Mass Lesions in CSF/Brain (cont'd)**

Subset	Pathogens	Preferred Therapy	Alternate Therapy
Neurocysticercosis	<i>Taenia solium</i>	Albendazole 400 mg* (PO) q12h × 2 weeks	Praziquantel 33 mg/kg (PO) q8h × 1 day, then 15 mg/kg (PO) q8h × 2–4 weeks ± cimetidine 400 mg (PO) q8h (to ↑ praziquantel levels)
Cerebral paragonimiasis (lung fluke)	<i>Paragonimus westermani</i>	Praziquantel 25 mg/kg (PO) q8h × 2 days	Bithionol 50 mg/kg (PO) q48h × 2 weeks  <b>or</b> Tridabendazole 10 mg/kg (PO) q24h × 2 days
Cerebral toxoplasmosis	<i>Toxoplasma gondii</i>	See pp. 331–332	
Chagas' disease (American trypanosomiasis)	<i>Trypanosoma brucei cruzi</i>	Nifurtimox 2 mg/kg (PO) q6h × 4 months	Benznidazole 3.5 mg/kg (PO) q12h × 2 months

\* If < 60 kg, give albendazole 7.5 mg/kg.

**Cerebral Nocardiosis**

**Clinical Presentation:** CNS mass lesion resembling brain tumor/abscess. Symptoms are highly variable, and result from local effects of granulomas/abscesses in CNS. Up to 40% of patients with systemic nocardiosis have associated mass lesions in CNS.

**Diagnostic Considerations:** Diagnosis by demonstrating *Nocardia* (gram positive, delicate, beaded, branching filaments) in brain biopsy specimens. Notify laboratory for modified acid fast specimen staining/aerobic cultures if suspect *Nocardia*. *Nocardia* are weakly acid-fast and aerobic.

**Pitfalls:** Usually not limited to brain. Look for *Nocardia* in skin, lungs or liver. Use in-vitro susceptibility data to guide therapy for refractory cases. IV regimens are recommended for critically ill patients. HIV patients require life-long suppression with TMP-SMX.

**Prognosis:** Related to health of host, degree of immunosuppression, and extent of lesions.

**Cerebral Amebiasis (*Entamoeba histolytica*)**

**Clinical Presentation:** Rare cause of brain abscess. Onset is frequently abrupt with rapid progression. Suspect in patients with a history of amebiasis and altered mental status/focal neurologic signs. If present, meningeal involvement resembles acute bacterial meningitis. CT/MRI shows focal lesions.

**Diagnostic Considerations:** Diagnosis by demonstrating *E. histolytica* trophozoites in wet preps or by trichrome stain from aspirated brain lesions under CT guidance. Worldwide distribution. Mass lesions may be single or multiple, and more commonly involve the left hemisphere. Most patients have concomitant liver ± lung abscesses.

**Pitfalls:** Trophozoites/cysts in stool are not diagnostic of CNS disease. *E. histolytica* serology is often positive, but is nonspecific. *E. histolytica* trophozoites are not present in CSF.

**Prognosis:** Related to size/location of CNS lesions.

**Primary Amebic Meningoencephalitis (*Naegleria fowleri*)** (see p. 23)**Granulomatous Amebic Encephalitis (*Acanthamoeba*)** (see p. 23)**Cerebral Echinococcosis (*Echinococcus granulosus*) Hydatid Cyst Disease**

**Clinical Presentation:** Most cysts are asymptomatic. Mass lesions may cause seizures, cranial nerve abnormalities, other focal neurologic symptoms.

**Diagnostic Considerations:** CT/MRI typically shows a single large cyst without edema or enhancement. Multiple cysts are rare. Diagnosis by demonstrating protoscolices in “hydatid sand” in cysts. Usually associated with liver/lung hydatid cysts.

**Pitfalls:** *E. granulosus* serology lacks specificity.

**Prognosis:** Related to size/location of CNS cysts. CSF eosinophilia is not a feature of CNS involvement. Treatment consists of surgical removal of total cyst after instilling cysticidal agent (hypertonic saline, iodophor, ethanol) into cyst plus albendazole.

**Cerebral Echinococcosis (*Echinococcus multilocularis*) Hydatid Cyst Disease**

**Clinical Presentation:** Frequently associated with hydatid bone cysts (may cause spinal cord compression), liver/lung cysts. Peripheral eosinophilia occurs in 50%, but eosinophils are not seen in the CSF.

**Diagnostic Considerations:** *E. multilocularis* ELISA is sensitive and specific.

**Pitfalls:** Praziquantel is ineffective for CNS hydatid cyst disease. Imaging studies suggest carcinoma/sarcoma. Diagnosis is frequently not made until brain biopsy.

**Prognosis:** If treatment is effective, improvement of CNS lesions is evident in 8 weeks. Brain/bone cysts are difficult to cure.

**Cerebral Gnathostomiasis (*Gnathostoma spinigerum*)**

**Clinical Presentation:** Nausea, vomiting, increased salivation, skin flushing, pruritus, urticaria, and upper abdominal pain 1–6 days after exposure. Cerebral form presents as eosinophilic meningitis with radiculomyeloencephalitis, with headache and severe sharp/shooting pains in extremities often followed by paraplegia and coma. Any cranial nerve may be involved. The most characteristic feature is changing/migratory neurological findings. Intense peripheral eosinophilia occurs in 90% of patients. CSF has eosinophilic pleocytosis and may have RBCs.

**Diagnostic Considerations:** In cases with ocular involvement, the worm may be seen in the anterior chamber of eye. Specific *Gnathostoma* serology of CSF is helpful in establishing the diagnosis. Acquired from infected cat/dog feces. Most cases occur in Southeast Asia also Central/South America (Mexico, Peru) and most recently in Africa (Botswana). Few other CNS infections have both RBCs and eosinophils in the CSF.

**Pitfalls:** Do not miss associated eye involvement or characteristic episodic non-pitting subcutaneous edema.

**Prognosis:** Related to invasion of medulla/brainstem.

**Cerebral Coenurosis (*Taenia multiceps*)**

**Clinical Presentation:** CNS mass lesion with seizures/cranial nerve abnormalities, often presenting as a posterior-fossa syndrome. Common sites of CNS involvement include paraventricular and basal subarachnoid spaces. Eosinophils not in CSF.

**Diagnostic Considerations:** Diagnosis by demonstrating protoscolices in brain cyst specimens. Worldwide distribution. Transmitted via dog feces.

**Pitfalls:** Do not miss associated ocular lesions, which mimic intraocular neoplasms/granulomas.

**Prognosis:** Related to size/extent of CNS lesions.

### **Neurocysticercosis (*Taenia solium*)**

**Clinical Presentation:** Chronic eosinophilic meningitis/mass lesions with seizures. Hydrocephalus is common. Spinal involvement may result in paraplegia. Cerebral cysts are usually multiple.

**Diagnostic Considerations:** CT/MRI shows multiple enhancing and non-enhancing unilocular cysts. Diagnosis by specific *T. solium* serology of serum/CSF or excision of cyst. Neurocysticercosis is the most common CNS parasite. Worldwide in distribution; most common in Eastern Europe, Asia, Latin America.

**Pitfalls:** Cranial nerve abnormalities are uncommon.

**Prognosis:** Related to extent/location of CNS lesions. Adjunctive therapy includes corticosteroids, anti-epileptics, and shunt for hydrocephalus.

### **Cerebral Paragonimiasis (*Paragonimus westermani*) Lung Fluke**

**Clinical Presentation:** Can resemble epilepsy, cerebral tumors, or brain embolism. Primary focus of infection is pulmonary, with pleuritic chest pain, cough, and night sweats. CNS findings are a manifestation of extrapulmonary (ectopic) organ involvement.

**Diagnostic Considerations:** Diagnosis by demonstrating operculated eggs in sputum, pleural fluid, or feces. Multiple sputum samples are needed to demonstrate *P. westermani* eggs. Charcot-Leyden crystals are seen in sputum. Endemic in Far East, India, Africa, and Central/South America.

**Pitfalls:** Extrapulmonary (ectopic) organ involvement (cerebral, subcutaneous, abdominal) is common. Up to 20% of patients have normal chest x-rays.

**Prognosis:** Related to size/location of CNS cysts and extent of lung involvement.

### **Cerebral Toxoplasmosis (*T. gondii*)** (see pp. 331–332)

### **Cerebral Cryptococcosis (*C. neoformans*)** (see pp. 329–330)

### **Chagas' Disease (*Trypanosoma brucei cruzi*) American Trypanosomiasis**

**Clinical Presentation:** Acute unilateral periorbital cellulitis (Romaña's sign) or regional adenopathy and edema of extremity at site of infected reduviid bug (Chagoma). Chronic disease manifests as myocarditis/heart block or megaesophagus, megaduodenum, megacolon. Hepatosplenomegaly is common. Overt CNS signs are frequently absent. CNS Chagas' disease typically have hypodense ring enhancing lesions with surrounding edema on head CT/MRI scans. If meningoencephalitis develops, the prognosis is very poor. In immunosuppressed patients (especially AIDS), recrudescence of disease occurs with development of *T. brucei cruzi* brain abscesses.

**Diagnostic Considerations:** Diagnosis in acute disease by demonstrating trypanosomes in wet prep of anticoagulated blood or stained buffy coat smears. Amastigote forms present intracellularly in monocytes/histiocytes in Giemsa-stained smear of bone marrow/lymph node aspirate, or by xenodiagnosis. Serology (mostly used for chronic disease) has limited value in endemic areas due to lack of specificity, but is useful in non-endemic areas. Common in Central/South America. Acquired from infected reduviid bugs, which infest mud/clay/stone parts of primitive dwellings. Infection in humans occurs in areas containing reduviids that defecate during or immediately after a blood meal.

**Diagnostic Considerations:** Screening test ELISA IFA; confirmatory test RIPA (radioimmuno precipitation assay).

**Pitfalls:** Do not overlook diagnosis in persons from endemic areas with unexplained heart block. For children ages 11–16 years, use nifurtimox 3.5 mg/kg (PO) q6h × 3 months. For children < 11 years, use nifurtimox 5 mg/kg (PO) q6h × 3 months.

**Prognosis:** Related to extent of GI cardiac or CNS involvement. The addition of gamma interferon to nifurtimox × 20 days may shorten the acute phase of the disease.

## Parasites, Fungi, Unusual Organisms in Lungs

### Pulmonary Cystic Lesions/Masses

Subset	Pathogens	Preferred Therapy	Alternate Therapy
Alveolar echinococcosis	Echinococcus multilocularis	<u>Operable cases</u> Wide surgical resection plus Albendazole 400 mg* (PO) q12h or Mebendazole 50 mg/ kg (PO) q24h until cured	<u>Inoperable cases:</u> Albendazole 400 mg* (PO) q12h × 1 month, then repeat therapy after 2 weeks × 3 cycles (i.e., 4 total months of albendazole)
Pulmonary amebiasis	Entamoeba histolytica	Metronidazole 750 mg (PO) q8h × 10 days	Tinidazole 800 mg (PO) q8h × 5 days
Pulmonary paragonimiasis (lung fluke)	Paragonimus westermani	Praziquantel 25 mg/kg (PO) q8h × 2 days	Bithionol 50 mg/kg (PO) q48h × 4 weeks  <b>or</b> Tridabendazole 10 mg/kg (PO) q24h × 2 days

\* If < 60 kg, give albendazole 7.5 mg/kg.

### Alveolar Echinococcosis (Echinococcus multilocularis)

**Clinical Presentation:** Slowly growing cysts remain asymptomatic for 5–20 years, until space-occupying effect elicits symptoms. Rupture/leak into bronchial tree can cause cough, chest pain, and hemoptysis.

**Diagnostic Considerations:** Diagnosis is suggested by typical “Swiss cheese calcification” findings on chest x-ray, and confirmed by specific *E. multilocularis* serology (which does not cross react with *E. granulosus*). Most common in Northern forest areas of Europe, Asia, North America, and Arctic. Acquired by ingestion of viable parasite eggs in food. Tapeworm-infected canines/cats or wild rodents are common vectors. Less common than infection with *E. granulosus*.

**Pitfalls:** Do not confuse central cavity lesions with squamous cell carcinoma.

**Prognosis:** Related to size/location of cysts.

### Pulmonary Amebiasis (Entamoeba histolytica)

**Clinical Presentation:** Cough, pelvic pain, fever, and right lung/pleural mass mimicking pneumonia or lung abscess. Bronchopleural fistulas may occur. Sputum has “liver-like” taste if cyst ruptures into bronchus. Bacterial co-infection is rare. Amebic lung lesions are associated with hepatic liver abscesses, and invariably involve the right lobe of lung/diaphragm.



**Diagnostic Considerations:** Diagnosis by aspiration of lungs cysts, which may be massive. Amebic serology is sensitive and specific. Worldwide distribution. Acquired by ingesting amebic cysts. Key to diagnosis is concomitant liver involvement; liver abscess presents years after initial diarrheal episode.

**Pitfalls:** Lung involvement is rarely the sole manifestation of amebic infection, and is usually due to direct extension of amebic liver abscess (10–20% of amebic liver abscesses penetrate through the diaphragm and into the lungs). Follow metronidazole with paromomycin 500 mg (PO) q8h × 7 days to eliminate intestinal focus.

**Prognosis:** Related to severity/extent of cysts.

### **Pulmonary Paragonimiasis (*Paragonimus westermani*) Lung Fluke**

**Clinical Presentation:** Mild infection; may be asymptomatic. Acute phase of infection is accompanied by abdominal pain, diarrhea and urticaria, followed by pleuritic chest pain/eosinophilic pleural effusion. Chronic symptoms occur within 6 months after exposure, with dyspnea/dry cough leading to productive cough ± hemoptysis. Complications include lung abscess, bronchiectasis, cough, and night sweats. Eosinophilia may be present acutely.

**Diagnostic Considerations:** Oriental lung fluke acquired by ingestion of freshwater crayfish/crabs. After penetration of the gut/peritoneal cavity, the fluke migrates through the diaphragm/pleural space and invades lung parenchyma. Incubation period is 2–20 days. Diagnosis by demonstrating operculated eggs in sputum, pleural fluid, or feces. Multiple sputum samples are needed to demonstrate *P. westermani* eggs. Charcot-Leyden crystals are seen in sputum, and characteristic chest x-ray findings of ring-shaped/crescent infiltrates with “thin-walled” cavities are evident in ~ 60%. Endemic in Asia, Africa, and Latin America. Chest x-ray findings take months to resolve.

**Pitfalls:** May have extrapulmonary (ectopic) organ involvement, e.g., cerebral, subcutaneous, abdominal. Up to 20% have normal chest x-rays. Commonest cause of eosinophilic pleural effusions in endemic areas. Diagnosis should be questioned if pleural effusion fluid does not have eosinophils.

**Prognosis:** Related to degree of lung damage, e.g., bronchiectasis and extrapulmonary organ involvement, especially CNS.

## **Pulmonary Coin Lesions**

Subset	Pathogens	Preferred Therapy	Alternate Therapy
Dog heartworm	<i>Dirofilaria immitis</i>	No therapy necessary	
Aspergilloma	<i>Aspergillus</i>	No therapy if asymptomatic. Surgery for massive hemoptysis	Itraconazole 200 mg (PO) solution q24h × 3–6 months <b>or</b> Voriconazole (see “usual dose,” p. 714) × 3–6 months

### **Dog Heartworm (*Dirofilaria immitis*)**

**Clinical Presentation:** Asymptomatic “coin lesion” after bite of infected mosquito transmits parasite from dogs to humans. Differential diagnosis includes granulomas and malignancy.

**Diagnostic Considerations:** Diagnosis by specific serology or pathological demonstration of organism in granuloma, usually when a coin lesion is biopsied to rule out malignancy. Worldwide distribution. Acquired from pet dogs. Dirofilariasis causes dog heartworm in carrier, but presents as a solitary lung nodule in humans.

**Pitfalls:** Often confused with malignancy.

**Prognosis:** Excellent.

### **Pulmonary Aspergilloma**

**Clinical Presentation:** Coin lesion(s) ± productive cough, hemoptysis, wheezing. May be asymptomatic. Usually occurs in pre-existing cavitory lung lesions, especially TB with cavity > 2 cm.

**Diagnostic Considerations:** Diagnosis by chest x-ray appearance of fungus ball in cavity and *Aspergillus* precipitins/biopsy or by demonstrating *Aspergillus* hyphae and “fruiting bodies” (conidiophores) in respiratory specimens. May present with “crescent sign” on chest x-ray (white fungus ball silhouetted against black crescent of the cavity).

**Pitfalls:** Role of itraconazole or voriconazole as therapy is unclear.

**Prognosis:** Related to degree of hemoptysis.

### **Pulmonary Infiltrates/Mass Lesions<sup>¶</sup>**

<b>Subset</b>	<b>Pathogens</b>	<b>Preferred Therapy</b>	<b>Alternate Therapy</b>
Pulmonary blastomycosis	<i>Blastomyces dermatitidis</i>	Mild illness: Itraconazole 200 mg (PO) TID × 3 days and then QD or BID for a total of 6–12 months (adequate serum levels should be confirmed). Moderately severe or severe illness: lipid amphotericin for 1–2 weeks followed by itraconazole as for mild illness.	Fluconazole has been disappointing; its use is limited to specialized settings such as CNS blastomycosis. Amphotericin B 0.7–1 mg/kg may be used as initial therapy instead of lipid amphotericin.
Pulmonary histoplasmosis	<i>Histoplasma capsulatum</i>	Mild illness: Therapy is not always needed, but symptoms lasting more than a month may be treated with Itraconazole 200 mg (PO) TID × 3 days and then QD or BID for a total of 6–12 weeks. Moderately severe or severe illness: lipid amphotericin for 1–2 weeks followed by Itraconazole as for mild illness. Chronic cavity histoplasmosis requires at least a year of therapy & confirmation of adequate blood levels.	Strategies are complex (See Wheat LJ et al. Clin Infect Dis 45:807–825, 2007).
Pulmonary paracoccidioidomycosis (South American blastomycosis)	<i>Paracoccidioides brasiliensis</i>	Itraconazole 200 mg (PO) q24h × 6 months <b>or</b> Ketoconazole 400 mg (PO) q24h × 18 months	Amphotericin B 0.5 mg/kg (IV) q24h until 1.5–2.5 grams given

<sup>¶</sup> See Color Atlas (center of book).

## Pulmonary Infiltrates/Mass Lesions (cont'd)

Subset	Pathogens	Preferred Therapy	Alternate Therapy
Pulmonary actinomycosis	Actinomyces israelii	Amoxicillin 1 gm (PO) q8h × 6 months <b>or</b> Doxycycline 100 mg (PO) q12h × 6 months	Clindamycin 300 mg (PO) q8h × 6 months <b>or</b> Chloramphenicol 500 mg (PO) q6h × 6 months
Pulmonary aspergillosis <i>BPA</i> <i>Acute invasive pneumonia/aspergillus</i> <i>Chronic pneumonia aspergillus</i>	Aspergillus	Systemic oral steroids	Itraconazole 200 mg (PO) q12h × 8 months
	Aspergillus	See p. 323	
	Aspergillus	Treat the same as on p. 165	Treat the same as on p. 165
Pulmonary sporotrichosis	Sporothrix schenckii	Lipid amphotericin (p. 525) (IV) q24h × 3 weeks <b>or</b> Amphotericin B 0.5 mg/kg (IV) q24h until 1–2 grams given	Itraconazole 200 mg (PO)* q12h until cured <b>or</b> Lipid amphotericin (p. 525) (IV) q24h until cured
Pulmonary coccidioidomycosis	Coccidioides immitis	Itraconazole 200 mg (PO)* q12h until cured <b>or</b> Fluconazole 800 mg (IV or PO) × 1 dose, then 400 mg (PO) q24h until cured	Amphotericin B 1 mg/kg (IV) q24h × 7 days† <b>or</b> Lipid amphotericin (p. 525) (IV) q24h × 7 days
Pulmonary nocardiosis	Nocardia asteroides	TMP–SMX 5–10 mg/kg/d (TMP) in 2–4 doses (IV) × 3–6 weeks, then 1 DS tablet (PO) q12h until cured	Minocycline 100 mg (PO) q12h until cured

\* Initiate therapy with Itraconazole 200 mg (IV) q12h × 7–14 days.

† Follow with Itraconazole 200 mg (PO) solution q12h until cured.

**Pulmonary Infiltrates/Mass Lesions (cont'd)**

Subset	Pathogens	Preferred Therapy	Alternate Therapy
Pulmonary cryptococcosis	Cryptococcus neoformans	Fluconazole 800 mg (IV or PO) × 1 dose, then 400 mg (PO) q24h until cured	Amphotericin B 0.5 mg/kg (IV) q24h until 1–2 grams given <b>or</b> Lipid amphotericin (p. 525) (IV) q24h × 3 weeks
Pulmonary zygomycosis (mucormycosis)	Rhizopus/ Mucor/Absidia	Lipid amphotericin (p. 525) (IV) q24h × 1–2 weeks <sup>†</sup> or × 3 wks <b>or</b> Amphotericin B 1–1.5 mg/kg (IV) q24h × 1–2 wks <sup>†</sup> or until 2–3 grams given	Voriconazole (see “usual dose,” p. 714) until cured <b>or</b> Isavuconazole 200 mg (IV) q8h × 48 hours, then 200 mg (IV/PO) q24h until cured <b>or</b> Itraconazole 200 mg (PO)* q12h until cured
Pulmonary pseudallescheriasis	Pseudallescheria boydii/ Scedosporium apiospermum	Voriconazole (see “usual dose,” p. 714) until cured	Itraconazole 200 mg (PO)* q12h until cured

BPA = bronchopulmonary aspergillosis.

\* Initiate therapy with itraconazole 200 mg (IV) q12h × 7–14 days.

† Follow with itraconazole 200 mg (PO) solution q12h until cured.

**Pulmonary Blastomycosis (Blastomyces dermatitidis)**

**Clinical Presentation:** Highly variable. May present as a chronic/non-resolving pneumonia with fever/cough and characteristic “right-sided perihilar infiltrate” ± small pleural effusion.

**Diagnostic Considerations:** Usual sites of dissemination include skin, bones and prostate, not CNS or adrenals. Blastomyces serology may XR (false +) with Aspergillus, Coccidioides, Histoplasma, or paracoccidioidomycosis. β 1,3 D-glucan –, aspergillus galactomannan –.

**Pitfalls:** Dissemination to extra-pulmonary sites may occur years after pneumonia.

**Prognosis:** Related to severity/extent of infection. One-third of cases are self-limited and do not require treatment.

**Pulmonary Histoplasmosis (Histoplasma capsulatum)**

**Clinical Presentation:** Acute primary infection presents as self-limiting flu-like illness with fever, headache, nonproductive cough, chills, and chest pain. Minority of patients become overtly ill with complicated respiratory or progressive pulmonary infection. Can cause arthralgias, E. nodosum, E. multiforme, or pericarditis. May occur in outbreaks. Chronic infection presents as chronic pneumonia resembling TB or chronic disseminated infection. Eosinophilia common.

**Diagnostic Considerations:** May be recovered from sputum or demonstrated in lung tissue specimens. Worldwide distribution, but most common in Central/South Central United States. Acute disseminated histoplasmosis suggests HIV. Intracellular (PMNs/monocytes) organisms may be seen in Wright/Giemsa stained peripheral blood smears/buffy coat smears in disseminated infection. Acute histoplasmosis with immunodiffusion (ID) assay M band precipitins; chronic histoplasmosis with ID assay M band precipitins or CF  $\geq 1:32$ . Histoplasma urinary antigen useful in diagnosis of acute/disseminated histoplasmosis. Histoplasma serology may XR (false +) with coccidiomycosis, blastomycosis, TB or sarcoidosis. Histoplasma urinary antigen may XR with acute/chronic coccidiomycosis (false + histoplasma urinary antigen may provide a clue to the presence of coccidiomycosis).  $\beta$  1,3 D-glucan +, aspergillus galactomannan +.

**Pitfalls:** Pleural effusion is uncommon. Do not treat old/inactive/minimal histoplasmosis, histoplasmosis pulmonary calcification, or histoplasmosis fibrosing mediastinitis. Differentiate from TB and kala-azar. Resembles kala-azar histologically, but rod-shaped kinetoplasts (kala-azar) are absent extracellularly.

**Prognosis:** Related to severity/extent of infection. No treatment is needed for self-limiting acute histoplasmosis presenting as flu-like illness. HIV patients should receive life-long suppressive therapy with itraconazole 200 mg (PO) solution q24h.

### **Pulmonary Paracoccidioidomycosis (South American Blastomycosis)**

**Clinical Presentation:** Typically presents as a chronic pneumonia syndrome with productive cough, blood-tinged sputum, dyspnea, and chest pain. May also develop fever, malaise, weight loss, mucosal ulcerations in/around mouth and nose, dysphagia, changes in voice, cutaneous lesions on face/limbs, or cervical adenopathy. Can disseminate to prostate, epididymis, kidneys, or adrenals.

**Diagnostic Considerations:** Characteristic "pilot wheel" shaped yeast in sputum. Diagnosis by culture and stain (Gomori) of organism from clinical specimen. Found only in Latin American. One-third of cases have only pulmonary involvement. Skin test is non-specific/non-diagnostic.

**Pitfalls:** No distinguishing radiologic features. No clinical adrenal insufficiency, in contrast to TB or histoplasmosis. Hilar adenopathy/pleural effusions are uncommon.

**Prognosis:** Related to severity/extent of infection. HIV require life-long suppression with TMP-SMX 1 DS tablet (PO) q24h or itraconazole 200 mg (PO) solution q24h.

### **Pulmonary Actinomycosis (Actinomyces israelii)**

**Clinical Presentation:** Indolent, slowly progressive infiltrates involving the pulmonary parenchyma  $\pm$  pleural space. Presents with fever, chest pain, weight loss. Cough/hemoptysis are less common. Chest wall sinuses frequently develop. Chest x-ray shows adjacent dense infiltrate. "Sulfur granules" are common in sinus drainage fluid.

**Diagnostic Considerations:** Diagnosis by stain/culture of drainage from sinuses or lung/bone biopsy specimens. Actinomycetes are non-acid fast and anaerobic to microaerophilic.

**Pitfalls:** No CNS lesions, but bone erosion is common with chest lesions. Prior antibiotic therapy may interfere with isolation of organism.

**Prognosis:** Excellent when treated until lesions resolve. Use IV regimen in critically ill patients, then switch to oral regimen.

### **Bronchopulmonary Aspergillosis (BPA/ABPA)**

**Clinical Presentation:** Migratory pulmonary infiltrates in chronic asthmatics. Eosinophilia is common, and sputum shows Charcot-Leyden crystals/brown flecks containing Aspergillus.

**Diagnostic Considerations:** Diagnosis by *Aspergillus* in sputum and high-titers of *Aspergillus* precipitins in serum. BPA is an allergic reaction in chronic asthmatics, *not* an infectious disease. Pulmonary infiltrates with peripheral eosinophilia in chronic asthmatics suggests the diagnosis.

**Pitfalls:** Correct diagnosis is important since therapy is steroids, not antifungals.

**Prognosis:** Related to severity/duration of asthma and promptness of steroid therapy.

### **Acute Invasive Aspergillus Pneumonia** (see p. 67)

#### **Chronic Aspergillus Pneumonia**

**Clinical Presentation:** Occurs in patients with AIDS, chronic granulomatous disease, alcoholism, diabetes, and those receiving steroids for chronic pulmonary disease. Usual features include chronic productive cough  $\pm$  hemoptysis, low-grade fever, weight loss, and malaise. Chronic *Aspergillus* pneumonia resembles TB, histoplasmosis, melioidosis.

**Diagnostic Considerations:** Diagnosis by lung biopsy demonstrating septate hyphae invading lung parenchyma. *Aspergillus* may be in sputum, but is not diagnostic of aspergillus pneumonia.  $\beta$  1,3 D-glucan +, aspergillus galactomannan +.

**Pitfalls:** May extend into chest wall, vertebral column, or brachial plexus.

**Prognosis:** Related to severity/extent of infection. Cavitation is a favorable prognostic sign.

#### **Pulmonary Sporotrichosis (*Sporothrix schenckii*)**

**Clinical Presentation:** Usually presents as productive cough, low-grade fever, and weight loss. Chest x-ray shows cavitary thin-walled lesions with associated infiltrate. Hemoptysis is unusual. Differential diagnosis includes other thin-walled cavitary lung lesions (e.g., coccidioidomycosis, atypical TB, paragonimiasis).

**Diagnostic Considerations:** Diagnosis by lung biopsy demonstrating invasive lung disease, not broncho-alveolar lavage. Usually a history of puncture/traumatic wound involving an extremity. May be associated with septic arthritis/osteomyelitis.

**Pitfalls:** Sporotrichosis in lungs implies disseminated disease. May need repeated attempts at culture.

**Prognosis:** Related to extent of infection/degree of immunosuppression.

#### **Pulmonary Coccidioidomycosis (*Coccidioides immitis*)**

**Clinical Presentation:** Usually presents as a solitary, peripheral, thin-walled cavitary lesion in early or later stage of primary infection. May present as a solitary pulmonary nodule. E. nodosum and bilateral hilar adenopathy are common (in contrast to sporotrichosis). Hemoptysis is unusual.

**Diagnostic Considerations:** Diagnosis by demonstration of spherules with endospores in sputum/BAL or biopsy specimens/Coccidioides serology. Increased incidence of dissemination in Filipinos, Blacks, and American Indians. Eosinophils in CSF in disseminated infection with CNS involvement. May be associated with chronic meningitis/osteomyelitis. CF IgG titer  $\geq$  1:32 diagnostic of active disease. Coccidioides CF may XR (false +) with Blastomyces, Histoplasma, Cryptococci, Torulopsis, or TB (XR minimal with EIA). (False + histoplasma urinary antigen may provide a clue to the presence of coccidiomycosis.)

**Pitfalls:** Dissemination is preceded by  $\downarrow$  Coccidioides titers/disappearance of Erythema nodosum.

**Prognosis:** Related to extent of infection/degree of immunosuppression.

#### **Pulmonary Nocardiosis (*Nocardia asteroides*)**

**Clinical Presentation:** Usually presents as a dense lower lobe lung mass  $\pm$  cavitation. May have associated mass lesions in CNS. Chest wall sinuses are more common with Actinomycosis.

**Diagnostic Considerations:** Diagnosis by demonstrating organisms by stain/culture of lung specimens. Nocardia are weakly acid-fast and aerobic.

**Pitfalls:** Use IV regimens in critically ill patients. HIV patients require life-long suppressive therapy with TMP-SMX or minocycline.

**Prognosis:** Related to extent of infection/degree of immunosuppression.

### **Pulmonary Cryptococcosis (Cryptococcus neoformans)**

**Clinical Presentation:** Individual focus of infection is usually inapparent/minimal when patient presents with disseminated cryptococcal infection. Pneumonia is typically a minor part of disseminated disease; CNS manifestations usually predominate (e.g., headache, subtle cognitive changes, occasional meningeal signs, focal neurological deficits).

**Diagnostic Considerations:** Diagnosis by serum cryptococcal antigen, by demonstrating encapsulated yeasts in sputum/lung specimens, or by culture of pulmonary specimens.  $\beta$  1,3 D-glucan – , aspergillus galactomannan – .

**Pitfalls:** Clinical presentation of isolated cryptococcal pneumonia is rare. In HIV, false + cryptococcal antigen may occur with BBL Port-A-Cul transport vials. Patients require life-long suppressive therapy with fluconazole.

**Prognosis:** Related to extent of dissemination/degree of immunosuppression.

### **Pulmonary Zygomycosis (Mucormycosis) (Rhizopus/Mucor/Absidia/Cunninghamella)**

**Clinical Presentation:** Progressive pneumonia with fever, dyspnea, hemoptysis and cough unresponsive to antibiotic therapy. Usually seen only in compromised hosts. Chest x-ray is not characteristic, but shows infiltrate with consolidation in > 50% of patients. Cavitation occurs in 40% as neutropenia resolves.

**Diagnostic Considerations:** Diagnosis by demonstrating branched non-septate “ribbon-like” hyphae with right angle branching in biopsy specimens in lung biopsy. Pleural effusion is not a feature of pulmonary mucormycosis.  $\beta$  1,3 D-glucan – , aspergillus galactomannan – .

**Pitfalls:** Causes rhinocerebral mucormycosis in diabetics, pneumonia in leukopenic compromised hosts.

**Prognosis:** Related to degree of immunosuppression and underlying disease. Angioinvasive with a propensity for dissemination.

### **Pulmonary Pseudallescheriasis (P. boydii/S. apiospermum)**

**Clinical Presentation:** Progressive pulmonary infiltrates indistinguishable from Aspergillosis or Mucor. Usually seen only in compromised hosts (e.g., prolonged neutropenia, high-dose steroids, bone marrow or solid organ transplants, AIDS). Manifests as cough, fever, pleuritic pain, and often hemoptysis. No characteristic chest x-ray appearance.

**Diagnostic Considerations:** Diagnosis by culture of organisms in lung biopsy. Hemoptysis is common in patients with cavitory lesions. CNS involvement is rare.

**Pitfalls:** One of few invasive fungi unresponsive to amphotericin B deoxycholate. Cause of sinusitis in diabetics, and pneumonia in leukopenic compromised hosts.

**Prognosis:** Related to severity/extent of infection and degree of immunosuppression. Cavitory lesions causing hemoptysis often require surgical excision. Disseminated infection is often fatal.

## Parasites, Fungi, Unusual Organisms in Heart

### Chagas' Disease (*Trypanosoma brucei cruzi*) American Trypanosomiasis

Subset	Pathogens	Preferred Therapy	Alternate Therapy
Chagas' disease (American trypanosomiasis)	<i>Trypanosoma brucei cruzi</i>	Benznidazole 2.5–3.5 mg/kg (PO) q12h × 60 days	Nifurtimox 2–3 mg/kg/day (PO) q6–8h × 90 days

## Parasites, Fungi, Unusual Organisms in the Liver

### Liver Flukes

Subset	Pathogens	Preferred Therapy	Alternate Therapy
Fascioliasis	<i>Fasciola hepatica</i> <i>Fasciola gigantica</i>	Tridabendazole 10 mg/kg (PO) × 1 or 2 doses	Bithionol 30–50 mg/kg (PO) q48h × 20–30 days <b>or</b> Nitazoxanide 500 mg (PO) q12h × 7 days
Clonorchiasis/ Opisthorchiasis	<i>Clonorchis sinensis</i> <i>Opisthorchis viverrini</i>	Praziquantel 25 mg/kg (PO) q8h × 2 days	<u><i>C. sinensis</i>/<i>O. viverrini</i></u> Albendazole 10 mg/kg (PO) q24h × 7 days <u><i>O. viverrini</i></u> ; Tribendimidine 400 mg (PO) × 1 dose

### Hepatic Fascioliasis (*F. hepatica*/*F. gigantica*)

**Clinical Presentation:** Frequently asymptomatic, but may present acutely with fever, right upper quadrant pain, nausea, diarrhea, wheezing, urticaria, hepatomegaly, eosinophilia, anemia. ELISA, IHA, CF, CIE serology helpful with acute diagnosis. Aspiration of biliary fluid may show flukes/eggs. Chronic disease is associated with gallstones, cholecystitis, cholangitis, liver abscess, generalized adenopathy. Subacute nodules, hydrocele, lung/brain abscess can be seen in ectopic forms.

**Diagnostic Consideration:** Diagnosis by *F. hepatica*/*F. gigantica* eggs in stool. Endemic in sheep-raising areas (sheep liver flukes). Acquired from freshwater plants (watercress). Not associated with cholangiocarcinoma.

**Pitfalls:** May present as Katayama syndrome resembling schistosomiasis, with high fever, eosinophilia, and hepatosplenomegaly. Unlike other trematodes, praziquantel is ineffective. Chronic *F. hepatica* often asymptomatic, but may present with intermittent biliary colic. No eosinophilia. Abdominal ultrasound/CT shows crescentic/leaf shaped defects in gallbladder/hepatic ducts.

**Prognosis:** Related to extent/location of liver damage.



### **Hepatic Clonorchiasis (*C. sinensis*)/Opisthorchiasis (*O. viverrini*)**

**Clinical Presentation:** Frequently asymptomatic, but may present 2–4 weeks after ingestion of fluke with fever, tender hepatomegaly, rash, and eosinophilia. Chronically presents as recurrent cholangitis, chronic cholecystitis, or pancreatitis. Associated with cholangiocarcinoma (unlike fascioliasis).

**Diagnostic Considerations:** Diagnosis by visualizing *C. sinensis*/*O. viverrini* (operculated lemon-shaped) eggs in stool. Clonorchiasis is acquired from ingesting raw/inadequately cooked infected freshwater (Cyprinoid) fish in Southeast Asia. Opisthorchiasis is acquired from ingesting raw/inadequately cooked infected freshwater fish/crayfish from Laos, Cambodia, or Thailand.

**Pitfalls:** Cholecystitis with eosinophilia should suggest clonorchiasis or ascariasis.

**Prognosis:** Related to extent/location of hepatic damage. Associated with cholangiocarcinoma.

### **Cystic Masses in Liver**

Subset	Pathogens	Preferred Therapy	Alternate Therapy
Hepatic amebiasis	Entamoeba histolytica	Metronidazole 750 mg (PO) q8h × 7–10 days	Tinidazole 2 gm/day (PO) in 3 divided doses × 3 days
Hepatic echinococcosis (hydatid cyst disease)	Echinococcus granulosus	<u>Operable</u> Surgical resection <b>plus</b> Albendazole 400 mg* (PO) q12h × 1–6 months	<u>Inoperable</u> Albendazole 400 mg* (PO) q12h × 1–6 months

\* If < 60 kg, give albendazole 7.5 mg/kg.

### **Hepatic Amebiasis (*Entamoeba histolytica*)**

**Clinical Presentation:** Presents insidiously with weight loss and night sweats or acutely ill with fever, nausea, vomiting, right upper quadrant pain. Typically, amebic liver abscesses are single, affect the right posterior lobe of liver, and do not show air/fluid levels. (In contrast, bacterial liver abscesses are usually multiple, and in all lobes of liver, and often show air/fluid levels.) Amebic liver abscesses do not calcify like hydatid cysts.

**Diagnostic Considerations:** Diagnosis by *E. histolytica* serology/*E. histolytica* in abscess wall. Worldwide distribution. Acquired by ingesting amebic cysts. Amebic liver abscess usually presents years after initial mild amebic dysenteric episode.

**Pitfalls:** Amebic abscess fluid (“anchovy paste”) contains no PMNs or amebas; amebas are found only in abscess wall. Eosinophilia is not a feature of amebiasis.

**Prognosis:** Related to health of host/extrapulmonary spread.

### **Hepatic Echinococcosis (*Echinococcus granulosus*) Hydatid Cyst Disease**

**Clinical Presentation:** Right upper quadrant pain/mass when cysts enlarge enough to cause symptoms. Hepatic cysts are unilocular in 70%, multilocular in 30%.

**Diagnostic Considerations:** Diagnosis by demonstrating *E. granulosus* scolices/hooklets in cyst/hydatid sand. Serology is unreliable. Worldwide distribution in sheep/cattle raising areas. Acquired by ingestion of eggs from dogs.

**Pitfalls:** Eosinophilia not a feature of hydatid cyst disease. Hydatid cysts are multifaceted, loculated, and calcified.

**Prognosis:** Related to location/extent of extrahepatic cysts. Large cysts are best treated by surgical removal after injection with hypertonic saline, alcohol, or iodophor to kill germinal layer/daughter cysts. Percutaneous drainage under ultrasound guidance plus albendazole may be effective.

## Hepatomegaly

Subset	Pathogens	Preferred Therapy	Alternate Therapy
Visceral leishmaniasis (Kala-azar)	<i>Leishmania donovani</i>	Antimony stibogluconate 20 mg/kg (IV/IM) q24h × 28 days <b>or</b> Lipid amphotericin (Immunocompetent adults: 3 mg/kg (IV) q24h on days 1–5, 14, and 21; Immunocompromised adults: 4 mg/kg (IV) q24h on days 1–5, 10, 17, 24, 31, and 38)*	Miltefosine 2.5 mg/kg* (PO) q24h × 28 days <b>or</b> Paromomycin 15 mg/kg (IM) q24h × 21 days
Indian (Bihar state) visceral leishmaniasis (Kala-azar)	<i>L. donovani</i>	AmBisome (L-Amb) <sup>†</sup> 5 mg/kg (IV) × 1 dose, then Miltefosine 50 mg (PO with food) q12h × 7–14 days	
Schistosomiasis	<i>Schistosoma mansoni</i>	Praziquantel 20 mg/kg (PO) q12h × 2 doses	Oxamniquine 15 mg/kg (PO) × 1 dose <sup>‡</sup>
	<i>Schistosoma japonicum</i>	Praziquantel 20 mg/kg (PO) q8h × 3 doses	None

\* Maximum 150 mg (PO)/day.

† Ultra short course therapy with AmBisome is 10 mg/kg (IV) × 2 days.

‡ In Africa, give 20 mg/kg (PO) q24h × 3 days.

### **Visceral Leishmaniasis (*Leishmania donovani*) Kala-azar**

**Clinical Presentation:** Subacute or chronic systemic cases manifest months to years after initial exposure to *Leishmania*, most often with fever, weight loss, anemia, hepatosplenomegaly ± generalized adenopathy. Long eyelashes in some. Laboratory abnormalities include leukopenia, anemia, and polyclonal gammopathy on SPEP. Incubation period is usually 3–8 months. May have atypical presentation in HIV (e.g., no splenomegaly). Acutely can mimic malaria with chills/temperature spikes. Post-kala-azar dermatitis may resemble leprosy, and is persistent/common on face.

**Diagnostic Considerations:** Double quotidian fever (double daily temperature spike) in persons from endemic areas with hepatosplenomegaly suggests the diagnosis. Kala-azar in children may have ↑ triglycerides/↓ HDL. Diagnosis by liver/bone marrow or buffy coat smear. Histologically, Kala-azar resembles histoplasmosis. Key diagnostic finding in tissue biopsy specimens is intracellular amastigotes with adjacent rod-shaped kinetoplasts. *L. donovani* serology is specific; immunochromatographic Anti-K39 strip test is also useful. Most common in Southern Europe, Middle East, Asia, Africa, South America. Facial lesion is a clue to the diagnosis.

**Pitfalls:** In acute cases, can mimic malaria with chills and temperature spikes, but no thrombocytopenia or atypical lymphocytes. Antimony resistance is common in India (Bihar state).

**Prognosis:** Related to degree of liver/spleen involvement.

**Hepatic Schistosomiasis (*Schistosoma mansoni*/*japonicum*)**

**Clinical Presentation:** May present acutely with Katayama fever (serum sickness-like illness with wheezing and eosinophilia) 4–8 weeks after exposure. May be accompanied or followed by fever/chills, headache, cough, abdominal pain, diarrhea, generalized lymphadenopathy, or hepatosplenomegaly. Laboratory abnormalities include leukocytosis, eosinophilia, and polyclonal gammopathy on SPEP. Resolves spontaneously after 2–4 weeks. After 10–15 years, may present chronically as hepatosplenic schistosomiasis, with pre-sinusoidal portal hypertension, hepatomegaly (L > R lobe enlargement), no jaundice, and intact liver function.

**Diagnostic Considerations:** Diagnosis by *S. mansoni*/*S. japonicum* eggs in stool/liver biopsy. Serology is good for acute (not chronic) schistosomiasis. CT/MRI of liver shows "turtle back" septal calcifications. Rare complications include cor pulmonale and protein-losing enteropathy. Increased incidence of hepatitis B/C and chronic Salmonella infections. Renal complications include glomerulonephritis and nephrotic syndrome.

**Pitfalls:** Chronic schistosomiasis is not associated with eosinophilia. *S. hematobium* does not infect the liver/spleen. Oxamniquine is contraindicated in pregnancy.

**Prognosis:** Related to egg burden.

## Parasites, Fungi, Unusual Organisms in Stool/Intestines

**Intestinal Protozoa**

Subset	Pathogens	Preferred Therapy	Alternate Therapy
Amebiasis	<i>E. histolytica</i>	See p. 88	
Giardiasis	<i>Giardia lamblia</i>	Tinidazole 2 gm (PO) × 1 dose <b>or</b> Metronidazole 250 mg (PO) q8h × 5–7 days	
Isospora	<i>Isospora belli</i>	TMP-SMX 1 DS tablet (PO) q12h × 10 days <b>or</b> Nitazoxanide 500 mg (PO) × 3 days (for HIV, see p. 334)	Ciprofloxacin 500 mg (PO) q12h × 7 days <b>or</b> Pyrimethamine 75 mg (PO) q24h + folinic acid 10 mg (PO) q24h × 2 weeks then q12h × 3 weeks
Dientamoebiasis	<i>Dientamoeba fragilis</i>	Iodoquinol 650 mg (PO) q8h × 20 days	Doxycycline 100 mg (PO) q12h × 10 days <b>or</b> Metronidazole 500–750 mg (PO) q8h × 10 days
Blastocystis	<i>Blastocystis hominis</i>	Nitazoxanide 500 mg (PO) q12h × 3 days	Iodoquinol 650 mg (PO) q8h × 20 days <b>or</b> TMP-SMX 1 DS tablet (PO) q12h × 7 days

**Intestinal Protozoa (cont'd)**

Subset	Pathogens	Preferred Therapy	Alternate Therapy
Cyclospora	Cyclospora	TMP-SMX 1 DS (PO) q12h × 10 days <b>or</b> Nitazoxanide 500 mg (PO) q12h × 3 days (for HIV, see p. 335)	
Cryptosporidiosis	Cryptosporidia	Nitazoxanide 500 mg (PO) q12h × 3 days (for HIV, see p. 334)	
Balantidiasis	Balantidium coli	Doxycycline 100 mg (PO) q12h × 10 days	Iodoquinol 650 mg (PO) q8h × 20 days <b>or</b> Metronidazole 750 mg (PO) q8h × 5 days

**Amebiasis (*Entamoeba histolytica*)** (see p. 91)**Giardiasis (*Giardia lamblia*)** (see p. 90)**Isosporiasis (*Isospora belli*)**

**Clinical Presentation:** Acute/subacute onset of diarrhea. *Isospora belli* is the only protozoa to cause diarrhea with eosinophils in stool.

**Diagnostic Considerations:** Diagnosis by demonstrating cysts in stool and organisms in intestinal biopsy specimen. Associated with HIV, immigration from Latin America, daycare centers, and mental institutions. If stool exam is negative, "string test"/duodenal aspirate and biopsy may be helpful.

**Pitfalls:** Difficult to eradicate; may last months. Multiple stool samples may be needed for diagnosis. Add folinic acid 10 mg (PO) q24 if pyrimethamine is used.

**Prognosis:** Related to adequacy of treatment/degree of immunosuppression.

**Dientamoebiasis (*Dientamoeba fragilis*)**

**Clinical Presentation:** Acute/subacute onset of diarrhea. No cyst stage. Lives only as trophozoite. Among intestinal protozoa, only *D. fragilis* and *I. belli* are associated with peripheral eosinophilia.

**Diagnostic Considerations:** Diagnosis by demonstrating trophozoites in stool/intestinal by trichrome stain. Mucus in diarrheal stools, not blood. May have abdominal pain. Diarrhea may last for months/years.

**Pitfalls:** Frequently associated with pinworm (*Enterobius vermicularis*) infection. Metronidazole failures common.

**Prognosis:** Related to adequacy of fluid replacement/underlying health of host.

**Blastocystis (*Blastocystis hominis*)**

**Clinical Presentation:** Acute/subacute onset of diarrhea.

**Diagnostic Considerations:** Diagnosis by demonstrating cysts and trophozoites in stool by trichrome stain. Trichrome stain reveals characteristic "halo" (slime capsule) in stool specimens.

**Pitfalls:** Uncommon GI pathogen. Consider as cause of diarrhea only after other pathogens excluded.

**Prognosis:** Related to adequacy of fluid replacement/underlying health of host.

**Cyclospora** (see p. 90; for HIV, see p. 335)**Cryptosporidiosis** (see p. 90; for HIV, see p. 334)

**Balantidiasis (Balantidium coli)**

**Clinical Presentation:** Acute/subacute onset of diarrhea. Fecal WBCs only with mucosal invasion. Largest intestinal protozoa and only ciliated protozoa to infect humans.

**Diagnostic Considerations:** Diagnosis by demonstrating organism in stool/intestinal biopsy specimen. Identifying features include darkly staining “kidney shaped” nucleus and large size. Fulminant dysentery seen only in debilitated/compromised hosts.

**Pitfalls:** Stools not bloody. Diarrhea may be intermittent.

**Prognosis:** Related to adequacy of fluid replacement/underlying health of host.

**Intestinal Nematodes (Roundworms)**

Subset	Pathogens	Preferred Therapy	Alternate Therapy
Capillariasis	Capillaria (Aonchotheca) philippinensis	Mebendazole 200 mg (PO) q12h × 20 days	Albendazole 400 mg (PO) q24h × 10 days
Angiostrongyliasis (rodent lung/intestinal worm)	Angiostrongylus cantonensis	Albendazole 400 mg (PO) q12h × 2–3 weeks <b>plus</b> corticosteroids	Mebendazole 200–400 mg (PO) q8h × 10 days
Hookworm	Necator americanus/ Ancylostoma duodenale	Albendazole 400 mg (PO) × 1 dose <b>or</b> Mebendazole 100 mg (PO) q12h × 3 days or 500 mg (PO) × 1 dose	Pyrantel pamoate 11 mg/kg (PO) q24h × 3 days (max. 1 gm/day)
Strongyloidiasis	Strongyloides stercoralis	Ivermectin 200 mcg/kg (PO) q24h × 2 days <b>or</b> Thiabendazole 25 mg/kg (PO) q12h × 2 days (max. 3 gm/day)	Albendazole 400 mg (PO) q24h × 3 days
Ascariasis	Ascaris lumbricoides	Albendazole 400 mg (PO) × 1 dose <b>plus</b> Pyrantel pamoate 11 mg/kg (PO) q24h × 3 days (max. 1 gm/day) <b>or</b> Mebendazole 100 mg (PO) q12h × 3 days or 500 mg (PO) × 1 dose	Ivermectin 200 mcg/kg (PO) × 1 dose*
Trichostrongyliasis	Trichostrongylus orientalis	Pyrantel pamoate 11 mg/kg (PO) × 1 dose (max. 1 gm)	Albendazole 400 mg (PO) × 1 dose <b>or</b> Mebendazole 100 mg (PO) q12h × 3 days

\* Avoid in young children and pregnant adults.

**Intestinal Nematodes (Roundworms) (cont'd)**

Subset	Pathogens	Preferred Therapy	Alternate Therapy
Pinworm	Enterobius vermicularis	Pyrantel pamoate 11 mg/kg (PO) × 1 dose (max. 1 gm); repeat in 2 weeks <b>or</b> Albendazole 400 mg (PO) × 1 dose; repeat in 2 weeks	Ivermectin 200 mcg/kg (PO) × 1 dose <b>or</b> Mebendazole 100 mg (PO) × 1 dose; repeat in 2 weeks
Whipworm	Trichuris trichiura	Mebendazole 100 mg (PO) q12h × 3–7 days	Albendazole 400 mg (PO) q24h × 3 days <b>or</b> Ivermectin 200 mcg/kg (PO) q24h × 3 days

**Capillariasis (Capillaria philippinensis)**

**Clinical Presentation:** Intermittent voluminous watery diarrhea ± malabsorption. Fever is uncommon.

**Diagnostic Considerations:** Diagnosis by demonstrating ova or parasite in stools. Resembles Trichuris, but *C. philippinensis* ova are larger and have a “pitted shell” with prominent polar plugs. Peripheral eosinophilia is uncommon until after therapy.

**Pitfalls:** Serology is positive in 85%, but cross-reacts with other parasites.

**Prognosis:** Related to severity of malabsorption/extra-intestinal disease.

**Angiostrongyliasis (A. cantonensis) Rodent Lung/Intestinal Worm**

**Clinical Presentation:** Presents as appendicitis (worm resides and deposits eggs in arteries/arterioles around ileocecum/appendix).

**Diagnostic Considerations:** Diagnosis by demonstrating organism in biopsied/excised tissue. May involve proximal small bowel, liver, CNS. When in CNS, a cause of eosinophilic meningitis.

**Pitfalls:** Can present as RLQ mass/fever resembling regional enteritis (Crohn's disease), but with eosinophilia and leukocytosis.

**Prognosis:** Related to severity of malabsorption and extra-intestinal disease.

**Hookworm (Necator americanus/Ancylostoma duodenale)**

**Clinical Presentation:** Pruritic, vesicular eruptions at site of filariform larval entry (“ground itch”). Pulmonary symptoms and transient eosinophilia may occur during migratory phase to intestines. Later, abdominal pain, diarrhea, weight loss, hypoalbuminemia, and anemia develop.

**Diagnostic Considerations:** Diagnosis by demonstrating eggs in stools. Larvae rare in fresh stool specimens (eggs may hatch if stools allowed to stand too long prior to examination). *N. americanus* can ingest 0.3 ml of blood/worm/day, much greater than *A. duodenale*. Anemia may be severe with heavy infestation (up to 100 mL/day).

**Pitfalls:** Eggs in fresh stool, not rhabditiform larvae. Eggs in hyperinfection syndrome.

**Prognosis:** Related to severity of anemia/malabsorption.

**Strongyloidiasis (Strongyloides stercoralis)**

**Clinical Presentation:** Pruritic, papular, erythematous rash. Pulmonary symptoms (cough, asthma) may occur during lung migration phase. May develop Loeffler's syndrome (pulmonary infiltrates with eosinophilia) or ARDS in heavy infections. Intestinal phase associated with colicky abdominal pain, diarrhea, and malabsorption.

**Diagnostic Considerations:** Diagnosis by demonstrating larvae in stool specimens/duodenal fluid. Usually asymptomatic in normal hosts, but causes “hyperinfection syndrome” in compromised hosts. CNS strongyloides (part of hyperinfection syndrome) should suggest diagnosis of HIV in non-immunosuppressed patients. Diarrhea/abdominal pain mimics regional enteritis (Crohn’s disease) or ulcerative colitis. Malabsorption is common and mimics tropical sprue. Anemia is usually mild.

**Pitfalls:** Usually rhabditiform larvae (not eggs) in stools.

**Prognosis:** Related to severity of malabsorption.

### **Ascariasis (*Ascaris lumbricoides*)**

**Clinical Presentation:** Pulmonary symptoms (cough, asthma) may occur during larval lung migration phase. May develop Loeffler’s syndrome (pulmonary infiltrates with eosinophilia), as with hookworm/*Strongyloides*. Intestinal symptoms develop late. Usually asymptomatic unless intestinal/biliary obstruction occurs. Can obstruct the appendix/pancreatic duct.

**Diagnostic Considerations:** Diagnosis by demonstrating eggs in stool specimens. Abdominal ultrasound can detect obstruction from adult worms. Most infections are asymptomatic; symptoms are related to “worm burden”/ectopic migration. Each female worm may produce up to 250,000 eggs/day.

**Pitfalls:** Lung involvement (bronchospasm, bronchopneumonia, lung abscess) is prominent in HIV.

**Prognosis:** Related to worm burden/extra-intestinal organ invasion.

### **Trichostrongyliasis (*Trichostrongylus orientalis*)**

**Clinical Presentation:** Mild intestinal symptoms with persistent eosinophilia.

**Diagnostic Considerations:** Diagnosis by demonstrating eggs in stool specimens. Most prevalent in the Middle East and Asia. Mild anemia. Eosinophilia is usually > 10%.

**Pitfalls:** Must differentiate eggs from hookworm, and rhabditiform larvae from *Strongyloides*. *T. orientalis* eggs have “pointed ends.”

**Prognosis:** Related to extent of disease/underlying health of host.

### **Pinworm (*Enterobius vermicularis*)**

**Clinical Presentation:** Primarily affects children. Perianal pruritus (ectopic migration) is the main symptom. Worm lives in the cecum, but patients do not have intestinal symptoms.

**Diagnostic Considerations:** Scotch tape of anus at night can be used to detect eggs left by migrating female worms (Scotch tape test).

**Pitfalls:** Abdominal pain and diarrhea should prompt search for *Dientamoeba fragilis*, since co-infection is common.

**Prognosis:** Excellent.

### **Whipworm (*Trichuris trichiura*)**

**Clinical Presentation:** May present as “chronic appendicitis.” Severe infestation may cause bloody diarrhea/abdominal pain (“*Trichuris* dysentery syndrome”), rectal prolapse.

**Diagnostic Considerations:** Diagnosis by demonstrating large eggs with bile-stained, triple-layered eggshell walls and doubly operculated transparent plugs. Most patients are asymptomatic or mildly anemic.

**Pitfalls:** Commonly co-exists with *Ascaris*, hookworm, or *E. histolytica*.

**Prognosis:** Related to severity/extent of dysentery. May need retreatment if heavy infection.

## Intestinal Cestodes (Tapeworms)

Subset	Pathogens	Preferred Therapy	Alternate Therapy
Beef tapeworm	<i>Taenia saginata</i>	Praziquantel 5–10 mg/kg (PO) × 1 dose	Niclosamide 2 gm (PO) × 1 dose <b>or</b> Nitazoxanide 500 mg (PO) q12h × 3 days
Pork tapeworm	<i>Taenia solium</i>	Praziquantel 5–10 mg/kg (PO) × 1 dose	Niclosamide 2 gm (PO) × 1 dose <b>or</b> Nitazoxanide 500 mg (PO) q12h × 3 days
Dwarf tapeworm	<i>Hymenolepis nana</i>	Praziquantel 25 mg/kg (PO) × 1 dose	Nitazoxanide 500 mg (PO) q12h × 3 days <b>or</b> Niclosamide 2 gm (PO) × 1 dose, then 1.5 gm q24h × 6 days
Fish tapeworm	<i>Diphyllobothrium latum</i>	Praziquantel 10 mg/kg (PO) × 1 dose	Niclosamide 2 gm (PO) × 1 dose <b>or</b> Nitazoxanide 500 mg (PO) q12h × 3 days

### **Beef Tapeworm (*Taenia saginata*)/Pork Tapeworm (*Taenia solium*)**

**Clinical Presentation:** Usually mild symptoms (weight loss, anemia), since most infections are caused by a single tapeworm. Worm segments passed between BMs with *T. saginata* and with BMs with *T. solium*.

**Diagnostic Considerations:** Diagnosis by demonstrating proglottids in stool. *Taenia* eggs in stool cannot be speciated; all are brown and spherical with a radially-striated inner shell. *T. saginata* may survive for 10 years, *T. solium* for 25 years.

**Pitfalls:** Severe cases may cause appendicitis, intestinal obstruction/perforation.

**Prognosis:** Related to severity of malabsorption/intestinal obstruction.

### **Dwarf Tapeworm (*Hymenolepis nana*)**

**Clinical Presentation:** Usually asymptomatic.

**Diagnostic Considerations:** Diagnosis by demonstrating typical eggs in stool, with two shells enclosing inner oncosphere with hooklets. ELISA is positive in 85%, but cross-reacts with *Taenia*/Cysticercosis. GI symptoms develop with stool egg counts > 15,000/gm.

**Pitfalls:** Abdominal pain and diarrhea in heavy infestations.

**Prognosis:** Related to severity of malabsorption/underlying health of host.

### **Fish Tapeworm (*Diphyllobothrium latum*)**

**Clinical Presentation:** Symptoms secondary to macrocytic anemia from vitamin B<sub>12</sub> deficiency. Most infestations are asymptomatic.

**Diagnostic Considerations:** Diagnosis by demonstrating operculated eggs and proglottids in stool.



**Pitfalls:** Vitamin B<sub>12</sub> deficiency anemia requires prolonged infection (> 3 years).

**Prognosis:** Related to severity of B<sub>12</sub> deficiency anemia/underlying health of host.

## Intestinal Trematodes (Flukes/Flatworms)

Subset	Pathogens	Preferred Therapy
Fasciolopsiasis	Fasciolopsis buski	Praziquantel 25 mg/kg (PO) q8h × 3 doses
Heterophyiasis	Heterophyes heterophyes Metagonimus yokogawai	Praziquantel 25 mg/kg (PO) q8h × 3 doses

### Fasciolopsiasis (Fasciolopsis buski)

**Clinical Presentation:** Diarrhea with copious mucus in stool. Most cases are asymptomatic.

**Diagnostic Considerations:** Diagnosis by demonstrating eggs in stool. May have eosinophilia, low-grade fever ± malabsorption. Intestinal obstruction is the most serious complication.

**Pitfalls:** Mimics peptic ulcer disease with upper abdominal pain relieved by food.

**Prognosis:** Related to severity/extent of malabsorption/intestinal obstruction.

### Heterophyiasis (Heterophyes heterophyes/Metagonimus yokogawai)

**Clinical Presentation:** Usually asymptomatic or mild intestinal symptoms. Embolization of eggs may result in myocarditis, myocardial fibrosis, or cerebral hemorrhage. Eosinophilia may be present.

**Diagnostic Considerations:** Diagnosis by demonstrating large operculated eggs in stool. Small intestine fluke.

**Pitfalls:** Difficult to differentiate from clonorchis sinensis.

**Prognosis:** Related to extent/severity of extra-intestinal dissemination to heart, lungs, CNS.

## Other Intestinal Infections

Subset	Pathogen	Preferred Therapy
Whipple's disease	Tropheryma whippelii	Ceftriaxone 2 gm (IV) q24h × 2 weeks* or Ceftriaxone 1 gm (IV) q24h × 2 weeks plus Streptomycin 1 gm (IM) q24h × 2 weeks*

\* Follow with TMP-SMX 1 DS tablet (PO) q12h × 1 year or Doxycycline 100 mg (PO) q12h × 1 year.

### Whipple's Disease (Tropheryma whippelii)

**Clinical Presentation:** Diarrhea, fever, encephalopathy/dementia, weight loss, polyarthritis, myocarditis, pericarditis, general lymphadenopathy ± malabsorption.

**Diagnostic Considerations:** Diagnosis by demonstrating organism by PAS staining macrophages in small bowel biopsy specimens.

**Pitfalls:** May present with dementia mimicking Alzheimer's disease, or FUO mimicking celiac disease or lymphoma. Optimum length of treatment is unknown. Relapses occur.

**Prognosis:** Related to severity/extent of extra-intestinal disease.

## Parasites, Fungi, Unusual Organisms in Skin/Muscle

### Infiltrative Skin/Subcutaneous Lesions

Subset	Pathogens	Preferred Therapy
Cutaneous leishmaniasis <i>Old World</i>	Leishmania major Leishmania tropica	Sodium stibogluconate 20 mg/kg (IV or IM) q24h × 20 days  <b>or</b> Pentamidine 2–3 mg/kg (IM) q48h × 4–7 days  <b>or</b>
	Leishmania mexicana Leishmania braziliensis	Lipid amphotericin 3 mg/kg (IV) q24h on days 1–5, 14, and 21  <b>or</b> Miltefosine 2.5 mg/kg (PO) q24h × 28 days (max. 150 mg/day)
Leprosy <i>Lepromatous</i>	Mycobacterium leprae	Dapsone 100 mg (PO) q24h × 1–2 years <b>plus</b> Clofazimine 50 mg (PO) q24h × 1–2 years <b>plus</b> Rifampin 600 mg (PO) monthly × 1–2 years
	<i>Non-lepromatous (tuberculoid)</i>	Mycobacterium leprae Dapsone 100 mg (PO) q24h × 6 months <b>plus</b> Rifampin 600 mg (PO) monthly × 6 months
Loa Loa (Loiasis)	L. loa	Diethylcarbamazine 2–3 mg/kg (PO) q8h × 2–3 weeks  <b>or</b> Albendazole 200 mg (PO) q12h × 3 weeks
Erythrasma	Corynebacterium minutissimum	Erythromycin 250 mg (PO) q6h × 2 weeks

#### **Cutaneous Leishmaniasis (Old World/New World)**

**Clinical Presentation:** Variable presentation. Typically, a nodule develops then ulcerates, with a raised/erythematous outer border and a central area of granulation tissue. May be single or multiple. Usually non-pruritic/non-painful. Occurs weeks after travel to endemic areas (New World leishmaniasis: Latin America; Old World leishmaniasis: Central Asia).

**Diagnostic Considerations:** Diagnosis by demonstrating Leishmania amastigotes in biopsy specimen.

**Pitfalls:** Most skin lesions undergo spontaneous resolution. However, treatment is advisable for lesions caused by *L. braziliensis* or related species causing mucosal leishmaniasis.

**Prognosis:** Excellent.

#### **Lepromatous Leprosy (Mycobacterium leprae)**

**Clinical Presentation:** Diffuse, symmetrical, red or brown skin lesions presenting as macules, papules, plaques, or nodules. May also present as diffuse thickening of skin, especially involving ear lobes, face, and extremities. Loss of eyebrows/body hair may occur.

**Diagnostic Considerations:** Diagnosis by demonstrating organism in acid fast stain of tissue specimens. Afebrile bacteremia is frequent, and buffy coat smears positive for *M. leprae*. Erythema nodosum and polyclonal gammopathy on SPEP are common, and lepromin skin test/PPD are negative (anergic). When present, peripheral neuropathy is often symmetrical and acral in distribution.

**Pitfalls:** Differential diagnosis is large. Consider leprosy in patients with unexplained skin diseases.

**Prognosis:** Good if treated early.

### **Non-Lepromatous (Tuberculoid) Leprosy (*Mycobacterium leprae*)**

**Clinical Presentation:** Small number of asymmetrical, hypopigmented skin lesions, which are often scaly with sharp borders and associated anesthesia. Asymmetric peripheral nerve trunk involvement is common.

**Diagnostic Considerations:** Diagnosis by demonstrating granulomas with few acid-fast bacilli. Differentiate from cutaneous leishmaniasis by skin biopsy. Lepromin skin test/PPD are positive and SPEP is normal (in contrast to lepromatous leprosy).

**Pitfalls:** Wide spectrum of presentations depending on immune status and duration of disease. Differential diagnosis is large. Consider leprosy in patients with unexplained skin diseases.

**Prognosis:** Good if treated early.

### **Erythrasma (*Corynebacterium minutissimum*)**

**Clinical Presentation:** Reddened/raised skin lesions on face/trunk. Not hot or pruritic.

**Diagnostic Considerations:** Differentiated from *Tinea versicolor* by culture. *C. minutissimum* skin lesions fluoresce red under UV light.

**Pitfalls:** Resembles *Tinea versicolor*, but lesions primarily involve the face, not trunk.

**Prognosis:** Excellent.

## **Infiltrative Skin Lesions ± Ulcers/Sinus Tracts/Abscesses**

Subset	Pathogens	Preferred Therapy
Cutaneous histoplasmosis	<i>Histoplasma capsulatum</i>	Treat as for disseminated histoplasmosis (p. 267) if part of systematic syndrome. If solely involves the skin, Itraconazole 200 mg (PO) q12h until resolution.
Cutaneous blastomycosis	<i>Blastomyces dermatitidis</i>	Treat the same as pulmonary infection (p. 267)
Cutaneous coccidioidomycosis	<i>Coccidioides immitis</i>	Treat the same as pulmonary infection (p. 268)
Cutaneous actinomycosis	<i>Actinomyces israelii</i>	Treat the same as pulmonary infection (p. 268)
Cutaneous nocardiosis	<i>Nocardia</i> sp.	Treat the same as pulmonary infection (p. 268)
Cutaneous amebiasis	<i>Entamoeba histolytica</i>	Treat the same as pulmonary infection (p. 265)

**Infiltrative Skin Lesions ± Ulcers/Sinus Tracts/Abscesses**

(cont'd)

Subset	Pathogens	Preferred Therapy
Cutaneous mycobacteria <i>Scrofula</i>	Mycobacterium tuberculosis	Treat same as pulmonary TB. See p. 53
	M. scrofulaceum	Surgical excision is curative
<i>M. haemophilum</i>	M. haemophilum	Surgical excision/TMP-SMX 1 DS tablet (PO) q12h <b>or</b> Minocycline 100 mg (PO) q12h <b>or</b> quinolone (PO) until cured
<i>M. chelonae</i>	M. chelonae	Clarithromycin 500 mg (PO) q12h × 6 months
<i>M. fortuitum</i>	M. fortuitum	Minocycline 100 mg (PO) q12h × 6–12 months <b>or</b> Doxycycline 100 mg (PO) q12h × 6–12 months <b>plus</b> TMP-SMX 1 DS tablet (PO) q12h × 6–12 months
<i>Swimming pool granuloma</i>	M. marinum	Clarithromycin XL 1 gm (PO) q24h × 3 months <b>plus</b> Rifampin 600 mg (PO) q24h × 3 months <b>or</b> TMP-SMX 1 DS tablet (PO) q12h × 3 months <b>plus</b> Ethambutol 15 mg/kg (PO) q24h × 3 months <b>or</b> Doxycycline 100 mg (PO) q12h × 3 months
<i>Buruli ulcer</i>	M. ulcerans	TMP-SMX 1 DS tablet (PO) q12h <b>plus</b> Ethambutol 15 mg/kg (PO) q24h × 6 weeks
<i>Cutaneous MAI</i>	Mycobacterium avium-intracellulare	Ethambutol 15 mg/kg (PO) q24h <b>plus</b> Azithromycin 1200 mg (PO) weekly × 6 months

**Cutaneous Histoplasmosis (Histoplasma capsulatum)****Clinical Presentation:** Chronic, raised, verrucous lesions.**Diagnostic Considerations:** Diagnosis by demonstrating organism by culture/tissue staining.**Pitfalls:** Skin nodules represent disseminated histoplasmosis, not isolated skin infection. Look for histoplasmosis elsewhere (e.g., lung, liver, bone marrow). Skin nodules (± chronic pneumonia) with osteolytic bone lesions suggests. African histoplasmosis due to H. capsulation var. duboisii.**Prognosis:** Related to extent of infection/degree of immunosuppression.**Cutaneous Blastomyces (Blastomyces dermatitidis)****Clinical Presentation:** Painless, erythematous, well-circumscribed, hyperkeratotic, crusted nodules or plaques that enlarge over time. Some may ulcerate and leave an undermined edge.**Diagnostic Considerations:** Diagnosis by demonstrating organism by culture/tissue staining. Blastomyces dermatitidis affects many organs.

**Pitfalls:** When found in skin, look for *Blastomyces* elsewhere (e.g., lungs, prostate).

**Prognosis:** Related to extent of infection/degree of immunosuppression.

### **Cutaneous Coccidioidomycosis (*Coccidioides immitis*)**

**Clinical Presentation:** Skin lesions take many forms, including raised verrucous lesions, cold subcutaneous abscesses, indolent ulcers, or small papules.

**Diagnostic Considerations:** Diagnosis by demonstrating organism by culture/tissue staining.

**Pitfalls:** Skin nodules represent disseminated coccidioidomycosis, not isolated skin infection. Look for *Coccidioides* elsewhere (e.g., CNS, bone, lungs).

**Prognosis:** Related to extent of infection/degree of immunosuppression.

### **Cutaneous Actinomycosis (*Actinomyces israelii*)**

**Clinical Presentation:** Erythematous, uneven, indurated, woody, hard, cervicofacial tumor. Localized single/multiple sinus tracts in chest wall, abdominal wall, or inguinal/pelvic area may be present.

**Diagnostic Considerations:** Diagnosis by demonstrating organism by culture/tissue staining.

**Pitfalls:** Look for underlying bone involvement.

**Prognosis:** Good with early/prolonged treatment.

### **Cutaneous Nocardia (*Nocardia brasiliensis*)**

**Clinical Presentation:** Subcutaneous abscesses may rupture to form chronically draining fistulas.

**Diagnostic Considerations:** Diagnosis by demonstrating organism by culture/tissue staining. May present as "Madura foot."

**Pitfalls:** Look for underlying immunosuppressive disorder.

**Prognosis:** Related to extent of infection/degree of immunosuppression.

### **Cutaneous Amebiasis (*Entamoeba histolytica*)**

**Clinical Presentation:** Ulcers with ragged edges, sinus tracts, amebomas, and strictures may develop around the anus/rectum or abdominal wall.

**Diagnostic Considerations:** Diagnosis by demonstrating organism by tissue staining.

**Pitfalls:** If abdominal sinus tract is present, look for underlying ameboma and evidence of infection in other organs (e.g., CNS, lung, liver).

**Prognosis:** Related to extent of infection/degree of organ damage.

### **Scrofula (*Mycobacterium tuberculosis*)**

**Clinical Presentation:** Cold, chronic, anterior cervical adenopathy ± sinus tracts. Usually in children.

**Diagnostic Considerations:** Diagnosis by acid-fast stain and culture of node/drainage.

**Pitfalls:** Cured by antibiotic therapy alone. No need for surgical excision.

**Prognosis:** Excellent with treatment.

### **Scrofula (*Mycobacterium scrofulaceum*)**

**Clinical Presentation:** Cold, chronic, anterior cervical adenopathy ± sinus tracts. Usually in adults.

**Diagnostic Considerations:** Diagnosis by acid-fast stain and culture of node/drainage.

**Pitfalls:** Highly resistant to anti-TB therapy.

**Prognosis:** Excellent with surgical excision.

### **Cutaneous Mycobacterium haemophilum**

**Clinical presentation:** Particularly common in immunosuppressed/HIV patients, presents as multiple, painful cutaneous ulcers/abscesses  $\pm$  draining fistulas.

**Diagnostic Considerations:** Diagnose by PCR, RT-PCR assay.

**Pitfalls:** Suspect *M. haemophilum* if drainage from ulcer/abscess is AFB smear + but fails to grow on AFB media. *M. haemophilum* grows only on AFB media supplemented with ferric containing compounds i.e., hemin, hemoglobin, or ferric ammonium citrate. Susceptible to minocycline, TMP-SMX, or quinolones, but most strains resistant to INH, EMB, RIF, PZA.

**Prognosis:** Good with surgical excision/prolonged treatment.

### **Cutaneous Mycobacterium chelonae**

**Clinical presentation:** Present as cold abscesses with draining fistulas.

**Diagnostic Considerations:** AF culture of wound drainage.

**Pitfalls:** Some strains clarithromycin resistant. May be susceptible to quinolones, doxycycline, or minocycline.

**Prognosis:** Good with effective therapy.

### **Cutaneous Mycobacterium fortuitum**

**Clinical Presentation:** Usually associated with chronic foreign body infection, especially infected breast implants. Associated with nail salon footbaths. Present as cold abscesses with draining fistulas.

**Diagnostic Considerations:** Diagnosis by demonstrating organism by acid fast smear and culture of drainage/infected prosthetic material.

**Pitfalls:** Highly resistant to anti-TB therapy, may be susceptible to quinolones or linezolid.

**Prognosis:** Good with surgical excision/prolonged treatment.

### **Swimming Pool Granuloma (Mycobacterium marinum)**

**Clinical Presentation:** Begins as erythema with tenderness at inoculation site, followed by a papule or violaceous nodule that ulcerates and drains pus. May have sporotrichoid spread. Presents as a skin lesion unresponsive to antibiotics after abrasive water exposure (e.g., fish tank, swimming pool/lake).

**Diagnostic Considerations:** Diagnosis by demonstrating organism by acid-fast smear/culture.

**Pitfalls:** Resistant to INH/pyrazinamide. Surgical excision of isolated lesions may be needed.

**Prognosis:** Good with prolonged therapy.

### **Buruli Ulcer (Mycobacterium ulcerans)**

**Clinical Presentation:** Begin as a firm, painless, movable, subcutaneous nodule. In 1–2 months, nodule becomes fluctuant, ulcerates, and develops an undermined edge. May have edema around lesion and in extremity (if involved).

**Diagnostic Considerations:** Diagnosis by acid-fast stain and culture of punch biopsy of ulcer rim. Patient is usually from Africa, but *M. ulcerans* also exists in Asia, Australia, and Central/South America.

**Pitfalls:** Steroids/skin grafting sometimes needed.

**Prognosis:** Good with surgical excision.

### **Cutaneous MAI (Mycobacterium avium-intracellulare)**

**Clinical Presentation:** Nodules, abscesses, ulcers, plaques, ecthyma and draining sinuses can occur, but are uncommon in normal hosts and usually only seen in immunosuppressed patients.

**Diagnostic Considerations:** Diagnosis by demonstrating organism by acid-fast staining and culture (for species identification) of tissue biopsy specimens.

**Pitfalls:** Usually represents disseminated infection. Look for non-cutaneous evidence of infection (e.g., lungs, bone marrow, liver/spleen).

**Prognosis:** Related to extent of organ damage/degree of immunosuppression.

## Skin Vesicles/Bullae

Subset	Pathogens	Preferred Therapy
Herpes simplex	Herpes simplex virus (HSV)	See p. 142 (for HIV, see p. 340)
Herpes zoster	Varicella zoster virus (VZV)	See p. 142 (for HIV, see p. 319, 343)

## Subcutaneous Serpiginous Lesions

Subset	Pathogens	Preferred Therapy
Cutaneous larva migrans (creeping eruption)	Ancylostoma braziliense	Ivermectin 200 mcg/kg (PO) × 1 dose <b>or</b> Albendazole 400 mg (PO) q24h × 3–7 days. Children may be treated topically with albendazole ointment (10%) q8h × 10 days
Guinea worm	Dracunculus medinensis	Surgical removal of worm near skin surface. Metronidazole 250 mg (PO) q8h × 10 days facilitates worm removal
Cutaneous gnathostomiasis	Gnathostoma spinigerum	Surgical removal <b>or</b> Albendazole 400 mg (PO) q12h × 3 weeks <b>or</b> Ivermectin 200 mcg/kg (PO) q24h × 2 days

### Cutaneous Larva Migrans (Ancylostoma braziliense) Creeping Eruption

**Clinical Presentation:** Intensely pruritic, migratory, subcutaneous, raised serpiginous lesions.

**Diagnostic Considerations:** Diagnosis by clinical appearance.

**Pitfalls:** Must be differentiated from “swimmer’s itch” caused by schistosomal cercariae.

**Prognosis:** Excellent with treatment.

### Guinea Worm (Dracunculus medinensis)

**Clinical Presentation:** Serpiginous, raised, subcutaneous tract overlying worm.

**Diagnostic Considerations:** Diagnosis by demonstrating Dracunculus worm when surgically removed.

**Pitfalls:** Resembles cutaneous larva migrans, but worm is visible below the skin and lesions are serpiginous with Dracunculus. May be complicated by painful arthritis.

**Prognosis:** Excellent with treatment. Soaking extremity in warm water promotes emergence/removal of worm. Metronidazole can also be used to decrease inflammation and facilitate worm removal. Mebendazole 200–400 mg (PO) q12h × 6 days may kill worms directly.

**Cutaneous Gnathostomiasis (*Gnathostoma spinigerum*)**

**Clinical Presentation:** Painful, intermittent, subcutaneous swelling with local edema, intense pruritus, and leukocytosis with eosinophilia. Acquired by eating undercooked fish, frogs, and other intermediate larvae-containing hosts. May be complicated by eosinophilic meningitis.

**Diagnostic Considerations:** Diagnosis by demonstrating *Gnathostoma* in tissue specimens. Relatively common infection in Thailand and parts of Japan, South America, and Southeast Asia.

**Prognosis:** Good if limited to the skin and surgically removed. Poor with CNS involvement. Albendazole is preferred.

**Skin Papules/Nodules/Abscesses**

Subset	Pathogens	Preferred Therapy
Bacillary angiomatosis	<i>Bartonella henselae</i> <i>Bartonella quintana</i>	Doxycycline 100 mg (PO) q12h <b>or</b> azithromycin 250 mg (PO) q24h <b>or</b> any quinolone (PO) × 8 weeks
Cutaneous Alternaria	<i>Alternaria alternata</i>	Amphotericin B 0.7–1.0 mg/kg (IV) q24h × 2–3 grams <sup>‡</sup> <u>Alternate therapy:</u> Voriconazole, itraconazole, posaconazole, caspofungin. (See Drug Summaries Chapter 11 for dosage.) Treat until cured.
Entomophthoromycosis	<i>E. basidiobolus</i> <i>E. conidiobolus</i>	Amphotericin B 0.7–1.0 mg/kg (IV) q24h × 1–2 grams <sup>‡</sup> <b>or</b> TMP–SMX 1 DS tablet (PO) q24h until cured
Chromomycosis	<i>Fonsecaea pedrosoi</i> , compactum <i>Phialophora verrucosa</i> , others	<u>Few small lesions:</u> Wide/deep surgical excision or cryosurgery with liquid nitrogen <u>Larger lesions:</u> Itraconazole 200 mg (PO) solution q24h until lesions regress ± cryosurgery
Cutaneous Fusarium	<i>Fusarium solani</i>	Amphotericin B 0.7–1.0 mg/kg (IV) q24h × 2–3 grams <sup>‡</sup> <b>or</b> Voriconazole (see “usual dose,” p. 714)
Cutaneous Penicillium	<i>Penicillium marneffei</i>	Amphotericin B 0.6 mg/kg (IV) q24h × 14 days, <sup>‡</sup> then Itraconazole 200 mg (IV) q12 × 2 days followed by 200 mg (PO) q12h × 10 weeks. For HIV, continue with Itraconazole 200 mg (PO) q24h indefinitely
Cutaneous Prototheca	<i>Prototheca wickerhamii</i>	Surgical excision. If excision is incomplete, add either Amphotericin B 0.7–1.0 mg/kg (IV) q24h × 2–3 grams <sup>‡</sup> <b>or</b> Itraconazole 200 mg (PO) <sup>†</sup> q12h until cured
Trichosporon	<i>Trichosporon beigelii</i>	Amphotericin B 0.7–1.0 mg/kg (IV) q24h × 2–3 grams <sup>‡</sup>

‡ Liposomal amphotericin (p. 525) may be a suitable alternative.

† Initiate therapy with Itraconazole 200 mg (IV) q12h × 7–14 days.



## Skin Papules/Nodules/Abscesses (cont'd)

Subset	Pathogens	Preferred Therapy
Cutaneous aspergillosis*	Aspergillus sp.	Voriconazole (see "usual dose," p. 714) <b>or</b> Isavuconazole 200 mg (IV) q8h × 48 hours, then 200 mg (IV/PO) q24h until cured <b>or</b> Itraconazole 200 mg (IV) q12h × 7–14 days, then 200 mg (PO) solution q12h until cured
Cutaneous zygomycosis (mucormycosis)	Mucor/Rhizopus/ Absidia	Treat as for pulmonary zygomycosis (see p. 269)
Cutaneous coccidioidomycosis	Coccidioides immitis	Fluconazole 800 mg (PO) × 1 dose, then 400 mg (PO) q24h until cured <b>or</b> Itraconazole 200 mg (PO) <sup>†</sup> q12h until cured <u>Alternate therapy</u> Amphotericin B 1 mg/kg (IV) q24h × 7 days, then 0.8 mg/kg (IV) q48h × 2–3 grams total dose <sup>‡</sup>
Cutaneous histoplasmosis	Histoplasma capsulatum	See p. 284
Cutaneous cryptococcosis	Cryptococcus neoformans	Amphotericin B 0.5 mg/kg (IV) q24h × 1–2 grams <sup>†</sup> <b>or</b> Lipid amphotericin (p. 525) (IV) q24h × 3 weeks, <b>then</b> Fluconazole 800 mg (PO) × 1 dose, then 400 mg (PO) q24h × 8–10 weeks
Cutaneous sporotrichosis	Sporothrix schenckii	Itraconazole 200 mg (PO). If HIV, chronic suppressive therapy may be needed q24h until 2–4 weeks after all lesions have resolved (typically 3–6 months in total).
Nodular/pustular candidiasis	Candida sp.	If occurs in moist skin areas (under breasts, groin, etc.), therapy is for mucocutaneous candidiasis (see p. 147). Nodular candidiasis occurring in non-moist skin areas in compromised hosts is a manifestation of disseminated disease and should be treated as such (see p. 148–149)
Cutaneous onchocerciasis	Onchocerca volvulus	Ivermectin 150 mcg/kg (PO) × 1 dose, repeated q 4–6 months until asymptomatic <b>plus</b> Doxycycline 100 mg (PO) q12h × 6 weeks

\* Cutaneous aspergillus is a manifestation of disseminated aspergillosis.

† Initiate therapy with Itraconazole 200 mg (IV) q12h × 7–14 days.

‡ Lipid amphotericin (p. 525) may be a suitable alternative.

**Bacillary Angiomatosis (*Bartonella henselae*/*B. quintana*) Peliosis Hepatis**

**Clinical Presentation:** Skin lesions resemble Kaposi's sarcoma. Liver lesions resemble CMV hepatitis in HIV patients.

**Diagnostic Considerations:** Diagnosis by demonstrating organism by stain/culture of skin lesions or by blood culture after lysis-centrifugation.

**Pitfalls:** Requires life-long suppressive therapy.

**Prognosis:** Related to extent of infection/degree of immunosuppression.

**Cutaneous *Alternaria* (*Alternaria alternata*)**

**Clinical Presentation:** Bluish/purple papules that are often painful and non-pruritic. Usually seen only in leukopenic compromised hosts.

**Diagnostic Considerations:** Diagnosis by demonstrating organism by stain/culture in tissue specimen.

**Pitfalls:** Skin lesions usually represent disseminated disease in compromised hosts, not local infection. Fluconazole and micafungin ineffective.

**Prognosis:** Poor and related to degree of immunosuppression.

**Cutaneous Entomophthoromycosis (*E. basidiobolus*/*E. conidiobolus*)**

**Clinical Presentation:** *E. conidiobolus* infection presents as swelling of nose, paranasal tissues and mouth, accompanied by nasal stuffiness, drainage, and sinus pain. Begins as swelling of inferior nasal turbinates and extends until generalized facial swelling occurs. Subcutaneous nodules can be palpated in tissue. *E. basidiobolus* infection presents as a non-painful, firm, slowly progressive, subcutaneous nodule of the trunk, arms, legs, or buttocks.

**Diagnostic Considerations:** Diagnosis by demonstrating organism by stain/culture in tissue specimen. Skin lesions usually represent disseminated disease in compromised hosts, not localized infection.

**Pitfalls:** Unlike *Mucor*, *E. basidiobolus* does not usually invade blood vessels, although tissue infarction/necrosis is occasionally seen in diabetics and immunocompromised patients.

**Prognosis:** May spontaneously resolve. Surgical removal of accessible nodules and reconstructive surgery may be helpful for disfigurement.

**Cutaneous Chromomycosis (*F. pedrosoi*/*F. compactum*, *P. verrucosa*, others)**

**Clinical Presentation:** Warty papule/nodule that enlarges slowly to form a scarred, verrucous plaque. May also begin as a pustule, plaque, or ulcer. Over time, typical papule/nodule ulcerates, and the center becomes dry/crusted with raised margins. Lesions can be pedunculated/cauliflower-like.

**Diagnostic Considerations:** Diagnosis by demonstrating organism by stain/culture in tissue specimen. Chromomycosis remains localized within cutaneous/subcutaneous tissues.

**Pitfalls:** May resemble other fungal diseases. Sclerotic bodies in tissue and exudate distinguish chromomycosis from other related fungal diseases.

**Prognosis:** Related to degree of organ damage.

**Cutaneous *Fusarium* (*Fusarium solani*)**

**Clinical Presentation:** Presents in compromised hosts as multiple papules or painful nodules, initially macular with central pallor, which then become raised, erythematous, and necrotic. Seen mostly in leukopenic compromised hosts (especially acute leukemia and bone marrow transplants). Also a cause of mycetoma/onychomycosis.

**Diagnostic Considerations:** Diagnosis by demonstrating organism by stain/culture from blood/tissue.  $\beta$  1,3 D-glucan + , aspergillus galactomannan + .

**Pitfalls:** Skin lesions usually represent disseminated disease, not localized infection.

**Prognosis:** Poor/fair. Related to degree of immunosuppression. Amphotericin B lipid formulation (p. 525), colony-stimulating granulocyte factor, and granulocyte transfusions may be useful.

### **Cutaneous Penicillium (Penicillium marneffeii)**

**Clinical Presentation:** Papules, pustules, nodules, ulcers, or abscesses. Mostly seen in HIV (requires life-long suppressive therapy with itraconazole).

**Diagnostic Considerations:** Diagnosis by demonstrating organism by stain/culture in tissue specimen. Affects residents/visitors of Southeast Asia/Southern China.

**Pitfalls:** Lesions commonly become umbilicated and resemble molluscum contagiosum. May present as disseminated infection.

**Prognosis:** Poor and related to degree of immunosuppression.

### **Cutaneous Prototheca (Prototheca wickermanii)**

**Clinical Presentation:** Most common presentation is a single plaque or papulonodular lesion of the skin or subcutaneous tissue. Lesions are usually painless, slowly progressive (enlarge over weeks to months without healing), well-circumscribed, and may become eczematoid/ulcerated.

**Diagnostic Considerations:** Diagnosis by demonstrating organism by stain/culture in tissue specimen. Skin lesions in HIV are not different from normal hosts.

**Pitfalls:** Lesions are usually painless and may resemble eczema.

**Prognosis:** Poor/fair. Related to degree of immunosuppression. Surgical excision has been successful.

### **Cutaneous Trichosporon (Trichosporon beigelii)**

**Clinical Presentation:** Seen mostly in leukopenic compromised hosts (especially in acute leukemia, but also in HIV, burn wounds, and organ transplants). Usually presents as multiple red-bluish/purple papules, which are often painful and non-pruritic.

**Diagnostic Considerations:** Diagnosis by demonstrating organism by stain/culture in tissue specimen.

**Pitfalls:** Skin lesions usually represent disseminated disease, not localized infection.

**Prognosis:** Related to extent of infection/degree of immunosuppression.

### **Cutaneous Aspergillosis (Aspergillus fumigatus)**

**Clinical Presentation:** Seen at site of IV catheter insertion or adhesive dressing applied to skin in leukopenic, compromised hosts. Lesion is similar to pyoderma gangrenosum. May also invade burn wounds and cause rapidly progressive necrotic lesions.

**Diagnostic Considerations:** Diagnosis by demonstrating organism by stain/culture in tissue specimen.  $\beta$  1,3 D-glucan + , aspergillus galactomannan + .

**Pitfalls:** Infiltrative/ulcerative skin lesions usually represent disseminated disease in compromised hosts, not localized infection. May cause invasive dermatitis/skin lesions in HIV.

**Prognosis:** Related to extent of infection/degree of immunosuppression.

### **Cutaneous Mucormycosis/Rhizopus/Absidia**

**Clinical Presentation:** Necrotic skin lesion secondary to vascular invasion. Involves epidermis and dermis. Black eschars are evident.

**Diagnostic Considerations:** Diagnosis by demonstrating broad, non-septate hyphae with branches at right angles by stain/culture in tissue specimen.

**Pitfalls:** Skin lesions usually represent disseminated disease in compromised hosts, not localized infection. Contaminated elastic bandages have been associated with cutaneous Mucor.

**Prognosis:** Related to extent of infection/degree of immunosuppression.

### **Cutaneous Coccidioidomycosis (*Coccidioides immitis*)**

**Clinical Presentation:** Skin lesions may take many forms, including verrucous granulomas, cold subcutaneous abscesses, indolent ulcers, or small papules.

**Diagnostic Considerations:** Diagnosis by demonstrating organism by stain/culture in tissue specimen.

**Pitfalls:** Skin lesions usually represent disseminated disease in compromised hosts, not local infection.

**Prognosis:** Related to extent of infection/degree of immunosuppression.

### **Cutaneous Histoplasmosis (*Histoplasma capsulatum*)**

**Clinical Presentation:** Common cutaneous findings include maculopapular eruption, petechiae, and ecchymosis. Histopathology reveals necrosis around superficial dermal vessels.

**Diagnostic Considerations:** Diagnosis by demonstrating organism by stain/culture in tissue specimen.  $\beta$  1,3 D-glucan +, aspergillus galactomannan +.

**Pitfalls:** Skin lesions usually represent disseminated disease in compromised hosts, not local infection.

**Prognosis:** Related to extent of infection/degree of immunosuppression.

### **Cutaneous Cryptococcosis (*Cryptococcus neoformans*)**

**Clinical Presentation:** May present as single or multiple papules, pustules, erythematous indurated plaques, soft subcutaneous masses, draining sinus tracts, or ulcers with undermined edges.

**Diagnostic Considerations:** Diagnosis by demonstrating spherules with endospores by stain/culture in tissue specimen.  $\beta$  1,3 D-glucan -, aspergillus galactomannan -.

**Pitfalls:** Skin lesions usually represent disseminated disease in compromised hosts, not localized infection. In AIDS patients, umbilicated papules resemble molluscum contagiosum. In organ transplants, cellulitis with necrotizing vasculitis may occur.

**Prognosis:** Related to extent of infection/degree of immunosuppression.

### **Cutaneous Sporotrichosis (*Sporothrix schenckii*)**

**Clinical Presentation:** Primary cutaneous lymphatic sporotrichosis starts as a small, firm, movable, subcutaneous nodule, which then becomes soft and breaks down to form a persistent, friable ulcer. Secondary lesions usually develop proximally along lymphatic channels, but do not involve lymph nodes. Plaque form does not spread locally.

**Diagnostic Considerations:** Diagnosis by demonstrating organism by stain/culture in tissue specimen. Cutaneous disease arises at sites of minor trauma with inoculation of fungus into skin. Skin lesions usually represent disseminated disease in compromised hosts, not localized infection.

**Pitfalls:** HIV patients with  $CD_4 < 200$  may have widespread lymphocutaneous disease that ulcerates and is associated with arthritis. Unusual in axilla due to increased temperature.

**Prognosis:** Related to extent of infection/degree of immunosuppression.

### **Nodular/Pustular Candidiasis (sepsis in chronic steroids, see p. 158)**

**Cutaneous Onchocerciasis (*Onchocerca volvulus*)**

**Clinical Presentation:** Early manifestation is pruritic, papular rash with altered pigmentation. Later, papules, scaling, edema, and depigmentation may develop. Nodules develop in deep dermis/subcutaneous tissue (especially over bony prominences) or in deeper sites near joints, muscles, bones. Endemic in Middle East, Central Africa, Central America.

**Diagnostic Considerations:** Diagnosis by serology/demonstration of microfilaria in thin snips of involved skin or in anterior chamber (by slit lamp) if eye involvement. Intra-dermal edema produces "peau d'orange" effect with pitting around hair follicles/sebaceous glands.

**Pitfalls:** Ivermectin is effective against microfilaria, not adult worms. Adults live ~ 18 years.

**Prognosis:** Related to location/extent of organ damage.

**Rickettsia (Fever/Petechial Skin Rash)**

Subset	Pathogens	Preferred Therapy	Alternate Therapy
Rocky Mountain spotted fever (RMSF)	<i>Rickettsia rickettsii</i> (tick borne)	Doxycycline 200 mg (IV or PO) q12h × 3 days, then 100 mg (IV or PO) × 4 days	Any quinolone (IV or PO) × 7 days <b>or</b> Chloramphenicol 500 mg (IV or PO) q6h × 7 days
Epidemic (louse-borne) typhus, flying squirrel typhus	<i>Rickettsia prowazekii</i>	Same as RMSF	Same as RMSF
Murine (flea-borne) typhus	<i>Rickettsia typhi</i>	Same as RMSF	Same as RMSF
Scrub (chigger/mite-borne) typhus (Tsutsugamushi fever)	<i>Rickettsia tsutsugamushi</i>	Same as RMSF, <b>or</b> in mild cases Azithromycin 500 mg (PO) × 1 dose	Rifampin 600–900 mg (PO) q24h × 7 days
Rickettsialpox	<i>Rickettsia akari</i>	Same as RMSF	Same as RMSF
Tick typhus fevers (Mediterranean spotted fever, Boutonneuse fever, Israeli spotted fever)	<i>Rickettsia conorii</i> <i>Rickettsia parkeri</i>	Same as RMSF	Same as RMSF
African tick bite fever	<i>Rickettsia africae</i>	Same as RMSF	Same as RMSF

**Rocky Mountain Spotted Fever (*Rickettsia rickettsia*) RMSF**

**Clinical Presentation:** Fever with relative bradycardia, severe frontal headache, severe myalgias of abdomen/back/legs 3–12 days after tick bite. Rash begins as erythematous macules on wrists and ankles 3–5 days after tick bite, and progresses to petechiae/palpable purpura with confluent areas of ecchymosis. Periorbital edema, conjunctival suffusion, acute deafness, and edema of the dorsum of the hands/feet are important early signs. Abdominal pain can mimic acute abdomen, and meningismus and headache can mimic meningitis. Hepatosplenomegaly, cough, and coma may develop late. Laboratory findings include normal leukocyte count, thrombocytopenia, ↑ LFTs, and pre-renal azotemia. Hypotension/shock may occur secondary to myocarditis, which is the most common causes of death. Primary vector in United States is the Dermacentor tick; primary animal reservoir is small wild animals.

**Diagnosis:** Primarily a clinical diagnosis requiring a high index of suspicion and early empiric therapy. Early/rapid diagnosis can be made by DFA of biopsy specimen of rash (Weil-Felix reactions: OX-19: + + +, OX-2: +, OX-K: -). Specific *R. rickettsii* IFA, complement fixation, ELISA antibody titers are confirmatory. Include RMSF in differential diagnosis of any patient with fever and potential tick exposure, especially during the summer months.

**Pitfalls:** Most cases occur in eastern and southeastern United States, not Rocky Mountain area. Many cases go unrecognized due to nonspecific early findings. Early antibiotic therapy may blunt/eliminate serologic response. Patients with signs/symptoms of RMSF but without a rash should be considered as having ehrlichiosis ("spotless RMSF") until proven otherwise. Early therapy is essential; begin empiric therapy as soon as RMSF is suspected. Other adjunctive measures may be required.

**Prognosis:** Late (after day 5) initiation of treatment increases the risk of death by 5-fold. Adverse prognostic factors include myocarditis and severe encephalitis.

**Epidemic (Louse-Borne) Typhus (*Rickettsia prowazekii*)**

**Clinical Presentation:** High fever with relative bradycardia, chills, headache, conjunctival suffusion, and myalgias. A macular, rubella-like, truncal rash develops in most on the fifth febrile day, which may become petechial and involve the extremities, but spares the palms/soles. Facial swelling/flushing occurs at end of first week, along with CNS symptoms (e.g., tinnitus, vertigo, delirium) and GI complaints (diarrhea, constipation, nausea, vomiting, abdominal pain). Hypotension, pneumonia, renal failure, gangrene, cerebral infarction may develop late. Laboratory findings include normal leukocyte count, thrombocytopenia, and ↑ serum creatinine/LFTs. Primary vector is the human body louse; primary reservoir is humans.

**Diagnosis:** Primarily a clinical diagnosis requiring a high index of suspicion and early empiric therapy (Weil-Felix reactions: OX-19: + + +, OX-2: +, OX-K: -). Specific *R. prowazekii* antibody titers are confirmatory. Consider epidemic (louse-borne) typhus in febrile impoverished persons infested with lice, especially in Africa and parts of Latin America. Rarely seen in the United States. Milder recrudescence form (Brill-Zinsser disease) is also rare.

**Pitfalls:** Many cases go unrecognized due to nonspecific early findings. Early therapy is essential.

**Prognosis:** Gangrene of nose, ear lobes, genitalia, toes, and fingers may develop in severe cases. Death occurs in 10–50% of untreated patients.

**Murine (Flea-Borne) Typhus (*Rickettsia typhi*)**

**Clinical Presentation:** Similar to epidemic typhus but less severe, with fever in most, and headache, myalgias, and a macular rash in half. Laboratory findings include a normal leukocyte count, mild thrombocytopenia, and mildly ↑ LFTs. Primary vector is the Asian rat flea (*Xenopsylla cheopis*); primary reservoir is the commensal rat (*Rattus* genus). Uncommon in United States; most cases from Texas, California, Florida, Hawaii.

**Diagnosis:** Primarily a clinical diagnosis requiring a high index of suspicion. More common during summer and fall (Weil-Felix reactions: OX-19: + + +, OX-2: +, OX-K: -). Specific *R. typhi* antibody titers are confirmatory.

**Pitfalls:** Rash becomes maculopapular, compared to epidemic typhus, which remains macular. A similar illness is caused by *R. felis*.

**Prognosis:** Good if treated early. Death occurs in < 1%.

**Scrub Typhus (*Rickettsia tsutsugamushi*) Tsutsugamushi Fever**

**Clinical Presentation:** Fever, chills, headache, myalgias, arthralgias, GI symptoms, other nonspecific complaints. Eschar at mite bite site (tache noire) ± regional adenopathy. A macular, truncal rash develops in most, usually in the first week, and may progress to involve the extremities and face, but spares the palms/soles. Vasculitis may lead to cardiopulmonary (ARDS) CNS (encephalitis) hematologic abnormalities during the second week. Hepatosplenomegaly is common. Laboratory abnormalities include normal/decreased WBC, relative lymphopenia, thrombocytopenia, ↑ serum transaminases. Primary vector/reservoir is the larval (chigger) trombiculid mite. Endemic areas include northern Australia, southeastern Asia, Indian subcontinent.

**Diagnosis:** Presumptive diagnosis is clinical (Weil-Felix reactions: OX-19: -, OX-2: -, OX-K: + + +). Specific *R. tsutsugamushi* serology is confirmatory.

**Pitfalls:** Incomplete therapy frequently results in relapse.

**Prognosis:** Excellent if treated early.

**Rickettsialpox (*Rickettsia akari*)**

**Clinical Presentation:** Milder illness than other rickettsioses, with initial eschar at bite site, high fever, and generalized rash (usually erythematous papules which become vesicular and spares the palms/soles). Fever peak is usually < 104°F, occurs 1–3 weeks after mite bite, and lasts ~ 1 week without therapy. Headache, photophobia, marked diaphoresis, sore throat, GI complaints (nausea/vomiting following initial headache) may occur. Leukopenia may be present. Primary vector is the mouse mite; primary animal reservoir is the house mouse. Rare in the United States.

**Diagnosis:** Presumptive diagnosis by clinical presentation (Weil-Felix reactions: OX-19: -, OX-2: -, OX-K: ±). Specific *R. akari* serology is confirmatory.

**Pitfalls:** Do not confuse with African tick-bite fever, which may also have a vesicular rash.

**Prognosis:** Excellent even without therapy.

**Tick Typhus Fevers (*Rickettsia conorii*, *Rickettsia parkeri*) Mediterranean Spotted Fever, Boutonnoise Fever, Israeli Fever**

**Clinical Presentation:** Similar to RMSF with fever, chills, myalgias, but less severe. Unlike RMSF, an eschar is usually present at the site of the tick bite ± regional adenopathy. Leukocyte count is normal and thrombocytopenia is common. Transmitted by the brown dog tick, *Rhipicephalus sanguineus*.

**Diagnosis:** Presumptive diagnosis by clinical presentation (Weil-Felix reactions: OX-19: +, OX-2: +, OX-K: +). Specific *R. conorii* serology is confirmatory.

**Pitfalls:** Suspect in travelers from endemic areas with a RMSF-like illness. Consider different diagnosis in absence of an eschar.

**Prognosis:** Good with early treatment. Prostration may be prolonged even with proper therapy.

**African Tick Bite Fever (*Rickettsia africae*)**

**Clinical Presentation:** Similar to murine typhus with fever, chills, myalgias, but regional adenopathy and multiple eschars are common. Incubation period ~ 6 days. *Amblyomma hebreum/variegatum* tick vectors frequently bite humans multiple times.

**Diagnosis:** Presumptive diagnosis by clinical presentation (Weil-Felix reactions: OX-19: +, OX-2: +, OX-K: +). Specific *R. africae* serology is confirmatory.

**Pitfalls:** Rash is transient, and may be vesicular or absent.

**Prognosis:** Good even without therapy; excellent with therapy.

**Other Skin Lesions**

Subset	Pathogens	Topical Therapy	PO Therapy
Tinea versicolor (pityriasis)	<i>Malassezia furfur</i> ( <i>Pityrosporum orbiculare</i> )	Clotrimazole cream (1%) or miconazole cream (2%) <b>or</b> Ketoconazole cream (2%) daily × 7 days	Ketoconazole 200 mg (PO) q24h × 7 days <b>or</b> Itraconazole 200 mg (PO) q24h × 7 days <b>or</b> Fluconazole 400 mg (PO) × 1 dose
Eosinophilic folliculitis	<i>Malassezia furfur</i> ( <i>Pityrosporum orbiculare</i> )	<u>Preferred Therapy</u> Ketoconazole cream (2%) topically × 2–3 weeks ± Ketoconazole 200 mg (PO) q24h × 2–3 weeks	

**Tinea Versicolor/Pityriasis (*Malassezia furfur*)**

**Clinical Presentation:** Hyper- or hypopigmented scaling papules (0.5–1 cm), which may coalesce into larger plaques. Most commonly affects the upper trunk and arms. May be asymptomatic or pruritic.

**Diagnostic Considerations:** *M. furfur* also causes eosinophilic folliculitis in HIV, and catheter-acquired sepsis mostly in neonates or immunosuppressed patients. Diagnosis is clinical.

**Pitfalls:** Skin pigmentation may take months to return to normal after adequate therapy.

**Prognosis:** Excellent.

**Eosinophilic Folliculitis (*Malassezia furfur*)**

**Clinical Presentation:** Intensely pruritic folliculitis, usually on lower extremities.

**Diagnostic Considerations:** Tissue biopsy shows eosinophilic folliculitis. Diagnosis by demonstrating organism by culture on Sabouraud's agar overlaid with olive oil.

**Pitfalls:** Resembles folliculitis, but lesions are concentrated on lower extremities (not on trunk as with cutaneous candidiasis).

**Prognosis:** Related to degree of immunosuppression. Use oral therapy if topical therapy fails.



## Myositis

Subset	Pathogens	Preferred Therapy
Chromomycosis	Cladosporium/Fonsecaea	Itraconazole 200 mg (PO) q24h until cured <b>or</b> Terbinafine 250 mg (PO) q24h until cured
Trichinosis	Trichinella spiralis	Albendazole 400 mg (PO) q12h × 8–14 days <b>or</b> Mebendazole 200–400 mg (PO) q8h × 3 days, then 400–500 mg q8h × 10 days

### **Chromomycosis (Cladosporium/Fonsecaea)**

**Clinical Presentation:** Subcutaneous/soft tissue nodules or verrucous lesions.

**Diagnostic Considerations:** Diagnosis by demonstrating organism by culture/tissue biopsy specimen.

**Pitfalls:** May resemble Madura foot or cause ulcerative lesions in muscle.

**Prognosis:** Related to degree of immunosuppression.

### **Trichinosis (Trichinella spiralis)**

**Clinical Presentation:** Muscle tenderness, low-grade fevers, peripheral eosinophilia, conjunctival suffusion.

**Diagnostic Considerations:** Diagnosis by Trichinella serology or by demonstrating larvae in muscle biopsy.

**Pitfalls:** ESR is very low (near zero), unlike other causes of myositis, which have elevated ESRs.

**Prognosis:** Excellent with early treatment. Short-term steroids may be useful during acute phase. Therapy is ineffective against calcified larvae in muscle.

---

## REFERENCES AND SUGGESTED READINGS

---

- Adjei GO, Goka BQ, Kurtzhals AL. Neurotoxicity of Artemisinin Derivatives. *Clin Infect Dis* 43:1616–17, 2006.
- Alexander BD, Pfaller MA. Contemporary Tools for the Diagnosis and Management of Invasive Mycoses. *Clin Infect Dis* 43:S15–S27, 2006.
- Ansari S, Perez L, Vergely O, et al. Illnesses in travelers returning from the tropics: a prospective study of 622 patients. *J Travel Med* 12:312–18, 2005.
- Avulunov A, Klaus S, Vardy D. Fluconazole for the treatment of cutaneous leishmaniasis. *N Engl J Med* 347:370–1, 2002.
- Ayeh-Kumi PF, Petri Jr. WA. Amebiasis: An Update on Diagnosis and Treatment. *Infections in Medicine* 23:301–10, 2006.
- Bisser S, N'Siesi FX, Lejon V, et al. Equivalence trial of melarsoprol and nifurtimox monotherapy and combination therapy for the treatment of second-stage Trypanosoma brucei gambiense sleeping sickness. *J Infect Dis* 195:322–9, 2007.
- Chayakulkeeree M, Ghannoum MA, Perfect JR. Zygomycosis: the re-emerging fungal infection. *Clin Infect Dis* 25:215–229, 2006.
- Chen XM, Keithly JS, Paya CV, et al. Cryptosporidiosis. *N Engl J Med* 346:1723–1731, 2002.
- Cheng AC, Currie BJ. Melioidosis: epidemiology, pathophysiology, and management. *Clin Microbiol Rev* 18:383–16, 2005.
- Cisneros Herreros JM, Cordero Matia E. Therapeutic armamentarium against systemic fungal infections. *Clin Microbiol and Infection* 12:S53–64, 2006.
- Croft SL, Yardley V. Chemotherapy of leishmaniasis. *Curr Pharm Des* 8:319–42, 2002.

- Cunha BA. Malaria vs. Typhoid Fever: A Diagnostic Dilemma? *Am J Med* 118:1442–43, 2005.
- DeGroot MA, Huitt G. Infections Due to Rapidly Growing Mycobacteria. *Clin Infect Dis* 42:1756–63, 2006.
- Galgiani JN, Ampel NM, Blair JE, Catanzaro A, et al. Coccidioidomycosis. *Clin Infect Dis* 41: 1217–23, 2005.
- Griffith, KS; Lewis, LS; Mali, S; Parise, ME. Treatment of Malaria in the United States. *JAMA* 297:2264–2277, 2007.
- Haque R, Huston CD, Hughes M et al. Amebiasis. *N Engl J Med* 348:1565–73, 2003.
- Herbrecht R, Natarajan A, Letscher-Bru V, et al. Invasive pulmonary aspergillosis. *Semin Respir Crit Care Med* 25:191–202, 2004.
- International Artemisinin Study Group. Artesunate combinations for treatment of malaria: meta-analysis. *Lancet* 363:9–17, 2004.
- Kremsner PG, Krishna S. Antimalarial combinations. *Lancet* 364:285–94, 2004.
- Montoya JG, Liesenfeld O. Toxoplasmosis. *Lancet* 363: 1965–76, 2004.
- Munoz P, Guinea J, Bouza E. Update on invasive aspergillosis: clinical and diagnostic aspects. *Clin Microbiol and Infection* 12:524–39, 2006.
- Murray HW. Treatment of visceral leishmaniasis in 2004. *Am J Trop Med Hyg* 71:787–94, 2004.
- Mushatt DM, Mihm LB, Dreisbach AW, et al. Antibiotic Dosing in Slow Extended Daily Dialysis. *Clin Infect Dis* 49:433–437, 2009.
- Olliaro PL, Guerin PJ, Gerstl S, et al. Treatment options for visceral leishmaniasis: a systematic review of clinical studies done in India, 1980–2004. *Lancet Infect Dis* 5:763–4, 2005.
- Ouellette M, Drummelsmith J, Papadopoulos B. Leishmaniasis: drugs in the clinic, resistance and new developments. *Drug Resist Updat* 7:257–66, 2004.
- Perea S, Patterson TF. Invasive *Aspergillus* infections in hematologic malignancy patients. *Semin Respir Infect* 17:99–105, 2002.
- Pirisi M, Salvador E, Bisoffi Z, et al. Unsuspected strongyloidiasis in hospitalized elderly patients with and without eosinophilia. *Clin Microbiol Infect* 12:787–92, 2006.
- Priotto G, Kasparian S, Mutombo W, et al. Nifurtimox-eflornithine combination therapy for second-stage African *Trypanosoma brucei gambiense* trypanosomiasis: a multicentre, randomized, phase III, noninferiority trial. *Lancet* 374:56–64, 2009.
- Ross AGP, Bartley PB, Sleight AC, et al. Schistosomiasis. *N Engl J Med* 346:1212–1220, 2002.
- Segal BH, Walsh TJ. Current Approaches to Diagnosis and Treatment of Invasive Aspergillosis. *Am J Respir Crit Care Med* 173:707–17, 2006.
- Sinha PK, Ranjan A, Singh VP, Das VNR, et al. Visceral leishmaniasis (kala-azar)—the Bihar (India) perspective. *Journal of Infection* 53:60–64, 2006.
- Sobel JD. Combination therapy for invasive mycosis: evaluation of past clinical trial designs. *Clin Infect Dis* 15; 39 Suppl 4:S224–7, 2004.
- Soto J. Miltefosine for new world cutaneous leishmaniasis. *Clin Infect Dis* 38:1266–72, 2004.
- Tortoli E. Clinical Manifestations of nontuberculous mycobacteria infections. *Clin Microbiol Infect* 15:906–910, 2009.
- Udall DN. Recent Updates on Onchocerciasis: Diagnosis and Treatment. *Clin Infect Dis* 44:53–60, 2007.
- Vardakas KZ, Michalopoulos A, Kiriakidou KG, et al. Candidaemia: Incidence, risk factors, characteristics and outcomes in immunocompetent critically ill patients. *Clinical Microbiology and Infectious Diseases* 15:286–295, 2009.
- White AC. New Developments in the Management of Neurocysticercosis. *J of Infectious Diseases* 199: 1261–1262, 2009.

#### GUIDELINES

- Chapman SW, Bradsher RW Jr, Campbell GD Jr, et al. Practice guidelines for the management of patients with blastomycosis. Infectious Diseases Society of America. *Clin Infect Dis* 30:679–83, 2000.
- Galgiani JN, Ampel NM, Catanzaro A, et al. Practice guidelines for the treatment of coccidioidomycosis. Infectious Diseases Society of America. *Clin Infect Dis* 30:658–61, 2000.
- Hill DR, Ericsson CD, Pearson RD, Keystone JS, et al. The Practice of Travel Medicine: Guidelines by the Infectious Diseases Society of America. *Clin Infect Dis* 43: 1499–539, 2006.
- Kauffman CA, Hajjeh R, Chapman SW. Practice guidelines for the management of patients with sporotrichosis. For the Mycoses Study Group. Infectious Diseases Society of America. *Clin Infect Dis* 30:684–7, 2000.
- Pappas PG, Rex JH, Sobel JD, Filler SG, et al. Guidelines for the Treatment of Candidiasis. *Clin Infect Dis* 38: 161–89, 2004.
- Perfect JR, Dismukes WE, Dromer F, et al. Clinical Practice Guidelines for the Management of Cryptococcal Disease: 2010 Update by the Infectious Disease Society of America. *Clin Infect Disease* 50:291–322, 2010.
- Sobel JD. Practice guidelines for the treatment of fungal infections. For the Mycoses Study Group. Infectious Diseases Society of America. *Clin Infect Dis* 30:652, 2000.

- Walsh TH, Anaissie EJ, Denning DW, et al. Treatment of Aspergillosis: Clinical Practice Guidelines of the Infectious Diseases Society of America. *Clin Infect Dis* 46: 327–60, 2008.
- Wheat J, Sarosi G, McKinsey D, Hamill R, et al. Practice guidelines for the management of patients with histoplasmosis. Infectious Diseases Society of America. *Clin Infect Dis* 30:688–95, 2000.
- Wormser GP, Dattwyler RJ, Shapiro ED, Halperin JJ, et al. The Clinical Assessment, Treatment, and Prevention of Lyme Disease, Human Granulocytic Anaplasmosis, and Babesiosis: Clinical Practice Guidelines by the Infectious Diseases Society of America. *Clin Infect Dis* 43:1089–134, 2006.
- TEXTBOOKS**
- Anaissie EJ, McGinnis MR, Pfaller MA (eds). *Clinical Mycology* (2<sup>nd</sup> Ed), Churchill Livingstone, New York, 2009.
- Arikan S, Rex JH. Antifungal Drugs. In: *Manual of Clinical Microbiology* (8<sup>th</sup> Ed), ASM Press, Washington, D.C., 2003.
- Bennett JE, Dolin R, Blaser MJ (eds). *Mandell, Douglas, and Bennett's Principles and Practice of Infectious Diseases* (8<sup>th</sup> Ed), Philadelphia, Elsevier, 2015.
- Bottone EJ (ed). *An Atlas of the Clinical Microbiology of Infectious Diseases, Volume 1. Bacterial Agents*. The Parthenon Publishing Group, Boca Raton, 2004.
- Bottone EJ (ed). *An Atlas of the Clinical Microbiology of Infectious Diseases, Volume 2. Viral, fungal, and parasitic agents*. Taylor and Francis, 2006.
- Bryskier A (ed). *Antimicrobial Agents*. ASM Press, Washington, D.C., 2005.
- Calderone RA (ed). *Candida and Candidiasis*. ASM Press, Washington D.C., 2002.
- Cook GC (ed). *Manson's Tropical Diseases* (24<sup>th</sup> Ed), WB Saunders Company, Ltd., London, 2014.
- Cunha BA (ed). *Tick-Borne Infectious Diseases*. Marcel Dekker, New York, 2000.
- Despommier DD, Gwadz RW, Hotez PJ, Knirsch CA (eds). *Parasitic Diseases* (4<sup>th</sup> Ed), Apple Trees Productions, LLC, New York, 2000.
- Garcia LS (ed). *Practical guide to Diagnostic Medical Parasitology*. (5<sup>th</sup> Ed), ASM Press, Washington, DC, 2007.
- Gorbach SL, Bartlett JG, Blacklow NR (eds). *Infectious Diseases* (3<sup>rd</sup> Ed), Philadelphia, Lippincott, Williams & Wilkins, 2004.
- Guerrant RL, Walker DH, Weller PF (eds). *Tropical Infectious Disease: Principles, Pathogens & Practice* (3<sup>rd</sup> Ed), Elsevier, Philadelphia, 2011.
- Gutierrez Y (ed). *Diagnostic Pathology of Parasitic Infections with Clinical Correlations* (2<sup>nd</sup> Ed), Oxford University Press, New York, 2000.
- Howard DH (ed). *Pathogenic Fungi in Humans and Animals* (2<sup>nd</sup> Ed), New York, Marcel Dekker, 2003.
- Jong E, Sanford C (eds). *The Travel and Tropical Medicine Manual* (4<sup>th</sup> Ed), Philadelphia: Saunders Elsevier, 2008.
- Keystone JS, Kozarsky PE, Freedman DO, Nothdurft HO, Connor B. *Travel Medicine* (2<sup>nd</sup> Ed), Philadelphia: Mosby Elsevier, 2008.
- Knipe DM, Howley PM (eds). *Fields, Virology* (5<sup>th</sup> Ed), Lippincott Williams & Wilkins, Philadelphia, 2008.
- Krauss H, Weber A, Appel M (eds). *Zoonoses. Infectious Diseases Transmissible from Animals to Humans* (3<sup>rd</sup> Ed), ASM Press, Washington, D.C. 2003.
- Maertens JA, Marr KA (eds). *Diagnosis of Fungal Infections*. Informa Healthcare, New York, 2007.
- Porterfield JS (ed). *Exotic Viral Infections*. Chapman & Hall Medical, London, 1995.
- Raoult D, Parola P. *Rickettsial Diseases*. Informa Healthcare, New York, 2007.
- Richman DD, Whitley RJ, Hayden FG (eds). *Clinical Virology* (2<sup>nd</sup> Ed), ASM Press, Washington, DC, 2002.
- Schollberg DM (ed). *Current Therapy of Infectious Diseases* (3<sup>rd</sup> Ed), Mosby St. Louis, 2008.
- Strickland GT (ed). *Hunter's Tropical Medicine and Emerging Infectious Diseases* (8<sup>th</sup> Ed), W.B. Saunders Company, Philadelphia, 2000.
- Sun T (ed). *Parasitic Disorders: Pathology, Diagnosis, and Management* (2<sup>nd</sup> Ed), Williams & Wilkins, Baltimore, 1999.
- Walker DH (ed). *Biology of Rickettsial Diseases*. CRC Press, Boca Raton, 2000.
- Warrell DA, Gilles HM. *Essential Malariology* (4<sup>th</sup> Ed), Oxford University Press, New York, 2002.
- Yu VL, Jr., Merigan TC, Jr., and Barriere SL (eds.). *Antimicrobial Therapy & Vaccines* (2<sup>nd</sup> Ed), Williams & Wilkins, Baltimore, MD, 2005.

## Chapter 5

**HIV Infection****Paul E. Sax, MD****Jean E. Hage, MD****Arthur Gran, MD****Jeffrey Baron, PharmD**

HIV Infection Overview .....	302
Stages of HIV Infection .....	303
Acute (Primary) HIV Infection .....	304
Approach to HIV Testing .....	305
Initial Assessment of HIV Infection .....	308
Indications for Treatment of HIV Infection .....	312
Antiretroviral Treatment .....	313
Antiretroviral Treatment Failure .....	316
Prophylaxis of Opportunistic Infections in HIV .....	319
Treatment of Opportunistic Infections in HIV .....	323
Treatment of Other Opportunistic Infections in HIV .....	338
HIV Coinfections (HBV/HCV) .....	344
References and Suggested Readings .....	351
For HIV Pre-Exposure Prophylaxis (PrEP) .....	365
For HIV Post-Exposure Prophylaxis (PEP) .....	366


---

**HIV INFECTION OVERVIEW**


---

**Paul E. Sax, M.D.**

Infection with Human Immunodeficiency Virus (HIV-1) leads to a chronic and without treatment usually fatal infection characterized by progressive immunodeficiency, a long clinical latency period, and opportunistic infections. The hallmark of HIV disease is infection and viral replication within T-lymphocytes expressing the CD<sub>4</sub> antigen (helper-inducer lymphocytes), a critical component of normal cell-mediated immunity. Qualitative defects in CD<sub>4</sub> responsiveness and progressive depletion in CD<sub>4</sub> cell counts increase the risk for opportunistic infections such as *Pneumocystis carinii* jiroveci pneumonia, and neoplasms such as lymphoma and Kaposi's sarcoma. HIV infection can also disrupt blood monocyte, tissue macrophage, and B-lymphocyte (humoral immunity) function, predisposing to infection with encapsulated bacteria. Direct attack of CD<sub>4</sub>-positive cells in the central and peripheral nervous system can cause HIV meningitis, peripheral neuropathy, and dementia. More than 1 million people in the United States and 30 million people worldwide are infected with HIV. Without treatment, the average time from acquisition of HIV to an AIDS-defining opportunistic infection is about 10 years; survival then averages 1–2 years. There is tremendous individual variability in these time intervals, with some patients progressing from acute HIV infection to death within 1–2 years, and others not manifesting HIV-related immunosuppression for > 20 years after HIV acquisition. Antiretroviral therapy and prophylaxis against opportunistic infections have markedly improved the overall prognosis of HIV disease. The approach to HIV infection is shown in Figure 5.1.




---

**Figure 5.1. Diagnosis, Evaluation, and Treatment of HIV Infection**

---

**STAGES OF HIV INFECTION**

---

- A. Viral Transmission.** HIV infection is acquired primarily by sexual intercourse (anal, vaginal, infrequently oral), exposure to contaminated blood (primarily needle transmission), or maternal-fetus (perinatal) transmission. Sexual practices with the highest risk of transmission include unprotected receptive anal intercourse (especially with mucosal tearing), unprotected receptive vaginal intercourse (especially during menses), and unprotected rectal/vaginal intercourse in the presence of genital ulcers (e.g., primary syphilis, genital herpes, chancroid). Lower risk sexual practices include insertive anal/vaginal intercourse and oral-genital contact. The risk of transmission after a single encounter with an HIV source has been estimated to be 1 in 150 with needle sharing, 1 in 300 with occupational percutaneous exposure, 1 in 300–1000 with receptive anal intercourse, 1 in 500–1250 with receptive vaginal intercourse, 1 in 1000–3000 with insertive vaginal intercourse, and 1 in 3000 with insertive anal intercourse. Transmission risk increases with the number of encounters and with higher HIV RNA plasma levels. The mode of transmission does not affect the natural history of HIV disease.
- B. Acute (Primary) HIV Infection (p. 304).** Acute HIV occurs 1–4 weeks after transmission, and is accompanied by a burst of viral replication with a decline in CD<sub>4</sub> cell count. Most patients manifest a symptomatic mononucleosis-like syndrome, which is often overlooked. Acute HIV infection is confirmed by a high HIV RNA in the absence of HIV antibody.
- C. Seroconversion.** Development of a positive HIV antibody test usually occurs within 4 weeks of acute infection, and invariably (with few exceptions) by 6 months.
- D. Asymptomatic HIV Infection** lasts a variable amount of time (average 8–10 years), and is accompanied by a gradual decline in CD<sub>4</sub> cell counts and a relatively stable HIV RNA level (sometimes referred to as the viral “set point”).
- E. Symptomatic HIV Infection.** Previously referred to as “AIDS Related Complex (ARC),” findings include thrush or vaginal candidiasis (persistent, frequent, or poorly responsive to treatment), cervical dysplasia/carcinoma in-situ, herpes zoster (recurrent episodes or involving multiple dermatomes), oral hairy leukoplakia, peripheral neuropathy, diarrhea, or constitutional symptoms (e.g., low-grade fevers, weight loss).
- F. AIDS** is defined by a CD<sub>4</sub> cell count < 200/mm<sup>3</sup>, a CD<sub>4</sub> cell percentage of total lymphocytes <14%, or one of several AIDS-related opportunistic infections. Common opportunistic infections include *Pneumocystis (carinii) jiroveci* pneumonia, cryptococcal meningitis, recurrent bacterial pneumonia, *Candida* esophagitis, CNS toxoplasmosis, tuberculosis, and non-Hodgkin’s lymphoma. Other AIDS indicators in HIV-infected patients include candidiasis of the bronchi, trachea, or lungs; disseminated/extrapulmonary coccidiomycosis, cryptococcosis, or histoplasmosis; chronic (>1 month) intestinal cryptosporidiosis or isosporiasis; Kaposi’s sarcoma; lymphoid interstitial pneumonia/pulmonary lymphoid hyperplasia; disseminated/extrapulmonary *Mycobacterium (avium-intracellulare, kansasii,*

other species) infection; progressive multifocal leukoencephalopathy (PML); recurrent Salmonella septicemia; or HIV wasting syndrome.

- G. Advanced HIV Disease** is diagnosed when the  $CD_4$  cell count is  $< 50/mm^3$ . Most AIDS-related deaths occur at this point. Common late stage opportunistic infections are caused by CMV disease (retinitis, colitis) or disseminated *Mycobacterium avium-intracellulare* (MAI).

---

### ACUTE (PRIMARY) HIV INFECTION

---

- A. Description.** Acute clinical illness associated with primary acquisition of HIV, occurring 1–4 weeks after viral transmission (range: 6 days to 6 weeks). Symptoms develop in 50–90%, but are often mistaken for the flu, mononucleosis, or other nonspecific viral syndrome. More severe symptoms may correlate with a higher viral set point and more rapid HIV disease progression. Even without therapy, most patients recover, reflecting development of a partially effective immune response and depletion of susceptible  $CD_4$  cells.
- B. Differential Diagnosis** includes **EBV, CMV**, viral hepatitis, enteroviral infection, 2<sup>o</sup> syphilis, toxoplasmosis, HSV with erythema multiforme, drug reaction, Behcet's disease, acute lupus.
- C. Signs and Symptoms** usually reflect hematogenous dissemination of virus to lymphoreticular and neurologic sites:
- Fever (97%).
  - Pharyngitis (73%). Typically non-exudative (unlike EBV, which is usually exudative).
  - Rash (77%). Maculopapular viral exanthem of the face and trunk is most common, but can involve the extremities, palms and soles.
  - Arthralgia/myalgia (58%).
  - Neurologic symptoms (12%). Headache is most common. Neuropathy, Bell's palsy, and meningoencephalitis are rare, but may predict worse outcome.
  - Oral/genital ulcerations, thrush, nausea, vomiting, diarrhea, weight loss.
- D. Laboratory Findings**
1. **CBC.** Lymphopenia followed by lymphocytosis (common). Atypical lymphocytosis is variable, but usually low level (unlike EBV, where atypical lymphocytosis may be 20–30% or higher). Thrombocytopenia occurs in some.
  2. **Elevated transaminases** in some but not all patients.
  3. **Depressed  $CD_4$  cell count.** Can rarely be low enough to induce opportunistic infections.
  4. **HIV antibody.** Usually negative, although persons with prolonged symptoms of acute HIV may have positive antibody tests if diagnosed late during the course of illness.

**E. Confirming the Diagnosis of Acute HIV Infection**

- 1. Obtain HIV antibody** after informed consent (if required by state law) to exclude prior disease.
- 2. Order viral load test (HIV RNA PCR)**, preferably RT-PCR. HIV RNA confirms acute HIV infection prior to seroconversion. Most individuals will have very high HIV RNA (>100,000 copies/mL). Be suspicious of a false-positive test if the HIV RNA is low (< 20,000 copies/mL). For any positive test, it is important to repeat HIV RNA and HIV antibody testing. p24 antigen can also be used to establish the diagnosis, but is less sensitive than HIV RNA PCR.
- 3. Order other tests/serologies if HIV RNA test is negative.** Order throat cultures for bacterial/viral respiratory pathogens, EBV VCA IgM/IgG, CMV IgM/IgG, HHV-6 IgM/IgG, and hepatitis serologies as appropriate to establish a diagnosis for patient's symptoms.

**F. Management of Acute HIV Infection**

- 1. Initiate antiretroviral therapy.** The rationale behind this change is to treat a greater proportion of people with HIV, regardless of disease stage, is based on accumulating evidence of the benefits of earlier initiation of HIV treatment and, conversely, the potential for uncontrolled viral replication and CD<sub>4</sub> depletion.
- 2. Obtain HIV resistance genotype** (see p. 317) because of a rising background prevalence of transmission of antiretroviral therapy-resistant virus. A genotype resistance test is preferred; therapy can be started pending results of the test.
- 3. Possible benefits for treatment of acute HIV infection.** Possible (but unproven) benefits include hastening symptom resolution, reducing viral transmission, lowering virologic "set point," and preserving virus-specific CD<sub>4</sub> responses.

---

**APPROACH TO HIV TESTING (Figure 5.2)**


---

**A. Standard HIV Antibody Tests.** Most patients produce antibody to HIV within 6–8 weeks of exposure; half will have a positive antibody test in 3–4 weeks, and nearly 100% will have detectable antibody by 6 months.

- 1. ELISA.** Usual screening test. All positives must be confirmed with Western blot or other more specific tests.
- 2. Western blot.** CDC criteria for interpretation: **positive:** at least two of the following bands: p24, gp41, gp160/120; **negative:** no bands; **indeterminate:** any HIV band, but does not meet criteria for positivity.
- 3. Test performance.** Standard method is ELISA screen with Western blot confirmation.
  - a. ELISA negative:** Western blot is not required (ELISA sensitivity 99.7%, specificity 98.5%). Obtain HIV RNA if acute HIV infection is suspected.
  - b. ELISA positive:** Confirm with Western blot. Probability that ELISA and Western blot are both false-positives is extremely low (< 1 per 140,000). Absence of p31 band could be a clue to a false positive Western blot.
  - c. Unexpected ELISA/Western blot:** Repeat test to exclude clerical/computer error.



- 4. Indeterminate Western Blot.** Common clinical problem, affecting 4–20% of reactive ELISAs. Usually due to a single p24 band or weak other bands. Causes include seroconversion in progress, advanced HIV disease with loss of antibody response,



**Figure 5.2. Approach to HIV Testing**

(-) = negative test; (+) = positive test.

\* Occurs 1–4 weeks after viral transmission. Most patients manifest a viral syndrome (fever, pharyngitis ± rash/arthralgias), which is often mistaken for the flu and therefore overlooked.

\*\* HIV RNA in acute HIV infection should be very high (usually > 100,000 copies/mL).

+ All positive ELISA tests must be confirmed by Western Blot; usually this is done automatically in clinical laboratories.

++ May be long-term non-progressor or laboratory error.

cross-reacting antibody from pregnancy, blood transfusions, organ transplantation, autoantibodies from collagen vascular disease, infection with HIV-2, influenza vaccination, or recipient of HIV vaccine. In low-risk patients, an indeterminate result almost never represents true HIV infection. Since seroconversion in progress is generally associated with high HIV RNA levels, the recommended approach is to order an HIV RNA test.

## B. Quantitative Plasma HIV RNA (HIV Viral Load Assays)

- 1. Description.** Measures amount of HIV RNA in plasma. High sensitivity of assays allows detection of virus in most patients not on antiviral therapy. Used to diagnose acute HIV infection and more commonly to monitor the response to antiretroviral therapy.
- 2. Uses of HIV RNA Assay**
  - a. Confirms diagnosis of acute HIV infection.** A high HIV RNA with a negative HIV antibody test confirms acute HIV infection prior to seroconversion.
  - b. Helpful in initial evaluation of HIV infection.** Establishes baseline HIV RNA and helps (along with CD<sub>4</sub> cell count) determine whether to initiate or defer therapy, as HIV RNA correlates with rate of CD<sub>4</sub> decline.
  - c. Monitors response to antiviral therapy.** HIV RNA changes rapidly decline 2–4 weeks after starting or changing effective antiretroviral therapy, with slower decline thereafter. Patients with the greatest HIV RNA response have the best clinical outcome. No change in HIV RNA suggests therapy will be ineffective.
  - d. Estimates risk for opportunistic infection.** For patients with similar CD<sub>4</sub> cell counts, the risk of opportunistic infections is higher with higher HIV RNAs.

## 3. Assays and Interpretation

- a. Tests, sensitivities, and dynamic range.** Three main assays, each with advantages and disadvantages, are widely used. Any assay can be used to diagnose acute HIV infection and guide/monitor therapy, but the same test should be used to follow patients longitudinally.
  - 1. RT-PCR Amplicor (Roche):** Sensitivity = 400 copies/mL; dynamic range = 400–750,000 copies/mL.
  - 2. RT-PCR Ultrasensitive 1.5 (Roche):** Sensitivity = 50 copies/mL; dynamic range = 50–75,000 copies/mL.
  - 3. bDNA Versant 3.0 (Bayer):** Sensitivity = 75 copies/mL; dynamic range = 50–500,000 copies/mL.
- b. Correlation between HIV RNA and CD<sub>4</sub>.** HIV RNA assays correlate inversely with CD<sub>4</sub> cell counts, but do so imperfectly (e.g., some patients with high CD<sub>4</sub> counts have relatively high HIV RNA levels, and vice versa.) For any given CD<sub>4</sub>, higher HIV RNA levels correlate with more rapid CD<sub>4</sub> decline. In response to antiretroviral therapy, changes in HIV RNA generally precede changes in CD<sub>4</sub> cell count.

- c. Significant change in HIV RNA assay** is defined by at least a 2-fold (0.3 log) change in viral RNA (accounts for normal variation in clinically stable patients), or a 3-fold (0.5 log) change in response to new antiretroviral therapy (accounts for intra-laboratory and patient variability). For example, if a HIV RNA result = 50,000 copies/mL, then the range of possible actual values = 25,000–100,000 copies/mL, and the value needed to demonstrate antiretroviral activity is  $\leq 17,000$  copies/mL.
- 4. Indications for HIV RNA Testing.** Usually performed in conjunction with CD<sub>4</sub> cell counts. Indicated for the diagnosis of acute HIV infection, and for initial evaluation of newly diagnosed HIV. Also recommended 2–8 weeks after initiation of antiretroviral therapy and every 3–4 months in all HIV patients.
- 5. When to Avoid HIV RNA Testing**
- a. During acute illnesses and immunizations.** Patients with acute infections (opportunistic infection, bacterial pneumonia, even HSV recurrences) may experience significant (> 5-fold) rises in HIV RNA, which return to baseline 1–2 months after recovery. Although data are conflicting, many studies show at least a transient increase in HIV RNA levels following influenza and other immunizations, which return to baseline after 1–2 months.
- b. When results of test would not influence therapy.** Frequent scenario in patients with advanced disease who have no antiretroviral options or cannot tolerate therapy.
- c. As a screening test for HIV infection,** except if acute (primary) HIV disease is suspected during the HIV antibody window (i.e., first 3–6 weeks after viral transmission).

---

## INITIAL ASSESSMENT OF HIV INFECTION

---

- A. Clinical Evaluation.** History and physical exam should focus on diagnoses associated with HIV infection. Compared to patients without HIV, the severity, frequency, and duration of these conditions are usually increased in HIV disease.
- 1. Dermatologic:** Severe herpes simplex (oral/anogenital); herpes zoster (especially recurrent, cranial nerve, or disseminated); molluscum contagiosum; staphylococcal abscesses; tinea nail infections; Kaposi's sarcoma (from HHV-8 infection); petechiae (from ITP); seborrheic dermatitis; new or worsening psoriasis; eosinophilic pustular folliculitis; severe cutaneous drug eruptions (especially sulfonamides).
- 2. Oropharyngeal:** Oral candidiasis; oral hairy leukoplakia (from EBV); Kaposi's sarcoma (most commonly on palate or gums); gingivitis/periodontitis; warts; aphthous ulcers (especially esophageal/perianal).

3. **Constitutional symptoms:** Fatigue, fevers, chronic diarrhea, weight loss.
4. **Lymphatic:** Persistent, generalized lymphadenopathy.
5. **Others:** Active TB (especially extrapulmonary); non-Hodgkin's lymphoma (especially CNS); unexplained leukopenia, anemia, thrombocytopenia (especially ITP); myopathy; miscellaneous neurologic conditions (cranial/peripheral neuropathies, Guillain-Barre syndrome, mononeuritis multiplex, aseptic meningitis, cognitive impairment).

#### B. Baseline Laboratory Testing (Table 5.1)

#### C. CD<sub>4</sub> Cell Count (lymphocyte subset analysis)

1. **Overview.** Acute HIV infection is characterized by a decline in CD<sub>4</sub> cell count, followed by a gradual rise associated with clinical recovery. Chronic HIV infection shows progressive declines (~ 50–80 cells/year) in CD<sub>4</sub> cell count without treatment, followed by more rapid decline 1–2 years prior to opportunistic infection (AIDS-defining diagnosis). Cell counts remain stable over 5–10 years in 5% of patients, while others may show rapid declines (> 300 cells/year). Since variability exists within individual patients and between laboratories, it is useful to *repeat any value before making management decisions*.
2. **Uses of CD<sub>4</sub> Cell Count**
  - a. **Gives context of degree of immunosuppression** for interpretation of symptoms/signs (Table 5.2).
  - b. **Used to guide therapy.** Guidelines support CD<sub>4</sub> < 500/mm<sup>3</sup> as the threshold for initiating treatment, regardless of HIV RNA or symptoms. For prophylaxis against PCP, toxoplasmosis, and MAI/CMV infection, CD<sub>4</sub> cell counts of 200/mm<sup>3</sup>, < 100/mm<sup>3</sup>, and < 50/mm<sup>3</sup> are used as threshold levels, respectively.
  - c. **Provides estimate of risk of opportunistic infection or death.** CD<sub>4</sub> cell counts < 50/mm<sup>3</sup> are associated with a markedly increased risk of death (median survival 1 year), although some patients with low counts survive > 3 years even without antiretroviral therapy. Prognosis is heavily influenced by HIV RNA, presence/history of opportunistic infections or neoplasms, performance status, and the immune reconstitution response to antiretroviral therapy.

#### D. HIV RNA Assay (HIV RNA PCR) (p. 307)

**Table 5.1. Baseline Laboratory Testing for HIV-Infected Patients**

Test	Rationale
Repeat HIV serology (ELISA/confirmatory Western blot)	Indicated for patients unable to document a prior positive test, and for "low risk" individuals with a positive test (to detect computer/clerical error). Repeat serology is now less important since HIV RNA testing provides an additional means of confirming HIV infection.

**Table 5.1. Baseline Laboratory Testing for HIV-Infected Patients** (cont'd)

Test	Rationale
CBC with differential, platelets	Detects cytopenias (e.g., ITP) seen in HIV. Needed to calculate CD <sub>4</sub> cell count.
Chemistry panel ("SMA 20") and fasting lipid panel	Detects renal dysfunction and electrolyte/LFT abnormalities, which may accompany HIV and associated infections (e.g., HIV nephropathy, HCV). Provides baseline lipid profile (many antiretroviral drugs can affect lipids).
CD <sub>4</sub> cell count	Determines the need for antiretroviral therapy and opportunistic infection (OI) prophylaxis. Best test for defining risk of OIs and prognosis.
HIV RNA assay ("viral load")	Provides a marker for the pace of HIV disease progression. Determines indication for and response to antiretroviral therapy.
Tuberculin skin test (standard 5 TU of PPD)	Detects latent TB infection and targets patients for preventive therapy. Anergy skin tests are no longer indicated due to poor predictive value. HIV is the most powerful co-factor for the development of active TB.
PAP smear	Risk of cervical cancer is nearly twice as high in HIV-positive women compared to uninfected controls.
HLA-B*5701	Needed if abacavir therapy is planned, to assess risk for severe abacavir hypersensitivity reactions.
Toxoplasmosis serology (IgG)	Identifies patients at risk for subsequent cerebral/systemic toxoplasmosis and the need for prophylaxis. Those with negative tests should be counseled on how to avoid infection.
Syphilis serology (VDRL or RPR)	Identifies co-infection with syphilis, which is epidemiologically-linked to HIV. Disease may have accelerated course in HIV patients.
Hepatitis C serology (anti-HCV)	Identifies HCV infection and usually chronic carriage. If positive, follow with HCV genotype and HCV viral load assay. If the patient is antibody-negative and at high-risk for hepatitis, order HCV RNA to exclude a false-negative result.
Hepatitis B serologies (HBsAb, HBcAb, HBsAg)	Identifies patients who are immune to hepatitis B (HBsAb) or chronic carriers (HBsAg). If all three are negative, hepatitis B vaccine is indicated.

**Table 5.1. Baseline Laboratory Testing for HIV-Infected Patients** (cont'd)

Test	Rationale
G6PD screen	Identifies patients at risk for dapsone or primaquine-associated hemolysis.
CMV serology (IgG)	Identifies patients who should receive CMV-negative or leukocyte-depleted blood if transfused.
VZV serology (IgG)	Identifies patients at risk for varicella (chickenpox), and those who should avoid contact with active varicella or herpes zoster patients. Serology-negative patients exposed to chickenpox should receive varicella-zoster immune globulin (VZIG).
Chest x-ray	Sometimes ordered as a baseline test for future comparisons. May detect healed granulomatous diseases/other processes. Indicated in all tuberculin skin test positive patients.

**Table 5.2. Use of CD<sub>4</sub> Cell Count for Interpretation of Patient Signs/Symptoms**

CD <sub>4</sub> Cell Count (cells/mm <sup>3</sup> )	Associated Conditions
> 500	Most illnesses are similar to those in HIV-negative patients. Some increased risk of bacterial infections (pneumococcal pneumonia, sinusitis), herpes zoster, tuberculosis, skin conditions.
200–500*	Bacterial infections (especially pneumococcal pneumonia, sinusitis), cutaneous Kaposi's sarcoma, vaginal candidiasis, ITP.
50–200*	Thrush, oral hairy leukoplakia, classic HIV-associated opportunistic infections (e.g., <i>P. carinii</i> jiroveci pneumonia, cryptococcal meningitis, toxoplasmosis). For patients receiving prophylaxis, most opportunistic infection do not occur until CD <sub>4</sub> cell counts fall significantly below 100/mm <sup>3</sup> .
< 50*	"Final common pathway" opportunistic infections (disseminated <i>M. avium-intracellulare</i> , CMV retinitis), HIV-associated wasting, neurologic disease (neuropathy, encephalopathy).

\* Patients remain at risk for all processes noted in earlier stages.

---

**INDICATIONS FOR TREATMENT OF HIV INFECTION**


---

**Table 5.3. Initiation of Antiretroviral Therapy in HIV-1 Infected Patients**

Clinical Condition and/or CD <sub>4</sub> count	Recommendation
Symptomatic HIV disease	ART strongly recommended regardless of CD <sub>4</sub> cell count
Pregnant women	
HIV-1 RNA > 100,000 copies/mm <sup>3</sup>	
Rapid decline in CD <sub>4</sub> cell count, >100 cells/mm <sup>3</sup> per year	
Acute hepatitis B coinfection	
Acute hepatitis C coinfection	
Active or high risk for cardiovascular disease	
HIV associated neuropathy	
Symptomatic primary HIV infection	
High risk for secondary HIV transmission (e.g., serodiscordant couples)	
Asymptomatic (any CD <sub>4</sub> cell count)	ART is recommended

ART = antiretroviral therapy.

Adapted from: Panel on Antiretroviral Guidelines for Adults and Adolescents. Department of Health and Human Services. Guidelines for the use of antiretroviral agents in HIV-infected adults and adolescents. <http://aidsinfo.nih.gov/ConsentFiles/AdultandAdolescentGL.pdf>, 2012.

## ANTIRETROVIRAL TREATMENT

Table 5.4. ART Therapy for Treatment-Naive Patients from DHHS Guidelines 2014

Regimen	Preferred Regimen	Alternate Regimen	Acceptable Regimen
NNRTI-based (1 NNRTI + 2 NRTIs)	Efavirenz (EFV) + Tenofovir (TDF) + Emtricitabine (FTC) (Atripla)	Efavirenz (EFV) (Sustiva) <b>plus</b> Abacavir (ABC) + Lamivudine (3TC) (Epzicom)  <b>or</b> Rilpivirine (RPV) + Tenofovir (TDF) + Emtricitabine (FTC) (Complera)  <i>These regimens are recommended only when pre-ART plasma HIV RNA &lt; 100,000 copies/ml</i>	None
PI-based (1 PI + 2 NRTIs)	Atazanavir (ATV) (Reyataz)/r <b>plus</b> Tenofovir (TDF) + Emtricitabine (FTC) (Truvada)  <b>or</b> Darunavir (DRV) (Prezista)/ <b>plus</b> Tenofovir (TDF) + Emtricitabine (FTC) (Truvada)	Atazanavir (ATV) (Reyataz)/r <b>plus</b> Abacavir (ABC) + Lamivudine (3TC) (Epzicom)  <i>This regimen is recommended only when pre-ART plasma HIV RNA &lt; 100,000 copies/ml</i>	Darunavir (DRV) (Prezista)/r <b>plus</b> Abacavir (ABC) + Lamivudine (3TC) (Epzicom)  <b>or</b> Lopinavir (LPV)/r (Kaletra) (once <sup>1</sup> or twice daily) <b>plus</b> Abacavir (ABC) + Lamivudine (3TC) (Epzicom)  <b>or</b> Lopinavir (LPV)/r (Kaletra) (once <sup>1</sup> or twice daily) <b>plus</b> Tenofovir (TDF) (Viread) <b>plus</b> Lamivudine (3TC) (Epivir)  Once daily LPV/r is not recommended for pregnant patients <sup>1</sup>
INSTI-based (1 INSTI + 2 NRTIs)	Raltegravir (RAL) (Isentress) <b>plus</b> Tenofovir (TDF) + Emtricitabine (FTC) (Truvada)  <b>or</b>	Raltegravir (RAL) (Isentress) <b>plus</b> Abacavir (ABC) + Lamivudine (3TC) (Epzicom)	None



**Table 5.4. ART Therapy for Treatment-Naive Patients from DHHS Guidelines 2014 (cont'd)**

Regimen	Preferred Regimen	Alternate Regimen	Acceptable Regimen
	Elvitegravir (EVG) + Cobicistat (COBI) + Tenofovir (TDF) + Emtricitabine (FTC) (Stribild)  <b>or</b> Dolutegravir (DTG) + Abacavir (ABC) + Lamivudine (3TC) (Triumeq)  <b>or</b> Dolutegravir (DTG) + (Tivicay) <b>plus</b> Tenofovir (TDF) + Emtricitabine (FTC) (Truvada)		

**Preferred Regimens for Pregnant Women**

Two-NRTI Backbone:

- Abacavir (ABC) + Lamivudine (3TC) (Epzicom)
- Tenofovir (TDF) + Emtricitabine (FTC) (Truvada) or Lamivudine (3TC) (Epivir)
- Zidovudine (ZDV) + Lamivudine (3TC) (Kaletra)—most experience for use in pregnancy

PI-based Regimens:

- Atazanavir (ATZ) (Reyataz)/r + a Preferred Two-NRTI Backbone (listed above)
- Lopinavir (LPV)/r (Kaletra) (BID) + a Preferred Two-NRT Backbone (listed above)

NNRTI Regimen:

- Efavirenz (EFV) (Sustiva) + a Preferred TWO-NRTI Backbone (listed above); may be initiated after the first 8 weeks of pregnancy

**Alternate Regimens**

PI-based:

- Darunavir (DRV) (Prezista)/r + a Preferred Two-NRTI Backbone (listed above)
- Saquinavir (SQV) (Invirase)/r + a Preferred Two-NRTI Backbone (listed above)

NNRTI-based:

- Nevirapine (NVP) (Viramune) + Preferred Two-NRTI Backbone (listed above)

INSTI-based:

- Raltegravir (RAL) (isentress) + a Preferred Two-NRTI Backbone (listed above)

- Efavirenz should not be used during the first trimester of pregnancy or in women trying to conceive or in those not using effective and consistent contraception.
- Lamivudine may substitute for emtricitabine or vice versa.
- Abacavir should not be used in patients who test positive for HLA-B\*5701 and should be used with caution in patients at high risk for cardiovascular disease or with pretreatment HIV-RNA > 100,000 copies/ml.
- Nevirapine should not be used in patients with moderate to severe hepatic impairment. It should not be used in women with pre-ARV CD<sub>4</sub> + > 250 cells/mm<sup>3</sup> or men with pre-ARV CD<sub>4</sub> + > 400 cells/mm<sup>3</sup>.
- Atazanavir/r should not be used in patients who requires . 20 mg omeprazole per day.
- Atazanavir/r is generally preferred over atazanavir. Unboosted atazanavir may be used when ritonavir boosting is not possible.
- Lopinavir/r qd is not recommended for pregnant women.

Adapted from: Panel on Antiretroviral Guidelines for Adults and Adolescents. Department of Health and Human Services. Guidelines for the use of antiretroviral agents in HIV-infected adults and adolescents. <http://aidsinfo.nih.gov/contentfiles/vguidelines/adultandadolescentgl.pdf>. 2014.

Adapted from: Recommendations for use of antiretroviral drugs in pregnant HIV-infected women for maternal health and interventions to reduce perinatal HIV transmission in the United States. <http://aidsinfo.nih.gov/contentfiles/vguidelines/perinatalgl.pdf>. 2014.

**Table 5.5. Recommended initial Antiretroviral Therapy from IAS-USA Guidelines**

	<b>Recommended regimen</b>	<b>Alternate regimen</b>	<b>Comments</b>
NNRTI-based	Efavirenz + Tenofovir + Emtricitabine	Nevirapine + Tenofovir + Emtricitabine	Abacavir only in HLA-B*5701-negative patients with HIV-1 RNA < 100000 copies/mL. Severe hepatotoxicity and rash with nevirapine are more common in initial therapy when CD4 cell count > 250/ $\mu$ L in women and > 400/ $\mu$ L in men
	Efavirenz + Abacavir + Lamivudine	Nevirapine + Abacavir + Lamivudine	
		Rilpivirine + Tenofovir + Emtricitabine	
		Rilpivirine + Abacavir + Lamivudine	
PI-based	Darunavir/r + Tenofovir + Emtricitabine	Darunavir/r + Abacavir + Lamivudine	Other alternative PIs include fosamprenavir/r and saquinavir/r but indications to use these options for initial treatment are rare
	Atazanavir/r + Tenofovir + Emtricitabine	Lopinavir/r + Tenofovir + Emtricitabine	
	Atazanavir/r + Abacavir + Lamivudine	Lopinavir/r + Abacavir + Lamivudine	
InSTI-based	Raltegravir + Tenofovir + Emtricitabine	Raltegravir + Abacavir + Lamivudine	Raltegravir is given twice daily. Experience with elvitegravir/combicistat/tenofovir/emtricitabine is limited to 48-week data
		Elvitegravir + Combicistat + Tenofovir + Emtricitabine	

Adapted from: Thompson MA, Aberg JA, Hoy JF, et al. Antiretroviral treatment of adult HIV infection: 2012 recommendations of the International Antiviral Society-USA Panel. *JAMA* 308:387–402, 2012.

**Selection of Optimal Antiretroviral Therapy.** Selection of the optimal initial antiretroviral regimen must take into consideration antiviral potency, tolerability, and safety. In the DHHS and IAS-USA Guidelines (Tables 5.4 and 5.5).

---

**ANTIRETROVIRAL TREATMENT FAILURE (Table 5.6)**

---

Antiretroviral treatment failure can be defined in various ways, as described below. Causes of treatment failure include inadequate adherence, preexisting drug resistance, regimen complexity, side effects, and suboptimal pharmacokinetics. All of these factors can lead to persistent viral replication and evolution of drug resistance. Poor medication adherence is the most common cause of treatment failure.

**A. Types of Treatment Failure**

- 1. Virologic Failure** is most strictly defined as the inability to achieve or maintain virologic suppression. In a treatment-naïve patient, the HIV RNA level should be < 400 copies/mL after 24 weeks or < 50 copies/mL by 48 weeks after starting therapy. Virologic rebound is seen when there is repeated detection of HIV RNA after virologic suppression in either treatment-naïve or treatment-experienced patients.
- 2. Immunologic Failure** can occur in the presence or absence of virologic failure and is defined as a failure to increase the CD<sub>4</sub> cell count by 25–50 cells/mm<sup>3</sup> above baseline during the first year of therapy, or as a decrease in CD<sub>4</sub> cell count to below baseline count while on therapy.
- 3. Clinical Failure** is the occurrence or recurrence of HIV-related events after at least 3 months on potent antiretroviral therapy, excluding events related to an immune reconstitution syndrome.
- 4. Usual Sequence of Treatment Failure.** Virologic failure usually occurs first, followed by immunologic failure, and finally by clinical progression. These events may be separated by months or years and may not occur in this order in all patients.

**B. Goals After Virologic Failure.** When patients have detectable HIV RNA on treatment, clinicians should attempt to identify the cause of their lack of response and set a treatment goal of achieving full virologic suppression (HIV RNA < 50 copies/mL). In addition to improving clinical and immunologic outcomes, this strategy will also prevent the selection of additional resistance mutations. Provided that medication adherence issues and regimen tolerability have been addressed, the regimen should be changed sooner than later. On the other hand, achieving an undetectable HIV RNA level in patients with an extensive prior treatment history may not be possible. The main goals in these patients should be partial suppression of HIV RNA below the pretreatment level to preserve immune function and prevent clinical progression.

**C. Resistance Testing and Selection of New Antiretroviral Therapy.** Genotypic assays characterize nucleotide sequences of the reverse transcriptase/protease portions of the virus, and identify resistance mutations associated with various drugs. Phenotypic assays

attempt to grow the virus in the presence of drugs, providing a more intuitively applicable measurement of resistance (similar to that done with bacteria). Compared to phenotypic assays, genotypic assays are faster (1–2 weeks vs. 2–4 weeks for results), less expensive (\$400 vs. \$1000), and have less inter-laboratory variability; however, mutations do not always correlate with resistance and results are difficult to interpret.

**Table 5.6. Management of Antiretroviral Treatment Failure**

Type of Failure	Recommended Approach	Comments
<i>Virologic failure Limited or intermediate prior treatment</i>	Assess for adherence and regimen tolerability. Obtain genotype resistance test. Select new regimen based on resistance test results and tolerability.	Usually associated with limited or no detectable resistance. If no resistance is found, consider re-testing for resistance 2–4 weeks after resuming antvirals. Stop NNRTI's if resistance is detected. Virologic suppression is likely if adherence is good.
<i>Extensive prior treatment</i>	Assess for adherence and regimen tolerability. Obtain resistance test – consider phenotype, “virtual phenotype,” or phenotype-genotype combination if level of resistance is likely to be high. Obtain viral tropism assay to assess possible use of CCR5 antagonist. Select new regimen using at least 2 new active agents; if 2 new active agents not available, continue a “holding” regimen.	In patients with resistance to NRTI's, NNRTI's and PI's, the new regimen should generally contain at least: (1) at least one drug from a new drug class (integrase inhibitor, CCR5 antagonist, or fusion inhibitor); (2) a boosted PI with activity against resistant viruses (tipranavir or darunavir); and (3) one or two NRTI's, one of them 3TC or FTC. A holding regimen should always contain 3TC or FTC plus a boosted PI; NNRTI's should never be used.
<i>Low-level HIV RNA (50–1000 copies)</i>	Assess for adherence, drug-drug interactions, intercurrent illness, recent immunizations. Repeat test in 3–4 weeks.	For low-level viremia followed by undetectable HIV RNA (“blip”), no treatment change is necessary. If HIV RNA is persistently detectable at > 500 copies/mL, obtain resistance test as described above, and treat accordingly. If HIV RNA is persistently detectable at 50–500 copies/mL, consider regimen “intensification” with use of an additional agent.

**Table 5.6. Management of Antiretroviral Treatment Failure** (cont'd)

Type of Failure	Recommended Approach	Comments
<u>Immunologic failure</u> <i>Detectable HIV RNA</i>	Assess for adherence and tolerability. If non-adherent, resume treatment after barriers to adherence are addressed. If adherent, obtain resistance testing and alter therapy as described above.	If HIV RNA is back to pre-treatment baseline, non-adherence is the most likely explanation.
<i>Suppressed HIV RNA</i>	Investigate for modifiable conditions that may be associated with impaired CD <sub>4</sub> response (chronic HCV, treatment with ZDV, TDF + ddI). If no modifiable conditions found, continue current regimen.	Prognosis for patients with suppressed HIV RNA and immunologic failure better than for those with comparable CD <sub>4</sub> cell counts and detectable viremia.
<u>Clinical failure</u> <i>Detectable HIV RNA</i>	Treat OI with appropriate anti-infective therapy. Assess for antiretroviral adherence and tolerability. Send resistance test and choose new regimen based on results of test and other treatment options.	OI's (IRIS excluded) most commonly occur in those not on antiretroviral therapy due to poor compliance and/or regimen tolerability.
<i>Suppressed HIV RNA</i>	Continue current antiretrovirals. Treat OI with appropriate anti-infective therapy. If symptoms persist and IRIS is likely, use adjunctive corticosteroids.	IRIS most likely when baseline CD <sub>4</sub> cell count is low (< 200/mm <sup>3</sup> ); onset usually weeks-to-months after starting a potent regimen. IRIS been reported with virtually all OI's. True clinical progression with suppressed HIV RNA is unusual; IRIS should not be considered a sign of treatment failure.

IRIS = immune reconstitution inflammatory syndrome, OI = opportunistic infection.

### PROPHYLAXIS OF OPPORTUNISTIC INFECTIONS IN HIV (Tables 5.7 and 5.8)

Patients with HIV disease are at risk for infectious complications not otherwise seen in immunocompetent patients. Such opportunistic infections occur in proportion to the severity of immune system dysfunction (reflected by CD<sub>4</sub> cell count depletion). While community acquired infections (e.g., pneumococcal pneumonia) can occur at any CD<sub>4</sub> cell count, "classic" HIV-related opportunistic infections (PCP, toxoplasmosis, cryptococcus, disseminated *M. avium-intracellulare*, CMV) do not occur until CD<sub>4</sub> cell counts are dramatically reduced. Specifically, it is rare to encounter PCP in HIV patients with CD<sub>4</sub> > 200/mm<sup>3</sup>, and CMV and disseminated MAI occur at median CD<sub>4</sub> cell counts < 50/mm<sup>3</sup>.

**Table 5.7. Overview of Prophylaxis (See Table 5.8 for details)**

Infection	Indication	Intervention
PCP	CD <sub>4</sub> < 200/mm <sup>3</sup>	TMP-SMX
TB ( <i>M. tuberculosis</i> )	PPD > 5 mm (current or past) or contact with active case	INH
Toxoplasma	IgG Ab (+) and CD <sub>4</sub> < 100/mm <sup>3</sup>	TMP-SMX
MAI	CD <sub>4</sub> < 50/mm <sup>3</sup>	Azithromycin or Clarithromycin
<i>S. pneumoniae</i>	CD <sub>4</sub> > 200/mm <sup>3</sup>	Pneumococcal vaccine
Hepatitis B (HBV)	Susceptible patients	Hepatitis B vaccine
Hepatitis A (HAV)	HCV (+) and HA Ab (-); HCV (-) and HA Ab (-) gay men and travelers to endemic areas, chronic liver disease	Hepatitis A vaccine
Influenza	All patients	Annual flu vaccine Swine influenza (H <sub>1</sub> N <sub>1</sub> ) vaccine
VZV	Exposure to chickenpox or shingles; no prior history	VZIG

Abbreviations: Ab = antibody; HAV = Hepatitis A virus; HCV = Hepatitis C virus; VZIG = varicella-zoster immune globulin; VZV = varicella-zoster virus; other abbreviations (p. xi).

**Table 5.8. Prophylaxis of Opportunistic Infections in HIV**

Infection	Indications and Prophylaxis	Comments
P. (carinii) jiroveci pneumonia (PCP)	<p><u>Indications:</u> CD<sub>4</sub> &lt; 200/mm<sup>3</sup>, oral thrush, constitutional symptoms, or previous history of PCP</p> <p><u>Preferred prophylaxis:</u> TMP-SMX 1 DS tablet (PO) q24h or 1 SS tablet (PO) q24h. 1 DS tablet (PO) 3x/week is also effective, but daily dosing may be slightly more effective</p> <p><u>Alternate prophylaxis:</u> Dapsone 100 mg (PO) q24h (preferred as second-line by most; more effective than aerosolized pentamidine when CD<sub>4</sub> cell count &lt; 100)</p> <p style="text-align: center;"><b>or</b></p> <p>Atovaquone 1500 mg (PO) q24h (comparably effective to dapsone and aerosolized pentamidine; more GI toxicity vs. dapsone, but less rash)</p> <p style="text-align: center;"><b>or</b></p> <p>Aerosolized pentamidine 300 mg via Respigard II nebulizer once monthly (exclude active pulmonary TB first to avoid nosocomial transmission)</p>	<p>Without prophylaxis, 80% of AIDS patients develop PCP, and 60–70% relapse within one year after the first episode. Prophylaxis with TMP-SMX also reduces the risk for toxoplasmosis and possibly bacterial infections. Among patients with prior non-life-threatening reactions to TMP-SMX, 55% can be successfully rechallenged with 1 SS tablet daily, and 80% can be rechallenged with gradual dose escalation using TMP-SMX elixir (8 mg TMP + 40 mg SMX/mL) given as 1 mL × 3 days, then 2 mL × 3 days, then 5 mL × 3 days, then 1 SS tablet (PO) q24h. Macrolide-regimens for MAI (azithromycin, clarithromycin) add to efficacy of PCP prophylaxis. Primary and secondary prophylaxis may be discontinued if CD<sub>4</sub> cell counts increase to &gt; 200 cells/mm<sup>3</sup> for 3 months or longer in response to antiretroviral therapy (i.e., immune reconstitution). Prophylaxis should be resumed if the CD<sub>4</sub> cell count decreases to &lt; 200/mm<sup>3</sup>.</p>
Toxoplasmosis	<p><u>Indications:</u> CD<sub>4</sub> &lt; 100/mm<sup>3</sup> with positive toxoplasmosis serology (IgG)</p> <p><u>Preferred prophylaxis:</u> TMP-SMX 1 DS tablet (PO) q24h</p> <p><u>Alternate prophylaxis:</u> Dapsone 50 mg (PO) q24h + pyrimethamine 50 mg (PO) weekly + folinic acid 25 mg (PO) weekly</p> <p style="text-align: center;"><b>or</b></p>	<p>Incidence of toxoplasmosis in seronegative patients is too low to warrant chemoprophylaxis. Primary prophylaxis can be discontinued if CD<sub>4</sub> cell counts increase to &gt; 200/mm<sup>3</sup> for at least 3 months in response to antiretroviral therapy. Secondary prophylaxis (chronic maintenance therapy) may be discontinued in patients who responded to initial therapy, remain asymptomatic, and whose CD<sub>4</sub> counts increase to &gt; 200/mm<sup>3</sup> for 6 months or longer in response to</p>

**Table 5.8. Prophylaxis of Opportunistic Infections in HIV** (cont'd)

Infection	Indications and Prophylaxis	Comments
	<p>Dapsone 100 mg/pyrimethamine 50 mg twice weekly (no folinic acid)</p> <p style="text-align: center;"><b>or</b></p> <p>Atovaquone 1500 mg (PO) q24h</p>	<p>antiretroviral therapy. Prophylaxis should be restarted if the CD<sub>4</sub> count decreases to &lt; 200/mm<sup>3</sup>. Some experts would obtain an MRI of the brain as part of the evaluation prior to stopping secondary prophylaxis.</p>
M. tuberculosis (TB)	<p><u>Indications:</u> PPD induration ≥ 5 mm or history of positive PPD without prior treatment, or close contact with active case of TB. Must exclude active disease (chest x-ray mandatory). Indicated at any CD<sub>4</sub> cell count</p> <p><u>Preferred prophylaxis:</u> INH 300 mg (PO) q24h × 9 months + pyridoxine 50 mg (PO) q24h × 9 months</p> <p><u>Alternate prophylaxis:</u> Rifampin 600 mg (PO) q24h × 2 months + pyrazinamide 20 mg/kg (PO) q24h × 2 months. Rifampin should not be given to patients receiving PI's</p>	<p>Consider prophylaxis for skin test negative patients when the probability of prior TB exposure is &gt; 10% (e.g., patients from developing countries, IV drug abusers in some cities, prisoners). However, a trial testing this strategy in the U.S. did not find a benefit for empiric prophylaxis. INH prophylaxis delayed progression to AIDS and prolonged life in Haitian cohort with positive PPD treated × 6 months. Rifampin plus pyrazinamide × 2 months was effective in a multinational clinical trial (but the combination may ↑ hepatotoxicity). Rifabutin may be substituted for rifampin in rifampin-containing regimens.</p>
M. avium-intracellulare (MAI)	<p><u>Indications:</u> CD<sub>4</sub> &lt; 50/mm<sup>3</sup></p> <p><u>Preferred prophylaxis:</u> Azithromycin 1200 mg (PO) once a week (fewest number of pills; fewest drug interactions; may add to efficacy of PCP prophylaxis)</p> <p style="text-align: center;"><b>or</b></p> <p>Clarithromycin 500 mg (PO) q12h (more effective than rifabutin; associated with survival advantage; resistance detected in some breakthrough cases)</p> <p><u>Alternate prophylaxis:</u> Rifabutin (less effective). See TB (p. 326) for dosing</p>	<p>Macrolide options (azithromycin, clarithromycin) preferable to rifabutin. Azithromycin is preferred for patients on protease inhibitors (fewer drug interactions). Primary prophylaxis may be discontinued if CD<sub>4</sub> cell counts increase to &gt; 100/mm<sup>3</sup> and HIV RNA suppresses for 3–6 months or longer in response to antiretroviral therapy. Secondary prophylaxis may be discontinued for CD<sub>4</sub> cell counts that increase to &gt; 100/mm<sup>3</sup> × 6 months or longer in response to antiretroviral therapy if patients have completed 12 months of MAI therapy and have no evidence of disease. Resume MAI prophylaxis for CD<sub>4</sub> &lt; 100/mm<sup>3</sup>.</p>



**Table 5.8. Prophylaxis of Opportunistic Infections in HIV (cont'd)**

Infection	Indications and Prophylaxis	Comments
Pneumococcus ( <i>S. pneumoniae</i> )	<u>Indications:</u> CD <sub>4</sub> > 200/mm <sup>3</sup> <u>Preferred prophylaxis:</u> Pneumococcal polysaccharide (23 valent) vaccine.* Re-vaccinate × 1 at 5 years	Incidence of invasive pneumococcal disease is > 100-fold higher in HIV patients. Efficacy of vaccine is variable in clinical studies.
Influenza (human seasonal)	<u>Indications:</u> Generally recommended for all patients <u>Preferred prophylaxis:</u> Influenza vaccine (inactivated whole virus and split virus vaccine)*	Give annually (optimally October–January). Some experts do not give vaccine if CD <sub>4</sub> is < 100/mm <sup>3</sup> (antibody response is poor). New intranasal live virus vaccine is contraindicated in immunosuppressed patients.
Hepatitis B (HBV)	<u>Indications:</u> All susceptible (anti-HBcAb negative and anti-HBsAg negative) patients <u>Preferred prophylaxis:</u> Hepatitis B recombinant DNA vaccine*	Response rate is lower than in HIV-negative controls. Repeat series if no response, especially if CD <sub>4</sub> was low during initial series and is now increased.
Hepatitis A (HAV)	<u>Indications:</u> All susceptible patients who are also infected with hepatitis C; HAV-susceptible seronegative gay men or travelers to endemic areas; chronic liver disease; illegal drug users <u>Preferred prophylaxis:</u> Hepatitis A vaccine*	Response rate is lower than in HIV-negative controls.
Measles, mumps, rubella	<u>Indications:</u> Patients born after 1957 and never vaccinated; patients vaccinated between 1963–1967 <u>Preferred prophylaxis:</u> MMR (measles, mumps, rubella) vaccine*	Single case of vaccine-strain measles pneumonia in severely immuno-compromised adult who received MMR; vaccine is therefore contraindicated in patients with severe immunodeficiency (CD <sub>4</sub> < 200).

\* Same dose as for normal hosts (see pp. 372–375). If possible, give vaccines early in course of HIV infection, while immune system may still respond. Alternatively, to increase the likelihood of response in patients with advanced HIV disease, vaccines may be administered after 6–12 months of effective antiretroviral therapy. Vaccines should be given when patients are clinically stable, not acutely ill (e.g., give during a routine office visit, rather than during hospitalization for an opportunistic infection). Live vaccines (e.g., oral polio, oral typhoid, Yellow fever) are generally contraindicated, but measles vaccine is well-tolerated in children, and MMR vaccine is recommended for adults as described above.

**Table 5.8. Prophylaxis of Opportunistic Infections in HIV** (cont'd)

Infection	Indications and Prophylaxis	Comments
H. influenzae	<u>Indications:</u> Not generally recommended for adults  <u>Preferred prophylaxis:</u> H. influenzae type B polysaccharide vaccine*	Incidence of H. influenzae disease is increased in HIV patients, but 65% are caused by non-type B strains. Unclear whether vaccine offers protection.
Travel vaccines*	<u>Indications:</u> Travel to endemic areas	All considered safe except oral polio, yellow fever, and live oral typhoid—each a live virus vaccine.

\* Same dose as for normal hosts (see pp. 372–375). If possible, give vaccines early in course of HIV infection, while immune system may still respond. Alternatively, to increase the likelihood of response in patients with advanced HIV disease, vaccines may be administered after 6–12 months of effective antiretroviral therapy. Vaccines should be given when patients are clinically stable, not acutely ill (e.g., give during a routine office visit, rather than during hospitalization for an opportunistic infection). Live vaccines (e.g., oral polio, oral typhoid, Yellow fever) are generally contraindicated, but measles vaccine is well-tolerated in children, and MMR vaccine is recommended for adults as described above.

## TREATMENT OF OPPORTUNISTIC INFECTIONS IN HIV

Antiretroviral therapy (ART) and specific antimicrobial prophylaxis regimens have led to a dramatic decline in HIV-related opportunistic infections. Today, opportunistic infections occur predominantly in patients not receiving ART (due to undiagnosed HIV infection or nonacceptance of therapy), or in the period after starting ART (due to lack of immune reconstitution or from eliciting a previously absent inflammatory host response). Despite high rates of virologic failure in clinical practice, the rate of opportunistic infections in patients compliant with ART remains low, presumably due to continued immunologic response despite virologic failure, a phenomenon that may be linked to impaired “fitness” (virulence) of resistant HIV strains. For patients on or off ART, the absolute CD<sub>4</sub> cell count provides the best marker of risk for opportunistic infections.

## Respiratory Tract Opportunistic Infections

### Aspergillosis, Invasive Pulmonary

Pathogen	Preferred Therapy	Alternate Therapy
Aspergillus fumigatus (rarely other species)	Voriconazole 400 mg (IV or PO) q12h × 2 days, then 200 mg (IV or PO) q12h until cured (typically 6–18 months)	Amphotericin B 1 mg/kg (IV) q24h until 2–3 gm total dose given (optimal duration of therapy poorly defined)  <b>or</b> Lipid amphotericin (Abelcet or Ambisome) 5 mg/kg/day (IV) until cured

**Clinical Presentation:** Pleuritic chest pain, hemoptysis, cough in a patient with advanced HIV disease. Additional risk factors include neutropenia and use of corticosteroids.

**Diagnostic Considerations:** Diagnosis by bronchoscopy with biopsy/culture. Open lung biopsy (usually video-assisted thorascopic surgery) is sometimes required. Radiographic appearance includes cavitation, nodules, sometimes focal consolidation. Dissemination to CNS may occur, and manifests as focal neurological deficits.

**Pitfalls:** Positive sputum culture for *Aspergillus* in advanced HIV disease should heighten awareness of possible infection.

**Therapeutic Considerations:** Decrease/discontinue corticosteroids, if possible. If present, treat neutropenia with granulocyte-colony stimulating factor (G-CSF) to achieve absolute neutrophil count > 1000/mm<sup>3</sup>. There are insufficient data to recommend chronic suppressive or maintenance therapy.

**Prognosis:** Poor unless immune deficits can be corrected.

## Bacterial Pneumonia

Usual Pathogens	Preferred Therapy	Comments
Streptococcus pneumoniae (most common) Haemophilus influenzae Pseudomonas aeruginosa	<p><b>Monotherapy with</b> Levofloxacin 750 mg (PO/IV) q24h or Moxifloxacin 400 mg (PO/IV) q24h × 7–14 days, depending on severity</p> <p><b>or combination therapy with</b> Ceftriaxone 1–2 gm (IV) q24h (or Cefotaxime 1 gm [IV] q8h) <b>plus</b> Azithromycin 500 gm q24h × 7–14 days</p>	For severe immunodeficiency (CD <sub>4</sub> < 100/mm <sup>3</sup> ), neutropenia, or a prior history of pseudomonas infection, broaden coverage to include <i>P. aeruginosa</i> and other gram-negative bacilli <u>by adding either</u> Ceftazidime 1 gm (IV) q8h <b>or</b> Cefepime 1 gm (IV) q12h <b>or</b> Ciprofloxacin 750 mg (PO) q12h or 400 mg (IV) q8h.

**Clinical Presentation:** HIV-infected patients with bacterial pneumonia present similar to those without HIV, with a relatively acute illness (over days) that is often associated with chills, rigors, pleuritic chest pain, and purulent sputum. Patients who have been ill over weeks to months are more likely have PCP, tuberculosis, or a fungal infection. Since bacterial pneumonia can occur at any CD<sub>4</sub> cell count, this infection is frequently the presenting symptom of HIV disease, prompting initial HIV testing and diagnosis.

**Diagnostic Considerations:** The most common pathogens are *Streptococcus pneumoniae*, followed by *Haemophilus influenzae*. The pathogens of atypical pneumonia (*Legionella pneumophila*, *Mycoplasma pneumoniae*, and *Chlamydia pneumoniae*) are rarely encountered, even with extensive laboratory investigation. A lobar infiltrate on chest radiography is a further predictor of bacterial pneumonia. Blood cultures should be obtained, as HIV patients have an increased rate of bacteremia compared to those without HIV.

**Pitfalls:** Sputum gram stain and culture are generally only helpful if collected prior to starting antibiotics, and only if a single organism predominates. HIV patients with bacterial pneumonia may rarely have a more subacute opportunistic infection concurrently, such as PCP or TB.

**Therapeutic Considerations:** Once improvement has occurred, a switch to oral therapy is generally safe. Patients with advanced HIV disease are at greater risk of bacteremic pneumonia due to gram-negative bacilli, and should be covered empirically for this condition.

**Prognosis:** Response to therapy is generally prompt and overall prognosis is good.

## Pneumocystis (carinii) jiroveci Pneumonia (PCP)

Subset	Preferred Therapy	Alternate Therapy
Severe disease (pO <sub>2</sub> < 70 mmHg, A-a gradient > 35)	TMP-SMX (5 mg/kg TMP) (IV) q6h × 21 days  <b>plus</b> Prednisone 40 mg (PO) q12h on days 1–5, then 40 mg (PO) q24h on days 6–10, then 20 mg (PO) q24h on days 11–21. Methylprednisolone (IV) can be substituted at 75% of prednisone dose	Pentamidine 4 mg/kg (IV) q24h (infused over ≥ 60 minutes) × 21 days <i>plus</i> prednisone × 21 days. (Dose reduction of pentamidine to 3 mg/kg [IV] q24h may reduce toxicity)  <b>or</b> Primaquine 30 mg (PO) q24h × 21 days <i>plus</i> clindamycin 600 mg (PO) q8h × 21 days
Mild or moderate disease (pO <sub>2</sub> > 70 mmHg, A-a gradient < 35)	TMP-SMX 2 DS tablets (PO) q6h × 21 days	Dapsone 100 mg (PO) q24h × 21 days <i>plus</i> TMP 5 mg/kg (PO) q8h × 21 days  <b>or</b> Primaquine 30 mg (PO) q24h × 21 days <i>plus</i> clindamycin 600 mg (PO) q8h × 21 days  <b>or</b> Atovaquone 750 mg (PO) q12h (with food) × 21 days

**Clinical Presentation:** Fever, cough, dyspnea; often indolent presentation. Progressive SOB usually over a week. Physical exam is usually normal. Chest x-ray is variable, but commonly shows a diffuse interstitial pattern. High A-a gradient with exercise desaturation.

**Diagnostic Considerations:** Diagnosis by immunofluorescent stain of induced sputum or bronchoscopy specimen. Highly elevated LDH.  $\beta$  1,3 D-glucan +, aspergillus galactomannan –.

**Pitfalls:** Slight worsening of symptoms is common after starting therapy, especially if not treated with steroids. Do not overlook superimposed bacterial pneumonia or other secondary infections, especially while on pentamidine. Patients receiving second-line agents for PCP prophylaxis—in particular aerosolized pentamidine—may present with atypical radiographic findings, including apical infiltrates, multiple small-walled cysts, pleural effusions, pneumothorax, or single/multiple nodules.

**Therapeutic Considerations:** Outpatient therapy is possible for mild disease, but only when close follow-up is assured. Adverse reactions to TMP-SMX (rash, fever, GI symptoms, hepatitis, hyperkalemia, leukopenia, hemolytic anemia) occur in 25–50% of patients, many of whom will need a second-line regimen to complete therapy (e.g., trimethoprim-dapsone or atovaquone). Unless an adverse reaction to TMP-SMX is particularly severe (e.g., Stevens-Johnson syndrome or other life-threatening problem), TMP-SMX may be considered for PCP prophylaxis, since prophylaxis requires a much lower dose (only 10–15% of treatment dose). Patients being treated for severe PCP with TMP-SMX who do not improve after one week may be switched to pentamidine, although there are no prospective data to confirm this approach. In general, patients receiving antiretroviral therapy when PCP develops should have their treatment continued, since intermittent antiretroviral therapy can lead to drug resistance. For newly-diagnosed or antiretroviral-naïve HIV patients, treatment of PCP may be completed before starting antiretroviral therapy. Steroids should be tapered, not discontinued

abruptly. Adjuvanted steroids increase the risk of thrush/herpes simplex infection, but probably not CMV, TB, or disseminated fungal infection.

**Prognosis:** Usually responds to treatment. Adverse prognostic factors include ↑ A-a gradient, hypoxemia, ↑ LDH.

## Pulmonary Tuberculosis (for isolates sensitive to INH and rifampin)

Pathogen	Patients NOT Receiving PI's or NNRTI's	Patients Receiving PI's or NNRTI's*
Mycobacterium tuberculosis (TB)	<u>Initial phase (8 weeks)</u> INH 300 mg (PO) q24h <b>plus</b> Rifampin† 600 mg (PO) q24h <b>plus</b> Pyrazinamide (PZA) 25 mg/kg (PO) q24h <b>plus</b> Ethambutol (EMB) 15–20 mg/kg (PO) q24h  <u>Continuation phase (18 weeks)</u> INH 300 mg (PO) q24h <b>plus</b> Rifampin† 600 mg (PO) q24h	<u>Initial phase (8 weeks)</u> INH 300 mg (PO) q24h <b>plus</b> Rifabutin* <b>plus</b> PZA 25 mg/kg (PO) q24h <b>plus</b> EMB 15 mg/kg (PO) q24h × 8 weeks.  <u>Continuation phase (18 weeks)</u> INH 300 mg (PO) q24h <b>plus</b> Rifabutin*

\* Rifabutin dose: If PI is nelfinavir, indinavir, amprenavir or fosamprenavir, then rifabutin dose is 150 mg (PO) q24h. If PI is ritonavir, lopinavir/ritonavir or atazanavir, then rifabutin dose is 150 mg (PO) 2–3 times weekly. If NNRTI is efavirenz, then rifabutin dose is 450 mg (PO) q24h or 600 mg (PO) 2–3 times weekly. If NNRTI is nevirapine, then rifabutin dose is 300 mg (PO) q24h. Rifabutin is **contraindicated** in patients receiving delavirdine rilpivirine, etravirine + ritonavir boosted PI or hard-gel saquinavir. Patients receiving PI's AND NNRTI's: as above, except adjust rifabutin to 300 mg (PO) q24h.

† For patients receiving triple NRTI regimens, substitute rifabutin 300 mg (PO) q24h for rifampin.

**Clinical Presentation:** May present atypically. HIV patients with high (> 500/mm<sup>3</sup>) CD<sub>4</sub> cell counts are more likely to have a typical pulmonary presentation, but patients with advanced HIV disease may have a diffuse interstitial pattern, hilar adenopathy, or a normal chest x-ray. Tuberculin skin testing (TST) is helpful if positive, but unreliable if negative due to anergy.

**Diagnostic Considerations:** In many urban areas, TB is one of the most common HIV-related respiratory illnesses. In other areas, HIV-related TB occurs infrequently except in immigrants or patients arriving from highly TB endemic areas. Maintain a high Index of suspicion for TB in HIV patients with unexplained fevers/pulmonary infiltrates.

**Pitfalls:** Extrapulmonary and pulmonary TB often coexist, especially in advanced HIV disease.

**Therapeutic Considerations:** Treatment by directly observed therapy (DOT) is strongly recommended for all HIV patients. If patients have cavitary disease or either positive sputum cultures or lack of clinical response at 2 months, total duration of therapy should be increased up to 9 months. If hepatic transaminases are elevated (AST > 3 times normal) before treatment initiation, treatment options include: (1) standard therapy with frequent monitoring; (2) rifamycin (rifampin or rifabutin) + EMB + PZA for 6 months; or

(3) INH + rifamycin + EMB for 2 months, then INH + rifamycin for 7 months. Once-weekly rifapentine is not recommended for HIV patients. Non-severe immune reconstitution inflammatory syndrome (IRIS) may be treated with nonsteroidal anti-inflammatory drugs (NSAIDs); severe cases should be treated with corticosteroids. In all cases of IRIS, antiretroviral therapy should be continued if possible. Monitor carefully for signs of rifabutin drug toxicity (arthralgias, uveitis, leukopenia).

**Prognosis:** Usually responds to treatment. Relapse rates are related to the degree of immunosuppression and local risk of re-exposure to TB.

## CNS Opportunistic Infections

### CMV Retinitis

Preferred Therapy, Duration of Therapy, Chronic Maintenance	Alternate Therapy	Other Options/Issues
<p><u>Preferred therapy for CMV retinitis</u>  <i>For immediate sight-threatening lesions</i>            Ganciclovir intraocular implant + valganciclovir 900 mg (PO) q12h for 14–21 days, then q24h daily            One dose of intravitreal ganciclovir may be administered immediately after diagnosis until ganciclovir implant can be placed</p> <p><i>For small peripheral lesions</i>            Valganciclovir 900 mg (PO) q12h × 14–21 days, then 900 mg (PO) q24h</p>	<p><u>Alternative therapy for CMV retinitis</u>            Ganciclovir 5 mg/kg (IV) q12h × 14–21 days, then 5 mg/kg (IV) q24h;  <b>or</b>            Ganciclovir 5 mg/kg (IV) q12h × 14–21 days, then valganciclovir 900 mg (PO) q24h  <b>or</b>            Foscarnet 60 mg/kg (IV) q8h or 90 mg/kg (IV) q12h × 14–21 days, then 90–120 mg/kg (IV) q24h;  <b>or</b>            Cidofovir 5 mg/kg (IV) q week × 2 weeks, then 5 mg/kg (IV) every other week with saline hydration before and after therapy plus probenecid 2 g (PO) 3 hours before the dose followed by 1 g (PO) 2 hours after the dose, and 1 g (PO) 8 hours after the dose (total of 4 g)  <b>Note:</b> This regimen should be avoided in patients with sulfa allergy because of cross hypersensitivity with probenecid</p>	<p>The choice of initial therapy for CMV retinitis should be individualized, based on location and severity of the lesion(s), level of immunosuppression, and other factors such as concomitant medications and ability to adhere to treatment</p> <p>Initial therapy in patients with CMV retinitis, esophagitis, colitis, and pneumonitis should include initiation or optimization of ART</p> <p>In patients with CMV neurological disease, localized morbidity might occur because of IRIS, a brief delay in initiation of ART until clinical improvement might be prudent</p> <p>Maintenance therapy for CMV retinitis can be safely discontinued in patients with inactive disease and sustained CD<sub>4</sub>+ count</p>

## CMV Retinitis (cont'd)

Preferred Therapy, Duration of Therapy, Chronic Maintenance	Alternate Therapy	Other Options/Issues
<p><u>Preferred chronic maintenance therapy (secondary prophylaxis) for CMV retinitis</u> Valganciclovir 900 mg (PO) daily</p> <p><b>or</b></p> <p>Ganciclovir implant (may be replaced every 6–8 months if CD<sub>4</sub>+ count remains &lt; 100 cells/<math>\mu</math>L) + Valganciclovir 900 mg (PO) q24h until immune recovery</p>	<p><u>Alternative chronic maintenance (secondary prophylaxis) for CMV retinitis</u> Ganciclovir 5 mg/kg IV 5–7 times weekly</p> <p><b>or</b></p> <p>Foscarnet 90–120 mg/kg (IV) q24h</p> <p><b>or</b></p> <p>Cidofovir 5 mg/kg (IV) of other week with saline hydration and probenecid (as above)</p>	<p>(&gt; 100 cells/mm<sup>3</sup> for <math>\geq</math> 3–6 months); consultation with ophthalmologist is advised</p> <p>Patients with CMV retinitis who discontinued maintenance therapy should undergo regular eye examination, optimally every 3 months, for early detection of relapse or immune recovery uveitis (IRU)</p> <p>IRU might develop in the setting of immune reconstitution.</p> <p><u>Treatment of IRU</u>: periocular corticosteroid or short courses of systemic steroid</p>

## CMV Encephalitis

Pathogen	Preferred Therapy	Alternate Therapy
Cytomegalovirus (CMV)	<p><u>Acute therapy</u> Ganciclovir (GCV) 5 mg/kg (IV) q12h until symptomatic improvement (typically &gt; 3 weeks). <i>For severe cases</i>, consider acute therapy with Ganciclovir 5 mg/kg (IV) q12h plus Foscarnet 90 mg/kg (IV) q24h until improvement</p> <p><u>Follow with lifelong suppressive therapy</u> Valganciclovir 900 mg (PO) q24h</p>	<p><u>Acute therapy</u> Foscarnet 60 mg/kg (IV) q8h or 90 mg/kg (IV) q12h <math>\times</math> 3 weeks</p> <p><u>Follow with lifelong suppressive therapy</u> Valganciclovir 900 mg (PO) q24h</p>

### CMV Retinitis

**Clinical Presentation:** Blurred vision, scotomata, field cuts common. Often bilateral, even when initial symptoms are unilateral.

**Diagnostic Considerations:** Diagnosis by characteristic hemorrhagic (“tomato soup and milk”) retinitis on funduscopic exam. Consult ophthalmology in suspected cases.

**Pitfalls:** May develop immune reconstitution vitritis after starting antiretroviral therapy.

**Therapeutic Considerations:** Oral valganciclovir is the preferred option for initial and maintenance therapy. Lifelong maintenance therapy for CMV retinitis is required for  $CD_4$  counts  $< 100/mm^3$ , but may be discontinued if  $CD_4$  counts increase to  $> 100\text{--}150/mm^3$  for 6 or more months in response to antiretroviral therapy (in consultation with ophthalmologist). Patients with CMV retinitis who discontinue therapy should undergo regular eye exams to monitor for relapse. Ganciclovir intraocular implants might need to be replaced every 6–8 months for patients who remain immunosuppressed with  $CD_4 < 100\text{--}150/mm^3$ . Immune recovery uveitis (IRU) may develop in the setting of immune reconstitution due to ART and be treated by ophthalmologist with periocular corticosteroid, sometimes systemic corticosteroid.

**Prognosis:** Good initial response to therapy. High relapse rate unless  $CD_4$  improves with antiretroviral therapy.

### **CMV Encephalitis**

**Clinical Presentation:** Encephalitis presents as fever, mental status changes, and headache evolving over 1–2 weeks. True meningismus is rare. CMV encephalitis occurs in advanced HIV disease ( $CD_4 < 50/mm^3$ ), often in patients with prior CMV retinitis. Polyradiculitis presents as rapidly evolving weakness/sensory disturbances in the lower extremities, often with bladder/bowel incontinence. Anesthesia in “saddle distribution” with ↓ sphincter tone possible.

**Diagnostic Considerations:** CSF may show lymphocytic or neutrophilic pleocytosis; glucose is often decreased. For CMV encephalitis, characteristic findings on brain MRI include confluent periventricular abnormalities with variable degrees of enhancement. Diagnosis is confirmed by CSF CMV PCR (preferred), CMV culture, or brain biopsy.

**Pitfalls:** For CMV encephalitis, a wide spectrum of radiographic findings are possible, including mass lesions (rare). Obtain ophthalmologic evaluation to exclude active retinitis. For polyradiculitis, obtain sagittal MRI of the spinal cord to exclude mass lesions, and CSF cytology to exclude lymphomatous involvement (can cause similar symptoms).

**Therapeutic Considerations:** For any established CMV disease, optimization of antiretroviral therapy is important along with initiating anti-CMV therapy. Ganciclovir plus foscarnet may be beneficial as initial therapy for severe cases. Consider discontinuation of valganciclovir maintenance therapy if  $CD_4$  increases to  $> 100\text{--}150/mm^3$  × 6 months or longer in response to antiretroviral therapy.

**Prognosis:** Unless  $CD_4$  cell count increases in response to antiretroviral therapy, response to anti-CMV treatment is usually transient, followed by progression of symptoms.

### **Cryptococcal Meningitis**

Pathogen	Preferred Therapy	Alternate Therapy
Cryptococcus neoformans	<p><u>Acute infection (induction therapy)</u> Amphotericin B 0.7 mg/kg (IV) q24h × 2 weeks ± flucytosine (5-FC) 25 mg/kg (PO) q6h × 2 weeks</p> <p style="text-align: center;"><b>or</b></p>	<p><u>Acute infection (induction therapy)</u> Amphotericin B 0.7 mg/kg/day (IV) × 2 weeks</p> <p style="text-align: center;"><b>or</b></p> <p>Fluconazole 400–800 mg (IV or PO) q24h × 6 weeks (less severe disease)</p> <p style="text-align: center;"><b>or</b></p>



## Cryptococcal Meningitis (cont'd)

Pathogen	Preferred Therapy	Alternate Therapy
	Lipid amphotericin 4 mg/kg (IV) q24h × 2 weeks ± flucytosine (5-FC) 25 mg/ kg (PO) q6h × 2 weeks  <u>Consolidation therapy</u> Fluconazole 400 mg (PO) q24h × 8 weeks or until CSF cultures are sterile  <u>Chronic maintenance therapy</u> <u>(secondary prophylaxis)</u> Fluconazole 200 mg (PO) q24h	Fluconazole 400–800 mg (IV or PO) q24h <i>plus</i> flucytosine (5-FC) 25 mg/kg (PO) q6h × 4–6 weeks  <u>Consolidation therapy</u> Itraconazole 200 mg (PO) q12h × 8 weeks or until CSF cultures are sterile  <u>Chronic maintenance therapy</u> Itraconazole 200 mg (PO) q24h for intolerance to fluconazole or failed fluconazole therapy

**Clinical Presentation:** Often indolent onset of fever, headache, subtle cognitive deficits. Occasional meningeal signs and focal neurologic findings, though non-specific presentation is most common.

**Diagnostic Considerations:** Diagnosis usually by cryptococcal antigen; India ink stain of CSF is less sensitive. Diagnosis is essentially excluded with a negative serum cryptococcal antigen (sensitivity of test in AIDS patients approaches 100%). If serum cryptococcal antigen is positive, CSF antigen may be negative in disseminated disease without spread to CNS/meninges. Brain imaging is often normal, but CSF analysis is usually abnormal with a markedly elevated opening pressure.

**Pitfalls:** Be sure to obtain a CSF opening pressure, since reduction of increased intracranial pressure is critical for successful treatment. Remove sufficient CSF during the initial lumbar puncture (LP) to reduce closing pressure to < 200 mm H<sub>2</sub>O or 50% of opening pressure. Increased intracranial pressure requires repeat daily lumbar punctures until CSF pressure stabilizes; persistently elevated pressure should prompt placement of a lumbar drain or ventriculo-peritoneal shunting. Adjunctive corticosteroids are not recommended.

**Therapeutic Considerations:** Optimal total dose/duration of amphotericin B prior to fluconazole switch is unknown (2–3 weeks is reasonable if patient is doing well). Treatment with 5-FC is optional; however, since 5-FC is associated with more rapid sterilization of CSF, it is reasonable to start 5-FC and then discontinue it for toxicity (neutropenia, nausea). Fluconazole is preferred over itraconazole for life-long maintenance therapy. Consider discontinuation of chronic maintenance therapy in patients who remain asymptomatic with CD<sub>4</sub> > 100–200/mm<sup>3</sup> for > 6 months due to ART.

**Prognosis:** Variable. Mortality up to 40%. Adverse prognostic factors include increased intracranial pressure, abnormal mental status.

## Progressive Multifocal Encephalopathy (PML)

Pathogen	Therapy
Reactivation of latent papovavirus (JC strain most common)	Effective antiretroviral therapy with immune reconstitution

**Clinical Presentation:** Hemiparesis, ataxia, aphasia, other focal neurologic defects, which may progress over weeks to months. Usually alert without headache or seizures on presentation.

**Diagnostic Considerations:** Demyelinating disease caused by reactivation of latent papovavirus (JC strain most common). Diagnosis by clinical presentation and MRI showing patchy demyelination of white matter  $\pm$  cerebellum/brainstem. JC virus PCR of CSF is useful for noninvasive diagnosis. In confusing or atypical presentation, biopsy may be needed to distinguish PML from other opportunistic infections, CNS lymphoma, or HIV encephalitis/encephalopathy.

**Pitfalls:** Primary HIV-related encephalopathy has a similar appearance on MRI.

**Therapeutic Considerations:** Most effective therapy is antiretroviral therapy with immune reconstitution. Some patients experience worsening neurologic symptoms once ART is initiated due to immune reconstitution induced inflammation. ART should be continued, with consideration of adjunctive steroids. Randomized controlled trials have evaluated cidofovir and vidarabine—neither is effective nor recommended.

**Prognosis:** Rapid progression of neurologic deficits over weeks to months is common. Best chance for survival is immune reconstitution in response to antiretroviral therapy, although some patients will have progressive disease despite immune recovery.

## Toxoplasma Encephalitis

Pathogen	Preferred Therapy	Alternate Therapy
Toxoplasma gondii	<p><u>Acute therapy (x 6–8 weeks until good clinical response)</u></p> <p>Pyrimethamine 200 mg (PO) <math>\times</math> 1 dose, then 50 mg (&lt; 60 kg body weight) or 75 mg (&gt; 60 kg) (PO) q24h</p> <p style="text-align: center;"><b>plus</b></p> <p>Sulfadiazine 1000 mg (&lt; 60 kg) or 1500 mg (&gt; 60 kg) (PO) q6h</p> <p style="text-align: center;"><b>plus</b></p> <p>Leucovorin 10–25 mg (PO) q24h</p> <p><i>For severely ill patients who cannot take oral medications, treat with TMP–SMX (5 mg/kg TMP and 25 mg/kg SMX) (IV) q12h</i></p>	<p><u>Acute therapy (x 6–8 weeks until good clinical response)</u></p> <p>Pyrimethamine 200 mg (PO) <math>\times</math> 1 dose, then 50 mg (&lt; 60 kg body weight) or 75 mg (&gt; 60 kg) (PO) q24h <i>plus</i> clindamycin 600 mg (IV or PO) q6h <i>plus</i> leucovorin 10 mg (PO) q24h</p> <p style="text-align: center;"><b>or</b></p> <p>TMP–SMX (5 mg/kg TMP (IV or PO) q12h</p> <p style="text-align: center;"><b>or</b></p> <p>Atovaquone 1500 mg (PO) q12h (with meals or nutritional supplement) <i>plus</i> pyrimethamine (as above)</p> <p style="text-align: center;"><b>or</b></p> <p>Atovaquone 1500 mg (PO) q12h (with meals or nutritional supplement) <i>plus</i> sulfadiazine 1000–1500 mg (PO) q6h</p> <p style="text-align: center;"><b>or</b></p> <p>Atovaquone 1500 mg (PO) q12h (with meals)</p> <p style="text-align: center;"><b>or</b></p> <p>Pyrimethamine (see above) <i>plus</i> leucovorin 10 mg (PO) q24h <i>plus</i> azithromycin 900–1200 mg (PO) q24h</p>

## Toxoplasma Encephalitis (cont'd)

Pathogen	Preferred Therapy	Alternate Therapy
	<p>Follow with lifelong suppressive therapy</p> <p>Sulfadiazine 500–1000 mg (PO) q6h</p> <p style="text-align: center;"><b>plus</b></p> <p>Pyrimethamine 50 mg (PO) q24h</p> <p style="text-align: center;"><b>plus</b></p> <p>Leucovorin 10–25 mg (PO) q24h</p>	<p>Lifelong suppressive therapy</p> <p>Clindamycin 600 mg (PO) q8h plus pyrimethamine 50 mg (PO) q24h <i>plus</i> leucovorin 10–25 mg (PO) q24h (2<sup>nd</sup> choice regimen)</p> <p style="text-align: center;"><b>or</b></p> <p>Atovaquone 750 mg (PO) q12h <math>\pm</math> pyrimethamine 25 mg (PO) q24h <i>plus</i> leucovorin 10 mg (PO) q24h (3<sup>rd</sup> choice regimen)</p>

**Clinical Presentation:** Wide spectrum of neurologic symptoms, including sensorimotor deficits, seizures, confusion, ataxia. Fever/headache are common.

**Diagnostic Considerations:** Diagnosis by characteristic radiographic appearance and response to empiric therapy in a for *T. gondii* seropositive patient.

**Pitfalls:** Use leucovorin (folinic acid) 10 mg (PO) daily with pyrimethamine-containing regimens, not folate/folic acid. Radiographic improvement may lag behind clinical response.

**Therapeutic Considerations:** Alternate agents include atovaquone, azithromycin, clarithromycin, minocycline (all with pyrimethamine if possible). Decadron 4 mg (PO or IV) q6h is useful for edema/mass effect. Chronic suppressive therapy can be discontinued if patients are free from signs and symptoms of disease and have a CD<sub>4</sub> cell count > 200/mm<sup>3</sup> for > 6 months due to ART.

**Prognosis:** Usually responds to treatment if able to tolerate drugs. Clinical response is evident by 1 week in 70%, by 2 weeks in 90%. Radiographic improvement is usually apparent by 2 weeks. Neurologic recovery is variable.

## Gastrointestinal Tract Opportunistic Infections

### Campylobacter (C. jejuni) Enteritis

Subset	Preferred Therapy
Mild disease	Might withhold therapy unless symptoms persist for several days
Moderate disease	<p>Ciprofloxacin 500 mg (PO) q12h <math>\times</math> 1 week</p> <p style="text-align: center;"><b>or</b></p> <p>Azithromycin 500 mg (PO) q24h <math>\times</math> 1 week</p>
Bacteremia	<p>Ciprofloxacin 500 mg (PO) q12h <math>\times</math> 2 weeks*</p> <p style="text-align: center;"><b>or</b></p> <p>Azithromycin 500 mg (PO) q24h <math>\times</math> 2 weeks*</p>

\* Consider addition of aminoglycoside in bacteremic patients.

**Clinical Presentation:** Acute onset of diarrhea, sometimes bloody; constitutional symptoms may be prominent.

**Diagnostic Considerations:** Diagnosis by stool culture; bacteremia may rarely occur, so blood cultures also indicated. Suspect campylobacter in AIDS patient with diarrhea and curved gram-negative rods in blood culture. Non-jejuni species may be more strongly correlated with bacteremia.

**Therapeutic Considerations:** Optimal therapy not well defined. Treat with quinolone or azithromycin; modify therapy based on susceptibility testing. Quinolone resistance can occur and correlates with treatment failure. Role of aminoglycoside is unclear.

**Prognosis:** Depends on underlying immune status; prognosis is generally good.

## Clostridium difficile Diarrhea/Colitis

Pathogen	Preferred Therapy	Alternate Therapy
<i>C. difficile</i>	Metronidazole 500 mg (PO) q8h x 10–14 days. Avoid use of other <i>C. difficile</i> associated antibiotics if possible.	Vancomycin 125 mg (PO) q6h x 10–14 days. Avoid use of <i>C. difficile</i> associated antibiotics if possible.

**Clinical Presentation:** Diarrhea and abdominal pain following antibiotic therapy. Diarrhea may be watery or bloody. Proton pump inhibitors increase the risk. Among antibiotics, clindamycin and beta-lactams are most frequent. Rarely due to aminoglycosides, linezolid, doxycycline, TMP–SMX, carbapenems, daptomycin, vancomycin.

**Diagnostic Considerations:** Most common cause of bacterial diarrhea in U.S. among HIV patients. Watery diarrhea with positive *C. difficile* toxin in stool specimen. *C. difficile* stool toxin test is sufficiently sensitive/specific. If positive, no need to retest until negative (endpoint is end of diarrhea); if negative, no need to retest (repeat tests will be negative). *C. difficile* colitis may be distinguished clinically from *C. difficile* diarrhea by temperature > 102°F, ↑ WBC, ↑ ESR, and/or abdominal pain. *C. difficile* virulent epidemic strain is type B1 (toxintype III), which produces 20-times the amount of toxin A/B compared to less virulent strains.

**Pitfalls:** In a patient with *C. difficile* diarrhea, *C. difficile* colitis is suggested by the presence of ↑ WBC, ↑ ESR, abdominal pain and temperature > 102°F; confirm diagnosis with CT of abdomen, which will show colonic wall thickening. New virulent strain of *C. difficile* may present with colitis with temperature ≤ 102°F, little/no ↑ WBC, and little/no abdominal pain; confirm diagnosis with CT/MRI of abdomen. *C. difficile* toxin may remain positive in stools for weeks following treatment; do not treat positive stool toxin unless patient has symptoms.

**Therapeutic Considerations:** Initiate therapy for mild disease with metronidazole; symptoms usually begin to improve within 2–3 days. For moderate or severe disease, or with evidence of colitis clinically (leukocytosis, fever, colonic thickening on CT scan), vancomycin has become the preferred agent in many centers due to concern for the more virulent strain, and based on the results of some observational studies suggesting vancomycin is more effective. The duration of therapy should be extended beyond 14 days if other systemic antibiotics must be continued. Relapse occurs in 10–25% of patients, and rates may be higher in patients with HIV due to the frequent need for other antimicrobial therapy. First relapses can be treated with a repeat of the initial regimen of metronidazole or vancomycin. For multiple relapses, a long-term taper is appropriate: week 1–125 mg 4x/day; week 2–125 mg 2x/day; week 3–125 mg once daily; week 4–125 mg every other day; weeks 5 and 6–125 mg every three days. Every effort should be made to resume a normal diet and to avoid other antibacterial therapies. Probiotic treatments (such as lactobacillus or *Saccharomyces boulardii*) have not yet been shown to reduce the risk of relapse in controlled clinical trials.

**Prognosis:** Prognosis with *C. difficile* colitis is related to severity of the colitis.

## CMV Esophagitis/Colitis

Infection	Preferred Therapy	Alternate Therapy
Initial infection	Ganciclovir 5 mg/kg (IV) q12h × 3–4 weeks or until signs and symptoms have resolved. <b>or</b> Valganciclovir 900 mg (PO) q12h can be used if able to tolerate oral intake. Maintenance therapy is generally not necessary but should be considered after relapses	Foscarnet 90 mg/kg (IV) q12h × 3–4 weeks or until signs and symptoms have resolved <b>or</b> Cidofovir 5 mg/kg (IV) q every other week with saline hydration and probenecid (see CMV retinitis)
Relapses	Valganciclovir 900 mg (PO) q24h indefinitely; consider discontinuation if $CD_4 > 200/mm^3$ for $\geq 6$ months on ART	

**Clinical Presentation:** Localizing symptoms, including odynophagia, abdominal pain, diarrhea, sometimes bloody stools.

**Diagnostic Considerations:** Diagnosis by finding CMV inclusions on biopsy. CMV can affect the entire GI tract, resulting in oral/esophageal ulcers, gastritis, and colitis (most common). CMV colitis varies greatly in severity, but typically causes fever, abdominal cramping, and sometimes bloody stools.

**Pitfalls:** CMV colitis may cause colonic perforation and should be considered in any AIDS patient presenting with an acute abdomen, especially if radiography demonstrates free intraperitoneal air.

**Therapeutic Considerations:** Duration of therapy is dependent on clinical response, typically 3–4 weeks. Consider chronic suppressive therapy for recurrent disease. Screen for CMV retinitis.

**Prognosis:** Relapse rate is greatly reduced with immune reconstitution due to antiretroviral therapy.

## Cryptosporidia Enteritis

Pathogen	Preferred Therapy	Alternate Therapy
Cryptosporidium sp.	Effective ART with immune reconstitution to $CD_4 > 100/mm^3$ can result in complete resolution of symptoms and clearance of infection	Nitazoxanide 500 mg (PO) q12h × 4–6 weeks <b>or</b> Paromomycin 1 gm (PO) q12h × 2–4 weeks

**Clinical Presentation:** High-volume watery diarrhea with weight loss and electrolyte disturbances, especially in advanced HIV disease.

**Diagnostic Considerations:** Spore-forming protozoa. Diagnosis by AFB smear of stool demonstrating characteristic oocyst. Malabsorption may occur.

**Pitfalls:** No fecal leukocytes; organisms are not visualized on standard ova and parasite exams (need to request special stains).

**Therapeutic Considerations:** Anecdotal reports of antimicrobial success. Newest agent nitazoxanide may be effective in some settings, but no increase in cure rate for nitazoxanide if  $CD_4 < 50/mm^3$ . Immune reconstitution in response to antiretroviral therapy is the most effective therapy, and may induce prolonged remissions and cure. Anti-diarrheal agents (Lomotil, Pepto-Bismol) are useful to control symptoms. Hyperalimentation may be required for severe cases.

**Prognosis:** Related to degree of immunosuppression/response to antiretroviral therapy.

## Isospora (*Isospora belli*) Enteritis

Subset	Preferred Therapy	Alternate Therapy
Acute infection	TMP 160 mg and SMX 800 mg (IV or PO) q6h × 10 days <b>or</b> TMP 320 mg and SMX 1600 mg (IV or PO) q12h × 10–14 days	Pyrimethamine 50–75 mg (PO) q24h <i>plus</i> leucovorin 5–10 mg (PO) q24h <b>or</b> Ciprofloxacin 500 mg (PO) q12h or other fluoroquinolones
Chronic maintenance therapy for CD <sub>4</sub> < 200 (secondary prophylaxis)	TMP 320 mg <i>plus</i> SMX 1600 mg (PO) q24h*	Pyrimethamine 25 mg (PO) q24h <i>plus</i> leucovorin 5–10 mg (PO) q24h*

\* Discontinuation of secondary prophylaxis may be considered if CD<sub>4</sub> > 200/mm<sup>3</sup> for > 3 months.

**Clinical Presentation:** Severe chronic diarrhea without fever/fecal leukocytes.

**Diagnostic Considerations:** Spore-forming protozoa (*Isospora belli*). Oocyst on AFB smear of stool larger than cryptosporidium (20–30 microns vs. 4–6 microns). More common in HIV patients from the tropics (e.g., Haiti). Less common than cryptosporidium or microsporidia. Malabsorption may occur.

**Pitfalls:** Multiple relapses are possible.

**Therapeutic Considerations:** Chronic suppressive therapy may be required if CD<sub>4</sub> does not increase.

**Prognosis:** Related to degree of immunosuppression/response to antiretroviral therapy.

**Comments:** Immune reconstitution with ART results in fewer relapses.

## Microsporidia Enteritis

Pathogen	Therapy*
Microsporidia other than <i>Enterocytozoon bienuesi</i>	Albendazole 400 mg (PO) q12h (continue until CD <sub>4</sub> > 200/mm <sup>3</sup> )
<i>Enterocytozoon bienuesi</i>	Nitazoxanide 1 gm (PO) q12h × 60 days <b>or</b> Fumagillin 60 mg (PO) q24h (not available in the U.S.)
<i>Trachipleistophora</i> or <i>Brachiola</i>	Itraconazole 400 mg (PO) q24h <i>plus</i> albendazole 400 mg (PO) q12h

\* Regardless of species, ART with immune reconstitution is a critical component of treatment.

**Clinical Presentation:** Intermittent chronic diarrhea without fever/fecal leukocytes.

**Diagnostic Considerations:** Spore-forming protozoa (*S. intestinalis*, *E. bienuesi*). Diagnosis by modified trichrome or fluorescent antibody stain of stool. Microsporidia can rarely disseminate to sinuses/cornea. Severe malabsorption may occur.

**Pitfalls:** Microsporidia is too small for detection by routine microscopic examination of stool.

**Therapeutic Considerations:** Albendazole is less effective for *E. bienuesi* than *S. intestinalis*, but speciation is usually not possible. May consider treatment discontinuation for CD<sub>4</sub> < 200/mm<sup>3</sup> if patient remains asymptomatic (no signs or symptoms of microsporidiosis). If ocular infection is present, continue treatment indefinitely.

**Prognosis:** Related to degree of immunosuppression/response to antiretroviral therapy.

## Oropharyngeal/Esophageal Candidiasis

Infection	Therapy	Fluconazole-Resistance
Oropharyngeal candidiasis (thrush)	<p><u>Preferred therapy</u> Fluconazole 100 mg (PO) q24h x 1–2 weeks</p> <p><u>Alternate therapy</u> Itraconazole oral solution 200 mg (PO) q24h x 1–2 weeks <b>or</b> Clotrimazole troches 10 mg (PO) 5x/day x 1–2 weeks <b>or</b> nystatin suspension 4–6 mL q6h or 1–2 flavored pastilles 4–5x/day x 1–2 weeks</p>	<p>Fluconazole at doses up to 800 mg (PO) q24h x 1–2 weeks</p> <p><b>or</b></p> <p>Caspofungin 70 mg (IV) on day 1, then 50 mg (IV) q24h x 1–2 weeks</p> <p><b>or</b></p> <p>Micafugin 150 mg (IV) q24h x 1–2 weeks</p>
Esophageal candidiasis	<p><u>Preferred therapy</u> Fluconazole 100 mg (up to 400 mg) (IV or PO) q24h x 1–2 weeks</p> <p><u>Alternate therapy</u> Itraconazole oral solution 200 mg (PO) q24h x 2–3 weeks <b>or</b> Voriconazole 200 mg (PO) q24h x 2–3 weeks <b>or</b> Caspofungin 50 mg (IV) q24h x 2–3 weeks</p>	<p>Amphotericin B 0.3 mg/kg (IV) q24h x 1–2 weeks</p> <p><b>or</b></p> <p>Lipid amphotericin 3–5 mg/kg (IV) q24h x 1–2 weeks</p>

### Oral Thrush (Candida)

**Clinical Presentation:** Dysphagia/odynophagia. More common/severe in advanced HIV disease.

**Diagnostic Considerations:** Pseudomembranous (most common), erythematous, and hyperplastic (leukoplakia) forms. Pseudomembranes (white plaques on inflamed base) on buccal mucosa/tongue/gingiva/palate scrape off easily, hyperplastic lesions do not. Diagnosis by clinical appearance  $\pm$  KOH/gram stain of scraping showing yeast/pseudomycelia. Other oral lesions in AIDS patients include herpes simplex, aphthous ulcers, Kaposi's sarcoma, oral hairy leukoplakia.

**Pitfalls:** Patients may be asymptomatic.

**Therapeutic Considerations:** Fluconazole is superior to topical therapy in preventing relapses of thrush and treating Candida esophagitis. Continuous treatment with fluconazole may lead to fluconazole-resistance, which is best treated initially with itraconazole suspension and, if no response, with IV caspofungin or amphotericin. Chronic suppressive therapy is usually only considered for severely immunosuppressed patients.

**Prognosis:** Improvement in symptoms are often seen within 24–48 hours.

### Candida Esophagitis

**Clinical Presentation:** Dysphagia/odynophagia, almost always in the setting of oropharyngeal thrush. Fever is uncommon.

**Diagnostic Considerations:** Most common cause of esophagitis in HIV disease. For persistent symptoms despite therapy, endoscopy with biopsy/culture is recommended to confirm diagnosis and assess azole-resistance.

**Pitfalls:** May extend into stomach. Other common causes of esophagitis include CMV, herpes simplex, and aphthous ulcers. Rarely, Kaposi's sarcoma, non-Hodgkin's lymphoma, zidovudine, dideoxycytidine, and other infections may cause esophageal symptoms.

**Therapeutic Considerations:** Systemic therapy is preferred over topical therapy. Failure to improve on empiric therapy mandates endoscopy to look for other causes, especially herpes viruses/aphthous ulcers. Consider maintenance therapy with fluconazole for frequent relapses, although the risk of fluconazole resistance is increased. Fluconazole-resistance is best treated initially with itraconazole suspension and, if no response, with IV caspofungin, micafungin, or amphotericin.

**Prognosis:** Relapse rate related to degree of immunosuppression.

## Salmonella Gastroenteritis (non-S. typhi)

Subset	Preferred Therapy	Alternate Therapy
Mild disease	Ciprofloxacin 750 mg (PO) q12h × 1–2 weeks	TMP–SMX 1 DS (PO) q12h × 2 weeks <b>or</b>
CD <sub>4</sub> < 200	Ciprofloxacin 750 mg (PO) q12h × 4–6 weeks	Ceftriaxone 2 gm (IV) q24h × 2 weeks <b>or</b>
Bacteremia	Ciprofloxacin 750 mg (PO) q12h × 4–6 weeks, then 500 mg (PO) q12h indefinitely	Cefotaxime 1 gm (IV) q8h × 2 weeks

**Clinical Presentation:** Patients with HIV are at markedly increased risk of developing salmonellosis. Three different presentations may be seen: (1) self-limited gastroenteritis, as typically seen in immunocompetent hosts; (2) a more severe and prolonged diarrheal disease, associated with fever, bloody diarrhea, and weight loss; or (3) Salmonella septicemia, which may present with or without gastrointestinal symptoms.

**Diagnostic Considerations:** The diagnosis is established through cultures of stool and blood. Given the high rate of bacteremia associated with Salmonella gastroenteritis—especially in advanced HIV disease—blood cultures should be obtained in any HIV patient presenting with diarrhea and fever.

**Pitfalls:** A distinctive feature of salmonella bacteremia in patients with AIDS is its propensity for relapse (rate > 20%).

**Therapeutic Considerations:** The mainstay of treatment is a fluoroquinolone; greatest experience is with ciprofloxacin, but newer quinolones (moxifloxacin, levofloxacin) may also be effective. For uncomplicated salmonellosis in an HIV patient with CD<sub>4</sub> > 200/mm<sup>3</sup>, 1–2 weeks of treatment is reasonable to reduce the risk of extraintestinal spread. For patients with advanced HIV disease (CD<sub>4</sub> < 200/mm<sup>3</sup>) or who have salmonella bacteremia, at least 4–6 weeks of treatment is required. Chronic suppressive therapy, given for several months or until antiretroviral therapy-induced immune reconstitution ensues, is indicated for patients who relapse after cessation of therapy. Consider using ZDV as part of the antiretroviral regimen (ZDV is active against Salmonella).

**Prognosis:** Usually responds well to treatment. Relapse rate in AIDS patients with bacteremia is > 20%.



## Shigella (Shigella sp.) Enteritis

Subset	Preferred Therapy	Alternate Therapy
No bacteremia	Fluoroquinolone (IV or PO) × 3–7 days	TMP–SMX 1 DS tablet (PO) q12h × 3–7 days  <b>or</b> Azithromycin 500 mg (PO) on day 1, then 250 mg (PO) q24h × 4 days
Bacteremia	Extend treatment duration to 14 days	Extend treatment duration to 14 days

**Clinical Presentation:** Acute onset of bloody diarrhea/mucus.

**Diagnostic Considerations:** Diagnosis by demonstrating organism in stool specimens. Shigella ulcers in colon are linear, serpiginous, and rarely lead to perforation. More common in gay men.

**Therapeutic Considerations:** Shigella dysentery is more acute/fulminating than amebic dysentery. Shigella has no carrier state, unlike Entamoeba. Shigella infections acquired outside of United States have high rates of TMP–SMX resistance. Therapy is indicated to shorten the duration of illness and to prevent spread of infection. Shigella has no carrier state.

**Prognosis:** Good if treated early. Severity of illness related to Shigella species: *S. dysenteriae* (most severe) > *S. flexneri* > *S. boydii*/*S. sonnei* (mildest).

### TREATMENT OF OTHER OPPORTUNISTIC INFECTIONS IN HIV

## Candida Vaginitis

Pathogen	Therapy
Candida albicans	Intravaginal miconazole suppository 200 mg q24h × 3 days or miconazole 3% × 7 days <b>or</b> Nystatin vaginal tablet 100,000U q24h × 14 days <b>or</b> Itraconazole 200 mg (PO) q12h × 1 day (or 200 mg q24h × 3 days) <b>or</b> Fluconazole 150 mg (PO) × 1 dose

**Clinical Presentation:** White, cheesy, vaginal discharge or vulvar rash ± itching/pain. Local infection not a sign of disseminated disease.

**Diagnostic Considerations:** Local infection. Not a manifestation of disseminated disease.

**Pitfalls:** Women with advanced AIDS receiving fluconazole may develop fluconazole-resistant Candida.

**Therapeutic Considerations:** For recurrence, consider maintenance with fluconazole 100–200 mg (PO) weekly.

**Prognosis:** Good response to therapy. Relapses are common.

## Coccidioidomycosis (*C. immitis*) Infection

Infection	Therapy*
Non-meningeal infection	<p><u>Acute therapy (diffuse pulmonary or disseminated disease)</u> Amphotericin B 0.5–1.0 mg/kg (IV) q24h until clinical improvement (usually 500–1000 mg total dose). Some specialists add an azole to amphotericin B therapy</p> <p><u>Acute therapy (milder disease)</u> Fluconazole 400–800 mg (PO) q24h or Itraconazole 200 mg (PO) q12h</p> <p><u>Chronic maintenance therapy (secondary prophylaxis)</u> <u>Preferred:</u> Fluconazole 400 mg (PO) q24h indefinitely; <u>alternative:</u> or Itraconazole 200 mg capsule (PO) q12h indefinitely</p>
Meningeal infection*	<p><u>Acute therapy</u> Fluconazole 400–800 mg (IV) or (PO) q24h. Intrathecal amphotericin B if no response to azole therapy</p> <p><u>Chronic maintenance therapy (secondary prophylaxis)</u> Fluconazole 400 mg (PO) q24h or itraconazole 200 mg capsule (PO) q12h indefinitely</p>

\* Therapy for meningeal infection should be lifelong with fluconazole 400–800 mg q24h. There are insufficient data to recommend discontinuation of chronic maintenance therapy in other settings.

**Clinical Presentation:** Typically a complication of advanced HIV infection ( $CD_4$  cell count  $< 200/mm^3$ ). Most patients present with disseminated disease, which can manifest as fever, diffuse pulmonary infiltrates, adenopathy, skin lesions (multiple forms – verrucous, cold abscesses, ulcers, nodules), and/or bone lesions. Approximately 10% will have spread to the CNS in the form of meningitis (fever, headache, altered mental status).

**Diagnostic Considerations:** Consider the diagnosis in any patient with advanced HIV-related immunosuppression who has been in a *C. immitis* endemic area (Southwestern US, northern Mexico) and presents with a systemic febrile syndrome. Diagnosis can be made by culture of the organism, visualization of characteristic spherules on histopathology, or a positive complement-fixation antibody ( $\geq 1:16$ ). In meningeal cases, CSF profile shows low glucose, high protein, and lymphocytic pleocytosis.

**Pitfalls:** Antibody titers often negative on presentation. CSF profile of meningitis can be similar to TB.

**Prognosis:** Related to extent of infection and degree of immunosuppression. Clinical response tends to be slow, especially with a high disease burden and advanced HIV disease. Meningeal disease is treated life-long regardless of  $CD_4$  recovery.

## Extrapulmonary Tuberculosis

Pathogen	Therapy
Mycobacterium tuberculosis	Treat the same as pulmonary TB (see p. 326). May require longer duration of therapy based on clinical response

**Clinical Presentation:** Multiple presentations possible (e.g., lymphadenitis, osteomyelitis, meningitis, hepatitis). Dissemination is more common in patients with low  $CD_4$  cell counts ( $< 100/mm^3$ ).

**Diagnostic Considerations:** Diagnosis by isolator blood cultures or tissue biopsy.

**Pitfalls:** Patients with disseminated disease frequently have pulmonary disease, which has implications for infection control.

**Therapeutic Considerations:** Response to therapy may be slower than in normal hosts.

**Prognosis:** Usually responsive to therapy.

## Herpes Simplex Virus (HSV) Disease

Infection	Preferred Therapy	Alternate Therapy
Orolabial lesions or initial/recurrent genital HSV	Famciclovir 500 mg (PO) q12h × 1–2 weeks <b>or</b> valacyclovir 1 gm (PO) q12h × 1–2 weeks <b>or</b> acyclovir 400 mg (PO) q8h × 1–2 weeks	<u>Acyclovir-resistant HSV</u> Foscarnet 60–100 mg/kg (IV) q12h until clinical response <b>or</b> Cidofovir 5 mg/kg (IV) weekly until clinical response
Moderate-to-severe mucocutaneous HSV	Initial therapy: Acyclovir 5 mg/kg (IV) q8h × 2–7 days. If improvement, switch to famciclovir 500 mg (PO) q12h <b>or</b> valacyclovir 1 gm (PO) q12h <b>or</b> acyclovir 400 mg (PO) q8h to complete 7–10 days	<u>Acyclovir-resistant HSV</u> Foscarnet 60–100 mg/kg (IV) q12h until clinical response <b>or</b> Cidofovir 5 mg/kg (IV) weekly until clinical response
HSV keratitis	Trifluridine 1% ophthalmic solution, one drop onto cornea q2h, not to exceed 9 drops per day and no longer than 21 days. Treatment in conjunction with ophthalmology consultation	<u>Acyclovir-resistant HSV</u> Foscarnet 60–100 mg/kg (IV) q12h until clinical response <b>or</b> Cidofovir 5 mg/kg (IV) weekly until clinical response
HSV encephalitis	Acyclovir 10 mg/kg (IV) q8h × 2–3 weeks <b>or</b> Valacyclovir 1 gm (PO) q6h × 2–3 weeks	<u>Acyclovir-resistant HSV</u> Foscarnet 60–100 mg/kg (IV) q12h until clinical response <b>or</b> Cidofovir 5 mg/kg (IV) weekly until response
Multiple mucocutaneous (oral or anogenital) relapses (chronic suppressive therapy)	Acyclovir 400 mg (PO) q12h <b>or</b> Famciclovir 250 mg (PO) q12h <b>or</b> Valacyclovir 500 mg (PO) q12h	Patients may be able to titrate dose downward to maintain response

### **Herpes Simplex (genital/oral)**

**Clinical Presentation:** Painful, grouped vesicles on an erythematous base that rupture, crust, and heal within 2 weeks. Lesions may be chronic, severe, ulcerative with advanced immunosuppression.

**Diagnostic Considerations:** Diagnosis by viral culture of swab from lesion base/roof of blister; alternative diagnostic techniques include Tzanck prep or immunofluorescence staining.

**Pitfalls:** Acyclovir prophylaxis is not required in patients receiving ganciclovir or foscarnet.

**Therapeutic Considerations:** In refractory cases, consider acyclovir resistance and treat with foscarnet. Topical trifluridine ophthalmic solution (Viroptic 1%) may be considered for direct application to small, localized areas of refractory disease; clean with hydrogen peroxide, then debride lightly with gauze, apply trifluridine, and cover with bacitracin/polymyxin ointment and nonadsorbent gauze; topical cidofovir (requires compounding) also may be tried. Chronic suppressive therapy with oral acyclovir, famciclovir, or valacyclovir may be indicated for patients with frequent recurrences, dosing similar to HIV-negative patients.

**Prognosis:** Responds well to treatment except in severely immunocompromised patients, in whom acyclovir resistance may develop. Prognosis for HSV meningitis is excellent.

### **Herpes Encephalitis (HSV-1)**

**Clinical Presentation:** Acute onset of fever and change in mental status.

**Diagnostic Considerations:** EEG is abnormal early (< 72 hours), showing unilateral temporal lobe abnormalities. Brain MRI is abnormal before CT scan, which may require several days before a temporal lobe focus is seen. Definitive diagnosis is by CSF PCR for HSV-1 DNA. Profound decrease in sensorium is characteristic of HSV meningoencephalitis. CSF may have PMN predominance and low glucose levels, unlike other viral causes of meningitis. A different clinical entity is HSV meningitis, which is usually associated with HSV-2 and can recur with lymphocytic meningitis.

**Pitfalls:** Rule out noninfectious causes of encephalopathy. Surprisingly, HIV encephalitis is a relatively rare cause of encephalitis in patients with HIV.

**Therapeutic Considerations:** Treat as soon as possible since neurological deficits may be mild and reversible early on, but severe and irreversible later.

**Prognosis:** Related to extent of brain injury and early antiviral therapy. Prognosis for HSV meningitis is excellent.

## **Histoplasmosis (*H. capsulatum*), Disseminated**

<b>Subset</b>	<b>Preferred Therapy</b>	<b>Alternate Therapy</b>
Acute phase (3–10 days or until clinically improved)	Liposomal amphotericin B 3 mg/kg × 1–2 weeks or other lipid associated formulations at 5 mg/kg	Amphotericin B doxycholate 0.7–1 mg/kg <b>or</b> Intravenous itraconazole (200 mg q12 × 4 doses then 200 mg q24)
Continuation phase	Itraconazole 200 (PO) TID × 3 days and then QD or BID for at least 12 months (adequate serum levels should be confirmed). Immunosuppressed patients may require lifelong suppressive therapy.	Itraconazole oral solution 200 mg (PO) q12h × 12 weeks <b>or</b> Fluconazole 800 mg (PO) q24h × 12 weeks
Meningitis	Amphotericin B deoxycholate 0.7 mg/kg (IV) q24h × 12–16 weeks <b>or</b> Liposomal amphotericin B 4 mg/kg (IV) q24h × 12–16 weeks	Fluconazole 800 mg (PO) q24h × 12 weeks

\* Duration of therapy dependent on response to therapy.

**Clinical Presentation:** Two general forms: Mild disease with fever/lymph node enlargement (e.g., cervical adenitis), or severe disease with fever, wasting  $\pm$  diarrhea/meningitis/GI ulcerations.

**Diagnostic Considerations:** Diagnosis by urine/serum histoplasmosis antigen, sometimes by culture of bone marrow/liver or isolator blood cultures. May occur in patients months to years after having lived/moved from an endemic area.

**Pitfalls:** Relapse is common after discontinuation of therapy. Cultures may take 7–21 days to turn positive.

**Therapeutic Considerations:** Initial therapy depends on severity of illness on presentation. Extremely sick patients should be started on amphotericin B deoxycholate, with duration of IV therapy dependent on response to treatment. Mildly ill patients can be started on itraconazole. All patients require chronic suppressive therapy, with possible discontinuation for immune reconstitution with  $CD_4$  counts  $> 100/mm^3$  for at least 6 months. HIV patients with  $CD_4 > 500/mm^3$  and acute pulmonary histoplasmosis might not require therapy, but a short course of itraconazole (4–8 weeks) is reasonable to prevent systemic spread.

**Prognosis:** Usually responds to treatment, except in fulminant cases.

## Mycobacterium avium-intracellulare (MAI)

Pathogen	Preferred Therapy	Alternate Therapy
Mycobacterium avium-intracellulare (MAI)	At least 2 drugs as initial therapy Clarithromycin 500 mg (PO) q12h plus ethambutol 15 mg/kg (PO) q24h (usually 800 mg or 1200 mg daily). Consider adding third drug, rifabutin 300 mg (PO) q24h, for patients with $CD_4 < 50/mm^3$ , high mycobacterial loads and severely symptomatic disease. Duration of therapy is <b>lifelong</b> , although consider discontinuation in asymptomatic patients with $> 12$ months therapy and $CD_4 > 100/mm^3$ for $> 6$ months in response to ART	Alternative to clarithromycin Azithromycin 500–600 mg (PO) q24h  Alternative 3 <sup>rd</sup> or 4 <sup>th</sup> drug for severe symptoms or disseminated disease Ciprofloxacin 500–750 mg (PO) q12h <b>or</b> Levofloxacin 500 mg (PO) q24h <b>or</b> Amikacin 10–15 mg/kg (IV) q24h

**Clinical Presentation:** Typically presents as a febrile wasting illness in advanced HIV disease ( $CD_4 < 50/mm^3$ ). Focal invasive disease is possible, especially in patients with advanced immunosuppression after starting antiretroviral therapy. Focal disease likely reflects restoration of pathogen-specific immune response to subclinical infection (“immune reconstitution inflammatory syndrome” [IRIS]), and typically manifests as lymphadenitis (mesenteric, cervical, thoracic) or rarely disease in the spine mimicking Pott’s disease. Immune reconstitution syndrome usually occurs within weeks to months after starting antiretroviral therapy for the first time, but may occur a year or more later.

**Diagnostic Considerations:** Diagnosis by isolation of organism from a normally sterile body site (blood, lymph node, bone marrow, liver biopsy). Lysis centrifugation (DuPont Isolator) is the preferred blood culture method. Anemia/ $\uparrow$  alkaline phosphatase are occasionally seen.

**Pitfalls:** Isolator blood cultures may be negative, especially in immune reconstitution inflammatory syndrome initially.

**Therapeutic Considerations:** Some studies suggest benefit for addition of rifabutin 300 mg (PO) q24h, others do not. Rifabutin may require dosage adjustment with NNRTI's and PI's. For concurrent use with nelfinavir, indinavir, or amprenavir, decrease rifabutin to 150 mg (PO) q24h. For concurrent use with ritonavir, decrease rifabutin to 150 mg (PO) 2–3x/week. For concurrent use with efavirenz, increase rifabutin to 450–600 mg (PO) q24h. The dose of PI's or NNRTI's may need to be increased by 20–25%. Monitor carefully for rifabutin drug toxicity (arthralgias, uveitis, leukopenia). Treat IRIS initially with NSAIDs; if symptoms persist, systemic corticosteroids (prednisone 20–40 mg daily) for 4–8 weeks can be used. Some patients will require a more prolonged course of corticosteroids with a slow taper over months. Azithromycin is often better tolerated than clarithromycin and has fewer drug-drug interactions. Optimal long-term management is unknown, though most studies suggest that treatment can be discontinued in asymptomatic patients with > 12 months of therapy and  $CD_4 > 100/mm^3$  for > 6 months.

**Prognosis:** Depends on immune reconstitution in response to antiretroviral therapy. Adverse prognostic factors include high-grade bacteremia or severe wasting.

## Varicella Zoster Virus (VZV)

Infection	Preferred Therapy
Primary VZV infection (chickenpox)	Acyclovir 10 mg/kg (IV) q8h × 7–10 days. Can start with or change to oral therapy with Valacyclovir 1 gm (PO) q8h or Famciclovir 500 mg (PO) q8h after defervescence if no evidence of visceral involvement exists
Local dermatomal herpes zoster	Famciclovir 500 mg (PO) q8h × 7–10 days <b>or</b> Valacyclovir 1 gm (PO) q8h × 7–10 days
Extensive cutaneous or visceral involvement	Acyclovir 10 mg/kg (IV) q8h until cutaneous and visceral disease has clearly resolved and the patient is clinically improved
Acute retinal necrosis	Acyclovir 10 mg/kg (IV) q8h until progression stops, then Valacyclovir 1 gm (PO) q8h × 6 weeks. Treat in conjunction with close ophthalmologic consultation

**Clinical Presentation:** Primary varicella (chickenpox) presents as clear vesicles on an erythematous base that heal with crusting and sometimes scarring. Zoster usually presents as painful tense vesicles on an erythematous base in a dermatomal distribution. In patients with HIV, primary varicella is more severe/prolonged, and zoster is more likely to involve multiple dermatomes/disseminate. VZV can rarely cause acute retinal necrosis, which requires close consultation with ophthalmology.

**Diagnostic Considerations:** Diagnosis is usually clinical. In atypical cases, immunofluorescence can be used to distinguish herpes zoster from herpes simplex.

**Pitfalls:** Extend treatment beyond 7–10 days if new vesicles are still forming after initial treatment period. Corticosteroids for dermatomal zoster are not recommended in HIV-positive patients.

**Therapeutic Considerations:** IV therapy is generally indicated for severe disease/cranial nerve zoster.

**Prognosis:** Usually responds slowly to treatment.

## HIV COINFECTIONS (HBV/HCV)

## Hepatitis B Virus (HBV)

Preferred Therapy, Duration of Therapy, Chronic Maintenance	Alternate Therapy	Other Options/Issues
<p><u>Therapy for patients who require ART</u></p> <p>Patients should be treated with agents active against both HIV and HBV or with agents with independent activity against each virus</p> <p>Consider tenofovir + emtricitabine as part of HIV and HBV regimen</p> <p><i>Lamivudine or emtricitabine-naïve patients</i>  [Lamivudine 150 mg PO BID (or 300 mg PO daily) or emtricitabine 200 mg PO daily] + tenofovir (TDF) 300 mg PO daily (+ additional agent[s] active against HIV)</p> <p><i>Lamivudine or emtricitabine-experienced patients with detectable HBV DNA (assume lamivudine-resistance)</i></p> <p><i>If not on TDF:</i> Add TDF 300 mg PO daily as part of an ART regimen + lamivudine or emtricitabine;</p> <p><b>or</b></p> <p>Adefovir 10 mg PO daily + lamivudine or emtricitabine + other combination ART;</p> <p><b>or</b></p>	<p><u>Treatment for patients who do not require ART</u></p> <p>Use agents with sole activity against HBV and with the least potential of selecting HIV resistance mutations</p> <p>Consider early initiation of ART, especially for patients with high HBV DNA <i>For patients with CD<sub>4</sub>+ count &gt; 350 cells/μL, HBeAg (-), HBV DNA &gt; 2,000 IU/mL (&gt; 20,000 copies/mL)</i> Adefovir 10 mg PO daily</p> <p><i>For patients with CD<sub>4</sub>+ count &gt; 350 cells/μL, HBeAg (+), HBV DNA &gt; 20,000 IU/mL (&gt; 200,000 copies/mL), and elevated ALT</i></p>	<p>Emtricitabine, entecavir, lamivudine, or tenofovir should not be used for the treatment of HBV infection in patients who are not receiving combination ART</p> <p>Among patients coinfectd with HIV, HBV, and HCV, consideration of starting ART should be the first priority. If ART is not required, an interferon-based regimen, which suppresses both HCV &amp; HBV, should be considered</p> <p>If IFN-based treatment for HCV has failed, treatment of chronic HBV with nucleoside or nucleotide analogs is recommended</p> <p>Cross-resistance to emtricitabine or telbivudine should be assumed in patients with suspected or proven 3TC resistance</p> <p>When changing ART regimens, continue agents with anti-HBV activity because of the risk of IRIS</p>

## Hepatitis B Virus (HBV) (cont'd)

Preferred Therapy, Duration of Therapy, Chronic Maintenance	Alternate Therapy	Other Options/Issues
Entecavir 1 mg PO daily can be considered in patients with complete HIV suppression (while on ART) who do not demonstrate YMDD (M204V/I) motif mutations in HBV DNA  Duration of therapy: Because of the high rates of relapse, certain specialists recommend continuing therapy indefinitely	Peginterferon alfa-2a 180 µg SQ weekly x 48 weeks—with careful follow-up of HBeAg conversion	If anti-HBV therapy is discontinued and a flare occurs, therapy should be reinstated, as it can be potentially life saving

Adapted from: Sax PE, Cohen CJ, Kuritzkes DR. HIV Essentials, 7<sup>th</sup> edition. Jones & Bartlett, Sudbury, MA, 2014.

**Epidemiology:** Hepatitis B virus (HBV) infection is relatively common in patients with HIV, with approximately 60% showing some evidence of prior exposure. Chronic hepatitis B infection interacts with HIV infection in several important ways:

- HBV increases the risk of liver-related death and hepatotoxicity from antiretroviral therapy
- 3TC, FTC, and tenofovir each have anti-HBV activity. Thus selection of antiretroviral therapy for patients with HBV can have clinical and resistance implications for HBV as well as HIV. This is most notable with 3TC and FTC, as a high proportion of coinfecting patients will develop HBV-associated resistance to these drugs after several years of therapy. This resistance reduces response to subsequent non-3TC or FTC anti-HBV therapy.
- Cessation of anti-HBV therapy may lead to exacerbations of underlying liver disease; in some cases, these flares have been fatal.
- Immune reconstitution may lead to worsening of liver status, presumably because HBV disease is immune mediated. This is sometimes associated with loss of HBeAg.
- Entecavir can no longer be recommended for HIV/HBV coinfecting patients, as it has anti-HIV activity and may select for HIV resistance mutation M184V. If needed, it should be used only with a fully suppressive HIV regimen.

**Diagnostic Consideration:** Obtain HBSAb, HBSAg, and HBCAb at baseline in all patients. If negative, hepatitis B vaccination is indicated. If chronic HBV infection (positive HBSAg) is identified, obtain HBeAg, HBeAb, and HBV DNA levels. As with HCV infection, vaccination with hepatitis A vaccine and counseling to avoid alcohol are important components of preventive care. **Isolated Hepatitis B Core Antibody:** Many patients with HIV have antibody to hepatitis B core (anti-HBc) but are negative for both HBSAg and HBSAb. This phenomenon appears to be more common in those with HCV coinfection (Clin Infect Dis 2003 36:1602–6). In this scenario, diagnostic considerations include: (1) recently acquired HBV, before development of HBSAb; (2) chronic HBV, with HBSAg below the levels of detection; (3) immunity to HBV, with HBSAb below the levels of detection; (4) false-positive anti-HBV core. As the incidence of HBV is relatively low in most populations and anti-HBc alone is usually a stable phenomenon over years, recent acquisition of HBV is rarely the explanation. We recommend checking HBV DNA in this situation:



If positive, this indicates chronic HBV; if negative, then low-level immunity or false-positive anti-HBV core remain as possible explanations; since distinguishing between these possibilities cannot be done, we recommend immunization with the hepatitis B vaccine series. It is useful to measure HBV serologic markers periodically in this population, as improvement in immune status due to ART may lead to increasing titers of HBSAb and subsequently confirm immunity.

**Therapeutic Considerations:** The optimal treatment for HBV infection is in evolution. Current guidelines suggest treatment of HBV in all patients with active HBV replication, defined as a detectable HBEAg or HBV DNA. Pending long-term studies defining optimal management, the recommendations set forth in the grid above are reasonable. Patients being treated with regimens for HBV should be monitored for ALT every 3–4 months. HBV DNA levels provide a good marker for efficacy of therapy and should be added to regular laboratory monitoring. The goal of therapy is to reduce HBV DNA to as low a level as possible, preferably below the limits of detection. The duration of HBV therapy is not well established; with development of HBEAb, while some individuals without HIV can stop therapy after reversion of HBEAg to negative, there are no clear stopping rules for HIV-coinfected patients.

## Hepatitis C Virus (HCV)

Subset	Pathogens	Therapy
HIV		<b>Genotype 1 (includes 1a &amp; 1b)</b> Preferred therapies: Ledipasvir <sup>6</sup> 90 mg/sofosbuvir 400 mg (Harvoni) (PO) q24h x 12 weeks Paritaprevir 150 mg/ritonavir 100 mg/ombitasvir 25 mg (PO) q24h + dasabuvir <sup>7</sup> 250 mg (Viekira Pak) (PO) q12h + ribavirin <sup>1,2</sup> x weeks <sup>3</sup> Sofosbuvir 400 mg (Savaldi) (PO) q24h + Simeprevir <sup>8</sup> 150 mg (Olysio) (PO) q24h ± ribavirin <sup>1</sup> x weeks <sup>3</sup>
		<b>Genotype 2</b> Preferred therapy: Sofosbuvir 400 mg (Savaldi) (PO) q24h + ribavirin <sup>1</sup> x 12 weeks (16 weeks in cirrhosis)
		<b>Genotype 3</b> Preferred therapies: Sofosbuvir 400 mg (Savaldi) (PO) q24h + ribavirin <sup>1</sup> x 24 weeks Alternate therapy: Sofosbuvir 400 mg (Savaldi) (PO) q24h + pegylated IFN alfa-2a/2b <sup>5</sup> ribavirin <sup>1</sup> x 12 weeks
		<b>Genotype 4</b> Preferred therapies: Ledipasvir <sup>6</sup> 90 mg/sofosbuvir 400 mg (Harvoni) (PO) q24h x 12 weeks Paritaprevir 150 mg/ritonavir 100 mg/ombitasvir 25 mg (Viekira Pak minus dasabuvir) (PO) q24h + ribavirin <sup>1</sup> x 12 weeks Sofosbuvir 400 mg (Sovaldi) (PO) q24h + ribavirin <sup>1</sup> x 24 weeks Alternative therapies: Sofosbuvir 400 mg (Sovaldi) (PO) q24h + pegylated IFN alfa-2a/2b <sup>5</sup> + ribavirin <sup>1</sup> x 12 weeks Sofosbuvir 400 mg (Sovaldi) (PO) q24h + simeprevir <sup>8</sup> 150 mg (Olysio) (PO) q24h ± ribavirin <sup>1</sup> x 12 weeks

## Hepatitis C Virus (HCV) (cont'd)

Subset	Pathogens	Therapy
	<b>Genotype 5</b>	Preferred therapy: Sofosbuvir 400 mg (Sovaidi) (PO) q24h + pegylated IFN alfa-2a/2b <sup>5</sup> + ribavirin <sup>1</sup> x 12 weeks Alternate therapy: Pegylated IFN alfa-2a/2b <sup>5</sup> + ribavirin <sup>1</sup> x 48 weeks
	<b>Genotype 6</b>	Preferred therapy: Ledipasvir <sup>6</sup> 90 mg/sofosbuvir 400 mg (Harvoni) (PO) q24h x 12 weeks Alternate therapy: Sofosbuvir 400 mg (Sovaidi) (PO) q24h + pegylated IFN alfa-2a/2b <sup>5</sup> + ribavirin <sup>1</sup> x 12 weeks

<sup>1</sup>Ribavirin should be dosed as following:

Weight < 75 kg: Ribavirin 1000 mg (PO) daily in 2 divided doses (400 mg in AM, 600 mg in PM)

Weight > 75 kg: Ribavirin 1200 mg (PO) daily in 2 divided doses (600 mg q12h)

<sup>2</sup>In Genotype 1b, ribavirin is only indicated in cirrhosis

<sup>3</sup>12 week therapy without cirrhosis; 24 week therapy with cirrhosis

<sup>4</sup>Simeprevir may be ineffective in patients with HCV genotype 1a Q80K polymorphism

<sup>5</sup>Pegylated IFN alfa-2a should be dosed 180 µg (SC) weekly. Pegylated IFN alfa-2b should be dosed 1.5 µg/kg (SC) weekly. Both types of pegylated IFN alfa are approved to treat genotype 1

<sup>6</sup>Ledipasvir increases tenofovir levels, increasing risk of nephrotoxicity; avoid combination when CrCl < 60 ml/min. Avoid combination with ritonavir-boosted regimens due to potentiation of this effect, unless regimen cannot be changed. Monitor for tenofovir nephrotoxicity.

<sup>7</sup>Due to concerns for significant interactions, paritaprevir/ritonavir/ombitasvir/dasabuvir, should only be co-administered with the following: raltegravir, dolutegravir, enfuvirtide, tenofovir, emtricitabine, lamivudine, and atazanavir. The dose of ritonavir used for boosting of HIV protease inhibitors may need to be adjusted (or held) when administered with paritaprevir/ritonavir/ombitasvir/dasabuvir & then restored when HCV treatment is completed. The HIV protease inhibitor should be administered at the same time as the fixed-dose HCV combination.

<sup>8</sup>Due to concerns for significant interactions, simeprevir should only be co-administered with the following: raltegravir, dolutegravir, rilpivirine, maraviroc, enfuvirtide, tenofovir, emtricitabine, lamivudine and abacavir.

Due to concerns for significant drug interactions, the following combinations are NOT recommended:

Ledipasvir/sofosbuvir and cobicistat & elvitegravir; sofosbuvir or ledipasvir/sofosbuvir with tipranavir; paritaprevir/ritonavir/ombitasvir/dasabuvir and efavirenz, rilpivirine, darunavir or ritonavir-boosted lopinavir; simeprevir and efavirenz, etravirine, nevirapine, cobicistat or any HIV protease inhibitors; ribavirin and didanosine, stavudine or zidovudine.

For patients with CD4 count < 200 cells/µl, Initiation of ART may be considered before HCV treatment.

Adapted from: American Association for the Study of Liver Diseases and Infectious Diseases Society of America. Recommendations for testing, managing, and treating hepatitis C. <http://www.hcvguidelines.org>. 2014.

**Epidemiology:** Hepatitis C virus (HCV) infection is transmitted primarily through blood exposure; sexual and perinatal transmission are also possible but less efficient. A notable exception is sexually transmitted HCV among gay men. Since modes of transmission of HIV and HCV are similar, there are high rates of HCV coinfection in HIV—an estimated 16% of HIV patients overall, including 80% or more of IDUs and 5–10% of gay men. Genotype 1 accounts for 75% of HCV in the United States. HIV accelerates the progression of chronic HCV infection to cirrhosis, liver failure, and hepatocellular carcinoma. Data are conflicting regarding the independent effect of HCV on HIV disease progression, but several studies have shown a markedly higher rate of antiretroviral therapy-induced hepatotoxicity in those with chronic HCV. In some series, liver failure from HCV is one of the leading causes of death in HIV/HCV coinfecting individuals.

**Clinical Presentation:** Persistently elevated liver transaminases; usually asymptomatic.

**Diagnostic Considerations:** All HIV-positive patients should be tested for HCV antibody. If the antibody test is negative but the likelihood of HCV infection is high (IDU, unexplained increase in LFTs), obtain an HCV RNA since false-negative antibody tests may occur, especially in advanced HIV disease. Since LFT elevation does not correlate well with underlying HCV activity, a liver biopsy is the best way to assess the degree of fibrosis and inflammation.

**Therapeutic Considerations:**

Once Diagnosis is Established. Advise patients to abstain from alcohol and administer vaccinations for hepatitis A and B (if non-immune). Also obtain HCV RNA levels with genotype assessment. HCV RNA levels do not have prognostic significance for underlying degree of liver disease, but higher levels make treatment for cure less likely. Genotype results also correlate with cure rates (reported cure rates: 60–75% for genotypes 2 and 3; 15–25% for genotype 1). Some clinicians elect to treat HCV without a liver biopsy due to the risks, costs, and discomfort of the test; the potential to underestimate the degree of HCV activity due to sampling error; and the high rate of treatment success for genotypes 2 and 3.


Optimal Patient Characteristics for HCV Treatment in HIV include no active psychiatric disease or substance abuse; stable HIV disease with undetectable HIV RNA and higher CD<sub>4</sub> cell count; receiving an antiretroviral regimen that does not contain ddI (in particular), d4T, or ZDV; and adherent to medications, follow-up visits, and blood test monitoring. Only a small proportion of patients will meet all these criteria; therefore, to maximize treatment effect, it is important to optimize clinical status prior to starting HCV therapy. Patients should be fully educated regarding the goals and risks of treatment, with provision of written information about side effects, local support groups, and whom to contact with questions. It is useful to administer the first dose of treatment in the office in order to provide instructions on injection techniques.

Choice of Drug Therapy. The treatment of choice for HCV infection is pegylated interferon plus ribavirin. All patients should also receive hepatitis A vaccine and counseling to avoid alcohol use.

- **Pegylated Interferon.** Two different formulations of pegylated interferon are available for treatment of HCV infection: (1) peginterferon alfa-2a (Pegasys), supplied as a pre-mixed solution and administered subcutaneously as a fixed dose of 180 mcg once a week; and (2) peginterferon alfa-2b (Peg-Intron), supplied as a powder that is reconstituted in saline and administered subcutaneously as a weight-based dose once a week. Efficacy and toxicity of these two peginterferon preparations appear to be similar. Three clinical trials have shown that pegylated interferon plus ribavirin is more effective than standard interferon plus ribavirin (N Engl J Med 2004;351:451–9; N Engl J Med 2004;351:438–50; JAMA 2004;292:2839–48). Interferon has numerous side effects, the most important of which are listed in Table 5.9.
- **Ribavirin.** Ribavirin is available in 200-mg capsules. Although the dose used in clinical trials was 800 mg daily (400 mg BID), it appears that a higher initial dose is associated with greater efficacy. Standard dose is now 400 mg qAM and 600 mg qPM for weight < 75 kg, and 600 mg qAM and

600 mg qPM for weight > 75 kg. Ribavirin causes hemolytic anemia that predictably leads to a measurable decline in hemoglobin; this stabilizes by week 4–8 of treatment. Hemolytic anemia may be exacerbated in patients receiving ZDV; if possible an alternative NRTI should be chosen, with preference for tenofovir or abacavir. For symptomatic anemia or patients with coexisting conditions exacerbated by anemia, erythropoietin is used to maintain hemoglobin levels > 10 gm/dL or higher as needed. The typical starting dose of erythropoietin is 40,000 units SQ once a week. ddI is contraindicated during ribavirin administration due to an increased risk of mitochondrial toxicity and hepatic decompensation (Clin Infect Dis 2004;38:e79–e80). d4T should also be avoided because of its potential for inducing mitochondrial toxicity itself. Finally, some data suggest that ABC may interact with ribavirin, reducing efficacy of HCV therapy. Another common side effect of ribavirin is GI distress, which can overlap with a similar effect of interferon. Ribavirin is pregnancy category X (potent teratogen), and can only be used in sexually active women of childbearing potential if they are using 2 forms of birth control. Pregnancy should be avoided until at least 6 months after stopping ribavirin.

**Monitoring.** The monitoring plan for HIV/HCV coinfecting patients consists of both safety and efficacy evaluations (Table 5.10). Results of HCV RNA testing are used to decide between completing 48 weeks of HCV therapy vs. discontinuing treatment at 12 or 24 weeks due to low probability of cure (Figure 5.3).



**Figure 5.3. Management of HCV Infection Based on HCV RNA Testing**

HCV RNA should be assessed at week 12. If the HCV viral load has not dropped by more than 2 logs (100-fold), then the likelihood of achieving a cure is extremely low. Many patients and providers will elect to discontinue therapy at this point, although theoretically low-dose peg-interferon may help improve liver fibrosis independent of its antiviral effect. (This strategy is being tested in clinical trials.) HCV RNA should be measured at weeks 24 and 48. If undetectable at week 48, additional measurements should be obtained at weeks 4, 12, and 24 after stopping treatment. An undetectable HCV RNA at week 24 post-treatment is the current standard for assessing a “sustained virologic response” (SVR), which can be equated with cure. Importantly, patients cured of HCV are susceptible to re-acquisition and should be cautioned about resuming high-risk behavior.

**Table 5.9. Adverse Effects Associated with Interferon**

Side Effect	Comments
Fatigue and flu-like symptoms (fever, chills, muscle aches, headache)	Dose at night so that some initial symptoms can be slept through. Symptoms may not peak until 48–78 hours after weekly dose and can be managed with acetaminophen (maximum 1300 mg/day) or ibuprofen plus good hydration. Fatigue is sometimes a manifestation of thyroid dysregulation or anemia; monitor thyroid function tests and CBC
Depression	Interferon can aggravate life-threatening psychiatric conditions. Low threshold for initiating anti-depressant therapy. Patients with a pre-existing history of depression or other psychiatric disease should be followed closely by mental health professionals during HCV treatment
Leukopenia, thrombocytopenia, anemia	Cytopenias are more common in HIV/HCV co-infection. Can treat with G-CSF 300 mcg 3x/week if absolute neutrophil count is < 500/mm <sup>3</sup> . If neutropenia persists, decrease dose of interferon. If platelet count falls to < 80,000/mm <sup>3</sup> , also consider decreasing interferon dose (some clinicians tolerate counts down to 50,000/mm <sup>3</sup> ). Low threshold for use of erythropoietin (EPO) 40,000 U/week for anemia (see section on ribavirin)
Mouth ulcers	Topical viscous lidocaine or sucralfate is helpful in some
Gastrointestinal symptoms	Nausea and anorexia are the most common, often leading to weight loss. Advise patients to eat several small meals daily rather than a few large meals
Hair loss	Reversible after completion of therapy

**Table 5.10. Monitoring Plan During Treatment for HCV Infection**

	Weeks of Treatment								
	Baseline	2	4	8	12	16	20	24	24–28
CBC	X	X	X	X	X	X	X	X	q4 weeks
LFTs + metabolic panel	X		X	X	X	X	X	X	q4 weeks
HCV RNA	X		X <sup>*</sup>		X <sup>*</sup>			X <sup>**</sup>	q12 weeks
HIV RNA + CD <sub>4</sub> profile	X				X <sup>†</sup>			X	q12 weeks
TSH	X				X			X	q12 weeks
Depression	X	X	X	X	X	X	X	X	ongoing

\* Patients who have not dropped  $\geq 2$  logs from baseline HCV RNA at 12 weeks (early virologic response) have < 3% chance of obtaining a sustained viral response (undetectable HCV RNA 6 months post-treatment). Optimally, the HCV RNA will be undetectable by week 12 (complete early virologic response). One proposed strategy for patients with low-level detectable HCV RNA at week 12 (partial early virologic response) is to lengthen the course of therapy to 72 weeks or longer to reduce the risk of relapse; this is currently under investigation.

\*\* If undetectable HCV RNA at 24 weeks, continue therapy for an additional 24 weeks (if known genotype 2 or 3 discuss option to stop, but some experts agree co-infected patients should continue treatment for 48 weeks to decrease risk of relapse). If HCV RNA is positive at 24 weeks, consider discontinuing HCV therapy. Even in patients without viral response, treatment may improve liver histology. Also see Figure 5.1.

<sup>†</sup> Anticipate decrease in absolute cell count but stable CD<sub>4</sub> %.

<sup>‡</sup> Check HCV RNA at week 4; if no decline in HIV RNA is seen, response is extremely unlikely. In contrast,  $\geq 1$  log decline can be very motivating and treatment should be continued. Best chance at cure is with an undetectable HIV RNA at 4 weeks, sometimes called a rapid virologic response.

**Table 5.10. Monitoring Plan During Treatment for HCV Infection (cont'd)**

	Weeks of Treatment								
	Baseline	2	4	8	12	16	20	24	
Ophthalmologic exam	X <sup>†</sup>				X			X	q12 weeks
PT	X					X			q12 weeks
Pregnancy test	Perform at regular intervals if appropriate								

<sup>†</sup> Necessary for patients with a history of retinopathy; IFN package insert recommends screening for all patients prior to treatment. Many clinicians choose to defer the initial exam and monitor for disturbances in vision and loss of color perception.

Adapted from: Sax PE, Cohen CJ, Kuritzkes DR. *HIV Essentials* (6<sup>th</sup> Ed), Jones & Bartlett, Sudbury, MA, 2013.

## REFERENCES AND SUGGESTED READINGS

- Aberg JA, Gallant JE, Ghanem KG, et al. Primary care guidelines for the management of persons infected with HIV: 2013 update by the HIV medicine association of the IDSA. *Clin Infect Dis* 58:1–10, 2014.
- Antiretroviral Postexposure Prophylaxis After Sexual, Injection-Drug Use, or Other Nonoccupational Exposure to HIV in the United States. *MMWR* 54(RR02): 1–20, 2005.
- Gandhi M, Gandhi RT. Single-Pill Combination Regimens for Treatment of HIV-1 Infection. *NEJM* 371:248-259, 2014.
- Guidelines for Treating Opportunistic Infections Among HIV-Infected Adults and Adolescents. [www.hivatis.org](http://www.hivatis.org). January 29, 2008.
- Gunthard HF, Aberg JA, Eron JJ, Hoy JF, et al. Antiretroviral Treatment of Adult HIV Infection. 2014 Recommendations of the International Antiviral Society USA Panel. *J Amer Med Assoc*. 312:410-425, 2014.
- Johnson LB, Saravolatz LD. The Quad pill, a once-daily combination therapy for HIV infection. *Clin Inf Dis* 58:93–8, 2014.
- Marrazzo JM, del Rio C, Holtgrave DR, Cohen MS, et al. HIV Prevention in Clinical Settings. 2014 Recommendations of the International Society USA Panel. *J Amer Med Assoc*. 312: 390-409, 2014.
- McLean HQ, Fiebelkorn AP, Temte JJ, et al. Guidelines for prevention and treatment of opportunistic infections in HIV-infected adults and adolescents. *AIDS Info* 6:1–416, 2013.
- Panel on Antiretroviral Guidelines for Adults and Adolescents. Guidelines for the use of antiretroviral agents in HIV-1-infected adults and adolescents. Department of Health and Human Services. <http://aidsinfo.nih.gov/contentfiles/lvguidelines/AdultandAdolescentGL.pdf>, 2013.
- Public Health Service Task Force Recommendations for Use of Antiretroviral Drugs in Pregnant HIV-1-Infected Women for Maternal Health and Interventions to Reduce Perinatal HIV-1 Transmission in the United States [www.hivatis.org](http://www.hivatis.org). November 2, 2007.
- Recommendation on Integrase Inhibitor Use in Antiretroviral Treatment-Naïve HIV Infected Individuals from the HHS Panel on Antiretroviral Guidelines for Adults and Adolescents. [http://aidsinfo.nih.gov/contentfiles/AdultARV\\_INSTIREcommendations.pdf](http://aidsinfo.nih.gov/contentfiles/AdultARV_INSTIREcommendations.pdf), 2013.
- Truong WR, Schafer JJ, Short WR. Once Daily, Single-Tablet Regimens for the Treatment of HIV-1 Infection. *Pharm & Therap*. 40:44-55, 2015.

**TEXTBOOKS**

- Dolin R, Masur H, Saag MS (eds). *AIDS Therapy* (3<sup>rd</sup> Ed), Churchill Livingstone, New York, 2008.
- Gorbach SL, Bartlett JG, Blacklow NR (eds). *Infectious Diseases* (3<sup>rd</sup> Ed), Philadelphia, Lippincott, Williams & Wilkins, 2004.
- Bennett JE, Dolin R, Blaser MJ (eds). *Mandell, Douglas, and Bennett's Principles and Practice of Infectious Diseases* (8<sup>th</sup> Ed), Philadelphia, Elsevier, 2015.
- Sax PE, Cohen CJ, Kuritzkes DR (eds). *HIV Essentials* (7<sup>th</sup> Ed), Jones & Bartlett, Sudbury, MA, 2014.
- Wormser GP (ed). *AIDS* (4<sup>th</sup> Ed), Elsevier, Philadelphia, 2004.

## Chapter 6

**Prophylaxis and Immunizations****Pierce Gardner, MD, John L. Brusck, MD****Jean E. Hage, MD, Ronald L. Nichols, MD****Staci A. Fischer, MD, Arthur Gran, MD****Muhammed Raza, MBBS, Burke A. Cunha, MD****Maria D. Mileno, MD**

Surgical Prophylaxis . . . . .	354
Post-Exposure Prophylaxis . . . . .	358
Chronic Medical Prophylaxis . . . . .	364
HIV Pre-Exposure Prophylaxis (PrEP) . . . . .	365
HIV Post-Exposure Prophylaxis (PEP) . . . . .	366
HIV Opportunistic Infections . . . . .	366
Transplant Prophylaxis . . . . .	367
Endocarditis Prophylaxis . . . . .	369
Travel Prophylaxis . . . . .	371
Malaria Prophylaxis . . . . .	375
Tetanus Prophylaxis . . . . .	377
Immunizations . . . . .	378
References and Suggested Readings . . . . .	382

\* For additional information on the prophylaxis of opportunistic infections in HIV see Chapter 5, p. 320.



---

**SURGICAL PROPHYLAXIS**


---

Antibiotic prophylaxis is designed to prevent infection for a defined period of time. **Prophylaxis is most likely to be effective when given for a short duration against a single pathogen with a known sensitivity pattern**, and least likely to be effective when given for a long duration against multiple organisms with varying/unpredictable sensitivity patterns (Table 6.1). It is a common misconception that antibiotics used for prophylaxis should not be used for therapy and vice versa. **The only difference between prophylaxis and therapy is the inoculum size and the duration of antibiotic administration:** In prophylaxis, there is no infection, so the inoculum is minimal/none and antibiotics are administered only for the duration of exposure/surgical procedure. With therapy, the inoculum is large (infection already exists), and antibiotics are continued until the infection is eradicated.

**Table 6.1. Factors Affecting the Efficacy of Surgical Antibiotic Prophylaxis**

Number of Organisms	Susceptibility Pattern	Duration of Protection	Efficacy of Prophylaxis
Single organism	Predictable	Short	Excellent
Multiple organisms	Predictable	Short	Excellent
Single organism	Unpredictable	Short	Good
Single organism	Predictable	Long	Good
Multiple organisms	Unpredictable	Long	Poor/none

**Antibiotic prophylaxis is designed to achieve effective antibiotic serum/tissue concentrations at the time of initial surgical incision, and is maintained throughout the “vulnerable period” of the procedure (i.e., time between skin incision and skin closure)** (Table 6.2). If prophylaxis is given too early, antibiotic levels will be suboptimal/non-existent when protection is needed. Properly timed pre-operative antibiotic prophylaxis is desirable for optimal effectiveness since **antibiotics given after skin closure are unlikely to be effective**. When no infection exists prior to surgery (clean/clean contaminated surgery), single-dose prophylaxis is preferred. When infection is present/likely prior to surgery (“dirty” surgery, e.g., perforated colon, TURP in the presence of positive urine cultures, repair of open fracture), antibiotics are given for > 1 day and represent early therapy, not true prophylaxis. Parenteral cephalosporins are commonly used for surgical prophylaxis, and ordinarily given as a bolus injection/rapid IV infusion 15–60 minutes prior to the procedure. Prophylaxis with vancomycin or gentamicin is given by slow IV infusion over 1–2 hours, starting ~1–2 hours prior to the procedure.

**Table 6.2. Surgical Prophylaxis**

Procedure	Usual Organisms	Preferred Prophylaxis	Alternate Prophylaxis	Comments
CNS shunt (VP/VA) placement, craniotomy, open CNS trauma	<i>S. epidermidis</i> (CoNS) <i>S. aureus</i> (MSSA)	<u>MRSA/MRSE unlikely</u> Ceftriaxone 1 gm (IV) × 1 dose  <u>MRSA/MRSE likely</u> Linezolid 600 mg (IV) × 1 dose	<u>MRSA/MRSE unlikely</u> Cefotaxime 2 gm (IV) × 1 dose <b>or</b> Ceftizoxime 2 gm (IV) × 1 dose  <u>MRSA/MRSE likely</u> Linezolid 600 mg (PO) × 1 dose <b>or</b> Vancomycin 1 gm (IV) × 1 dose <b>or</b> Minocycline 200 mg (IV) × 1 dose	Administer immediately prior to procedure. Vancomycin protects against wound infections, but may not prevent CNS infections. Give vancomycin slowly IV over 1 hour prior to procedure.
Thoracic (non-cardiac) surgery	<i>S. aureus</i> (MSSA)	Cefazolin 1 gm (IV) × 1 dose <b>or</b> Ceftriaxone 1 gm (IV) × 1 dose	Cefotaxime 2 gm (IV) × 1 dose <b>or</b> Ceftizoxime 2 gm (IV) × 1 dose	Administer immediately prior to procedure.
Cardiac valve replacement surgery	<i>S. epidermidis</i> (MSSE/MRSE) <i>S. aureus</i> (MSSA/MRSA) Enterobacter	Vancomycin 1 gm (IV) × 1 dose <b>plus</b> Gentamicin 120 mg (IV) × 1 dose	Linezolid 600 mg (IV) × 1 dose <b>plus</b> Gentamicin 120 mg (IV) × 1 dose	Administer vancomycin and gentamicin slowly IV over 1 hour prior to procedure.
Coronary artery bypass graft (CABG) surgery	<i>S. aureus</i> (MSSA)	Cefazolin 2 gm (IV) × 1 dose <b>or</b> Ceftriaxone 1 gm (IV) × 1 dose	Cefotaxime 2 gm (IV) × 1 dose <b>or</b> Ceftizoxime 2 gm (IV) × 1 dose	Administer immediately prior to procedure. Except for ceftriaxone, repeat dose intraoperatively for procedures lasting > 3 hours.
Biliary tract surgery	<i>E. coli</i> <i>Klebsiella</i> <i>E. faecalis</i> (VSE)	Meropenem 1 gm (IV) × 1 dose <b>or</b> Piperacillin 4 gm (IV) × 1 dose	Ampicillin/sulbactam 3 gm (IV) × 1 dose	Administer immediately prior to procedure (anaerobic coverage unnecessary).

**Table 6.2. Surgical Prophylaxis** (cont'd)

Procedure	Usual Organisms	Preferred Prophylaxis	Alternate Prophylaxis	Comments
Hepatic surgery	E. coli Klebsiella E. faecalis (VSE) B. fragilis	Ampicillin/sulbactam 3 gm (IV) × 1 dose <b>or</b> Piperacillin 4 gm (IV) × 1 dose	Meropenem 1 gm (IV) × 1 dose <b>or</b> Moxifloxacin 400 mg (IV) × 1 dose	Administer immediately prior to procedure.
Stomach, upper small bowel surgery	S. aureus (MSSA) Group A streptococci	Ceftriaxone 1 gm (IV) × 1 dose <b>or</b> Cefazolin 1 gm (IV) × 1 dose	Cefotaxime 2 gm (IV) × 1 dose <b>or</b> Ceftizoxime 2 gm (IV) × 1 dose	Administer immediately prior to procedure (anaerobic coverage unnecessary).
Distal small bowel, colon surgery	E. coli Klebsiella B. fragilis	<u>Oral</u> Neomycin* <b>plus either</b> Erythromycin base* <b>or</b> Metronidazole* <u>Parenteral</u> Ertapenem 1 gm (IV) × 1 dose	Piperacillin 3 gm (IV) × 1 dose <b>or</b> Cefoxitin 2 gm (IV) × 1 dose <b>or combination therapy with</b> Metronidazole 1 gm (IV) × 1 dose <b>plus either</b> Ceftriaxone 1 gm (IV) × 1 dose <b>or</b> Levofloxacin 500 mg (IV) × 1 dose <b>or</b> Gentamicin 240 mg (IV) × 1 dose	Administer immediately prior to procedure. Give gentamicin slowly IV over 1 hour.
Pelvic (OB/GYN) surgery	Aerobic GNBs Anaerobic streptococci B. fragilis	Ceftriaxone 1 gm (IV) × 1 dose <b>plus</b> Metronidazole 1 gm (IV) × 1 dose	Cefotetan 2 gm (IV) × 1 dose <b>or</b> Cefoxitin 2 gm (IV) × 1 dose <b>or</b> Ceftizoxime 2 gm (IV) × 1 dose	Administer immediately prior to procedure.

\* After appropriate diet and catharsis give either neomycin 1 gm (PO) plus erythromycin base 1 gm (PO) at 1 pm, 2 pm, and 11 pm, or give neomycin 2 gm (PO) plus metronidazole 2 gm (PO) at 7 pm and 11 pm the day before an 8 am operation.

**Table 6.2. Surgical Prophylaxis** (cont'd)

Procedure	Usual Organisms	Preferred Prophylaxis	Alternate Prophylaxis	Comments
Orthopedic prosthetic implant surgery (total hip/knee replacement)	<i>S. epidermidis</i> (CoNS) <i>S. aureus</i> (MSSA)	<u>MRSA/MRSE unlikely</u> Cefazolin 2 gm (IV) × 1 dose  <u>MRSA/MRSE likely</u> Vancomycin 1 gm (IV) × 1 dose	<u>MRSA/MRSE unlikely</u> Ceftriaxone 1 gm (IV) × 1 dose  <u>MRSA/MRSE likely</u> Linezolid 600 mg (IV) × 1 dose	Administer immediately prior to procedure. Post-operative doses are ineffective and unnecessary.
Arthroscopy	<i>S. aureus</i> (MSSA)	Cefazolin 1 gm (IV) × 1 dose <b>or</b> Ceftriaxone 1 gm (IV) × 1 dose	Cefotaxime 2 gm (IV) × 1 dose <b>or</b> Ceftizoxime 2 gm (IV) × 1 dose	Pre-procedure prophylaxis is usually unnecessary in clean surgical procedures.
Orthopedic surgery (open fracture)	<i>S. aureus</i> (MSSA) Aerobic GNBs	Ceftriaxone 1 gm (IV) × 1 week	Clindamycin 600 mg (IV) q8h × 1 week <b>plus</b> Gentamicin 240 mg (IV) q24h × 1 week	Represents early therapy, not true prophylaxis. Duration of post-op antibiotics depends on severity of infection.
Urological implant surgery	<i>S. aureus</i> (MSSA) Aerobic GNBs bacilli	Ceftriaxone 1 gm (IV) × 1 dose	Cefotaxime 2 gm (IV) × 1 dose <b>or</b> Ceftizoxime 2 gm (IV) × 1 dose	Administer immediately prior to procedure.
TURP, cystoscopy	<i>P. aeruginosa</i> <i>P. cepacia</i> <i>P. maltophilia</i> <i>E. faecalis</i> (VRE) Aerobic GNBs	Ciprofloxacin 400 mg (IV) × 1 dose <b>or</b> Piperacillin 4 gm (IV) × 1 dose	Levofloxacin 500 mg (IV) × 1 dose <b>or</b> Gatifloxacin 400 mg (IV) × 1 dose	Prophylaxis given to TURP patients with positive pre-op urine cultures. Represents early therapy, not true prophylaxis. No prophylaxis required for TURP if pre-op urine culture is negative.
	<i>E. faecium</i> (VRE)	Linezolid 600 mg (IV) × 1 dose	Quinupristin/dalfopristin 7.5 mg/kg (IV) × 1 dose	

MSSA/MRSA = methicillin-sensitive/resistant *S. aureus*; MSSE/MRSE = methicillin-sensitive/resistant *S. epidermidis*.

---

**POST-EXPOSURE PROPHYLAXIS**


---

Some infectious diseases can be prevented by post-exposure prophylaxis (PEP). To be maximally effective, PEP should be administered within 24 hours of the exposure, since the effectiveness of prophylaxis > 24 hours after exposure decreases. PEP is usually reserved for persons with close/intimate contact with an infected individual. Casual contact does not warrant PEP.

**Table 6.3. Post-Exposure Prophylaxis**

Exposure	Usual Organisms	Preferred Prophylaxis	Alternate Prophylaxis	Comments
Meningitis	N. meningitidis	Any quinolone (PO) × 1 dose	Minocycline 100 mg (PO) q12h × 2 days <b>or</b> Rifampin 600 mg (PO) q12h × 2 days	Must be administered within 24 hours of close face-to-face exposure to be effective. Otherwise, observe and treat if infection develops.
	H. influenzae	Rifampin 600 mg (PO) q24h × 3 days	Any quinolone (PO) × 3 days	Must be administered within 24 hours of close face-to-face exposure to be effective. H. influenzae requires 3 days of prophylaxis.
Influenza	Influenza A or B	Oseltamivir (Tamiflu) 75 mg (PO) q24h for duration of outbreak (or at least 7 days after close contact to an infected person). <sup>5</sup>	Rimantadine 100 mg (PO) q12h* for duration of outbreak (or at least 7–10 days after close contact to infected person) <b>or</b> Amantidine 200 mg (PO) q24h <sup>†</sup> for 7–10 days or outbreak	Give to non-immunized contacts and high-risk contacts even if immunized. Begin at onset of outbreak or within 3 days of close contact to an infected person. Oseltamivir is active against both influenza A and B; rimantadine and amantidine are only active against influenza A. Oseltamivir may be ineffective due to resistance.

<sup>5</sup> Dose for CrCl > 30–60 mL/min = 30 mg (PO) q24h; for CrCl > 10–30 mL/min = 30 mg (PO) q48h.

**Table 6.3. Post-Exposure Prophylaxis (cont'd)**

Exposure	Usual Organisms	Preferred Prophylaxis	Alternate Prophylaxis	Comments
Avian influenza	Influenza A (H <sub>5</sub> N <sub>1</sub> )	Oseltamivir (as for viral influenza, above)	Rimantadine <b>or</b> Amantadine <sup>†</sup> (as for viral influenza, above)	Oseltamivir may be ineffective due to resistance.
Swine influenza	Influenza A (H <sub>1</sub> N <sub>1</sub> )	Oseltamivir above (as for viral influenza, above)	None	Amantadine/rimantadine ineffective Continue PEP for swine influenza (H <sub>1</sub> N <sub>1</sub> ) × 10 days post-close exposure. <sup>‡</sup>
Pertussis	B. pertussis	Erythromycin 500 mg (PO) q6h × 2 weeks	TMP-SMX 1 SS tablet (PO) q12h × 2 weeks <b>or</b> Levofloxacin 500 mg (PO) q24h × 2 weeks	Administer as soon as possible after exposure. Effectiveness is greatly reduced after 24 hours.
Diphtheria	C. diphtheriae	Erythromycin 500 mg (PO) q6h × 1 week <b>or</b> Benzathine penicillin 1.2 mu (IM) × 1 dose	Azithromycin 500 mg (PO) q24h × 3 days	Administer as soon as possible after exposure. Effectiveness is greatly reduced after 24 hours.
TB	M. tuberculosis	Rifampin 600 mg (PO) q24h × 4 months <sup>§</sup> <b>or</b> INH 300 mg (PO) q24h × 9 months	Rifapentine 900 mg (PO) q week <b>plus</b> INH 900 mg (PO) q week × 3 months (D.O.T.)	For INH, monitor SGOT/SGPT weekly × 4, then monthly. Mild elevations are common and resolve spontaneously. D/C INH if SGOT/SGPT levels ≥ 5 × normal.

\* For elderly, severe liver dysfunction, or CrCl < 10 cc/min, give 100 mg (PO) q24h.

† For age ≥ 65 years, give 100 mg (PO) q24h. For renal dysfunction, give 200 mg (PO) load followed by 100 mg q24h (CrCl 30–50 ml/min), 100 mg q48h (CrCl 15–29 ml/min), or 200 mg weekly (CrCl < 15 ml/min).

‡ *Pediatric prophylaxis:* < 15 kg: 30 mg PO qd for 10 days, > 15–24 kg: 45 mg PO qd for 10 days, 24–40 kg: 60 mg PO qd for 10 days, > 40 kg: same as adults.

§ INH preferred if patient uses contact lens, has HIV, or is taking medications that may interact with rifampin.

**Table 6.3. Post-Exposure Prophylaxis** (cont'd)

Exposure	Usual Organisms	Preferred Prophylaxis	Alternate Prophylaxis	Comments
Gonorrhea (GC)	N. gonorrhoeae	Ceftriaxone 125 mg (IM) × 1 dose	Spectinomycin 2 gm (IM) × 1 dose <b>or</b> Any oral quinolone × 1 dose	Administer as soon as possible after sexual exposure (≤ 72 hours). Ceftriaxone also treats incubating syphilis.
Syphilis	T. pallidum	Benzathine penicillin 2.4 mu (IM) × 1 dose	Doxycycline 100 mg (PO) q12h × 1 week	Administer as soon as possible after sexual exposure. Obtain HIV serology.
Chancroid	H. ducreyi	Ceftriaxone 250 mg (IM) × 1 dose	Azithromycin 1 gm (PO) × 1 dose <b>or</b> Any oral quinolone × 3 days	Administer as soon as possible after sexual exposure. Obtain HIV and syphilis serologies.
Non-gonococcal urethritis (NGU)	C. trachomatis U. urealyticum M. genitalium	Azithromycin 1 gm (PO) × 1 dose <b>or</b> Doxycycline 100 mg (PO) q12h × 1 week	Any oral quinolone × 1 week	Administer as soon as possible after sexual exposure. Also test for gonorrhea/Ureaplasma.
Varicella (chicken-pox)	VZV	<u>Preferred:</u> For exposure < 72 hours, give varicella-zoster immune globulin (VZIG) 625 mcg (IM) × 1 dose to immuno-compromised hosts and pregnant women (esp. with respiratory conditions). For others or exposure > 72 hours, consider acyclovir 800 mg (PO) 5x/day × 5–10 days <u>Alternate:</u> Varicella vaccine 0.5 mL (SC) × 1 dose. Repeat in 4 weeks		Administer as soon as possible after exposure (≤ 72 hours). Varicella vaccine is a live attenuated vaccine and should not be given to immunocompromised or pregnant patients. If varicella develops, start acyclovir treatment immediately.

**Table 6.3. Post-Exposure Prophylaxis (cont'd)**

Exposure	Usual Organisms	Preferred Prophylaxis	Alternate Prophylaxis	Comments
Hepatitis A (HAV)	Hepatitis A virus	HAV vaccine 1 mL (IM) × 1 dose	Immune serum globulin (IG) 0.02 mL/kg (IM) × 1 dose	Give HAV vaccine alone if within 14 days after exposure.
Hepatitis B (HBV)	Hepatitis B virus	<u>Unvaccinated</u> Hepatitis B immune globulin (HBIG) 0.06 mL/kg (IM) × 1 dose <b>plus</b> HBV vaccine (40 mcg HBsAg/mL) deep deltoid (IM) at 0, 1, 6 months (can use 10-mcg dose in healthy adults < 40 years)	<u>Previously vaccinated</u> <i>Known responder</i> (anti-HBsAg antibody levels ≥ 10 IU/mL): No treatment. <i>Known non-responder</i> (anti-HBsAg antibody levels < 10 IU/mL): Treat as if unvaccinated. <i>Antibody status unknown</i> : Obtain HBsAg antibody levels to determine immunity status. If testing is not possible or results are not available within 24 hours of exposure, give HBIG plus 1 dose of HBV vaccine (booster).	
Hepatitis C (HCV)	Hepatitis C virus	None		
Rocky Mountain spotted fever	<i>R. rickettsia</i>	Doxycycline 100 mg (PO) q12h × 1 week	Any oral quinolone × 1 week	Administer prophylaxis after removal of Dermacentor tick.
Lyme disease	<i>B. burgdorferi</i>	Doxycycline 200 mg (PO) × 1 dose	Amoxicillin* 1 gm (PO) q8h × 3 days <b>or</b> Any oral 1 <sup>st</sup> gen. cephalosporin* × 3 days <b>or</b> Azithromycin* 500 mg (PO) q24h × 3 days	If tick is in place ≥ 72 hours or is grossly engorged, prophylaxis may be given after tick is removed. Otherwise, prophylaxis is usually not recommended.

HDCV = human diploid cell vaccine, HRIG = human rabies immune globulin, PCEC = purified chick embryo cells, RVA = rabies vaccine absorbed.

\* All or as much of the full dose of HRIG should be injected into the wound, and the remaining vaccine should be injected IM into the deltoid. Do not give HRIG at the same site or through the same syringe with PCEC, RVA, or HDCV.



**Table 6.3. Post-Exposure Prophylaxis** (cont'd)

Exposure	Usual Organisms	Preferred Prophylaxis	Alternate Prophylaxis	Comments
		* Although experience is limited, single-dose prophylaxis with these agents is probably also effective.		
Zoonotic diseases (plague, anthrax)	B. anthracis Y. pestis	Doxycycline 100 mg (PO) q12h for duration of exposure	Any oral quinolone for duration of exposure	Continued for the duration of a naturally-acquired exposure/outbreak. See p. 363 for bioterrorist plague/anthrax recommendations.
Rabies	Rabies virus	<u>No Previous Immunization</u> HRIG 20 IU/kg* <b>plus either</b> PCEC 1 mL (IM) in deltoid <b>or</b> RVA 1 mL (IM) in deltoid <b>or</b> HDCV 1 mL (IM) in deltoid  PCEC, RVA, HDCV given on days 0, 3, 7, 14, and 28 post-exposure	<u>Previous Immunization</u> PCEC 1 mL (IM) in deltoid on days 0 and 3 <b>or</b> RVA 1 mL (IM) in deltoid on days 0 and 3 <b>or</b> HDCV 1 mL (IM) in deltoid on days 0 and 3	Following unprovoked or suspicious dog or cat bite, immediately begin prophylaxis if animal develops rabies during a 10-day observation period. If dog or cat is suspected of being rabid, begin vaccination sequence immediately. Raccoon, skunk, bat, fox, and most wild carnivore bites should be regarded as rabid, and bite victims should be vaccinated against rabies immediately (contact local health department regarding rabies potential of animals in your area). All potential rabies wounds should immediately be thoroughly cleaned with soap and water. Do not inject rabies vaccine IV (may cause hypotension/shock). Serum sickness may occur with HDCV.

HDCV = human diploid cell vaccine, HRIG = human rabies immune globulin, PCEC = purified chick embryo cells, RVA = rabies vaccine absorbed.

\* All or as much of the full dose of HRIG should be injected into the wound, and the remaining vaccine should be injected IM into the deltoid. Do not give HRIG at the same site or through the same syringe with PCEC, RVA, or HDCV.

**Table 6.3. Post-Exposure Prophylaxis** (cont'd)

Exposure	Usual Organisms	Preferred Prophylaxis	Alternate Prophylaxis	Comments
<b>BIOTERRORIST AGENTS</b>				
Anthrax <i>Inhalation/ cutaneous</i>	<i>B. anthracis</i>	Doxycycline 100 mg (PO) q12h × 60 days <b>or</b> Ciprofloxacin 500 mg (PO) q12h × 60 days <b>or</b> Levofloxacin 500 mg (PO) q24h × 60 days	Amoxicillin 1 gm (PO) q8h × 60 days	Duration of anthrax PEP based on longest incubation period of inhaled spores in nares.
Tularemia pneumonia	<i>F. tularensis</i>	Doxycycline 100 mg (PO) q12h × 2 weeks	Ciprofloxacin 500 mg (PO) q12h × 2 weeks <b>or</b> Levofloxacin 500 mg (PO) q24h × 2 weeks	Duration of PEP for tularemia is 2 weeks, not 1 week as for plague.
Pneumonic plague	<i>Y. pestis</i>	Doxycycline 100 mg (PO) q12h × 7 days <b>or</b> Ciprofloxacin 500 mg (PO) q12h × 7 days <b>or</b> Levofloxacin 500 mg (PO) q24h × 7 days	Chloramphenicol 500 mg (PO) q6h × 7 days	Pneumonic plague should be considered bioterrorism since most natural cases of plague are bubonic plague.
Smallpox	Variola virus	Smallpox vaccine ≤ 4 days after exposure	None	Smallpox vaccine is protective when diluted 1:5.
Viral hemorrhagic fever	Lassa fever	Ribavirin loading dose: 35mg/kg (PO) (not to exceed 2.5 g), then 15 mg/kg (PO) (not to exceed 1g) q8h × 10 days	None	Decrease maintenance dose to 7.5 mg/kg (PO) q8h if CrCl < 50/ml/min.

### CHRONIC MEDICAL PROPHYLAXIS

Some infectious diseases are prone to recurrence/relapse and may benefit from intermittent or chronic suppressive therapy. The goal of suppressive therapy is to minimize the frequency/severity of recurrent infectious episodes.

**Table 6.4. Chronic Medical Prophylaxis**

Disorder	Usual Organisms	Preferred Prophylaxis	Alternate Prophylaxis	Comments
Asplenia/ impaired splenic function	<i>S. pneumoniae</i> <i>H. influenzae</i> <i>N. meningitidis</i>	Amoxicillin 1 gm (PO) q24h indefinitely	Respiratory quinolone* (PO) q24h indefinitely	Long-term prophylaxis effective. Vaccines may be given but are not always protective. Use amoxicillin in children.
UTIs (recurrent)	Gram-negative bacilli Enterococci	Nitrofurantoin 100 mg (PO) q24h × 6 months	Amoxicillin 500 mg (PO) q24h × 6 mos <b>or</b> TMP-SMX 1 SS tablet (PO) q24h × 6 months	Prophylaxis for reinfection UTIs (≥ 3 per year). Relapse UTIs should be investigated for stones, abscesses, or structural problems.
Asymptomatic bacteriuria in pregnancy	Gram-negative bacilli	Nitrofurantoin 100 mg (PO) q24h × 1 week	Amoxicillin 1 gm (PO) q24h × 1 week	Prophylaxis prevents symptomatic infections.
Prophylaxis of fungal infections in neutropenic patients <sup>†</sup>	<i>Candida albicans</i> Non-albicans <i>Candida</i> <i>Aspergillus</i>	Posaconazole 200 mg (PO) q8h <b>or</b> Itraconazole 200 mg (PO) q12h		Prophylaxis given until neutropenia resolves (absolute neutrophil count ≥ 500 cells/mm <sup>3</sup> ).
Recurrent genital herpes (< 6 episodes/ year)	<i>H. simplex</i> (HSV-2)	Famciclovir 125 mg (PO) q12h × 5 days <b>or</b> Valacyclovir 500 mg (PO) q24h × 5 days	Acyclovir 200 mg (PO) 5x/day × 5 days	Begin therapy as soon as lesions appear. (For HIV disease, see Ch. 5.) Famciclovir 1 gm (PO) q12h × 1 day ↓ lesion progression by 2 days.
Recurrent genital herpes (> 6 episodes/ year)	<i>H. simplex</i> (HSV-2)	Famciclovir 250 mg (PO) q12h × 1 year <b>or</b> Valacyclovir 1 gm (PO) q24h × 1 year	Acyclovir 400 mg (PO) q12h × 1 year	Suppressive therapy is indicated for frequent recurrences. (For HIV disease, see Ch. 5.)

**Table 6.4. Chronic Medical Prophylaxis (cont'd)**

Disorder	Usual Organisms	Preferred Prophylaxis	Alternate Prophylaxis	Comments
Acute exacerbation of chronic bronchitis (AECB)	<i>S. pneumoniae</i> H. <i>influenzae</i> <i>M. catarrhalis</i>	Moxifloxacin 400 mg or levofloxacin 500 mg or gatifloxacin 400 mg or gemifloxacin 320 mg (PO) q24h × 5 days <b>or</b> Amoxicillin/clavulanic acid XR 2 tablets (PO) q12h × 5 days <b>or</b> Clarithromycin XL 1 gm (PO) q24h × 5 days <b>or</b> Doxycycline 100 mg (PO) q12h × 5 days <b>or</b> Azithromycin 500 mg (PO) × 3 days		Treat each episode individually.
Acute rheumatic fever (ARF)	Group A streptococci	Benzathine penicillin 1.2 mu (IM) monthly until age 30	Amoxicillin 500 mg (PO) q24h <b>or</b> Azithromycin 500 mg (PO) q72h until age 30	Group A streptococcal pharyngitis and acute rheumatic fever are uncommon after age 30.
Neonatal Group B streptococcal (GBS) infection (primary prevention)	Group B streptococci	Ampicillin 2 gm (IV) q4h at onset of labor until delivery	Clindamycin 600 mg (IV) q8h at onset of labor until delivery <b>or</b> Vancomycin 1 gm (IV) q12h at onset of labor until delivery	Indications: previous infant with GBS infection, maternal GBS colonization/infection during pregnancy, vaginal/rectal culture of GBS after week 35 of gestation, delivery ≤ week 37 of gestation without labor/ruptured membranes, ruptured membranes ≥ 12 hrs, or intrapartum temp ≥ 100.4°F.

\* Levofloxacin 500 mg (PO) or gatifloxacin 400 mg (PO) q24h or moxifloxacin 400 mg (PO).

† During induction chemotherapy for acute myelogenous leukemia (AML) or myelodysplastic syndrome (MDS).

**Table 6.5. HIV Pre-Exposure Prophylaxis (PrEP)**

Disorder	Usual Organisms	Preferred Prophylaxis	Alternate Prophylaxis	Comments
Repeated potential HIV exposures	HIV	Truvada 1 tablet (PO) q24h × 90 days (recheck HIV status)		Avoid if CrCl < 60 ml/min. Avoid if breast feeding.

**Table 6.6. HIV Post-Exposure Prophylaxis (PEP)**

Exposure	Preferred regimen	Recommended regimen	Alternate recommended regimen	Comments
HIV	<p><b>Truvada 1</b> tablet (PO) q24h [tenofovir (TDF) 300 mg (PO) q24h + emtricitabine (FTC) 200 mg (PO) q24h] × 4 weeks</p> <p><b>plus either</b></p> <p>Raltegravir (RAL) 400 mg (PO) q12h × 4 weeks</p> <p><b>or</b></p> <p>Dolutegravir (DTG) 50 mg (PO) q24h × 4 weeks</p>	<p><b>Truvada 1</b> tablet (PO) q24h [tenofovir (TDF) 300 mg (PO) q24h + emtricitabine (FTC) 200 mg (PO) q24h] × 4 weeks</p> <p><b>plus</b></p> <p><i>(one of the following)</i></p> <p>Darunavir (DRV) 800 mg (PO) q24h × 4 weeks</p> <p><b>or</b></p> <p>Atazanavir (ATZ) 300 mg (PO) q24h × 4 weeks</p> <p><b>or</b></p> <p>Fosamprenavir (FPV) 1400 mg (PO) q24h × 4 weeks</p> <p><b>plus</b></p> <p>Ritonavir (RTV) 100 mg (PO) q24h × 4 weeks</p>	<p><b>Truvada 1</b> tablet (PO) q24h [tenofovir (TDF) 300 mg (PO) q24h + emtricitabine (FTC) 200 mg (PO) q24h] × 4 weeks</p> <p><b>plus</b></p> <p>Zidovudine (ZDV) 300 mg (PO) q12h × 4 weeks</p> <p><b>or</b></p> <p><b>Truvada 1</b> tablet (PO) q24h [tenofovir (TDF) 300 mg (PO) q24h + emtricitabine (FTC) 200 mg (PO) q24h] × 4 weeks</p> <p><b>plus</b></p> <p><b>Kaletra</b> 2 tablets (PO) q12h [lopinavir (LPV) 200 mg (PO) q12h + ritonavir (RTV) 50 mg (PO) q12h] × 4 weeks</p>	...

**Table 6.7. HIV: Prophylaxis of Opportunistic Infections**

Disorder	Usual organism	Indication for Prophylaxis	Intervention
HIV	PCP	CD <sub>4</sub> < 200/mm <sup>3</sup>	TMP-SMX: 1 SS tablet or 1 DS tablet (PO) q24h <b>or</b> Atovaquone 1500 mg (PO) q24h
	TB	PPD > 5 mm (current or past) or contact with active case	INH: 300 mg (PO) q24h × 9 months Rifabutin: Dose based on concomitant ART × 4 months
	Toxoplasmosis	IgG (+) and CD <sub>4</sub> < 100/mm <sup>3</sup>	TMP-SMX: 1 DS tablet (PO) q24h
	MAI	CD <sub>4</sub> < 50/mm <sup>3</sup>	Azithromycin: 1200 mg (PO) q week
	S.pneumoniae	CD <sub>4</sub> > 200/mm <sup>3</sup>	Pneumococcal vaccine
	Hepatitis B (HBV)	Susceptible patients	Hepatitis B vaccine
	Hepatitis A (HAV)	Susceptible patients	Hepatitis A vaccine
	Influenza	All patients	Annual influenza vaccine
	VZV	CD <sub>4</sub> > 200/mm <sup>3</sup> , VZV antibody negative	VZV vaccine

**Table 6.8. Transplant Prophylaxis (BMT/SOT)**

Exposure	Usual organism	Preferred prophylaxis	Alternate prophylaxis	Comments
Herpes simplex (HSV)	Herpes simplex	<u>HSV → &lt; 6 recurrences/year</u> Valacyclovir 500 mg (PO) q24h x 30 days <b>or</b> Famciclovir 1 g (PO) q12 x 30 days	Acyclovir 400 mg (PO) q8h x 30 days post-BMT	<u>Acute prophylaxis</u> Nearly all post-transplant HSV infections occur < 1 month. Transplants receiving CMV prophylaxis with ganciclovir, or valganciclovir are protected against HSV (see CMV prophylaxis).
		<u>HSV → &gt; 6 recurrences/year</u> Valacyclovir 1 gm (PO) q24h x 90 days <b>or</b> Famciclovir 250 mg (PO) q12 x 30 days	Acyclovir 400 mg (PO) q8h x 90 days post-BMT	<u>Chronic prophylaxis</u> Valacyclovir 500 mg (PO) q12h <b>or</b> Famciclovir 500 mg (PO) q12h <b>or</b> Acyclovir 400 mg (PO) q8h
Varicella zoster virus (VZV)	Varicella zoster virus	<u>VZV seropositive recipient</u> Valacyclovir 500 mg (PO) q12h post-transplant x 4–24 months	Acyclovir 400 mg (PO) q8h post-transplant x 4–24 months	Post-transplant VZV infections occur later than HSV. > 95% of adults are VZV seropositive. In transplants, most VZV infections are due to reactivation rather than primary infection. Patients may develop shingles after prophylaxis is discontinued.
		<u>VZV seronegative recipient (VZV seropositive graft)</u> Valacyclovir 1 g (PO) q8h post-transplant x 4–24 months	Acyclovir 800mg (PO) 5x/day post-transplant x 4–24 months	Transplant recipients on CMV prophylaxis with ganciclovir or valganciclovir do not require VZV prophylaxis.

**Table 6.8. Transplant Prophylaxis (BMT/SOT)** (cont'd)

Exposure	Usual organism	Preferred prophylaxis	Alternate prophylaxis	Comments
Cytomegalovirus (CMV)	Transplants (SOT) D+/R-, D+/R+, D-/R+	Valganciclovir 900 mg (PO) q24h × 3–6 months	Ganciclovir 5 mg/kg (IV) q24h × 3–6 months	Alternately, preemptive therapy may be used. When quantitative CMV PCR/pp 65 antigenemia levels become +/↑.  D-/R- patients should receive CMV negative blood and leukocyte depleted RBCs but do not require antiviral prophylaxis.
PCP	SOT/BMT (with GVHD)	TMP-SMX 1 DS tablet (PO) q week × 12 months	TMP-SMX 1 SS tablet (PO) q24h × 12 months <b>or</b> Atovaquone 1500 mg (PO) q24h × 12 months (with a meal)	See Drug Summaries for drug-drug interactions.
Toxoplasmosis	Heart transplants	TMP-SMX 1 DS tablet (PO) q24h	Atovaquone 1500 mg (PO) q24h (take with a meal)	See Drug Summaries for drug-drug interactions.
<i>Candida sp.</i>	Liver/pancreas transplants	Fluconazole 400 mg (IV/PO) q24h × 4 weeks	Posaconazole 200 mg (PO) q8h × 4 weeks	See Drug Summaries for drug-drug interactions.
<i>Aspergillus sp.</i>	Allogeneic BMT/lung transplants	Voriconazole 200 mg (PO) q12h	Posaconazole 200 mg (PO) q8h	Until engrafted (BMT); duration in lung transplants unclear. See Drug Summaries for drug-drug interactions.

GVHD = graft vs. host disease.

### ENDOCARDITIS PROPHYLAXIS

***Endocarditis prophylaxis is now recommended only for previous endocarditis, prosthetic cardiac valve or prosthetic cardiac valve material, congenital heart disease (CHD), unrepaired cyanotic CHD, including palliative shunts and conduits, completely repaired congenital heart defects with prosthetic material/devices (placed by surgery or by catheter intervention during the first 6 months post-procedure), repaired CHD with residual defects adjacent to the site of prosthetic patch/device, and cardiac transplantation with cardiac valvulopathy.***

**Table 6.9. Previous Indications for Infective Endocarditis (IE) Prophylaxis\***

Subset	Prophylaxis Recommended (Column A)	Prophylaxis Not Recommended (Column B)
Cardiac conditions	<ul style="list-style-type: none"> <li>• Ostium primum ASD</li> <li>• Prosthetic heart valves, including bioprosthetic and homograft valves</li> <li>• Previous infective endocarditis</li> <li>• Most congenital cardiac malformations</li> <li>• Rheumatic valve disease</li> <li>• Hypertrophic cardiomyopathy</li> <li>• MVP with significant valvular regurgitation</li> <li>• Calcific aortic stenosis</li> </ul>	<ul style="list-style-type: none"> <li>• Isolated ostium secundum ASD</li> <li>• Surgical repair without residue beyond 6 months of ostium secundum ASD or PDA</li> <li>• Previous coronary artery bypass surgery</li> <li>• MVP without valvular regurgitation</li> <li>• Physiologic, functional, or innocent murmurs</li> <li>• Previous Kawasaki's cardiac disease or rheumatic fever without valve disease</li> </ul>
Procedures	<ul style="list-style-type: none"> <li>• Dental procedures known to induce gingival/mucosal bleeding, including dental cleaning</li> <li>• Tonsillectomy or adenoidectomy</li> <li>• Surgical operations involving intestinal or respiratory mucosa</li> <li>• Cystoscopy or urethral dilation</li> <li>• Urethral catheterization or urinary tract surgery if UTI is present</li> <li>• Prostate surgery</li> <li>• I &amp; D of infected tissue</li> </ul>	<ul style="list-style-type: none"> <li>• Dental procedures not likely to induce gingival bleeding</li> <li>• Tympanostomy tube insertion</li> <li>• Flexible bronchoscopy ± biopsy</li> <li>• Endotracheal intubation</li> <li>• Endoscopy ± gastrointestinal biopsy</li> <li>• Cesarean section</li> <li>• D &amp; C, IUD insertion/removal, or therapeutic abortion in the absence of infection</li> <li>• Cardiac pacemaker/defibrillator insertion</li> </ul>

\* Wilson W, et al. Prevention of Infective Endocarditis. *Circulation* 116:1736–1754, 2007.



**Table 6.9. Previous Indications for Infective Endocarditis (IE) Prophylaxis\*** (cont'd)

Subset	Prophylaxis Recommended (Column A)	Prophylaxis Not Recommended (Column B)
	<ul style="list-style-type: none"> <li>• Biopsies of infected respiratory mucosa or infected skin/soft tissues</li> <li>• Any surgical procedure involving an infected field</li> </ul>	<ul style="list-style-type: none"> <li>• Coronary stent implantation</li> <li>• Percutaneous transluminal coronary angioplasty (PTCA)</li> <li>• Cardiac catheterization</li> </ul>

ASD = atrial septal defect, D & C = dilatation and curettage, I & D = incision and drain, IUD = intrauterine device, MVP = mitral valve prolapse, PDA = patent ductus arteriosus, UTI = urinary tract infection.

\* Prophylaxis is indicated for patients with cardiac conditions in Column A undergoing procedures in Column A. Prophylaxis is not recommended for patients or procedures in Column B. See Tables 6.9 and 6.10 for prophylaxis regimens for above-the-waist and below-the-waist procedures, respectively.

**Table 6.10. Endocarditis Prophylaxis for Above-the-Waist (Dental, Oral, Esophageal, Respiratory Tract) Procedures\***

Prophylaxis	Reaction to Penicillin	Antibiotic Regimen
Oral prophylaxis	None	Amoxicillin 2 gm (PO) 1 hour pre-procedure <sup>†</sup>
	Non-anaphylactoid	Cephalexin 1 gm (PO) 1 hour pre-procedure
	Anaphylactoid	Clindamycin 300 mg (PO) 1 hour pre-procedure <sup>††</sup>
IV prophylaxis	None	Ampicillin 2 gm (IV) 30 minutes pre-procedure
	Non-anaphylactoid	Cefazolin 1 gm (IV) 15 minutes pre-procedure
	Anaphylactoid	Clindamycin 600 mg (IV) 30 minutes pre-procedure

\* Endocarditis prophylaxis is directed against viridans streptococci, the usual SBE pathogen above the waist. Macrolide regimens are less effective than other regimens; clarithromycin/azithromycin regimens (500 mg PO 1 hour pre-procedure) are of unproven efficacy.

† Some recommend a 3 gm dose of amoxicillin, which is excessive given the sensitivity of viridans streptococci to amoxicillin.

†† Some recommend a 600 mg dose of clindamycin, but a 300 mg dose gives adequate blood levels and is better tolerated (less diarrhea).

**Table 6.11. Endocarditis Prophylaxis for Below-the-Waist (Genitourinary, Gastrointestinal) Procedures Involving an Infected Field\*\***

Prophylaxis	Reaction to Penicillin	Antibiotic Regimen
Oral prophylaxis	None	Amoxicillin 2 gm (PO) 1 hour pre-procedure
	Non-anaphylactoid, anaphylactoid	Linezolid 600 mg (PO) 1 hour pre-procedure

**Table 6.11. Endocarditis Prophylaxis for Below-the-Waist (Genitourinary, Gastrointestinal) Procedures Involving an Infected Field\*\* (cont'd)**

Prophylaxis	Reaction to Penicillin	Antibiotic Regimen
IV prophylaxis	None	Ampicillin 2 gm (IV) 30 minutes pre-procedure <b>plus</b> Gentamicin 80 mg (IM) or (IV) over 1 hour 60 minutes pre-procedure
	Non-anaphylactoid, anaphylactoid	Vancomycin 1 gm (IV) over 1 hour 60 minutes pre-procedure <b>plus</b> Gentamicin 80 mg (IM) or (IV) over 1 hour 60 minutes pre-procedure

\* Endocarditis prophylaxis is directed against *E. faecalis*, the usual SBE pathogen below the waist.

† Seto TB. The case for infectious endocarditis prophylaxis. *Arch Intern Med* 167:327–330, 2007.

Harrison JL, Hoen B, Prendergast BD. Antibiotic Prophylaxis for Infective Endocarditis. *Lancet* 371: 1317–1319, 2008.

**Table 6.12. Q Fever Endocarditis Prophylaxis**

Prophylaxis	Antibiotic Regimen
PO prophylaxis (for significant valvulopathy* in acute Q fever)	Doxycycline 100 mg (PO) of q12 h × 12 months <b>Plus</b> Hydroxychloroquine 200 mg (PO) q8h × 12 months

\* Significant valvulopathy: history of rheumatic fever, aortic bicuspid valve,  $\geq 2$  valve stenosis/regurgitation, MVP, remodeled/thickened valve.

## TRAVEL PROPHYLAXIS

**Travelers may acquire infectious diseases from ingestion of fecally-contaminated water/food, exchange of infected body secretions, inhalation of aerosolized droplets, direct inoculation via insect bites, or from close contact with infected birds/animals.**

Recommendations to prevent infection in travelers consist of general travel precautions (Table 6.12), and specific travel prophylaxis regimens (Table 6.13).

**Table 6.13. General Infectious Disease Travel Precautions**

Exposure	Risk	Precautions
Unsafe water (fecally-contaminated)	Diarrhea/dysentery, viral hepatitis (HAV)	<ul style="list-style-type: none"> <li>Avoid ingestion of unbottled/unpotable water. Be sure bottled water has an unbroken seal and has not been opened/refilled with tap water.</li> <li>Avoid ice cubes made from water of uncertain of origin/handling, and drinking from unclean glasses.</li> <li>Drink only pasteurized bottled drinks. Be sure bottles/cans are opened by you or in your presence.</li> </ul>

**Table 6.13. General Infectious Disease Travel Precautions (cont'd)**

Exposure	Risk	Precautions
		<ul style="list-style-type: none"> <li>Avoid drinking unpasteurized/warm milk; beer, wine, and pure alcoholic beverages are safe.</li> <li>Eat only canned fruit or fresh fruit peeled by you or in your presence with clean utensils.</li> <li>Avoid eating soft cheeses.</li> <li>Avoid eating raw tomatoes/uncooked vegetables that may have been exposed to contaminated water.</li> <li>Avoid using hotel water for tooth brushing/rinsing unless certain of purity. Many hotels use common lines for bath/sink water that is unsuitable for drinking.</li> <li>Avoid wading/swimming/bathing in lakes or rivers.</li> </ul>
Food-borne (fecally-contaminated)	Diarrhea/dysentery	<ul style="list-style-type: none"> <li>Eat only seafood/poultry/meats that are freshly cooked and served hot. Avoid eating at roadside stands or small local restaurants with questionable sanitary practices.</li> <li>"Boil it, peel it, or forget it."</li> </ul>
Body fluid secretions	Viral hepatitis (HBV, HCV, etc.), STDs, HIV/other retroviruses	<ul style="list-style-type: none"> <li>Do not share utensils/glasses/straws or engage in "risky behaviors" involving body secretion exchange.</li> <li>Avoid blood transfusion (use blood expanders instead).</li> <li>Treat dental problems before travel.</li> </ul>
Animal bite	Animal bite-associated infections	<ul style="list-style-type: none"> <li>Do not pet/play with stray dogs/cats. Rabies and other infections are common in wild (and some urban) animals.</li> </ul>
Flying insects	Malaria, arthropod-borne infections	<ul style="list-style-type: none"> <li>Avoid flying/biting insects by wearing dark protective clothing (long sleeves/pants) and using insect repellent on clothes/exposed skin, especially during evening hours.</li> <li>Minimize dawn-to-dusk outdoor exposure.</li> <li>Use screens/mosquito nets when possible.</li> <li>Do not use perfume, after shave, or scented deodorants/toiletries that will attract flying insects.</li> </ul>

**Table 6.14. Travel Prophylaxis Regimens**

Exposure	Usual Pathogens	Prophylaxis Regimens	Comments
	E. coli Salmonella Shigella Non-cholera vibrios V. cholerae Aeromonas Plesiomonas Rotavirus Norwalk virus Giardia lamblia Campylobacter	Doxycycline 100 mg (PO) q24h for duration of exposure <b>or</b> Any quinolone (PO) for duration of exposure <b>or</b> TMP-SMX 1 SS tablet (PO) q24h for duration of exposure	Observe without prophylaxis and treat mild diarrhea symptomatically with loperamide (2 mg). Persons with medical conditions adversely affected by dehydration caused by diarrhea may begin prophylaxis after arrival in country and continue for 1 day after returning home. Should severe diarrhea/dysentery occur, continue/switch to a quinolone, maximize oral hydration, and see a physician if possible.

**Table 6.14. Travel Prophylaxis Regimens** (cont'd)

Exposure	Usual Pathogens	Prophylaxis Regimens	Comments
Traveler's diarrhea	Yersinia Cryptosporidium Cyclospora Enteroviruses Amebiasis		Anti-spasmodics may be used for symptomatic relief of mild, watery diarrhea but are contraindicated in severe diarrhea/dysentery. Bismuth subsalicylate is less effective than antibiotic prophylaxis. Most cases are due to enterotoxigenic <i>E. coli</i> . TMP-SMX is active against some bacterial pathogens and <i>Cyclospora</i> , but not against <i>E. histolytica</i> or enteroviral pathogens (e.g., Rotavirus, Norwalk agent). Doxycycline is active against most bacterial pathogens and <i>E. histolytica</i> , but misses <i>Campylobacter</i> , <i>Cryptosporidium</i> , <i>Cyclospora</i> , <i>Giardia</i> , and enteroviral pathogens. Rifaxamin only active against <i>E. coli</i> . Antibiotics are inactive against viral/parasitic pathogens causing diarrhea.
Meningococcal meningitis	<i>N. meningitidis</i>	<u>Pre-travel prophylaxis</u> Meningococcal conjugate vaccine 0.5 mL (IM) $\geq$ 1 month prior to travel to endemic/epidemic areas  <u>Post-exposure prophylaxis</u> See p. 358	Acquired via close face-to-face contact (airborne aerosol/droplet exposure). Vaccine is highly protective against <i>N. meningitidis</i> serotypes A, C, Y, and W-135, but misses serotype B. For areas with serogroup B may use meningococcal group B vaccine. Alternately, consider chemoprophylaxis.
Hepatitis A (HAV)	Hepatitis A virus	HAV vaccine 1 mL (IM) prior to travel, then follow with a one-time booster 3, 6 months later	HAV vaccine is better than immune globulin for prophylaxis. Take care to avoid direct/indirect ingestion of fecally contaminated water. HAV vaccine is recommended for travel to all developing countries. Protective antibody titers develop after 2 weeks.
Typhoid fever	<i>S. typhi</i>	ViCPS vaccine 0.5 mL (IM). Booster every 2 years for repeat travelers	For the oral vaccine, do not co-administer with antibiotics. Contraindicated in compromised hosts and children < 6 years old.

**Table 6.14. Travel Prophylaxis Regimens** (cont'd)

Exposure	Usual Pathogens	Prophylaxis Regimens	Comments
		<p><b>or</b></p> <p>Oral Ty21a vaccine 1 capsule (PO) q48h × 4 doses. Booster every 5 years for repeat travelers</p>	<p>Take oral capsules with cold water. Degree of protective immunity is limited with vaccine. Some prefer chemoprophylaxis the same as for Traveler's diarrhea (p. 373).</p>
Yellow fever	Yellow fever virus	Yellow fever vaccine 0.5 mL (SC). Booster every 10 years for repeat travelers	<p>Vaccine is often required for travel to or from Tropical South America or Tropical Central Africa. Administer 1 month apart from other live vaccines. Contraindicated in children &lt; 4 months old; caution in children &lt; 1 year old. Reactions may occur in persons with egg allergies. Immunity is probably life long, but a booster every 10 years is needed for vaccination certification by some countries.</p>
Japanese encephalitis (JE)	Japanese encephalitis virus	JE vaccine 1 mL (SC) on days 0, 7, and 14 or 30. Booster schedule not established	<p>Recommended for travelers planning prolonged (&gt; 3 week) visits during the rainy season to rural, endemic areas of Asia (e.g., Eastern Russia, Indian subcontinent, China, Southeast Asia, Thailand, Korea, Laos, Cambodia, Vietnam, Malaysia, Philippines). Administer 3, 2 weeks before exposure. Children &lt; 3 years may be given 0.5 mL (SC) on same schedule as adults.</p>
Rabies	Rabies virus	HDCV, PCEC, or RVA 1 mL (IM) on days 0, 7, and 21 or 28 prior to travel	<p>Avoid contact with wild dogs/ animals during travel. Dose of rabies vaccine for adults and children are the same. A booster dose prior to travel is recommended if antibody levels are measured and are low.</p>

**Table 6.14. Travel Prophylaxis Regimens** (cont'd)

Exposure	Usual Pathogens	Prophylaxis Regimens	Comments
		<p style="text-align: center;"><b>or</b></p> HDCV 0.1 mL (ID) on days 0, 7, and 21 or 28 prior to travel	
Tetanus Diphtheria Pertussis	C. tetani C. diphtheriae B. pertussis	Tdap 0.5 ml (IM)	Tdap preferred to Td since Tdap also boosts pertussis immunity.

**Table 6.15. Malaria Prophylaxis**

Drug	Area of Exposure	Adult Dose	Comments
Atovaquone/ proguanil (Malarone)	Prophylaxis in all areas	Adult tablets contain 250 mg atovaquone and 100 mg proguanil hydrochloride. 1 adult tablet orally, daily	Begin 1–2 days before travel to malarious areas. Take daily at the same time each day while in the malarious area and for 7 days after leaving such areas. Contraindicated in persons with severe renal impairment (creatinine clearance < 30 mL/min). Atovaquone/proguanil should be taken with food or a milk drink. Not recommended prophylaxis for pregnant women, and women breastfeeding.
Chloroquine phosphate (Aralen and generic)	Prophylaxis only in areas with chloroquine-sensitive malaria	300 mg base (500 mg salt) orally, once/week	Begin 1–2 weeks before travel to malarious areas. Take weekly on the same day of the week while in the malarious area and for 4 weeks after leaving such areas.
Doxycycline	Prophylaxis in all areas	100 mg orally, (with food) daily	Begin 1–2 days before travel to malarious areas. Take daily at the same time each day while in the malarious area and for 4 weeks after leaving such areas. Contraindicated in children < 8 years and in pregnancy.

**Table 6.15. Malaria Prophylaxis** (cont'd)

Drug	Area of Exposure	Adult Dose	Comments
Hydroxy-chloroquine sulfate (Plaquenil)	An alternative to chloroquine for prophylaxis only in areas with chloroquine-sensitive malaria	310 mg base (400 mg salt) orally, once/week	Begin 1–2 weeks before travel to malarious areas. Take weekly on the same day of the week while in the malarious area and for 4 weeks after leaving such areas.
Mefloquine (Larium)	Prophylaxis in areas with mefloquine-sensitive malaria (mefloquine resistance in SE Asia)	228 mg base (250 mg salt) orally, once/week	Begin at least 2 weeks before travel to malarious areas. Take weekly on the same day of the week while in the malarious area and for 4 weeks after leaving such areas. Contraindicated in persons allergic to mefloquine or related compounds (e.g., quinine, quinidine) and in persons with active depression, a recent history of depression, generalized anxiety disorder, psychosis, schizophrenia, other major psychiatric disorders, or seizures. Use with caution in persons with psychiatric disturbances or a previous history of depression. Not recommended for persons with cardiac conduction abnormalities.
Primaquine <sup>†</sup>	Prophylaxis for short-duration travel to areas with principally <i>P. vivax</i>	30 mg base (52.6 mg salt) orally, daily	Begin 1–2 days before travel to malarious areas. Take daily at the same time each day while in the malarious area and for 7 days after leaving such areas.  Contraindicated in persons with G6PD deficiency, pregnancy and breastfeeding unless the infant being breastfed has a documented normal G6PD level.

**Table 6.15. Malaria Prophylaxis** (cont'd)

Drug	Area of Exposure	Adult Dose	Comments
Primaquine	Used for presumptive antirelapse therapy (terminal prophylaxis) to decrease the risk for relapses of <i>P. vivax</i> and <i>P. ovale</i>	30 mg base (52.6 mg salt) orally, once/day for 14 days after departure from the malarious area.	Indicated for persons who have had prolonged exposure to <i>P. vivax</i> and <i>P. ovale</i> or both. Contraindicated in persons with G6PD <sup>+</sup> deficiency. Also contraindicated during pregnancy and lactation unless the infant being breastfed has a documented normal G6PD level.

HDCV = human diploid cell vaccine, PCEC = purified chick embryo cell vaccine, RVA = rabies vaccine adsorbed, RIG = rabies immune globulin.

† Glucose-6-phosphate dehydrogenase. Those who take primaquine should have a normal G6PD level before starting the medication.

## TETANUS PROPHYLAXIS

**Current information suggests that immunity lasts for decades/life-time after tetanus immunization.** A tetanus booster should not be routinely given for minor wounds, but is recommended for wounds with high tetanus potential (e.g., massive crush wounds, soil-contaminated wounds, or deep puncture wounds).

**Table 6.16. Tetanus Prophylaxis in Routine Wound Management**

History of Adsorbed Tetanus Toxoid	Wound Type	Recommendations
Unknown or < 3 doses	Clean, minor wounds	Td <sup>†</sup> or Tdap <sup>‡</sup>
	Tetanus-prone wounds <sup>†</sup>	(Td <sup>†</sup> or Tdap <sup>‡</sup> ) plus TIG
≥ 3 doses	Clean, minor wounds	No prophylaxis needed
	Tetanus-prone wounds <sup>†</sup>	Td <sup>†</sup> if > 10 years since last dose*

DT = diphtheria and tetanus toxoids adsorbed (pediatrics), DTP = diphtheria and tetanus toxoids and pertussis vaccine adsorbed, Td = tetanus and diphtheria toxoids adsorbed (adult), TIG = tetanus immune globulin, Tdap = tetanus and diphtheria toxoids and pertussis vaccine adsorbed.

† For example, massive crush wounds; wounds contaminated with dirt, soil, feces, or saliva; deep puncture wounds; or significant burn wounds or frostbite.

‡ For children < 7 years, DTP (DT if pertussis is contraindicated) is preferred to tetanus toxoid alone. For children ≥ 7 years old and adults, Tdap is preferred to Td or tetanus toxoid alone.

\* More frequent booster doses are unnecessary and can increase side effects. Protection lasts > 20 yrs. Adapted from: Centers for Disease Control and Prevention. MMWR Rep 40 (RR-10):1–28. 1991.



## IMMUNIZATIONS

**Immunizations are designed to reduce infections in large populations, and may prevent/decrease the severity of infection in non-immunized individuals.** Compromised hosts with altered immune systems may not develop protective antibody titers to antigenic components of various vaccines. **Immunizations are not fully protective, but are recommended** (depending on the vaccine) for most normal hosts, since some protection is better than none.

**Table 6.17. Adult Immunizations**

Vaccine	Indications	Dosage	Comments
Bacille Calmette Guérin (BCG)	Possibly beneficial for adults at high-risk of multiple-drug resistant tuberculosis.	Primary: 1 dose (intradermal). Booster not recommended.	Live attenuated vaccine induces a positive PPD which may remain positive for years/life. Contraindicated in immunocompromised hosts. Injection site infection or disseminated infection are rare.
Hemophilus influenzae (type B)	For those at increased risk for invasive disease, e.g., functional or anatomic asplenia, HIV, immunoglobulin deficiency, complement deficiency (C <sub>1-3</sub> ), stem cell transplants, chemotherapy or radiation therapy.	Primary: 0.5 mL dose (IM). Booster not recommended.	Capsular polysaccharide conjugated to diphtheria toxoid. Benefit uncertain. Safety in pregnancy unknown. Mild local reaction in 10%.
Hepatitis A (HAV)	All children beginning age 12 to 23 months and adults at increased risk of HAV.	Primary: 1 mL dose (IM). One-time booster $\geq$ 6 months later. Booster not routinely recommended.	Inactivated whole virus. Pregnancy risk not fully evaluated. Mild soreness at injection site. Occasional headache/malaise.
Hepatitis B (HBV)	Household/sexual contact with carrier, IV drug use, multiple sex partners (heterosexual), homosexual male activity, blood product recipients, hemodialysis, occupational exposure to blood,	Primary (3 dose series): Recombivax 10 mcg (1 mL) or Engerix-B 20 mcg (1 mL) IM in deltoid at 0, 1, and 6 months. Alternate schedule for Engerix-B: 4 dose series at 0, 1, 2, and 12 months. Booster	Recombinant vaccine comprised of hepatitis B surface antigen. For compromised hosts (including dialysis patients), use specially packaged Recombivax 40-mcg doses (1 mL vial containing 40 mcg/mL). HBsAb titers should be obtained 6 months after 3-dose primary series. Those with non-protective titers ( $\leq$ 10 mIU/mL) should receive 1 dose monthly

**Table 6.17. Adult Immunizations** (cont'd)

Vaccine	Indications	Dosage	Comments
	residents/staff of institutions for developmentally disabled, prison inmates, residence $\geq 6$ months in areas of high endemicity, others at high risk.	not routinely recommended.	up to a maximum of 3 doses and retest for HbsAb titers. Pregnancy not a contraindication in high-risk females. Mild local reaction in 10–20%. Occasional fever, headache, fatigue, nausea. Twinrix 1 mL (IM) (combination of Hepatitis A inactivated vaccine and Hepatitis B recombinant vaccine) is available for adults on a 0, 1, and 6-month schedule or 0, 7 days, 21–30 days, and 12 month schedules.
Herpes zoster (VZV)	Adults $\geq 60$ years to reduce the frequency of shingles/ prevent post-herpetic neuralgia. Use in those with previous H. zoster is not yet defined. Protection best in 60–69 year group; efficacy decreases with increasing age.	Primary: 0.65 mL (SC). Need for revaccination not yet defined. Vaccine must be stored frozen and used within 30 minutes after thawing.	Duration of protection is at least 4 years. Injection site reactions in 48%. Contraindicated in immunocompromised hosts (with immunosuppressive disorder or receiving immunosuppressive drug) or untreated TB. Not indicated for therapy of H. zoster or post-herpetic neuralgia.
Influenza	All adults.	Annual vaccine. Single 0.5 mL dose (IM) before flu season is optimal, but can be given anytime during flu season. (2 A strains + 1 B strain)	High dose (HD) vaccine has 4 $\times$ the antigen content as the standard dose (SD) vaccine. Quadrivalent (2 A strains + 2 B strain) and trivalent inactivated whole and split virus vaccines available. Contraindications include anaphylaxis to eggs or sensitivity to thimerosal. Mild local reaction common. Malaise/myalgias in some. For pregnancy, administer in 2 <sup>nd</sup> or 3 <sup>rd</sup> trimester during flu season. High titer (HD) inactivated vaccine indicated for adults $\geq 65$ years, but enhanced protective efficacy (vs. SD) not yet demonstrated.

**Table 6.17. Adult Immunizations** (cont'd)

Vaccine	Indications	Dosage	Comments
Measles	Adults born after 1956 without live-virus immunization or measles diagnosed by a physician or immunologic test. Also indicated for revaccination of persons given killed measles vaccine between 1963–67.	Primary: 0.5 mL dose (SC). A second dose ( $\geq 1$ month later) is recommended for certain adults at increased risk of exposure (e.g., healthcare workers, travelers to developing countries). No routine booster.	Live virus vaccine (usually given in MMR). Contraindicated in compromised hosts, pregnancy, history of anaphylaxis to eggs or neomycin. Ineffective if given 3–11 months after blood products. Side effects include low-grade fever 5–21 days after vaccination, rash, and local reactions in if previously immunized with killed vaccine (1963–67).
Meningococcus (invasive disease)	Patients with splenic dysfunction or with complement defects ( $C_{7-9}$ ) laboratory workers. May be given to 1 <sup>st</sup> year college students living in dormitories.	Meningococcal conjugate vaccine 0.5 mL (IM). Primary: 0.5 mL (IM) then again at months 2 and 6 (3 dose series)	Also used in epidemic control of <i>N. meningitidis</i> serogroups A, C, Y, and W-135. For serogroup B use meningococcal group B vaccine.
Mumps	Non-immune adults.	Primary: 0.5 mL dose (SC). No routine booster.	Live attenuated vaccine (usually given in MMR). Contraindicated in immunocompromised hosts, pregnancy, history of anaphylaxis to eggs or neomycin.
Papilloma (human) Virus (HPV)	Females up to 26 years of age. Contraindicated in pregnancy. Suggested for males up to 26 years of age.	Primary: 0.5 mL (IM). Second dose: 2 months after 1 <sup>st</sup> dose. Third dose: 6 months after 1 <sup>st</sup> dose.	Quadrivalent HPV vaccine to prevent cervical, vulvar, vaginal and cancers caused by HPV types 6, 11, 16, 18. 9 valent HPV vaccine includes HPV types 52, 58, 31, 33, 45, 6, 11, 16, 18.
Pertussis	Use Tdap instead of Td in booster dose.	A Single Tdap booster dose 0.5 mL (IM).	Recommended since adults may get pertussis or transmit it to susceptible infants.

**Table 6.17. Adult Immunizations** (cont'd)

Vaccine	Indications	Dosage	Comments
Pneumo-coccus (S. Pneumoniae)	Immunocompetent hosts $\geq 65$ years old, or $> 19$ years old with diabetes, CSF leaks, or chronic cardiac, pulmonary or liver disease. Also for immunocompromised hosts $> 19$ years old with functional/anatomic asplenia,* leukemia, lymphoma, multiple myeloma, widespread malignancy, chronic renal failure, bone marrow/organ transplant, or on immunosuppressive/steroid therapy.	PPSV-23 PCV-13 Primary: 0.5 mL dose (IM). No booster. Primary: 0.5 mL dose (SC or IM). A one-time booster at 5 years is recommended for immuno-compromised hosts $> 2$ years old and for those who received the vaccine before age 65 for high-risk conditions.	<i>Pneumococcal vaccine naïve persons aged <math>\geq 65</math>. Give PCV-13, then give PPSV-23 6-12 months later (minimal interval 8 weeks).</i> <i>Persons who previously received PPSV-23 at age <math>\geq 65</math>. Give PCV-13 <math>&gt; 1</math> year later after PPSV-23 given.</i> <i>Persons who previously received PPSV-23 before age 65 years who are now age <math>\geq 65</math>. Give PCV-13 when <math>\geq 65</math> years give <math>&gt; 1</math> year later (minimum interval between sequential administration of PCV-13 and PPSV-23 is 8 weeks). If this window is missed, PPSV-23 can be given 6-12 months after PCV-13.</i>
Rubella	Non-immune adults, particularly women of childbearing age.	Primary: 0.5 mL dose (SC). No routine booster.	Live virus (RA 27/3 strain) vaccine (usually given in MMR). Contraindicated in immunocompromised hosts, pregnancy, history of anaphylactic reaction to neomycin. Joint pains and transient arthralgias in up to 40%, beginning 3–25 days after vaccination and lasting 1–11 days; arthritis in $< 2\%$ .
Tetanus-diphtheria	Adults	Primary: Td two 0.5 mL doses (IM), 1–2 months apart; third dose 6–12 months after second dose. Booster: a single Tdap 0.5 mL (IM) is recommended. Tdap should be substituted for one of the three Td doses.	Adsorbed toxoid vaccine. Contraindicated if hypersensitivity/neurological reaction or severe local reaction to previous doses. Side effects include local reactions, occasional fever, systemic symptoms, Arthus-like reaction in persons with multiple previous boosters, and systemic allergy (rare).

PCV-13 = 13 valent pneumococcal conjugate vaccine (contains strains: 1, 3, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F, 23F)  
PPSV-23 = 23 valent pneumococcal polysaccharide vaccine (contains strains: 1, 2, 3, 4, 5, 6B, 7F, 8, 9N, 9V, 10A, 11A, 12F, 14, 15B, 17F, 18C, 19F, 19A, 20, 22F, 23F, 33F).

\*Pneumococcal polysaccharide vaccine may be given before splenectomy, but is in effective post-splenectomy, the conjugate vaccine is more effective.

**Table 6.17. Adult Immunizations** (cont'd)

Vaccine	Indications	Dosage	Comments
Varicella (VZV) chickenpox	Non-immune adolescents and adults, especially healthcare workers and others likely to be exposed.	Primary: Two 0.5 mL doses (SC), 4–8 weeks apart. Vaccine must be stored frozen and used within 30 minutes after thawing and reconstitution. No routine booster.	Live attenuated vaccine. Contraindications include pregnancy, active untreated TB, immunocompromised host, malignancy of bone marrow or lymphatic system, anaphylactic reaction to gelatin/neomycin, or blood product recipient within previous 6 months (may prevent development of protective antibody). Mild febrile illness in 10%. Injection site symptoms in 25–30% (local rash in 3%). Mild diffuse rash in 5%.

---

## REFERENCES AND SUGGESTED READINGS

---

- Andrews WW, Kimberlin DF, Whitley R, Cliver S, Ramsey PS. Valacyclovir therapy to reduce recurrent genital herpes in pregnant women. *Am J Obstet Gynecol* 194:774–81, 2006.
- Antrum RM, Solomkin JS. A review of antibiotic prophylaxis for open fractures. *Orthop Rev* 16:246–54, 1987.
- Baum SG. Oseltamivir and the influenza alphabet. *Clin Infect Dis* 43:445–6, 2006.
- Brantley JS, Hicks L, Sra K, Tyring SK. Valacyclovir for the treatment of genital herpes. *Expert Rev Anti-Infect Ther* 4:367–76, 2006.
- Bryson DJ, Gulihar A, Aujla RS, Taylor GJ. The hip fracture best practice tariff: early surgery and the implications for MRSA screening and antibiotic prophylaxis. *Eur J Orthop Surg Trauma* 25:123–127, 2015.
- Camann W, Tuomala R. Antibiotic prophylaxis for cesarean delivery: always before skin incision! *Int J Obstet Anesth* 20:1–2, 2011.
- Caple J. Varicella-Zoster Virus vaccine: a review of its use in the prevention of Herpes Zoster in older adults. *Drugs of Today* 42:249–54, 2006.
- Carratala J. Role of antibiotic prophylaxis for the prevention of intravascular catheter-related infection. *Clin Microbiol Infect* 7 (Suppl 4):83–90, 2001.
- Centers for Disease Control and Prevention. Recommended adult immunization schedule for Adults 19 years or older—United States, 2012. *MMWR*, 63(05):110–112, 2014.
- Chen LH, Wilson ME, Schlagenhauf P. Prevention of Malaria in Long-term Travelers. *JAMA* 296:2234–44, 2006.
- Chattopadhyay B. Splenectomy pneumococcal vaccination and antibiotic prophylaxis. *Br J Hosp Med* 41:172–4, 1989.
- Clifford V, Daley A. Antibiotic prophylaxis in obstetric and gynaecological procedures: a review. *Aust N Z J Obstet Gynaecol*. 52:412–419, 2012.
- Colizza S, Rossi S. Antibiotic prophylaxis and treatment of surgical abdominal sepsis. *J Chemother* 13:193–201, 2001.
- Craig AS, Schaffner W. Prevention of hepatitis A with the hepatitis A vaccine. *N Engl J Med* 350:476–81, 2004.

- Cunha BA, Gossling HR, Nightingale CH, et al. Penetration of cefazolin and cefradine into bone in patients undergoing total knee arthroplasty. *Infection* 2: 80–84, 1984.
- Cunha BA. Antibiotic tissue penetration. *Bulletin of the New York Academy of Medicine* 59:443–449, 1983.
- Cunha BA, Ristuccia A, Jonas M, et al. Penetration of ceftizoxime and cefazolin into bile and gallbladder wall. *Journal of Antimicrobial Chemotherapy* 10: 117–120, 1982.
- Cunha BA, Pyrttek LJ, Quintiliani R. Prophylactic antibiotics in cholecystectomy. *Lancet* 1:207–8, 1979.
- Cunha BA, Gossling HR, Nightingale C, et al. Penetration characteristics of cefazolin, cephalothin, and cephadrine into bone in patients undergoing total hip replacement. *Journal of Bone and Surgery* 59: 856–859, 1977.
- Darouiche RG. Treatment of infections associated with surgical implants. *N Engl J Med* 350:1422–9, 2004.
- De Lall F. Antibiotic prophylaxis in orthopedic prosthetic surgery. *J Chemother* 13 Spec No 1:48–53, 2001.
- De la Camara R. Antifungal prophylaxis in haematology patients. 12:565–76, 2006.
- Diaz-Pedroche C, Lumbreras C, San Juan R, et al. Valganciclovir preemptive therapy for the prevention of cytomegalovirus disease in high-risk seropositive solid-organ transplant recipients. *Transplantation* 82: 30–5, 2006.
- Dietrich ES, Bieser U, Frank U, et al. Ceftriaxone versus other cephalosporins for perioperative antibiotic prophylaxis: a meta analysis of 43 randomized controlled trials. *Chemotherapy* 48:49–56, 2002.
- Drew RH, Perfect JR. Use of Polyenes for Prophylaxis for Invasive Fungal Infections. *Infections in Medicine* 23:512–24, 2006.
- Esposito S. Is single-dose antibiotic prophylaxis sufficient for any surgical procedure? *J Chemother* 11:556, 2000.
- Faix, RG. Immunization during pregnancy. *Clin Obstet Gynecol* 45:42–58, 2002.
- Fennessy BG, O'Sullivan MJ, Fulton GJ, et al. Prospective Study of Use of Perioperative Antimicrobial Therapy in General Surgery. *Surgical Infections* 7:355–60, 2006.
- Fritze D, Englesbe MJ, Campbell DA Jr. Oral antibiotics to prevent surgical site infections following colon surgery. *Adv Surg* 45:141–153, 2011.
- Fry DE. Colon preparation and surgical site infection. *Am J Surg* 202:225–232, 2011.
- Gardner P, Eickhoff T. Immunization in adults in the 1990s. *Curr Clin To Infect Dis* 15:271–300, 1995.
- Gardner P, Peter G. Recommended schedules for routine immunization of children and adults. *Infect Dis Clin North Am* 15:1–8, 2001.
- Gardner P, Schaffner W. Immunization of adults. *N Engl J Med* 328:1252–8, 1993.
- Grabe M. Perioperative antibiotic prophylaxis in urology. *Curr Opin Urol* 11:81–5, 2001.
- Guaschino S, De Santo D, De Seta F. New perspectives in antibiotic prophylaxis for obstetric and gynecological surgery. *J Hosp Infect* 50 (Suppl A):S513–6, 2002.
- Haines SJ. Antibiotic prophylaxis in neurosurgery. The controlled trials. *Neurosurg Clin N Am* 3:355–8, 1992.
- Hartman BJ. Selective aspects of infective endocarditis: considerations on diagnosis, risk factors, treatment and prophylaxis. *Adv Cardiol* 39:195–202, 2002.
- Hasin T, Davidovitch N, Cohen R, et al. Postexposure Treatment with Doxycycline for the Prevention of Tick-Borne Relapsing Fever. *N Engl J Med* 355:147–55, 2006.
- Hawn MT, Richman JS, Vick CC, et al. Timing of surgical antibiotic prophylaxis and the risk of surgical site infection. *JAMA* 147:649–657, 2013.
- Higuchi Y, Takesue Y, Yamada Y, et al. A single-dose regimen for antimicrobial prophylaxis to prevent perioperative infection in urological clean and clean-contaminated surgery. *J Infect Chemother* 17:219–223, 2011.
- Hill DR, Ericsson CD, Pearson RD, Keystone JS, et al. The Practice of Travel Medicine: Guidelines by the Infectious Diseases Society of America. *Clin Infect Dis* 43:1499–539, 2006.
- Hornberger H, Robertus K. Cost-Effectiveness of a Vaccine To Prevent Herpes Zoster and Postherpetic Neuralgia in Older Adults. 145:317–25, 2006.
- Howdieshell TF, Heffernan D, Dipiro JT. Surgical Infection Society Guidelines for Vaccination after Traumatic Injury. *Surgical Infections* 7:275–303, 2006.
- Jackson LA, Neuzil KM, Yu O et al. Effectiveness of pneumococcal polysaccharide vaccine in older adults. *N Engl J Med* 348:1747–55, 2003.
- Kambaroudis AG, Papadopoulos S, Christodoulidou M, et al. Perioperative use of antibiotics in intra-abdominal surgical infections. *Surg Infect (Larchmt)* 11:535–544, 2011.
- Khoury JA, Storch GA, Bohl DL, et al. Prophylactic versus preemptive oral valganciclovir for the management of cytomegalovirus infection in adult renal transplant recipients. *Am J Transplant* 6:2134–43, 2006.
- Klein A, Miller KB, Sprague K, et al. A randomized, double-blind, placebo controlled trial of valganciclovir prophylaxis to prevent zoster recurrence from months 4 to 24 after BMT. *Bone Marrow Transplant*, 26: 1–6, 2010.

- Kotton CN, Kumar D, Caliendo AM, et al. International consensus guidelines on the management cytomegalovirus in solid organ transplantation. *Transplantation* 89:779–795, 2010.
- Laftavi MR, Rostami R, Feng L, et al. The role of antibiotic prophylaxis in the new era of immunosuppression. *Transplant Proc* 43:533–535, 2011.
- Lewis RT. Oral versus systemic antibiotic prophylaxis in elective colon surgery: a randomized study and meta-analysis send a message from the 1990s. *Can J Surg* 45:173–80, 2002.
- Million M, et al. Evolution from Acute Q Fever to Endocarditis Is Associated with Underlying Valvulopathy and Age and Can Be Prevented by Prolonged Antibiotic Treatment. *Clin Infect Dis* 57:836–844, 2013.
- Musher DM, Rueda-Jaimes AM, Graviss EA, Rodriguez-Baradas MC. Effect of Pneumococcal Vaccination: A Comparison of Vaccination Rates in Patients with Bacteremic and Nonbacteremic Pneumococcal Pneumonia. *Clin Infect Dis* 43:1004–08, 2006.
- New York State Department of Health AIDS Institute. HIV Prophylaxis following occupational exposure. October 2012 <http://www.hivguidelines.org/clinical-guidelines/post-exposure-prophylaxis/hiv-prophylaxis-following-occupational-exposure/>, 2012.
- Norton TD, Skeete F, Dubrovskaya Y, et al. Orthopedic surgical site infections: analysis of causative bacteria and implications for antibiotic stewardship. *Am J Orthop* 43: e89–92, 2014.
- Paul M, Yahav D, Fraser A, et al. Empirical antibiotic monotherapy for febrile neutropenia: systematic review and meta-analysis of randomized controlled trials. *J Antimicrob Chemother* 57:176–89, 2006.
- Pham H, Geraci SA, Burton MJ. Adult immunizations: update on recommendations. *Am J Med* 124:698–201, 2011.
- Phillips P, Chan K, Hogg R, et al. Azithromycin prophylaxis for *Mycobacterium avium* complex during the era of highly active antiretroviral therapy: evaluation of a provincial program. *Clin Infect Dis* 34:371–8, 2002.
- Pratesi C, Russo D, Dorigo W, et al. Antibiotic prophylaxis in clean surgery: vascular surgery. *J Chemother* 13 (Spec 1):123–8, 2001.
- Rex JH, Sobel JD. Prophylactic antifungal therapy in the intensive care unit. *Clin Infect Dis* 32:191–200, 2001.
- Rupprecht CE, Gibbons RV. Prophylaxis against rabies. *N Engl J Med* 351:2626–35, 2004.
- Salkind AR, Rao KC. Antibiotic prophylaxis to prevent surgical site infections. *Am Fam Physician* 83: 585–590, 2011.
- Segreti J. Is antibiotic prophylaxis necessary for preventing prosthetic device infection. *Infect Dis Clin North Am* 13:871–7, 1999.
- Seto TB. The case for infectious endocarditis prophylaxis. *Arch Intern Med* 167:327–330, 2007.
- Seymour RA, Whitworth JM. Antibiotic prophylaxis for endocarditis, prosthetic joints, and surgery. *Dent Clin North Am* 46:635–51, 2002.
- Small TN, Cowan MJ. Immunization of hematopoietic stem cell transplant recipients against vaccinepreventable diseases. *Expert Rev Clin Immunol* 7:193–203, 2011.
- Snydman DR. Cytomegalovirus prevention strategies: The case for prophylaxis. *Am J Transplant* 9:1254, 2009.
- Snydman DR. Prevention of cytomegalovirus (CMV) infection and CMV disease in recipients of solid organ transplants: the case for prophylaxis. *Clin Infect Dis* 40:709–12, 2005.
- Sobel JD, Rex JH. Invasive candidiasis: Turning risk into a practical prevention policy? *Clin Infect Dis* 33: 187–190, 2001.
- Spira AM. Yellow Fever: An Update on Risks, Presentation, and Prevention. *Infections in Medicine* 23: 385–89, 2006.
- Syndman DR. Use of valganciclovir for prevention and treatment of cytomegalovirus disease. *Clin Infect Dis*, 46:28–29, 2008.
- Taub DD, Ershler WB, Janowski M, et al. Immunity from Smallpox Vaccine Persists for Decades: A Longitudinal Study. *Am J Med* 121:1058–1064, 2008.
- Vazquez M, LaRussa PS, Gershon AA, et al. The effectiveness of the varicella vaccine in clinical practice. *N Engl J Med* 344:955–60, 2001.
- Victor JC, Mourfi AS, Sundina TY, et al. Hepatitis A vaccine versus immune globulin for postexposure prophylaxis. *N Engl J Med* 357:1685–1694, 2007.
- Wingard JR. Antifungal chemoprophylaxis after blood and marrow transplantation. *Clin Infect Dis* 34: 1386–90, 2002.
- Winston DJ, Busuttill RW. Randomized controlled trial of oral itraconazole solution versus intravenous/oral fluconazole for prevention of fungal infections in liver transplant recipients. *Transplantation* 74:688–95, 2002.
- Wistrom J, Norrby R. Antibiotic prophylaxis of travellers' diarrhoea. *Scand J Infect Dis Suppl* 70:111–29, 1990.
- Zelenitsky SA, Ariano RE, Harding GK, et al. Antibiotic pharmacodynamics in surgical prophylaxis: an association between intraoperative antibiotic concentrations and efficacy. *Antimicrob Agents Chemother* 46:3026–30, 2002.

**GUIDELINES**

- CDC. Prevention of Perinatal Group B streptococcal disease. *MMWR*, 59:1–31, 2010.
- Hauser CJ, Adams CA, Eachempati SR. Prophylactic antibiotic use in open fractures: An evidence-based guideline. *Surgical Infections* 7:379–405, 2006.
- Kim DK, Bridges CB, Harriman KH. Advisory committee on immunization practices recommended immunization schedule for adults aged 19 years or older – United States 2015. *Morbidity and Mortality Weekly Report* 64:91, 2015.
- Medical Letter Antimicrobial prophylaxis for surgery. Guidelines. *Medical Letter* 7:47–50, 2009.
- Strikas RA. Advisory committee on immunization practices recommended immunization schedules for persons aged 0 through 18 years – United States, 2015. *Morbidity and Mortality Weekly Report* 64:93, 2015.
- Williams WW, Lu P, O'Halloran AO, Bridges CB, et al. Vaccination coverage among adults, excluding influenza vaccination – United States, 2013. *Morbidity and Mortality Weekly Report* 63:95–102, 2015.

**TEXTBOOKS**

- Antibiotic Prophylaxis for Surgical Site Infection Prevention in Adults (4<sup>th</sup> Ed), Institute for Clinical Systems Improvement, October, 2010.
- Brunette GW, Kozarsky PE, Magill AJ, et al. *CDC Health Information for International Travel 2010*, Mosby Elsevier, 2010.
- Brunette GW, Kozarsky P, Magill A. *The Yellow Book CDC Health Information for International Travel*. Oxford University Press, 2014.
- Brusch JL, *Endocarditis Essentials*. Jones & Bartlett, Sudbury, MA, 2011.
- Gorbach SL, Bartlett JG, Blacklow NR Infectious (eds). *Diseases* (3<sup>rd</sup> Ed), Lippincott, Williams & Wilkins, Philadelphia, 2004.
- Hawker J, Begg N, Blair, et al. *Communicable Disease Control Handbook* (2<sup>nd</sup> Ed), Malden, Massachusetts, Blackwell Publishing, 2005.
- Heymann DL. *Control of Communicable Disease Manual* (19<sup>th</sup> Ed), American Public Health Association, 2008.
- Keystone JS, Kozarski PE, Freedman DO, et al. *Travel Medicine*. Mosby, Edinburgh, 2004.
- Koff R. *Hepatitis Essentials*. Jones & Bartlett, Sudbury, MA, 2011.
- Mandell GL, Bennett JE, Blaser MJ (eds). *Mandell, Douglas, and Bennett's Principles and Practice of Infectious Diseases* (8<sup>th</sup> Ed), Philadelphia, Elsevier, 2015.
- Marrazzo JM, del Rio C, Holtgrave DR. HIV Prevention in Clinical Settings. Recommendations of the International Society USA Panel. *JAMA*. 312: 390-409, 2014.
- Plotkin SA, Orenstein WA *Vaccines* (5<sup>th</sup> Ed), Philadelphia, W.B. Saunders, 2004.
- Sax PE, Cohen CJ, Kuritzkes DR. *HIV Essentials* (7<sup>th</sup> Ed), Jones & Bartlett, Sudbury, MA, 2014.
- Yu VL, Merigan Jr, TC, Barriere SL. *Antimicrobial Therapy and Vaccines* (2<sup>nd</sup> Ed), Williams & Wilkins, Baltimore, 2005.



## Chapter 7

## Pediatric Infectious Diseases and Pediatric Drug Summaries

Leonard R. Krilov, MD

George H. McCracken, Jr, MD

Empiric Therapy of CNS Infections .....	387
Empiric Therapy of HEENT Infections .....	391
Empiric Therapy of Lower Respiratory Tract Infections .....	396
Empiric Therapy of Vascular Infections .....	402
Empiric Therapy of Gastrointestinal Infections .....	403
Empiric Therapy of Bone and Joint Infections .....	407
Empiric Therapy of Skin and Soft Tissue Infections .....	411
Common Pediatric Antimicrobial Drugs .....	414
References and Suggested Readings .....	423
Treatment of Malaria in Children in the US .....	726

This chapter pertains to infectious diseases and antimicrobial agents in the pediatric population. It is organized by clinical syndrome, patient subset, and in some cases, specific organism. Clinical summaries immediately follow each treatment grid. Therapeutic recommendations are based on antimicrobial effectiveness, reliability, cost, safety, and resistance potential. Antimicrobial agents and duration of therapy are listed in the treatment grids; corresponding drug dosages are provided on pages 414–422 and represent the usual dosages for normal renal and hepatic function. Drug dosages in infants/children are generally based on weight, up to adult dosage as maximum. **For any treatment category, i.e., preferred IV therapy, alternate IV therapy, PO therapy recommended drugs are equally effective and not ranked by priority.** Please refer to other pediatric drug references and the manufacturer's package inserts for dosage adjustments, side effects, drug interactions, and other important prescribing information.

"IV-to-PO Switch" in the last column of the shaded title bar in each treatment grid indicates the clinical syndrome can be treated either by IV therapy alone or IV followed by PO therapy, but

not by PO therapy alone. "PO Therapy or IV-to-PO Switch" indicates the clinical syndrome can be treated by IV therapy alone, PO therapy alone, or IV followed by PO therapy (unless otherwise indicated in the footnotes under each treatment grid). Most patients on IV therapy able to take oral medications should be switched to PO equivalent therapy after clinical improvement.

## Empiric Therapy of CNS Infections

### Acute Bacterial Meningitis

Subset (age)	Usual Pathogens	Preferred IV Therapy <sup>†</sup>	Alternate IV Therapy <sup>†</sup>
Neonate (0–30 days)	Group B streptococci E. coli Listeria monocytogenes Other gram-negatives (e.g., Citrobacter, Serratia) and gram-positives (e.g., Enterococci)	Ampicillin <b>plus</b> Gentamicin*	Ampicillin <b>plus</b> Cefotaxime*
1–3 months	Overlap neonate and > 3 months	Vancomycin <b>plus</b> <b>either</b> Cefotaxime <b>or</b> Ceftriaxone*	Ampicillin <b>plus</b> <b>either</b> Cefotaxime <b>or</b> Ceftriaxone*
> 3 months	S. pneumoniae N. meningitidis H. influenzae B (very rare) Non-typable H. influenzae (rare)	<u>Before culture results:</u> Vancomycin <b>plus</b> <b>either</b> Ceftriaxone <b>or</b> Cefotaxime*  <u>After culture results</u> Discontinue vancomycin if penicillin-susceptible S. pneumoniae, N. meningitidis, or H. influenzae is isolated	Meropenem*  <u>Severe penicillin allergy</u> Vancomycin <b>plus</b> <b>either</b> Rifampin <b>or</b> Chloramphenicol*

## Acute Bacterial Meningitis (cont'd)

Subset (age)	Usual Pathogens	Preferred IV Therapy†	Alternate IV Therapy‡
CNS shunt infection  (Treat initially for <i>S. epidermidis</i> ; if later identified as MSSA, treat accordingly)	<i>S. epidermidis</i> <i>S. aureus</i> (MRSA)	Vancomycin <b>with or without</b> Rifampin 10 mg/kg (IV or PO) q12h × 7–10 days after shunt removal	Linezolid × 7–10 days after shunt removal  <u>If MSSA isolated</u> Nafcillin <b>with or without</b> Rifampin 10 mg/kg (PO) q12h × 7–10 days after shunt removal
	<i>S. aureus</i> (MSSA)	Nafcillin × 7–10 days after shunt removal	Vancomycin × 7–10 days after shunt removal
	Enterobacteriaceae	Cefotaxime <b>or</b> Ceftriaxone × 7–10 days after shunt removal	Meropenem × 7–10 days after shunt removal

MRSA/MSSA = methicillin-resistant/sensitive *S. aureus*. Duration of therapy represents total time IV or IV + PO. Most patients on IV therapy able to take PO meds should be switched to PO therapy after clinical improvement.

† See pp. 414–422 for drug dosages.

\* Duration of therapy: Group B strep ≥ 14 days (with ventriculitis 4 weeks); *E. coli*, *Citrobacter*, *Serratia* ≥ 21 days; *Listeria* 10–14 days; *Enterococcus* ≥ 14 days; *S. pneumoniae* 10–14 days; *H. influenzae* ≥ 10 days; *N. meningitidis* 4–7 days.

## Acute Bacterial Meningitis (Normal Hosts)

**Clinical Presentation:** Fever, headache, stiff neck (or bulging anterior fontanelle in infants), irritability, vomiting, lethargy, evolving over hours to 1–2 days. The younger the child, the more nonspecific the presentation (e.g., irritability, lethargy, fever, poor feeding).

**Diagnostic Considerations:** Diagnosis by CSF chemistries, gram stain, culture. CSF findings are similar to those in adults: pleocytosis (100–5000 WBCs/mm<sup>3</sup>; predominately PMNs), ↑ protein, ↓↓ glucose. Normal CSF values differ by age:

	WBC/mm <sup>3</sup> (range); %PMNs	Protein, mg/dL (range)	Glucose, mg/dL (range)
Preterm	9 (0–25); 57% PMNs	115 (65–150)	50 (24–63)
Term newborn	8 (0–22); 61% PMNs	90 (20–170)	52 (34–119)
Child	0 (0–7); 0% PMNs	(5–40)	(40–80)

Modified from *The Harriet Lane Handbook*, 20<sup>th</sup> edition.

**Pitfalls:** The younger the child, the more nonspecific the presentation. With prior antibiotic use (e.g., oral therapy for otitis media), partially-treated meningitis (CSF lactic acid levels 4–6 nm/L) is a concern. Especially in the summer season, rapid viral diagnosis (especially enteroviral PCR) can exclude ABM and prevent unnecessary hospitalization and antibiotic use. HSV PCR can also be helpful in excluding ABM.

**Therapeutic Considerations:** Meningitic doses of antibiotics are required for the *entire* course of treatment. Repeat lumbar puncture (LP) is indicated at 24–48 hours if not responding clinically or in situations where a resistant organism is a concern (e.g., penicillin-resistant *S. pneumoniae*, or neonate with gram-negative bacillary meningitis). Dexamethasone (0.3 mg/kg IV q12h × 2 days) may reduce neurologic sequelae (e.g., hearing loss) in children ≥ 6 weeks of age when given before/with the first dose of antibiotics.

**Prognosis:** Varies with causative agent and age at presentation. Overall mortality < 5% in US (higher in developing countries). Morbidity 30–35% with hearing loss the most common neurologic sequella. Incidence of neurologic sequelae: *S. pneumoniae* > *H. influenzae* > *N. meningitidis*. Use of *H. influenzae* (type B) and pneumococcal conjugate vaccines in routine childhood immunizations have greatly reduced the incidence of meningitis caused by these pathogens. Neonates are at increased risk of mortality and major neurologic sequelae, and there is a significant incidence of brain abscesses in neonates with gram-negative meningitis (especially *Citrobacter* and *Serratia* infections).

### **Acute Bacterial Meningitis (CNS Shunt Infections)**

**Clinical Presentation:** Indolent (lethargy, irritability, vomiting) or acute (high fever, depressed mental status).

**Diagnostic Considerations:** Diagnosis by CSF gram stain/culture, often obtained by shunt tap.

**Pitfalls:** Blood cultures are usually negative for shunt pathogens and CSF WBCs may be low.

**Therapeutic Considerations:** High risk of nafcillin resistance with coagulase-negative staphylococcal infection. The addition of rifampin to vancomycin may improve clearance of bacteria. Removal of prosthetic material is usually necessary to achieve a cure.

**Prognosis:** Good with removal of prosthetic material.

## **Encephalitis**

Subset	Usual Pathogens	IV Therapy <sup>†</sup>	IV-to-PO Switch <sup>†</sup>
Herpes	HSV-1	Acyclovir × 14–21 days	Treat IV only
Arbovirus	WNV, SLE, Powassan encephalitis, EEE, WEE, VEE, JE, CE	No specific therapy	
Mycoplasma	<i>M. pneumoniae</i>	Doxycycline <b>or</b> Minocycline × 2–4 weeks	Doxycycline <b>or</b> minocycline × 2–4 weeks

## Encephalitis (cont'd)

Subset	Usual Pathogens	IV Therapy <sup>†</sup>	IV-to-PO Switch <sup>†</sup>
Respiratory viruses	Influenza, Enteroviruses, Measles	No specific therapy	

WNV = West Nile Virus, SLE = St. Louis encephalitis, EEE = Eastern equine encephalitis, WEE = Western equine encephalitis, VEE = Venezuelan equine encephalitis, JE = Japanese encephalitis, CE = California encephalitis.

<sup>†</sup> See pp. 414–422 for drug dosages.

### Herpes Encephalitis (HSV-1)

**Clinical Presentation:** Acute onset of fever and mental status/behavioral changes. High fever with intractable seizures and profoundly depressed sensorium may dominate the clinical picture.

**Diagnostic Considerations:** Diagnosis by CSF PCR for HSV-1. CSF findings include ↑ RBC/WBC, ↓ glucose, and ↑↑ protein. Classic EEG changes with temporal lobe spikes may be present early. Brain biopsy is rarely, if ever, indicated.

**Pitfalls:** MRI/CT scan can be normal initially. Delays in diagnosis and therapy adversely affect outcome.

**Therapeutic Considerations:** HSV is the only treatable viral encephalitis in normal hosts.

**Prognosis:** Antiviral therapy improves survival based on level of consciousness at presentation. Mortality is 10–20%, and major neurologic sequelae are frequent in survivors.

### Arboviral Encephalitis

**Clinical Presentation:** Acute onset of fever, headache, altered mental status. Usually seasonal (more common in summer/fall), based on vector/travel history. Can be very severe with high mortality. Symptomatic West Nile encephalitis is rare in children.

**Diagnostic Considerations:** Serological studies are the primary means of diagnosis. PCR "encephalitis panels" are being developed for clinical use.

**Therapeutic Considerations:** No specific antiviral therapy. Supportive therapy is crucial.

**Pitfalls:** Be sure to elicit potential exposures/travel history in children with fever/alter mental status.

**Prognosis:** Varies with agent. Severe residual deficits and high mortality rates can occur, especially with Eastern Equine encephalitis.

### Mycoplasma Encephalitis

**Clinical Presentation:** Acute onset of fever and mental status changes. Prodromal cough, sore throat, respiratory symptoms may occur.

**Diagnostic Considerations:** CSF shows mild pleocytosis with a predominance of mononuclear cells, normal or mildly ↓ glucose, and mildly ↑ protein. Mycoplasma IgG/IgM titers are elevated.

**Pitfalls:** CSF findings can be confused with viral encephalitis.

**Therapeutic Considerations:** Often self-limited illness even without antibiotic therapy. Doxycycline or minocycline penetrate CNS and can be used in children > 8 years of age. Macrolides may treat pulmonary infection but do not penetrate CNS.

**Prognosis:** Good. Neurologic sequelae are rare.

### Respiratory Virus Encephalitis

**Clinical Presentation:** Encephalitis may occur during or immediately after an acute respiratory illness. Influenza encephalitis, although rare, may be especially severe with a number of deaths reported every year in previously healthy children and adolescents.

**Diagnostic Considerations:** Rarely, adenoviruses have been recovered from CSF or brain in cases of encephalitis. Encephalitis has also been described very rarely in association with RSV and parainfluenza infections as well.

**Pitfalls:** Most cases of encephalitis remain undiagnosed, in part due to failure to consider and test for respiratory viruses.

**Therapeutic Considerations:** No evidence to date suggests a benefit of anti-influenza therapy in influenza encephalitis.

**Prognosis:** Generally self-limited with recovery, but death has been reported with influenza encephalitis. Adenovirus, measles, and mumps encephalitis in immunocompromised hosts may also be severe.

## Empiric Therapy of HEENT Infections

### Periorbital (Preseptal) Cellulitis/Orbital Cellulitis

Subset	Usual Pathogens	Preferred IV Therapy <sup>†</sup>	Alternate IV Therapy <sup>†</sup>	IV-to-PO Switch <sup>†</sup>
Periorbital cellulitis	S. pneumoniae H. influenzae M. catarrhalis S. aureus	<b>Combination therapy with Nafcillin* plus either Ceftriaxone or Cefotaxime</b> × 10–14 days	Ampicillin-sulbactam × 10–14 days	Amoxicillin/clavulanate <b>or</b> Cefuroxime <b>or</b> Cefpodoxime <b>or</b> Cefdinir × 10–14 days
Orbital cellulitis	S. pneumoniae H. influenzae M. catarrhalis S. aureus Anaerobes Group A streptococci	<b>Combination therapy with Nafcillin plus Ceftriaxone</b> × 10–14 days	Piperacillin/tazobactam <b>or</b> Ampicillin-sulbactam <b>or</b> Ticarcillin/clavulanate × 10–14 days	Amoxicillin/clavulanate <b>or</b> Cefuroxime <b>or</b> Cefpodoxime <b>or</b> Cefdinir × 10–14 days

Duration of therapy represents total time IV or IV + PO. Most patients on IV therapy able to take PO meds should be switched to PO therapy after clinical improvement.

\* Clindamycin or vancomycin if CA-MRSA likely.

† See pp. 414–422 for drug dosages.

**Clinical Presentation:** Periorbital and orbital cellulitis are bacterial infections. Fever, lid swelling, and erythema around the eye often in conjunction with acute sinusitis. In periorbital cellulitis the infection is

anterior to the orbital septum. Orbital cellulitis involves the orbit proper, extraocular muscles/nerves, and possibly the orbital nerve. Proptosis and limitation of ocular mobility define orbital cellulitis.

**Diagnostic Considerations:** CT scan is used to differentiate preseptal from periorbital cellulitis and identify the extent of orbital involvement when present.

**Pitfalls:** Failure to recognize orbital involvement leading to optic nerve damage or CNS extension/cavernous sinus thrombosis. CT scan cannot differentiate phlegmon from abscess.

**Therapeutic Considerations:** Orbital cellulitis is more emergent than periorbital cellulitis and should be treated with IV antibiotics initially. Surgical drainage may be indicated if a well defined abscess is present or in more severe disease.

**Prognosis:** Good with prompt antimicrobial therapy and ophthalmologic surgery if needed.

## Sinusitis

Subset	Usual Pathogens	IV Therapy <sup>†</sup> (Hospitalized)	PO Therapy or IV-to-PO Switch <sup>†</sup> (Ambulatory)
Acute	<i>S. pneumoniae</i> <i>H. influenzae</i> <i>M. catarrhalis</i>	Ceftriaxone <b>or</b> cefuroxime × 1–2 weeks	Amoxicillin <b>or</b> Amoxicillin/ clavulanic acid <b>or</b> 2 <sup>nd</sup> or 3 <sup>rd</sup> generation cephalosporin <b>or</b> Clarithromycin* × 10–14 days <b>or</b> Azithromycin* × 5 days
Chronic	Same as acute + oral anaerobes	Requires prolonged antimicrobial therapy (2–4 weeks)	

Duration of therapy represents total time IV, PO, or IV + PO. Most patients on IV therapy able to take PO meds should be switched to PO therapy soon after clinical improvement (usually < 72 hrs).

\* Macrolides may be less effective and should be reserved for penicillin- and cephalosporin-allergic patients.

† See pp. 414–422 for drug dosages.

**Clinical Presentation:** Nasal discharge and cough frequently with headache, facial pain, and low-grade fever lasting > 10–14 days. Can also present acutely with high fever (≥ 104°F) and purulent nasal discharge ± intense headache for ≥ 3 days.

**Diagnostic Considerations:** Acute sinusitis is a clinical diagnosis. Imaging studies are not routinely indicated. Overlap with acute viral infection and allergic symptoms may make diagnosis difficult.

**Pitfalls:** Transillumination, sinus tenderness to percussion, and color of nasal mucus are not reliable indicators of sinusitis.

**Therapeutic Considerations:** Microbiology/antibiotics are similar to acute otitis media, but duration of therapy is 10–14 days. Failure to respond to initial antibiotic therapy suggests a resistant pathogen or an alternative diagnosis. There are insufficient data to support long-course antibiotic treatment. Rarely, quinolones may be considered as 3<sup>rd</sup> line alternative.

**Prognosis:** Good. For frequent recurrences, consider radiologic studies and ENT/allergy consultation.

## Otitis Externa

Subset	Usual Pathogens	Topical Therapy
"Swimmer's ear"	<i>Pseudomonas</i> sp. Enterobacteriaceae <i>S. aureus</i>	Polymyxin B plus neomycin <b>plus</b> hydrocortisone (eardrops) q6h × 7–10 days  <b>or</b> Ciprofloxacin (otic solution) q12h × 7–10 days  <b>plus</b> Dexamethasone or hydrocortisone ear drops  <b>or</b> Ofloxacin (otic solution) q12h × 7–10 days

**Clinical Presentation:** Ear pain, itching, and sensation of fullness. Pain is exacerbated by tugging on the pinna or tragus of the outer ear. Purulent discharge may be visible in the external ear canal. Fever is generally absent.

**Diagnostic Considerations:** Otitis externa is a clinical diagnosis. A recent history of swimming or cleaning with cotton swabs is often elicited. Malignant otitis externa, as seen in elderly adults with diabetes mellitus, is extremely rare in children but may be seen in immunocompromised hosts.

**Pitfalls:** Failure to recognize a ruptured tympanic membrane may lead to a misdiagnosis of otitis externa based on purulence in the canal.

**Therapeutic Considerations:** Local cleansing (e.g., 2% acetic acid) and topical therapy with corticosteroid-polymyxin B-neomycin suspension is usually sufficient. Oral antibiotics should be considered for fever/cervical adenitis.

**Prognosis:** Excellent. Cleansing with 2% acetic acid drops after swimming prevents recurrences.

## Acute Otitis Media

Subset	Usual Pathogens	IV/IM Therapy <sup>†</sup>	PO Therapy <sup>†</sup>
Initial uncomplicated bacterial infection	<i>S. pneumoniae</i> <i>H. influenzae</i> <i>M. catarrhalis</i>	Ceftriaxone × 1 dose	Amoxicillin × 10 days <b>or</b> Azithromycin <sup>§</sup> (1-, 3-, or 5-day regimen) <b>or</b> Erythromycin-sulfisoxazole × 10 days** <b>or</b> TMP-SMX × 10 days**
Treatment failure or resistant organism*	MRSP Beta-lactamase positive <i>H. influenzae</i>	Ceftriaxone q24h × 3 doses	Amoxicillin/clavulanic acid <b>or</b> cephalosporin × 10 days**

DRSP = drug-resistant *S. pneumoniae*. Pediatric doses are provided; acute otitis media is uncommon in adults. For chronic otitis media, prolonged antimicrobial therapy is required.

† See pp. 414–422 for drug dosages.

\* Treatment failure = persistent symptoms and otoscopy abnormalities 48–72 hours after starting initial antimicrobial therapy. For risk factors for DRSP, see Therapeutic Considerations, below. If still fails after recommended therapy, consider clindamycin for resistant *S. pneumoniae* or tympanocentesis for gram stain and culture.

\*\* In children > 6 years of age with mild-moderate acute otitis media, a 5–7 day course of antimicrobial therapy may be adequate.

†† ES-600 = 600 mg amoxicillin/5 mL.

‡ 10-day course with either cefuroxime axetil 15 mg/kg (PO) q12h or cefdinir 7 mg/kg (PO) q12h or 14 mg/kg (PO) q24h or cefpodoxime 5 mg/kg (PO) q12h may be used.

§ Macrolides and TMP-SMX may be less effective and should be reserved for penicillin or cephalosporin allergic patients.



**Clinical Presentation:** Fever, otalgia, hearing loss. Nonspecific presentation is more common in younger children (irritability, fever). Key to diagnosis is examination of the tympanic membrane. Acute otitis media requires evidence of inflammation *and* effusion. Uncommon in adults.

**Diagnostic Considerations:** Diagnosis is made by finding an opaque, hyperemic, bulging tympanic membrane with loss of landmarks and decreased mobility on pneumatic otoscopy.

**Pitfalls:** Otitis media with effusion (i.e., tympanic membrane retracted or in normal position with decreased mobility or mobility with negative pressure; fluid present behind the drum but normal in color) usually resolves spontaneously and should not be treated with antibiotics.

**Therapeutic Considerations:** American Academy of Pediatrics guidelines suggest initial observation without antibiotics for non-severe otitis media or in children > 6 month of age. Risk factors for infection with drug-resistant *S. pneumoniae* (DRSP) include antibiotic therapy in past 30 days, failure to respond within 48–72 hours of therapy, day care attendance, and antimicrobial prophylaxis. Quinolones not approved for therapy.

**Prognosis:** Excellent, but tends to recur. Chronic otitis, cholesteatomas, mastoiditis are rare complications. Tympanostomy tubes/adenoidectomy for frequent recurrences of otitis media are the leading surgical procedures in children.

## Mastoiditis

Subset	Usual Pathogens	Preferred IV Therapy <sup>†</sup>	Alternate IV Therapy <sup>†</sup>	PO Therapy or IV-to-PO Switch <sup>†</sup>
Acute	<i>S. pneumoniae</i> <i>S. aureus</i> Group A streptococci <i>H. influenzae</i>	Nafcillin <b>or</b> Clindamycin <b>or</b> Vancomycin (if CA-MRSA suspected) <b>plus either</b> Ceftriaxone <b>or</b> Cefotaxime × 10–14 days	Ampicillin-sulbactam × 10–14 days	Amoxicillin/clavulanate <b>or</b> Cefpodoxime <b>or</b> Cefdinir <b>or</b> Cefuroxime axetil × 10–14 days
Chronic	Polymicrobial, including <i>P. aeruginosa</i> , <i>S. aureus</i> , anaerobes, Enterobacteriaceae	Piperacillin/Tazobactam <b>or</b> Ticarcillin/clavulanate × 10–14 days	Meropenem <b>or</b> Imipenem × 10–14 days	None

Duration of therapy represents total time IV, IV + PO, or PO. Most patients on IV therapy able to take PO meds should be switched to PO therapy after clinical improvement.

† See pp. 414–422 for drug dosages.

**Clinical Presentation:** Fever and otalgia with postauricular swelling/erythema pushing the ear superiorly and laterally. The presentation may be more subtle (e.g., less toxic, less swelling, Bell's palsy alone) in older children partially treated with antibiotics. Concomitant otitis media is rare.

**Diagnostic Considerations:** Acute mastoiditis is diagnosed clinically, but CT scan is definitive. Tympanocentesis through intact ear drum for aspirate and insertion of tympanostomy tube are helpful for microbiology and drainage, respectively. Chronic mastoiditis is often polymicrobial, including anaerobes and *P. aeruginosa*. Tuberculosis rarely presents as chronic mastoiditis.

**Pitfalls:** Do not overlook mastoiditis in older child with unresponsive otitis. Orbital involvement may lead to optic nerve damage or CNS extension/cavernous sinus thrombosis.

**Therapeutic Considerations:** Treatment is based on microbiology and requires at least 3 weeks of antibiotics.

**Prognosis:** Good with early treatment.

## Pharyngitis

Subset	Usual Pathogens	IV or IM Therapy <sup>†</sup>	PO Therapy or IV-to-PO Switch <sup>‡</sup>
Exudative (culture)	Group A streptococci	Benzathine penicillin IM x 1 dose	Penicillin V <b>or</b> Amoxicillin x 10 days. <i>Alternate:</i> Azithromycin 12 mg/kg/day x 5 days <b>or</b> Cephalexin <b>or</b> Cefadroxil <b>or</b> Erythromycin <b>or</b> Clarithromycin <b>or</b> Clindamycin x 10 days
Asymptomatic carrier	Group A streptococci	No treatment indicated	No treatment indicated
Persistent/recurrent disease	Group A streptococci	Clindamycin	Amoxicillin/clavulanate x 10 days <b>or combination therapy with either</b> Penicillin V <b>or</b> Amoxicillin x 10 days plus Rifampin added on days 7–10
Exudative, sexually active	N. gonorrhoeae	Ceftriaxone (IM) x 1 dose	
Lemierre's Syndrome (jugular vein septic thrombophlebitis) <sup>†</sup>	Fusobacterium necrophorum	Clindamycin (IV) <b>or</b> Penicillin G (IV) x 4–6 weeks	Clindamycin <b>or</b> Penicillin VK x 4–6 weeks
Vesicular, ulcerative	Enteroviruses HSV 1 or 2	Primary HSV: Acyclovir x 5–7 days	Primary HSV: Acyclovir <b>or</b> Valacyclovir x 5–7 days

Duration of therapy represents total time IV, IM, IV + PO, or PO. Most patients on IV therapy able to take PO meds should be switched to PO therapy after clinical improvement.

<sup>†</sup> See pp. 414–422 for drug dosages.

<sup>‡</sup> Treat only IV or IV-to-PO switch.

**Clinical Presentation:** Acute sore throat and fever with tender cervical lymphadenitis. Primary clinical consideration is differentiating Group A streptococci (GAS) from viral/other causes (e.g., adenovirus, enterovirus, respiratory viruses, other strep groups [C, G], Arcanobacterium hemolyticum, M. pneumoniae, C. pneumoniae, EBV). GAS is less likely with concomitant coryza, conjunctivitis, hoarseness, cough, acute stomatitis, discrete oral ulcerations, or diarrhea—children with these manifestations should not be cultured routinely.

**Diagnostic Considerations:** Laboratory testing for GAS is recommended, since clinical differentiation of viral pharyngitis from GAS is not possible. Rapid tests for GAS antigens are reliable with excellent specificities, but due to variable sensitivities of the assays, a negative rapid test should be confirmed by a throat culture. The accuracy of antigen and culture tests is dependent on obtaining a good throat swab containing pharyngeal/tonsillar secretions.

**Pitfalls:** Be sure to obtain a good throat swab. Post-treatment testing is generally not recommended unless the patient is at high risk for rheumatic fever (e.g., family history, ongoing outbreak) or is still symptomatic after 10 days of therapy. Asymptomatic GAS carriers do not require antibiotic therapy, but identifying a “true carrier” may be difficult. Eradication of GAS carriage should be considered in the following situations: an outbreak of acute rheumatic fever or post-streptococcal glomerulonephritis; an outbreak of GAS in a closed community; a family history of rheumatic fever; multiple episodes of GAS infection within the family for many weeks despite therapy; family anxiety about the presence of GAS; or tonsillectomy is being considered based on persistent carriage. Eradication is achieved using the same antimicrobial regimen as for “persistent/recurrent disease” (see treatment grid p. 395).

**Therapeutic Considerations:** Penicillin V (or amoxicillin) is the drug of choice for GAS pharyngitis. Erythromycin is still considered the drug of choice for penicillin-allergic individuals, although macrolide-resistant GAS strains are being reported. First-generation cephalosporins are also useful alternatives. Broader spectrum agents, although likely effective, should not be used routinely. Macrolides or sulfonamides may not eradicate GAS pharyngitis. When eradication of carriage is indicated amoxicillin, amoxicillin/clavulanate or clindamycin alone or penicillin plus rifampin may be useful.

**Prognosis:** Excellent. Rheumatic fever is rare in the US.

## Empiric Therapy of Lower Respiratory Tract Infections

### Community-Acquired Pneumonia

Subset (age)	Usual Pathogens	IV Therapy <sup>†</sup>	PO Therapy or IV-to-PO Switch <sup>†</sup>
Community-acquired pneumonia <i>Birth to 20 days</i>	Group B streptococci, Gram-negative enteric bacteria	Ampicillin <b>Plus either</b> Gentamicin <b>or</b> Cefotaxime × 10–21 days	Not applicable

Duration of therapy represents total time IV or IV, PO, or IV + PO. Most patients on IV therapy able to take PO meds should be switched to PO therapy after clinical improvement.

† See pp. 414–422 for drug dosages.

## Community-Acquired Pneumonia (cont'd)

Subset (age)	Usual Pathogens	IV Therapy <sup>†</sup>	PO Therapy or IV-to-PO Switch <sup>‡</sup>
3 weeks to 3 months	RSV Parainfluenza Human metapneumovirus (hmpv)	None (supportive care)	None
	C. trachomatis S. pneumoniae B. pertussis S. aureus	Cefotaxime <b>or</b> Ceftriaxone × 10–14 days*. Alternative: Ampicillin <b>or</b> Clindamycin*	<u>Afebrile</u> : Erythromycin × 14 days <b>or</b> Azithromycin × 5–7 days. <u>Lobar, febrile</u> : Amoxicillin <b>or</b> amoxicillin/clavulanic acid <b>or</b> Cefdinir <b>or</b> Cefuroxime <b>or</b> Cefpodoxime × 10–14 days
> 3 months to < 5 years	Viruses (RSV, parainfluenza, influenza, adenovirus, rhinoviruses)	<u>RSV</u> : consider ribavirin. <sup>§</sup> For infants at highest risk for severe RSV infection, consider palivizumab 15 mg/kg/month × 1–2 seasons for prevention. <u>Influenza</u> : Oseltamivir (influenza A, B). <sup>¶</sup> Routine immunization for all children > 6 months	
	S. pneumoniae H. influenzae M. pneumoniae	Ampicillin** <b>or</b> Ceftriaxone <b>or</b> Cefotaxime × 10–14 days*	Amoxicillin <b>or</b> Amoxicillin/clavulanate <b>or</b> Clarithromycin <b>or</b> Azithromycin × 10–14 days
5–15 years	M. pneumoniae C. pneumoniae S. pneumoniae	Ceftriaxone <b>plus</b> a macrolide × 10–14 days*	Amoxicillin <b>plus</b> a macrolide <b>or</b> Doxycycline (age > 8 years) × 10–14 days
Pertussis	Bordetella spp. <sup>‡</sup> certain Adenoviruses M. pneumoniae C. trachomatis C. pneumoniae	Azithromycin × 5 days <b>or</b> Erythromycin × 14 days	Erythromycin estolate × 14 days <b>or</b> Clarithromycin × 7 days <b>or</b> Azithromycin × 5 days. If macrolide-intolerant: TMP-SMX × 14 days
Tuberculosis	M. tuberculosis	See pp. 400–401	See pp. 400–401

Duration of therapy represents total time IV or IV, PO, or IV + PO. Most patients on IV therapy able to take PO meds should be switched to PO therapy after clinical improvement.

<sup>†</sup> See pp. 414–422 for drug dosages.

\* If chronic cough of more insidious onset, consider adding IV or PO macrolide (azithromycin, clarithromycin, erythromycin) to cover Pertussis/C. trachomatis (3 weeks–3 months of age), Mycoplasma (> 5 years of age), or Mycoplasma/C. pneumoniae (5–15 years of age).

\*\* If fully immunized against H. influenzae and S. pneumoniae.

<sup>‡</sup> B. pertussis, B. parapertussis, B. bronchiseptica.

<sup>§</sup> Use should be limited to the most severely ill with RSV, i.e., transplant patients.

<sup>¶</sup> Influenza A strains now circulating are amantadine resistant.

### **Community-Acquired Pneumonia**

**Clinical Presentation:** Fever  $\pm$  dyspnea, cough, tachypnea with infiltrates on chest x-ray.

**Diagnostic Considerations:** Usual pathogens differ by age. In neonates, pneumonia is typically diffuse and part of early-onset sepsis. In young infants, there is significant overlap between signs and symptoms of bronchiolitis (primarily due to RSVU) and pneumonia. Severe pneumonia is usually due to bacterial infection, although the organism is frequently not isolated (e.g., blood cultures are positive in only 10–20% of children < 2 years of age with bacterial pneumonia). In young infants, *Chlamydia trachomatis* can be detected by or culture of nasopharyngeal (NP) secretions. *Mycoplasma pneumoniae*, suggested by high titer cold agglutinins > 1:64. Diagnosis by nucleic acid amplification testing or  $\uparrow$  IgM titers. Although cases in children < 5 years have been reported. Respiratory viruses (RSV, influenza, adenoviruses, parainfluenza viruses, hMPV) can also be detected by PCR of throat/nasal secretions. If child lives in area with high prevalence of tuberculosis, consider tuberculosis in the differential diagnosis of primary pneumonia.

**Pitfalls:** Reliance on upper airway specimen for gram stain/culture leads to misdiagnosis and mistreatment, as true deep sputum specimen is rarely obtainable in children.

**Therapeutic Considerations:** Therapy is primarily empiric based on child's age, clinical/epidemiologic features and chest x-ray findings. *Mycoplasma* requires 2–3 weeks of treatment; *C. pneumoniae* may require up to 4 weeks of treatment. Routine use of pneumococcal vaccines has decreased the incidence of pneumococcal pneumonia.

**Prognosis:** Varies with pathogen, clinical condition at presentation, and underlying health status. Prognosis is worse in children with chronic lung disease, congenital heart disease, immunodeficiency, neuromuscular disease, hemoglobinopathy.

### **Lower Respiratory Tract Infections due to Respiratory Viruses**

**Clinical Presentation:** Viruses cause the majority of lower respiratory tract infections (LRTIs) in children. Respiratory syncytial virus (RSV) is the leading cause of LRTI in young infants, manifesting as bronchiolitis/viral pneumonia and causing annual mid-winter epidemics. The risk of secondary bacterial infection (other than possibly otitis media) is very low. Fever is typically low grade and usually improves over 3–5 days, even if hospitalized. Influenza viruses are another major cause of winter epidemic LRTIs in children of all ages. Hospitalization rates in infants under one year of age rival those in the > 65 year old population. Characteristic findings include high fever and diffuse inflammation of the airways. Young infants may have prominent GI symptoms as well. Secondary bacterial infection (otitis media, pneumonia, sepsis) are frequent complications of influenza. Primary influenza pneumonia, encephalopathy, and myocarditis are rare, severe complications. Other respiratory viruses associated with LRTIs in children include parainfluenza type 3 (viral pneumonia), parainfluenza types 1 and 2 (croup), adenoviruses, and human metapneumovirus (hMPV). Influenza in children resembles that in adults, but in children, lymphocytosis is usual. Atypical lymphocytes are common in children, but not in adults. Thrombocytopenia is common in both children and adults.

**Diagnostic Considerations:** Viral syndromes are often diagnosed based on clinical assessment alone. For confirmation or in more severe cases, rapid diagnosis by PCR.

**Pitfalls:** Routine use of corticosteroids or bronchodilators in RSV bronchiolitis are not supported by clinical evidence. Overuse of the diagnosis of "flu" (e.g., stomach flu, summer flu) has led to diluted appreciation for true influenza and its severity. Influenza vaccine (RIDT) has been traditionally underutilized in children. As in adults, a negative rapid influenza diagnostic test does not rule out influenza. ILI with negative RIDTs should be retested by PCR for respiratory viruses.

**Therapeutic Considerations:** For most viral LRTIs treatment is primarily supportive (e.g., adequate hydration, fever control, supplemental oxygen for severe illness). Ribavirin is approved for RSV infection but is very rarely used due to uncertain clinical benefit, high cost, and cumbersome method of administration (prolonged aerosol). For infants at greatest risk of severe RSV disease (e.g., premature infants, infants with underlying chronic lung disease or congenital heart disease), monthly prophylaxis with palivizumab (synagis) 15 mg/kg/month (IM) decreases RSV hospitalization rates. Per American Academy of Pediatrics guidelines, palivizumab is indicated for infants with chronic lung disease and  $\leq 32$  weeks gestational age (GA) or congenital heart disease who are 12 months of age at start of RSV season. For premature infants, palivizumab is considered for premature infants  $< 29$  weeks GA for a maximum of 5 doses through the winter season are recommended.

Annual influenza vaccination is indicated for all individuals  $> 6$  months of age. All circulating influenza A strains are resistant to amantadine. The neuraminidase inhibitors oseltamivir and zanamivir have pediatric indications for treatment and prophylaxis of influenza (Table 7.1).

**Table 7.1. Influenza Antiviral Medication Dosing Recommendations**

Antiviral		Treatment	Chemoprophylaxis
<b>Oseltamivir*</b>			
<b>Adults</b>		75-mg capsule twice per day 3 5 days	75-mg capsule once per day 3 10 days
<b>Children</b> (age, 12 months or older), weight:	15 kg or less	60 mg per day divided into 2 doses	30 mg once per day
	15–23 kg	90 mg per day divided into 2 doses	45 mg once per day
	24–40 kg	120 mg per day divided into 2 doses	60 mg once per day
	$> 40$ kg	150 mg per day divided into 2 doses	75 mg once per day
<b>Zanamivir</b>			
<b>Adults</b>		Two 5-mg inhalations (10 mg total) twice per day	Two 5-mg inhalations (10 mg total) once per day
<b>Children</b>		Two 5-mg inhalations (10 mg total) twice per day (age, 7 years or older)	Two 5-mg inhalations (10 mg total) once per day (age, 5 years or older)

\* CDC recommends oseltamivir 3 mg/kg (PO) q12h  $\times$  5 days for full term infants  $< 12$  months of age. Lower doses recommended for pre-term infants.

**Prognosis:** Most children with viral LRTIs do well and recover without sequelae. Infants hospitalized with RSV infection have higher rates of wheezing episodes over the next 10 years. The highest rates of hospitalization from influenza occur in children  $< 2$  years of age and in the elderly.

### **Pertussis**

**Clinical Presentation:** Upper respiratory tract symptoms (congestion, rhinorrhea) over 1–2 weeks (catarrhal stage) progressing to paroxysms of cough (paroxysmal stage) lasting 2–4 weeks, often with a characteristic inspiratory whoop, followed by a convalescent stage lasting 1–2 weeks during which

cough paroxysms decrease in frequency and severity. Fever is low grade or absent. In children < 6 months, whoop is frequently absent and apnea may occur. Duration of classic pertussis is 6–10 weeks. Older children and adults may present with persistent cough (without whoop) lasting 2–6 weeks. Complications include seizures, secondary bacterial pneumonia, encephalopathy, death; risk of complications is greatest in children < 1 year.

**Diagnostic Considerations:** Diagnosis is usually based on nature of cough and duration of symptoms. Laboratory diagnosis may be difficult. A positive culture for *Bordetella pertussis* from a nasopharyngeal swab inoculated on fresh selective media is diagnostic, but the organism is difficult to recover after 3–4 weeks of illness. PCR of nasal secretions is the best rapid diagnostic test for pertussis.

**Pitfalls:** Be sure to consider pertussis in older children and adults with prolonged coughing illness. Family contacts of index case should receive post-exposure antimicrobial prophylaxis. Virtually all children should be vaccinated against pertussis. A single booster dose of acellular pertussis is recommended at 11–12 years of age (additional guidelines for catch-up for older adolescents and adults up to 64 years of age). Pregnant women should be vaccinated during the 2<sup>nd</sup>–3<sup>rd</sup> trimester of each pregnancy to protect themselves and provide passive immunity to their infants. Relative precautions to further pertussis immunization include: seizure within 3 days of a dose; persistent, severe, inconsolable crying for  $\geq 3$  hours within 48 hours of a dose; collapse or shock-like state within 48 hours of a dose; or temperature of  $\geq 40.5^\circ\text{C}$  without other cause within 48 hours of a dose.

**Therapeutic Considerations:** Infants < 6 months frequently require hospitalization. By the paroxysmal stage, antibiotics have minimal effect on the course of the illness but are still indicated to decrease transmission. An association has been made between oral erythromycin and infantile hypertrophic pyloric stenosis in infants < 6 weeks of age; consider an alternative macrolide (azithromycin or clarithromycin) in these cases.

**Prognosis:** Good. Despite the prolonged course, long-term pulmonary sequelae have not been described after pertussis. Children < 1 year are at greatest risk of morbidity and mortality, although mortality rates remain very low.

## Tuberculosis

Subset	Pathogen	PO or IM Therapy (see footnote for drug dosages)
Latent infection (positive PPD, clinically well, negative chest x-ray)	<i>M. tuberculosis</i>	INH $\times$ 9 months <b>or</b> Rifampin $\times$ 6 months (if INH-resistant) <b>or</b> INH unavailable and child at risk
Pulmonary and extrapulmonary TB (except meningitis)	<i>M. tuberculosis</i>	INH <b>plus</b> Rifampin <b>plus</b> PZA $\times$ 2 months followed by INH <b>plus</b> Rifampin $\times$ 4 months <sup>†</sup>

Duration of therapy represents total time IV or IV, PO, or IV + PO. Most patients on IV therapy able to take PO meds should be switched to PO therapy after clinical improvement.

<sup>†</sup> If drug resistance is a concern, EMB or streptomycin (IM) is added (4-drug regimen in areas where MDR TB is prevalent) to the initial regimen until drug susceptibilities are determined.

**Tuberculosis** (cont'd)

Subset	Pathogen	PO or IM Therapy (see footnote for drug dosages)
Meningitis	M. tuberculosis	INH <b>plus</b> Rifampin <b>plus</b> PZA <b>plus</b> either Streptomycin (IM) <b>or</b> Ethionamide × 2 months, followed by INH <b>plus</b> Rifampin × 7–10 months (i.e., 9–12 months total therapy) <sup>†</sup>
Congenital	M. tuberculosis	INH <b>plus</b> rifampin <b>plus</b> PZA <b>plus</b> streptomycin (IM)

Duration of therapy represents total time IV or IV, PO, or IV + PO. Most patients on IV therapy able to take PO meds should be switched to PO therapy after clinical improvement.

‡ Plus steroids.

TB Drug	Daily dosage (mg/kg)	Twice weekly dosage (mg/kg)
Isoniazid (INH)	10–15	20–30
Rifampin (RIF)	10–20	10–20
Ethambutol (EMB)	15–25	50
Pyrazinamide (PZA)	20–40	50
Streptomycin	20–40 (IM)	–
Ethionamide	15–20 (in 2–3 divided doses/day)	–

Alternative drugs (capreomycin, ciprofloxacin, levofloxacin, cycloserine, kanamycin, para-aminosalicylic acid) are used less commonly and should be administered in consultation with an expert in the treatment of tuberculosis.

**Tuberculosis**

**Clinical Presentation:** Most children diagnosed with tuberculosis have asymptomatic infection detected by tuberculin skin testing. Symptomatic disease typically presents 1–6 months after infection with fever, growth delay, weight loss, night sweats, and cough. Extrapulmonary involvement may present with meningitis, lymphadenopathy, bone, joint, or skin involvement.

**Diagnostic Considerations:** A positive tuberculin skin test indicates likely infection with M. tuberculosis. Tuberculin reactivity develops 2–12 weeks after infection. Children < 8 years old cannot produce sputum for AFB smear and culture. Interferon gamma release assays (IGRAs) may be used to diagnose TB in children > 5 years. Testing recommended only for children or increased risk, e.g., HIV, immigrants from TB endemic areas clinical and those with clinical findings suggestive of TB.

**Pitfalls:** Tuberculosis may be missed if not considering the diagnosis in children at increased epidemiological risk for exposure. Tuberculous meningitis often presents insidiously with nonspecific irritability and lethargy weeks prior to the development of frank neurological defects.



**Therapeutic Considerations:** Choice of initial therapy depends on stage of disease and likelihood of resistant organisms (based on index case, geographical region of acquisition). For HIV-infected patients, duration of therapy is prolonged to  $\geq 12$  months. For tuberculosis meningitis and miliary disease, the addition of corticosteroids to anti-TB therapy is beneficial.

**Prognosis:** Varies with extent of disease, drug resistance and underlying immune status, but is generally good for pulmonary disease in children. Bone infection may result in orthopedic sequelae (e.g., Pott's disease of the spine with vertebral collapse). The prognosis for meningitis is guarded once focal neurological deficits and persistent depression of mental status occur.

## Empiric Therapy of Vascular Infections

### Central Venous Catheter (CVC) Infections (Broviac, Hickman, Mediport)

Subset	Usual Pathogens	Preferred IV Therapy <sup>†</sup>	Alternate IV Therapy <sup>†</sup>
Empiric; immunocompromised host	<i>S. aureus</i> <sup>‡</sup> Enterobacteriaceae <i>Pseudomonas</i> Viridans Streptococci Enterococcus	Meropenem <b>or</b> Piperacillin/tazobactam <b>or</b> cefepime $\times 7-14$ days	Piperacillin/tazobactam <b>plus</b> an aminoglycoside $\times 7-14$ days
Isolate-based	<i>S. aureus</i> <i>MSSA</i>	Nafcillin <b>or</b> Oxacillin for $\geq 2$ weeks	Cefazolin <b>or</b> Vancomycin (preferred if severe beta-lactam allergy) for $\geq 2$ weeks
	<i>MRSA</i>	Vancomycin <b>with or without</b> Gentamicin for $\geq 2$ weeks	Linezolid <b>or</b> Quinupristin/dalfopristin for $\geq 2$ weeks (addition of Rifampin may be of benefit, for MRSA)
	Enterobacteriaceae <i>Pseudomonas</i>	Piperacillin/tazobactam <b>or</b> Ticarcillin/clavulanate <b>with or without</b> an aminoglycoside* for $\geq 2$ weeks	Meropenem <b>or</b> Imipenem <b>with or without</b> an aminoglycoside* for $\geq 2$ weeks
	ESBL-producing gram-negative bacilli	Meropenem <b>or</b> Ertapenem <b>with or without</b> an aminoglycoside* for $\geq 2$ weeks	—

## Central Venous Catheter Infections (Broviac, Hickman, Mediport)

(cont'd)

Subset	Usual Pathogens	Preferred IV Therapy <sup>†</sup>	Alternate IV Therapy <sup>†</sup>
	Candida	Amphotericin B × 2–6 weeks <sup>‡</sup>	Fluconazole <b>or</b> Caspofungin <b>or</b> liposomal amphotericin B (if renal dysfunction) × 2–6 weeks <sup>‡</sup>

MSSA/MRSA = methicillin-sensitive/resistant *S. aureus*. Duration of therapy represents total time IV.

† See pp. 414–422 for drug dosages.

\* Gentamicin, tobramycin, or amikacin.

‡ Based on promptness of line removal, clearance of blood cultures, evidence of metastatic foci.

¶ If severely ill or suspicion of MRSA based on local epidemiology, include vancomycin in initial regimen pending culture results.

**Clinical Presentation:** Fever ± site tenderness, erythema.

**Diagnostic considerations:** Quantitative blood cultures from the peripheral blood/vascular catheter are best used to make the diagnosis. However in clinical practice these are not often obtained, and the diagnosis is based on culture results in conjunction with one or more of the following features: local phlebitis or inflammation at the catheter insertion site; embolic disease distal to the catheter; sepsis refractory to appropriate therapy; resolution of fever after device removal; or clustered infections caused by infusion-associated organisms.

**Pitfalls:** It may be difficult to differentiate infection from contamination in blood cultures, especially with coagulase-negative staphylococci. Multiple positive cultures with the same organism and/or clinical features noted above suggest infection, not colonization. Semiquantitative culture of the catheter tip yielding ≥ 15 colonies may also be useful in confirming the diagnosis but requires removal of the device.

**Therapeutic Considerations:** Indications for catheter removal include septic shock, tunnel infection, failure to respond to treatment within 48–72 hours, or infection with *Candida*, atypical mycobacteria, or possibly *S. aureus*. Otherwise attempt to retain the catheter while treating with antibiotics. Localized exit site infections (erythema, induration, tenderness, purulence) within 2 cm of the exit site should be treated topically (e.g., Neosporin, Bacitracin, Bactroban) in conjunction with systemic therapy.

**Prognosis:** Major complications (septic emboli, endocarditis, vasculitis) are rare with aggressive therapy. Recrudescence infection can occur after therapy/apparent clearance and can often be successfully treated with additional courses of antibiotics; persistence ultimately requires line removal.

## Empiric Therapy of Gastrointestinal Infections

### Acute Diarrheal Syndromes (Gastroenteritis)

Subset	Usual Pathogens	Preferred Therapy <sup>†</sup>	Alternate Therapy <sup>†</sup>
Community-acquired	Viruses (Rotavirus, Norwalk agent, enteric adenovirus, enteroviruses)	No specific therapy indicated	No specific therapy indicated

**Acute Diarrheal Syndromes (Gastroenteritis)** (cont'd)

Subset	Usual Pathogens	Preferred Therapy <sup>†</sup>	Alternate Therapy <sup>†</sup>
	Salmonella non-typhi	Ceftriaxone (IV) <b>or</b> Cefotaxime (IV) × 10–14 days*	TMP–SMX (IV or PO) <b>or</b> Amoxicillin (PO) <b>or</b> cefixime (PO) × 10–14 days
	Shigella	Ceftriaxone (IV) <b>or</b> Azithromycin (PO) × 5 days	TMP–SMX (PO) <b>or</b> Cefixime (PO) <b>or</b> Ampicillin (PO) × 5 days
	Campylobacter	Erythromycin (PO) × 7 days <b>or</b> Azithromycin (PO) × 5 days	Doxycycline (PO) (> 8 year old) × 7 days
	Yersinia enterocolitica	TMP–SMX (PO) × 5–7 days	Cefotaxime (IV) <b>or</b> Doxycycline (PO) × 5–7 days
Traveler's diarrhea	E. coli	TMP–SMX (PO) × 3 days	Azithromycin (PO) × 3 days
Typhoid (enteric) fever	Salmonella typhi	Ceftriaxone (IV) <b>or</b> Cefotaxime (IV) × 10–14 days	TMP–SMX (IV or PO) <b>or</b> Amoxicillin (PO) <b>or</b> Cefixime (PO) × 10–14 days
Antibiotic-associated colitis	Clostridium difficile	Metronidazole (PO) × 7–10 days <b>or</b> Nitazoxanide (PO) × 3 days	Vancomycin (PO) × 7–10 days
Chronic watery diarrhea	Giardia lamblia <sup>‡</sup>	Metronidazole (PO) × 5–7 days <b>or</b> Nitazoxanide (PO) × 3 days	Tinidazole (PO) × 1 dose <b>or</b> Furazolidone (PO) × 7–10 days <b>or</b> Albendazole × 5–7 days
	Cryptosporidia <sup>‡</sup>	Nitazoxanide (PO) × 3 days	Human immunoglobulin (PO) <b>or</b> bovine colostrum (PO) for immunocompromised hosts
Acute dysentery	E. histolytica	Metronidazole (PO) × 10 days <b>followed by either</b> Iodoquinol (PO) × 20 days <b>or</b> Paromomycin (PO) × 7 days	Tinidazole (PO) × 3–5 days <b>followed by either</b> Iodoquinol (PO) × 20 days <b>or</b> Paromomycin (PO) × 7 days

## Acute Diarrheal Syndromes (Gastroenteritis) (cont'd)

Subset	Usual Pathogens	Preferred Therapy <sup>†</sup>	Alternate Therapy <sup>†</sup>
	Shigella**	Ceftriaxone (IV) <b>or</b> Azithromycin (PO) × 5 days	TMP-SMX (PO) <b>or</b> Cefixime (PO) <b>or</b> Ampicillin (PO) × 5 days

<sup>†</sup> See pp. 414–422 for drug dosages.

\* Therapy only indicated in child < 3 to 6 months of age, immunocompromised host, or toxic appearing child.

\*\* Mild cases acquire no antibiotic therapy. However, antibiotic therapy shortens durations of symptoms and by decreasing the duration of diarrhea limits potential syneral.

<sup>†</sup> May also present as acute watery diarrhea.

### Acute Gastroenteritis (Community-Acquired)

**Clinical Presentation:** Typically presents with the acute onset of diarrhea with fever. This is not an indication for antibiotic therapy unless illness is severe ( $\geq 6$  unformed stools/day, fever  $\geq 102^\circ$  F, bloody stools). Travel history regarding risk for potential *E. coli* and parasitic exposures is important.

**Diagnostic Considerations:** In the absence of blood in the stools viruses are the most common cause of acute community-acquired gastroenteritis. Rotaviruses are the most common cause of acute gastroenteritis in 4–24 month old children. Enteric adenoviruses, Norwalk-like virus, enteroviruses and astroviruses are common causes of gastroenteritis in older children. Commercially available antigen tests using ELISA or latex agglutination techniques are readily available to detect rotavirus. Testing for the other viral agents may not be as readily available. Inflammatory changes (presence of white blood cells and/or blood) in the stool are more consistent with bacterial infection. When requesting stool cultures, it may be necessary to order special conditions/media for the detection of yersinia or *E. coli* O157.

**Pitfalls:** Antimotility drugs may worsen the course of illness in children with colitis. Empiric therapy with antibiotics may prolong the carriage of *Salmonella* or increase the risk for developing hemolytic uremic syndrome (HUS) with *E. coli* O157 infection. The benefit of antibiotics in treating diarrhea caused by yersinia is also unproven. Thus, antibiotic therapy is not routinely indicated prior to culture results in most instances of mild diarrheal disease, especially as most such infections are self-limited.

**Therapeutic Considerations:** As above, pending cultures in the absence of severe symptoms or dysentery antibiotics may not be indicated. Overall, the fluoroquinolones have the most complete spectrum for pathogens causing bacterial diarrhea but are not presently approved for use in children < 18 years of age.

**Prognosis:** Good. Up to 5–10% of children with *E. coli* O157 are at risk for hemolytic-uremic syndrome.

### Giardiasis (*Giardia lamblia*)

**Clinical Presentation:** *Giardia lamblia* is the most common parasite causing diarrheal illness in children. Giardiasis may present as acute watery diarrhea with abdominal pain and bloating or as a chronic, intermittent illness with foul smelling stools, abdominal distension, and anorexia.

**Diagnostic Considerations:** Trophozoites or cysts of *Giardia lamblia* can be identified in direct examination of infected stools with 75%–95% sensitivity on a single specimen. Testing 3 or more stools further

increases sensitivity of detection. If giardiasis is suspected with negative stool tests, examination of duodenal contents (Entero- or string test) may be helpful.

**Pitfalls:** Asymptomatic infection is commonly seen in children in day care; therefore, indications to treat must take into account stool testing results *and* clinical findings.

**Therapeutic Considerations:** Treatment failures occur commonly, and retreatment with the same initial drug is recommended. Furazolidone is the only pediatric liquid available for treating giardiasis < 3 years of age.

**Prognosis:** Good.

### **Cryptosporidiosis**

**Clinical Presentation:** Usually presents as fever, vomiting, and non-bloody, watery diarrhea. Infection may also be asymptomatic. More severe and chronic infection is seen in immunocompromised patients (e.g., HIV infection). Cryptosporidia parasite is resistant to chlorine and maybe transmitted in swimming pools. Transmission can also occur from livestock, and a major outbreak through contamination of a public water supply has been reported.

**Diagnostic Considerations:** Cryptosporidium cysts are detected by microscopic examination of Kinyoun-stained stool specimens using a sucrose floatation method or formalin-ethyl acetate method to concentrate oocysts. This test is not part of routine stool ova and parasite examination and must be specifically requested. Shedding is intermittent; therefore, 3 stool samples should be submitted for optimal detection.

**Pitfalls:** The oocysts of cryptosporidium are small and may be missed by an inexperienced exam-iner. A commercially available ELISA test is available but may have false-positive and false-negative results.

**Therapeutic Considerations:** Treatment failures are frequent. In immunocompromised hosts oral human immune globulin and bovine colostrum are beneficial. Improvement in CD<sub>4</sub> counts with antiretroviral therapy in HIV-infected patients shortens the clinical illness.

**Prognosis:** Good. Recovery may take months.

### **Amebiasis (Entamoeba histolytica)**

**Clinical Presentation:** E. histolytica can lead to a spectrum of clinical illnesses from asymptomatic infection to acute dysentery to fulminant colitis. Disseminated disease, primarily manifest as hepatic abscesses, can also occur. E. histolytica is most prevalent in developing countries and is transmitted by the fecal-oral route.

**Diagnostic Considerations:** Trophozoites or cysts of E. histolytica can be identified in stool specimens. In more severe disease (amebic colitis, hepatic abscesses), serum antibodies can be detected.

**Pitfalls:** Treatment with steroids or antimotility drugs can worsen symptoms and should not be used.

**Therapeutic Considerations:** Treatment is two-staged to eliminate tissue-invading trophozoites and organisms in the intestinal lumen. Surgical drainage of large hepatic abscesses may be beneficial.

**Prognosis:** Good.

### **Clostridium difficile Colitis**

**Clinical Presentation:** Classically occurs in a child receiving antibiotic therapy and presents as diarrhea, cramping, bloody/mucous stools, abdominal tenderness, fever, and toxicity. It may also present weeks after a course of antibiotics.

**Diagnostic Considerations:** *C. difficile* toxin can be detected with commercially available immunoassays. Endoscopic finding of pseudomembranous colitis is the definitive method of diagnosis although rarely indicated.

**Pitfalls:** *C. difficile* may be normal flora in infants < 1 year of age and probably does not cause illness in this age group. The finding of *C. difficile* toxin in an infant should not be equated with cause and additional evaluations should be pursued. Antimotility drugs may worsen symptoms and should be avoided.

**Therapeutic Considerations:** Cessation of antibiotics is recommended if possible in the presence of significant *C. difficile* colitis. Patients with severe symptoms or persistent diarrhea  $\times$  2–3 days after discontinuing antibiotics should then be treated with oral metronidazole or oral vancomycin. Up to ~25% relapse after a single course; re-treatment using the same initial antibiotic is recommended. For persistent infection alternate therapies such as nitazoxanide, rifaximin, tinidazole, oral immune globulin therapy, toxin binders or repopulating intestinal flora may be considered.

**Prognosis:** Good.

## Empiric Therapy of Bone and Joint Infections

### Septic Arthritis

Subset	Usual Pathogens	Preferred IV Therapy <sup>†*</sup>	Alternate IV Therapy <sup>†*</sup>	IV-to-PO Switch <sup>†</sup>
Newborns ( $\leq$ 3 months)	<i>S. aureus</i> <sup>‡</sup> Group B streptococci Enterobacteriaceae <i>N. gonorrhoeae</i>	<b>Combination therapy with Nafcillin or Vancomycin plus either Ceftriaxone or Cefotaxime*</b>	<b>Combination therapy with Vancomycin or Clindamycin plus either ceftriaxone or cefotaxime or gentamicin or tobramycin*</b>	Not applicable
Child ( $>$ 3 months to 14 years)	<i>S. aureus</i> <sup>‡</sup> Group A streptococci <i>S. pneumoniae</i> Gram-negative bacilli <i>N. meningitidis</i> ( <i>H. influenzae</i> type b if unvaccinated) <i>Kingella</i>	<b>Combination therapy with Nafcillin or Vancomycin plus either Ceftriaxone or Cefotaxime*</b>	<b>Combination therapy with Vancomycin or Clindamycin plus either Ceftriaxone or Cefotaxime*</b>	Dicloxacillin or Cephalexin or Clindamycin (PO) after 1 week of IV therapy*

<sup>†</sup> See pp. 414–422 for drug dosages.

\* Total duration of therapy for non-gonococcal septic arthritis  $\geq$  3 weeks based on clinical response.

<sup>‡</sup> If CA-MRSA is suspected based on clinical presentation/local epidemiology, consider empiric coverage for CA-MRSA with clindamycin, TMP-SMX, doxycycline, or vancomycin pending culture results.

## Septic Arthritis (cont'd)

Subset	Usual Pathogens	Preferred IV Therapy**	Alternate IV Therapy**	IV-to-PO Switch†
Adolescents (sexually active)	As above plus <i>N. gonorrhoeae</i> (typically 2 or 3 joints involved)	<b>Combination therapy with</b> Nafcillin <b>or</b> Vancomycin <b>plus either</b> Ceftriaxone <b>or</b> Cefotaxime*	If GC isolated and <u>penicillin allergy</u> : Azithromycin	GC arthritis with prompt response to IV therapy may switch to Cefixime to complete 7 days of total therapy

Duration of therapy represents total time IV or IV + PO. Most patients on IV therapy able to take PO meds should be switched to PO therapy after clinical improvement. Taper to individual drug therapy once organism isolated and sensitivities are available.

† See pp. 414–422 for drug dosages.

\* Total duration of therapy for non-gonococcal septic arthritis ≥ 3 weeks based on clinical response.

‡ If CA-MRSA is suspected based on clinical presentation/local epidemiology, consider empiric coverage for CA-MRSA with clindamycin, TMP-SMX, doxycycline, or vancomycin pending culture results.

**Clinical Presentation:** Varies with age of the child. Presentation is often nonspecific in infants (i.e., fever, poor feeding, tachycardia). Physical exam findings can be subtle: asymmetrical tissue folds, unilateral swelling of an extremity, subtle changes in limb/joint position. In older children, signs and symptoms are more localized to the involved joint. Commonly affected joints: hips, elbows, knees.

**Diagnostic Considerations:** Joint aspiration (large-bore needle) shows 50,000–250,000 WBCs/mm<sup>3</sup> with a marked predominance of PMNs. Gram stain/culture of joint fluid confirm the diagnosis. Toxic synovitis, Lyme arthritis, rheumatoid arthritis, traumatic arthritis, and sympathetic effusion from adjacent osteomyelitis can mimic septic arthritis on presentation, but joint fluid has fewer WBCs and cultures are negative. Multiple joint involvement is more typical of rheumatic/Lyme disease. In young infants, persistence of the nutrient artery can lead to osteomyelitis and septic arthritis.

**Pitfalls:** Septic arthritis of the hip is an emergency concomitant condition requiring prompt joint aspiration and irrigation to minimize the risk of femoral head necrosis, which may occur within 24 hours. Toxic synovitis of the hip, which is treated with anti-inflammatory agents and observation, typically causes less fever, pain, and leukocytosis than septic arthritis; however, at times the two can not be differentiated and aspiration of the joint is indicated. In sexually active adolescents, consider *N. gonorrhoeae* and culture aspirate appropriately.

**Therapeutic Considerations:** Therapy is recommended for at least 3–4 weeks (IV followed by oral therapy), for a total antibiotic course of 4–6 weeks based on clinical response and laboratory parameters (WBC, ESR, CRP). Intra-articular therapy is not helpful. Empiric coverage is broader than for osteomyelitis in children.

**Prognosis:** Good with prompt joint aspiration and at least 3 weeks of antibiotics.

## Acute Osteomyelitis, Osteochondritis, Diskitis

Subset	Usual Pathogens	Preferred IV Therapy†	Alternate IV Therapy†	IV-to-PO Switch or PO Therapy†
Acute osteomyelitis Newborn (0–3 months)	S. aureus <sup>§</sup> Gram-negative bacilli Group B streptococci	<b>Combination therapy with Nafcillin or Oxacillin plus either Cefotaxime or Ceftriaxone</b> × 4–6 weeks	Vancomycin <b>plus either</b> Cefotaxime <b>or</b> Ceftriaxone × 4–6 weeks	Not applicable
Osteochondritis	P. aeruginosa S. aureus <sup>§</sup>	Ticarcillin clavulanate × 7–10 days after surgery <b>or combination therapy with Nafcillin plus Ceftazidime</b> × 7–10 days after surgery	Piperacillin/tazobactam <b>or</b> Ciprofloxacin × 7–10 days after surgery	Ciprofloxacin** × 7–10 days after surgery
Diskitis	S. aureus, K. kingae Enterobacteriaceae S. pneumoniae S. epidermidis	<b>PO Therapy:</b> Cephalexin <b>or</b> Cefadroxil <b>or</b> Clindamycin × 3–4 weeks or ESR returns to normal††		

Duration of therapy represents total time IV, PO, or IV + PO. Most patients on IV therapy able to take PO meds should be switched to PO therapy soon after clinical improvement.

† **See pp. 414–422 for drug dosages.**

\*\* Not approved for children but might consider in adolescent.

\* Treat only IV or IV-to-PO switch.

†† Add gram-negative coverage only if culture proven.

§ If CA-MRSA is suspected based on clinical presentation/local epidemiology, consider empiric coverage for CA-MRSA with clindamycin, TMP-SMX, doxycycline, or vancomycin pending culture results.

### Acute Osteomyelitis

**Clinical Presentation:** Acute onset of fever and pain/decreased movement around the infected area. Can be difficult to localize, especially in younger children. Occurs primarily via hematogenous spread (vs. direct inoculation) to metaphysis of long bones.

**Diagnostic Considerations:** Acute phase reactants are elevated (ESR, CRP, WBC). X-rays may not reveal osteolytic lesions for > 7 days, but soft tissue swelling and periosteal reaction may be seen as early as



3 days. Bone scan/MRI reveal changes in the first 24 hours, but bone scans are insensitive in neonates. Blood cultures may be positive, especially in younger children. Definitive diagnosis by bone aspirate for gram stain and culture, but empiric therapy is often initiated based on clinical history and the presence of an acute lytic lesion on MRI, bone scan, or x-ray.

**Pitfalls:** Bony changes on x-ray may not be present initially. It may be difficult to differentiate cellulitis from osteomyelitis, even on bone scan.

**Therapeutic Considerations:** Initiate empiric therapy by IV route. Treatment is required  $\geq 4$  weeks, but children with a prompt clinical response can complete therapy with high-dose oral antibiotics. With adequate treatment, CPR and ESR normalize over 2 weeks and 4 weeks, respectively.

**Prognosis:** Good with 4–6 weeks of total therapy. Long-term growth of affected bone may be impaired.

### **Osteochondritis of the Foot (“Puncture Wound Osteomyelitis”)**

**Clinical Presentation:** Tenderness, erythema, and swelling several days to weeks after a nail puncture wound through a sneaker/tennis shoe (*Pseudomonas* found in foam layer between sole and lining of shoe). Fever and other systemic complaints are rare. Develops in 1–2% of puncture wounds of foot.

**Diagnostic Considerations:** *P. aeruginosa* is the primary pathogen, although infection with *S. aureus* is also a concern.

**Pitfalls:** Antibiotics with surgical debridement of necrotic cartilage.

**Therapeutic Considerations:** With prompt debridement, 7–10 days of antibiotic therapy is usually sufficient.

**Prognosis:** Good with debridement and antibiotics.

### **Diskitis**

**Clinical Presentation:** Typically involves the lumbar region and occurs in children  $< 6$  years old. Presents with gradual onset (over weeks) of irritability and refusal to walk. Fever is low-grade or absent. Older children may be able to localize pain to back, hip, or abdomen. Pain is exacerbated by motion of the spine and can be localized by percussion of the vertebral bodies.

**Diagnostic Considerations:** ESR is elevated, but WBC count is normal. X-rays are usually normal initially, but later reveal disk space narrowing and sclerosis of the vertebrae. Increased disk space uptake can be seen on bone scan. MRI is very sensitive for assessing disk space involvement.

**Therapeutic Considerations:** Diskitis is probably a low-grade bacterial infection, but the role of antibiotic therapy is unclear. Bed rest and anti-inflammatory medications are the mainstays of treatment. Immobilization may be required for severe cases. Oral antibiotics are given for 3–4 weeks or until the ESR returns to normal.

**Pitfalls:** Difficult to isolate an organism, even with needle aspiration of disk space. *S. aureus* is the most common pathogen, but diskitis can also be caused by coagulase-negative staphylococcus, *K. kingae*, coliforms, *S. pneumoniae*.

**Prognosis:** Fusion of involved vertebrae may occur as the infection resolves. Otherwise, outcome is generally good.

## Empiric Therapy of Skin and Soft Tissue Infections

### Skin and Soft Tissue Infections

Subset	Usual Pathogens	Preferred IV Therapy <sup>†</sup>	Alternate IV Therapy <sup>†</sup>	PO Therapy or IV-to-PO Switch <sup>†</sup>
Cellulitis, impetigo	<i>S. aureus</i> Group A streptococci	Cefazolin <b>or</b> nafcillin <b>or</b> oxacillin × 7–10 days	Clindamycin <sup>†</sup> × 7–10 days	Cephalexin <b>or</b> Cefadroxil <b>or</b> Dicloxacillin <b>or</b> Clindamycin <b>or</b> amoxicillin/clavulanate <b>or</b> Erythromycin <b>or</b> Azithromycin × 7–10 days
Severe pyoderma, abscesses	CA-MRSA	Vancomycin <b>plus</b> Clindamycin × 7–14 days	Vancomycin <b>or</b> Linezolid × 7–14 days	TMP-SMX <b>or</b> Doxycycline × 7–14 days**
Animal bite wounds (dog/cat)	Group A streptococci <i>P. multocida</i> Capnocytophaga <i>S. aureus</i>	Ampicillin-sulbactam × 7–10 days	Piperacillin <b>or</b> Ticarcillin × 7–10 days  <u>Penicillin allergy:</u> Clindamycin <b>plus</b> TMP-SMX × 7–10 days (dog bites); Doxycycline <b>or</b> Cefuroxime × 7–10 days (cat bites)	Amoxicillin/clavulanate <b>or</b> Doxycycline × 7–10 days
Human bite wounds	Oral anaerobes <i>E. corrodens</i> Group A streptococci <i>S. aureus</i>	Ampicillin-sulbactam × 5–7 days	Piperacillin <b>or</b> Ticarcillin × 5–7 days  <u>Penicillin allergic patient:</u> Clindamycin <b>plus</b> TMP-SMX × 5–7 days	Amoxicillin/clavulanate <b>or</b> Doxycycline × 5–7 days
Cat scratch disease (CSD)*	<i>Bartonella henselae</i>	Gentamicin × 10–14 days	—	Azithromycin × 5 days <b>or</b> TMP-SMX <b>or</b> Ciprofloxacin <b>or</b> Rifampin × 10–14 days

† See pp. 414–422 for drug dosages.

\*\* In children > 8 years of age.

\* No well-controlled trials of antibiotic treatment for CSD to demonstrate benefit.

‡ Preferred therapy for patients in geographic regions with a high prevalence of MRSA or for those with penicillin/cephalosporin allergy.

## Skin and Soft Tissue Infections (cont'd)

Subset	Pathogens	IV Therapy <sup>†</sup>	PO Therapy or IV-to-PO Switch <sup>†</sup>
Chicken pox <i>Immuno-compromised host</i>	VZV	Acyclovir × 7–10 days	Acyclovir <b>or</b> Valacyclovir × 7–10 days
<i>Immuno-competent host</i>	VZV	Acyclovir × 5 days	Acyclovir <b>or</b> Valacyclovir × 5 days
H. zoster (Shingles)	VZV	Same as for chicken pox	Acyclovir <b>or</b> Valacyclovir × 10 days (for individuals ≥ 12 years of age)

Duration of therapy represents total time IV, PO, or IV + PO. Most patients on IV therapy able to take PO meds should be switched to PO therapy soon after clinical improvement.

† See pp. 414–422 for drug dosages.

### Cellulitis

**Clinical Presentation:** Erythema, warmth, and tenderness of skin. Impetigo is characterized by a vesiculopapular rash with honey-colored discharge.

**Diagnostic Considerations:** Primarily a clinical diagnosis. Group A streptococci is the primary pathogens in healthy children. Cellulitis, alone without a pustular component, is caused by streptococci (not staphylococci).

**Pitfalls:** Differential diagnosis of cellulitis may include hypersensitivity to insect bites. Herpetic whitlow (HSV) may be mistaken for a bacterial skin or paronychia infection.

**Therapeutic Considerations:** First generation cephalosporin or semi-synthetic penicillin with anti-staphylococcal activity (i.e., dicloxacillin, nafcillin, oxacillin) are drugs of choice. Increasing incidence of community-acquired MRSA may affect treatment decisions.

**Prognosis:** Excellent. Impetigo may only require topical treatment (Mupirocin).

### Bite Wounds

**Clinical Presentation:** 80% of animal bite wounds in children are from dogs, and 15%–50% of dog bites become infected. More than 50% of cat bites become infected, and due to their long teeth, there is an increased risk of inoculation into bone/joints with development of osteomyelitis/septic arthritis. Human bite wounds are most prone to infection, and 75%–90% of all human bites become infected.

**Diagnostic Considerations:** Clinical diagnosis. Culture of wound exudate may yield organism.

**Pitfalls:** Failure to assess depth of infection, especially with cat bites, may result in late identification of bone/joint infection. Macrolides are ineffective against *P. multocida*.

**Therapeutic Considerations:** It is important to cover oral anaerobes, *S. aureus*, and Group A streptococci in human bite wounds. *P. multocida* is an important pathogen in cat and dog bite wounds. Facial/hand lesions require plastic surgery evaluation. Recurrent debridement may be necessary, especially with human bite wound infections of hand. Assess tetanus immunization status for all bite wounds

(p. 377). For human bites consider the risk of HIV and hepatitis B. For dog bites consider the risk of rabies. Antimicrobial therapy initiated within 8 hours of a bite wound and administered for 2–3 days may decrease the rate of infection.

**Prognosis:** Good with early debridement and antibiotics.

### **Cat Scratch Disease (CSD)**

**Clinical Presentation:** Classic presentation is a papular lesion at site of cat scratch with lymphadenitis in the draining region (axillary, epitrochlear, inguinal, cervical most commonly). Frequently associated with fever/malaise 1–2 weeks after scratch. Infection can present with conjunctivitis and ipsilateral preauricular lymph node (Parinaud oculoglandular syndrome). Unusual presentations in normal hosts include encephalitis, hepatitis, microabscesses in liver/spleen, fever of unknown origin, osteolytic lesions.

**Diagnostic Considerations:** Most often secondary to kitten scratch with *Bartonella henselae*. Diagnosed using specific serology for antibodies to *B. henselae*.

**Pitfalls:** Failure to obtain history of kitten exposure. Surgical excision of lymph node is generally not necessary.

**Therapeutic Considerations:** Most lesions are self-limited and resolve over 2–4 months. If lymph nodes are fluctuant, I&D may be indicated. Antibiotic therapy may be helpful in severe cases with hepatosplenomegaly. Doxycycline, erythromycin, or azithromycin are helpful in immunocompromised hosts.

**Prognosis:** Very good with spontaneous resolution over 2–4 months.

### **Chicken Pox/Shingles (VZV)**

**Clinical Presentation:** Primary illness is chicken pox, a generalized pruritic, vesicular rash with fever that erupts in crops of lesions over 3–5 days followed by crusting and recovery. Complications include bacterial superinfection of skin lesions, sepsis, cerebellar ataxia, thrombocytopenia, hepatitis, pneumonia and encephalitis. The disease tends to be more severe in adolescents and adults, particularly if immunocompromised. Primary infection early in pregnancy can rarely result in varicella embryopathy with limb atrophy and CNS malformations in the neonate. Reactivation disease (shingles) may occur in children and in normal hosts and remains localized to a single dermatome. Post-herpetic neuralgia occurs less often in children than adults. Reactivation disease in immunocompromised hosts can spread and re-disseminate.

**Diagnostic Considerations:** The characteristic eruption of chicken pox occurs in waves—multiple stages appear at the same time, from new papules and vesicles to more advanced larger crusted lesions—and is unique to varicella. Direct fluorescent antibody staining of a scraped lesion or PCR can confirm the diagnosis.

**Pitfalls:** Initially lesions may be primarily papular, and if the diagnosis is not considered, exposure of others can occur.

**Therapeutic Considerations:** Antiviral therapy with acyclovir, if started within 24 hours of rash, should be considered for children > 12 years of age, those on steroid or salicylate therapy, and those with underlying chronic pulmonary, skin, or immunosuppressive states. Oral administration is acceptable, although IV therapy may be preferred for immunocompromised hosts at risk of disseminated disease. More severe varicella has been observed in individuals acquiring the infection from a household contact, presumably due to a higher inoculum with closer contact; non-immune household contacts may be considered for acyclovir therapy at onset of rash in child. Additionally,

individuals  $\geq 13$  years may develop more extensive varicella than younger children. Immunocompromised children or pregnant women without a history of varicella or immunization may benefit from prophylaxis with varicella zoster immune globulin (VZIG) within 96 hours of varicella exposure. Newborns whose mothers develop chicken pox within 5 days before or 48 hours after delivery and exposed premature infants are also candidates for VZIG. A live-attenuated varicella vaccine has been licensed since 1995 for use in individuals  $\geq 12$  months of age who have not had chicken pox. A two-dose vaccine schedule is recommended for children  $\geq 12$  months of age.

**Prognosis:** Overall prognosis is good with complete recovery and minimal risk of scarring unless immunosuppressed host with disseminated disease. Although rare, Group A streptococcal toxic shock syndrome (manifest as cellulitis or in conjunction with necrotizing fasciitis complicating varicella skin lesions) and Group A streptococcal septicemia, which can occur in the absence of apparent secondarily infected skin lesions, may be fatal complications of varicella in normal children.

## Common Pediatric Antimicrobial Drugs

Drug	Dosage in Neonates	Dosage in Infants/Children*
Acyclovir	20 mg/kg (IV) q8h $\times$ 14–21 days. Dosing interval may need to be increased for infants < 34 weeks post-maturational age (GA + CA) or if significant renal impairment or liver failure followed by <u>Chronic suppression:</u> 75 mg/kg (PO) q12h $\times$ 6 months	<u>HSV encephalitis:</u> 10 mg/kg (IV) q8h $\times$ 14–21 days <u>Primary HSV infection:</u> 10–20 mg/kg (PO) q6h $\times$ 5–10 days <b>or</b> 5 mg/kg/dose (IV) q8h $\times$ 5 days <u>Varicella in immunocompromised hosts:</u> 10 mg/kg (IV) q8h $\times$ 7–10 days <u>Varicella in immunocompetent hosts:</u> 20 mg/kg (PO) q6h $\times$ 5 days (maximum 800 mg/dose)
Albendazole	Not applicable	400 mg (PO) q24h
Amikacin**	<u>During first week of life</u> dosing is based on gestational age (administer IV dose over 30 min) <ul style="list-style-type: none"> <li>• <math>\leq 27</math> weeks (or asphyxia, PDA, or indomethacin): 18 mg/kg (IV) q48h</li> <li>• 28–30 weeks: 18 mg/kg (IV) q36h</li> <li>• 31–33 weeks: 16 mg/kg (IV) q36h</li> <li>• <math>\geq 34</math> weeks: 15 mg/kg (IV) q24h</li> </ul>	5–7.5 mg/kg (IV or IM) q8h

Drug	Dosage in Neonates			Dosage in Infants/Children*
	<i>After first week of life:</i> Initial dose of 15 mg/kg, then draw serum concentrations 30 min after end of infusion (peak) and 12–24 hours later (trough) to determine dosing interval. Aim for peak of 20–30 mcg/mL and trough of 2–5 mcg/mL.			
Amoxicillin	Not indicated			22.5–45 mg/kg (PO) q12h
Amoxicillin-clavulanate	Not indicated			22.5–45 mg/kg (of amoxicillin component) (PO) q12h
Amphotericin B (conventional)	0.5–1 mg/kg (IV over 2–6 hours) q24–48h (Some authorities recommend an initial test dose of 0.1–0.5 mg/kg IV over 2–6 hours)			
Ampicillin	25–50 mg/kg/dose (IV or IM). <u>Severe Group B streptococcal sepsis:</u> 100 mg/kg/dose. Dosing interval is based on gestational age (GA) and chronological age (CA):			25–50 mg/kg (IV or IM) q6h
	<b>GA + CA (weeks)</b>	<b>CA (days)</b>	<b>Interval (hours)</b>	
	≤ 29	0–28 > 28	12 8	
	30–36	0–14 > 14	12 8	
	≥ 37	0–7 > 7	12 8	
Ampicillin-sulbactam	Not indicated			25–50 mg/kg (of ampicillin component) (IV) q6h
Azithromycin	Not indicated			Otitis media/sinusitis: 30 mg/kg (PO) × 1 dose <b>or</b> 10 mg/kg (PO) q24h × 3 days <b>or</b> 10 mg/kg (PO) on day 1 followed by 5 mg/kg (PO) q24h on days 2–5 <u>Pharyngitis/tonsillitis:</u> 12 mg/kg (PO) q24h × 5 days

\* Dosages are generally based on weight (mg/kg), up to adult dose as maximum.

\*\* Drug can be given IM but absorption may be variable.

Drug	Dosage in Neonates	Dosage in Infants/Children*
		Community-acquired pneumonia (not indicated for moderate or severe disease): 10 mg/kg (PO) × 5 days <b>or</b> 10 mg/kg (IV or PO) on day 1 followed by 5 mg/kg (IV or PO) q24h on days 2–5 Skin/soft tissue infections (including Cat Scratch Disease): 10 mg/kg (PO) on day 1 followed by 5 mg/kg (PO) q24h on days 2–5
Aztreonam	30 mg/kg (IV or IM). See <i>ampicillin</i> for dosing interval (p. 415)	30 mg/kg (IV or IM) q6–8h
Caspofungin	70 mg/m <sup>2</sup> loading dose, then 25 mg/m <sup>2</sup> (IV) q24h	70 mg/m <sup>2</sup> loading dose, then 50 mg/m <sup>2</sup> (IV) q24h (> 3 months)
Cefadroxil	Not indicated	15 mg/kg (PO) q12h
Cefazolin	25 mg/kg (IV or IM). See <i>ampicillin</i> for dosing interval (p. 415)	25–100 mg/kg/day (IV or IM) divided q6–q8h
Cefdinir	Not indicated	7 mg/kg (PO) q12h or 14 mg/kg (PO) q24h
Cefepime	50 mg/kg (IV) q12h	33.3–50 mg/kg (IV or IM) q8h
Cefotaxime	50 mg/kg (IV or IM). (25 mg/kg/dose is adequate for gonococcal infection). See <i>ampicillin</i> for dosing interval (p. 415)	25–50 mg/kg (IV or IM) q6–8h
Cefotetan	Not indicated	20–40 mg/kg (IV or IM) q12h
Cefoxitin	25–33 mg/kg/dose (IV or IM). See <i>ampicillin</i> for dosing interval (p. 415)	80–160 mg/kg/day (IV or IM) divided q4–8h
Cefpodoxime	Not indicated	5 mg/kg (PO) q12h
Cefprozil	Not indicated	15 mg/kg (PO) q12h
Ceftazidime	30 mg/kg/dose (IV or IM). See <i>ampicillin</i> for dosing interval (p. 415)	25–50 mg/kg (IV or IM) q8h
Ceftibuten	Not indicated	9 mg/kg (PO) q24h
Ceftizoxime	Not indicated	50 mg/kg (IV or IM) q6–8h

\* Dosages are generally based on weight (mg/kg), up to adult dose as maximum.

Drug	Dosage in Neonates			Dosage in Infants/Children*
Ceftriaxone <sup>‡</sup>	<u>Sepsis and disseminated gonococcal infection</u> : 50 mg/kg (IV or IM) q24h <u>Meningitis</u> : 100 mg/kg loading dose followed by 80 mg/kg (IV or IM) q24h <u>Uncomplicated gonococcal ophthalmia</u> : 50 mg/kg (maximum 125 mg) as a single dose (IV or IM)			50 mg/kg (IV or IM) q24h. Meningitis: 50 mg/kg (IV or IM) q12h or 100 mg/kg (IV or IM) q24h <u>Acute otitis media</u> : 50 mg/kg (IM) × 1 dose (or 3 doses IM q24h in high-risk patients)
Cefuroxime	Not indicated			10–15 mg/kg (PO) q12h 25–50 mg/kg (IV or IM) q8h
Cephalexin	Not indicated			6.25–25 mg/kg (PO) q6h
Cephalothin	Not indicated			25 mg/kg (IV or IM) q4–6h
Clarithromycin	Not indicated			7.5 mg/kg (PO) q12h
Clindamycin	5.0–7.5 mg (IV or PO). Dosing interval is based on gestational age (GA) and chronological age (CA)			5–10 mg/kg (IV or IM) q6–8h or 10–30 mg/kg/day (PO) divided q6–8h
	<b>GA + CA (weeks)</b>	<b>CA (days)</b>	<b>Interval (hours)</b>	
	< 29	0–28 > 28	12 8	
	30–36	0–14 > 14	12 8	
	37–44	0–7 > 7	8 6	
Dicloxacillin	Not indicated			6.25–12.5 mg/kg (PO) q6h
Doxycycline	Contraindicated			> 45 kg: 100 mg (PO) q12h ≤ 45 kg: 1.1–2.5 mg/kg (PO) q12h Use only in children > 8 years (unless RMSF) 1–2 mg/kg (IV) q12–24h

\* Dosages are generally based on weight (mg/kg), up to adult dose as maximum.

‡ Do not use in presence of hyperbilirubinemia.



Drug	Dosage in Neonates			Dosage in Infants/Children*
Erythromycin	<u>Chlamydia pneumoniae/conjunctivitis or pertussis</u> : 12.5 mg/kg (PO) q6h (E. estolate preferred) <u>Other infections</u> : E. estolate 10 mg/kg (PO) q8h <b>or</b> E. ethylsuccinate 10 mg/kg (PO) q6h <u>Severe infections and PO not possible</u> : 5–10 mg/kg (IV over $\geq$ 60 min) q6h			10–12.5 mg/kg (PO) q6–8h 5–12.5 mg/kg (IV) q6h
Ertapenem	Not indicated			15 mg/kg (IV) q12h (not to exceed 1 gm/day)
Ethambutol	See p. 401			See p. 401
Ethionamide	See p. 401			See p. 401
Fluconazole	<u>Systemic infection or meningitis</u> : 12 mg/kg (IV over 30 min or PO) $\times$ 1 dose, then 6 mg/kg/dose (IV or PO) with dosing interval based on gestational age (GA) and chronological age (CA) (below) <u>Prophylaxis</u> (e.g., extremely low birth weight infants in NICU with high rates of fungal disease): 3 mg/kg/dose (IV or PO) according to dosing interval grid (below) <u>Thrush</u> : 6 mg/kg (PO) $\times$ 1 dose, then 3 mg/kg (PO) q24h			10 mg/kg (IV or PO) loading dose followed by 12 mg/kg (IV or PO) q24h
	<b>GA + CA (weeks)</b>	<b>CA (days)</b>	<b>Interval (weeks)</b>	
	$\leq$ 29	0–14 > 14	72 48	
	30–36	0–14 > 14	48 24	
	37–44	0–7 > 7	48 24	

\* Dosages are generally based on weight (mg/kg), up to adult dose as maximum.

Drug	Dosage in Neonates	Dosage in Infants/Children*
Gentamicin**	<p><i>During first week of life</i> dosing is based on gestational age (administer IV dose over 30 min):</p> <ul style="list-style-type: none"> <li>• ≤ 29 weeks (or asphyxia, PDA, or indomethacin): 5 mg/kg (IV) q48h</li> <li>• 30–33 weeks: 4.5 mg/kg (IV) q48h</li> <li>• 34–37 weeks: 4 mg/kg (IV) q36h</li> <li>• ≥ 38 weeks: 4 mg/kg (IV) q24h</li> </ul> <p><i>After first week of life:</i> Initial dose of 4 mg/kg, then draw serum concentrations 30 min after end of infusion (peak) and 12–24 hours later (trough) to determine dosing interval. Aim for peak of 5–12 mcg/mL and trough of 0.5–1 mcg/mL</p>	<p>2–2.5 mg/kg (IV or IM) q8h  <b>or</b>            4.5–7.5 mg/kg (IV) q24h</p>
Imipenem	20–25 mg/kg (IV) q12h	15–25 mg/kg (IV or IM) q6h
Iodoquinol	No information	10–13.3 mg/kg (PO) q8h
Isoniazid	See p. 401	See p. 401
Linezolid	10 mg/kg (IV) q12h	10 mg/kg (IV) q8h
Liposomal/ lipid complex Amphotericin preparations	1–5 mg/kg (IV over 2 hours) q24h	3–6 mg/kg (IV) q24h
Meropenem	20 mg/kg (IV) q12h	10 mg/kg (IV) q8h (skin); 20 mg/kg (IV) q8h (intraabdominal); 40 mg/kg (IV) q8h (meningitis)
Methenamine mandelate	Not indicated	15–25 mg (PO) q6–8h
Metronidazole	15 mg/kg (IV or PO) × 1 dose, then 7.5 mg/kg/dose (IV or PO) with dosing interval based on gestational age (GA) and chronological age (CA):	5–12.5 mg/kg (PO) q8h 15 mg/kg (IV) × 1 dose followed by 7.5 mg/kg (IV) q6h

\* Dosages are generally based on weight (mg/kg), up to adult dose as maximum.

\*\* Drug can be given IM but absorption may be variable.

Drug	Dosage in Neonates			Dosage in Infants/Children*
	GA + CA (weeks)	CA (days)	Interval (hours)	
	≤ 29	0–28 > 28	24 48	
	30–36	0–14 > 14	24 12	
	37–44	0–7 > 7	24 12	
Micafungin	10 mg/kg (IV) of q24h			4–12 mg/kg (IV) q24 (higher dose for patients < 8 year of age)
Mupirocin	Not indicated			Nasal cream: ½ of single use tube into nostril q12h × 5 days; Cream: apply q8h × 5–10 days
Nafcillin	25–50 mg/kg/dose (IV). See <i>ampicillin</i> for dosing interval (p. 415)			12.5–50 mg/kg (IV or IM) q6h
Nitazoxanide	Not applicable			Children 1–3 years old: 100 mg (PO) q12h; children 4–11 years old: 200 mg (PO) q12h
Nitrofurantoin	Not indicated			<u>UTI</u> : 1.25–1.75 mg/kg (PO) q6h <u>UTI prophylaxis</u> : 1–2 mg/kg (PO) q24h
Nystatin	<u>Oral</u> : 1 mL (preterm) to 2 mL (term) of 100,000 U/mL suspension applied with swab to each side of mouth q6h until 3 days after resolution of lesions <u>Topical</u> : Apply ointment or cream to affected area q6h until 3 days after resolution of lesions.			<u>Suspension</u> : 4–6 mL swish and swallow 4×/day <u>Troche</u> : 1–2 troches 4–5×/ day
Oxacillin	25–50 mg/kg/dose (IV). See <i>ampicillin</i> for dosing interval (p. 415)			25–50 mg/kg (IV or IM) q6h
Paromomycin	Not applicable			10 mg/kg (PO) q8h
Penicillin G	25,000–50,000 IU/kg/dose (IV). See <i>ampicillin</i> for dosing interval (p. 415) <u>Meningitis</u> : 75,000–100,000 IU/kg			12,500–75,000 U/kg (IV or IM) q4–6h

\* Dosages are generally based on weight (mg/kg), up to adult dose as maximum.

Drug	Dosage in Neonates	Dosage in Infants/Children*
	(IV) in meningitis (IV). <sup>†</sup> Q8–12h based on GA + CA. See <i>ampicillin</i> for dosing interval (p. 415). Crystalline penicillin G: IM: procaine penicillin G q24 hours. <u>Congenital syphilis</u> : Aqueous penicillin G 50,000 IU/kg (slow IV push) q12h × 7 days, then q8h to complete 10–14 days <b>or</b> procaine penicillin G 50,000 IU/kg (IM) q24h × 10–14 days	
Penicillin V	Not indicated	6.25–12.5 mg/kg (PO) q6–8h
Piperacillin	50–100 mg/kg/dose (IV or IM). See <i>ampicillin</i> for dosing interval (p. 415)	25–75 mg/kg (IV or IM) q6h; may increase to q4h in severe infection, especially with <i>pseudomonas</i>
Piperacillin-tazobactam	Not indicated	100–300 mg/kg/day (IV) (of piperacillin component) divided q6–8h
Pyrazinamide	See p. 401	See p. 401
Quinupristin/dalfopristin	No information	7.5 mg/kg (IV) q12h
Rifampin	10–20 mg/kg (PO) q24h <b>or</b> 5–10 mg/kg (IV) q24h	20 mg/kg (PO) q24h <b>or</b> 10 mg/kg (PO) q12h 10–20 mg/kg/day (IV) divided q12–24h
Streptomycin	See p. 401	See p. 401
Sulfisoxazole	Contraindicated	30–35 mg/kg (PO) q6h <u>Otitis media prophylaxis</u> : 37.5 mg/kg (PO) q12h
Tetracycline	Contraindicated	5–12.5 mg/kg (PO) q6h. Use only in children > 8 years
Ticarcillin	75–100 mg/kg/dose (IV). See <i>ampicillin</i> for dosing interval (p. 415)	25–75 mg/kg (IV or IM) q6h
Ticarcillin-clavulanate	75–100 mg/kg/dose (of ticarcillin component) (IV). See <i>ampicillin</i> for dosing interval (p. 415)	25–75 mg/kg (of ticarcillin component) (IV or IM) q6h
Tinidazole	Not applicable	50–60 mg/kg (PO) q24h
Tobramycin	Same as gentamicin (p. 419)	2–2.5 mg/kg (IV or IM) q8h or 4.5–7.5 mg/kg (IV) of q24h

\* Dosages are generally based on weight (mg/kg), up to adult dose as maximum.

Drug	Dosage in Neonates			Dosage in Infants/Children*
Trimethoprim-sulfamethoxazole (TMP-SMX)	Contraindicated			UTI: 4–5 mg/kg (of trimethoprim component) (PO) q12h <u>Pneumocystis carinii pneumonia (PCP)</u> 5 mg/kg (PO) q6h (typically after initial IV therapy) <u>UTI prophylaxis:</u> 2–4 mg/kg (PO) q24h <u>IV dosing</u> <u>PCP or severe infection:</u> 5 mg/kg (of trimethoprim component) (IV) q6h <u>Minor infections:</u> 4–6 mg/kg (of trimethoprim component) (IV) q12h
Vancomycin	<u>Bacteremia:</u> 10 mg/kg/dose (IV) <u>Meningitis:</u> 15 mg/kg/dose (IV). Administer IV dose over 60 min. Dosing interval is based on gestational age (GA) and chronological age (CA):			10–20 mg/kg (IV) q6h
	<b>GA + CA (weeks)</b>	<b>CA (days)</b>	<b>Interval (weeks)</b>	
	≤ 29	0–14 > 14	18 12	
	30–36	0–14 > 14	12 8	
	37–44	0–7 > 7	12 8	
Voriconazole	Not applicable			7 mg/kg (IV) q12h 8 mg/kg (PO) q12h × 1 day, then 7 mg/kg (PO) q12h

\* Dosages are generally based on weight (mg/kg), up to adult dose as maximum.

## REFERENCES AND SUGGESTED READINGS

- Betts R, Glasmacher A, Maertens J, et al. Efficacy of caspofungin against invasive *Candida* or invasive *Aspergillus* infections in neutropenic patients. *Cancer* 106:466–73, 2006.
- Bisno AL, Kaplan EL. Strep throat over and over: How frequent? How real? *Mayo Clin Proc* 81:1153–4, 2006.
- CDC. Prevention of measles, rubella, congenital rubella syndrome and mumps 2013. *MMWR* 62:1–34, 2013.
- CDC. Use of rapid influenza diagnostic tests for patients with influenza-like illness during the novel H<sub>1</sub>N<sub>1</sub> influenza virus (Swine Flu) outbreak. *Swine Influenza Bulletin* June 12, 2009.
- Cohen-Wolkowicz M, Poindexter B, Bidegain M, et al. Safety and effectiveness of meropenem in infants with suspected or complicated intra-abdominal infections. *Clin Infect Dis* 55:1495–1502, 2012.
- Committee of Infectious Disease. Prevention of varicella: Recommendations for use of varicella vaccine in children, including a recommendation for a routine 2-dose varicella immunization schedule. *Pediatrics* 120:221–231, 2007.
- Ellis JM, Kuti JL, Nicolau DP. Pharmacodynamic evaluation of meropenem and cefotaxime for pediatric meningitis: a report from the OPTAMA program. *Pediatr Drugs* 8:131–8, 2006.
- Fox JW, Marcon MJ, Bonsu BK. Diagnosis of streptococcal pharyngitis by detection of *Streptococcus pyogenes* in posterior pharyngeal vs oral cavity specimens. *Journal of Clinical Microbiology* 44:2593–94, 2006.
- Groll AH, Attarbasehi A, Schuster FR, et al. Treatment with caspofungin in immunocompromised paediatric patients: a multicentre survey. *Antimicrob Chemother* 57:527–35, 2006.
- Hidron AI, Low CE, Honig EG, et al. Emergence of community-acquired methicillin-resistant *Staphylococcus aureus* strain USA300 as a cause of necrotizing community-onset pneumonia. *Lancet Infect Dis* 9:384–392, 2009.
- Kaplan SL, Afghani B, Lopez P, et al. Linezolid for the treatment of methicillin-resistant *Staphylococcus aureus* infections in children. *Pediatric Infect Disease* 22:S178–185, 2003.
- Lehtinen P, Jartti T, Virkki R, et al. Bacterial coinfections in children with viral wheezing. *Eur J Clin Microbiol Infect Dis* 25:463–9, 2006.
- Lorente L, Lorenzo L, Martin MM, Jimenez A, Mora ML. Meropenem by continuous versus intermittent infusion in ventilator-associated pneumonia due to gram-negative bacilli. *Ann Pharmacother* 40:219–23, 2006.
- Mathys J, DeMeyere M. Acute pharyngitis: no reliability of rapid streptococcal tests and clinical findings. *Arch Intern Med* 166:2285–86, 2006.
- Saiman L, Goldfarb J, Kaplan SL, et al. Safety and tolerability of linezolid in children. *Pediatric Infect Disease* 22:S193–200, 2003.

## GUIDELINES

- ACIP Recommended childhood and adolescent immunization schedule—United States, 2007. *MMWR* 62:2–8, 2013.
- Bradley J, Byington C, Shah S, et al. The management of community-acquired pneumonia in infants and children older than 3 months of age: clinical practice guidelines by the Pediatric Infectious Diseases Society and the Infectious Diseases Society of America. *Clinical Infectious Diseases*. 53:617–630, 2011.
- CDC Prevention and Control of Seasonal Influenza with Vaccines. Recommendations of the Advisory Committee on Immunization Practices (ACIP); *MMWR* 63:961–967, 2014.
- CDC. Antiviral Agents for Treatment and Chemoprophylaxis of Influenza, *MMWR* 60:1–24, 2011.
- CDC. Prevention of perinatal group B streptococcal disease. Revised guidelines from CDC, 2010. *MMWR* 59:1–32, 2010.
- Committee of infectious Disease. Prevention of varicella: Recommendations for use of varicella vaccine in children, including a recommendation for a routine 2-dose varicella immunization schedule. *Pediatrics* 120:221–231, 2007.
- Lieberthal AS, Carroll AE, Clomaitree T, et al. Clinical practice guideline for the diagnosis and management of acute otitis media. *Pediatrics* 131:e964–e999, 2013.
- Solomkin JS, Mazuski JE, Bradley JS, et al. Diagnosis and management of complicated intra-abdominal infection in adults and children: Guidelines by the Surgical Infection Society and the Infectious Diseases Society of America. *Clin Infect Dis* 50:133–164, 2010.
- Tunkel AR, Glaser CA, Block KC, et al. The management of encephalitis: Clinical practice guidelines by the Infectious Diseases Society of America. *Clin Infect Dis* 47:303–327, 2008.

Wald ER, Applegate KE, Bordley C, et al. Clinical Practice Guidelines for the diagnosis and management of acute bacterial sinusitis in children aged 1 to 18 years. *Pediatrics*. 132:e262-e280; 2013.

#### TEXTBOOKS

Bradley JS, Nelson JD, Kimberlin DW, et al (eds). *Nelson's Pediatric Antimicrobial Therapy* (20<sup>th</sup> Ed), American Academy of Pediatrics, Elk Grove Village IL, 2014.

Cherry J, Demmler GJ, Kaplan SL, et al. *Feigin and Cherry's Textbook of Pediatric Infectious Diseases* (7<sup>th</sup> Ed), Saunders, Philadelphia PA, 2010.

Long SS, Pickering LK, Prober CG (eds). *Principles and Practice of Pediatric Infectious Diseases* (4<sup>th</sup> Ed), Saunders, Philadelphia, 2012.

Pickering LK (ed). *Red Book: 2012 Report of the Committee on Infectious Diseases* (29<sup>th</sup> Ed), American Academy of Pediatrics, Elk Grove Village, IL, 2012.

Remington JS, Klein JO, Wilson CB, et al. *Infectious Diseases of the Fetus and Newborn* (7<sup>th</sup> Ed), Saunders, Philadelphia PA, 2010.

Shah S. *Pediatric Infectious Disease Practice*. New York: McGraw-Hill, 2009.

Steele RW. *Clinical Handbook of Pediatric Infectious Disease* (3<sup>rd</sup> Ed), Informa Healthcare, New York, 2007.

## Chapter 8


**CHEST X-RAY ATLAS****Burke A. Cunha, MD****Douglas S. Katz, MD****Robert Moore, MD****Daniel S. Siegal, MD****Chest X-Ray Patterns**

Unilateral Focal Segmental/Lobar Infiltrate Without Effusion .....	426
Unilateral Focal Segmental/Lobar Infiltrate With Effusion .....	431
Unilateral Ill-Defined Infiltrates Without Effusion .....	436
Unilateral Ill-Defined Infiltrates With Effusion .....	443
Bilateral Infiltrates Without Effusion .....	448
Bilateral Infiltrates With Effusion .....	458
Cavitary Infiltrates (Thick-Walled) .....	463
Cavitary Infiltrates (Thin-Walled) .....	470
References and Suggested Readings .....	473

This atlas has been developed to assist in the management of patients who present with respiratory symptoms and chest x-ray abnormalities. Eight common chest x-ray patterns are provided. Common infections and noninfectious etiologies are followed by usual clinical features, which can be used to identify the disorder and help guide empiric/specific therapy.



## UNILATERAL FOCAL SEGMENTAL/LOBAR INFILTRATE WITHOUT EFFUSION



### Infectious Causes

Causes	Features (may have some, none, or all)			
	History	Physical	Laboratory	Chest X-Ray
<i>S. pneumoniae</i>	Elderly, smokers, COPD, ↓ humoral immunity (multiple myeloma, SLE, CLL, hyposplenism).	Fever, chills, no relative bradycardia. Chest signs related to extent of consolidation.	↑ WBC, ↓ platelets (overwhelming infection/hyposplenism), normal LFTs. Sputum with abundant PMNs and gram-positive diplococci. Blood cultures usually positive.	Consolidation usually limited to one lobe (RLL most common) ± air bronchogram. Pleural effusion very uncommon. Empyema not uncommon. No cavitation.
<i>H. influenzae</i>	Recent contact with <i>H. influenzae</i> URI.	Fever, chills.	↑ WBC. Sputum/pleural effusion with gram-negative pleomorphic bacilli. Blood cultures often positive.	Usually RLL with small/moderate effusion. No empyema. No cavitation.

## Infectious Causes (cont'd)

Causes	Features (may have some, none, or all)			
	History	Physical	Laboratory	Chest X-Ray
M. catarrhalis	Chronic/heavy smoker, COPD.	Nonspecific.	Sputum with gram-negative/variable diplococci. Blood cultures negative.	Usually lower lobe ± consolidation. No pleural effusion or cavitation.
K. pneumoniae	Nosocomial pneumonia or history of alcoholism in patient with community-acquired pneumonia.	Fever, chills, signs of alcoholic cirrhosis, signs of consolidation over involved lobe.	↑ WBC, ↓ platelets, ↑ SGOT/SGPT (2° to alcoholism). "Red currant jelly" sputum with PMNs and plump gram-negative encapsulated bacilli.	"Bulging fissure" sign secondary to expanded lobar volume. Empyema rather than pleural effusion. Cavitation in 5–7 days (thick walled).
Legionella	Recent contact with Legionella containing water. Usually elderly. May have watery diarrhea, abdominal pain, mental confusion.	Fever/chills, relative bradycardia. Hepatic/splenic enlargement goes against the diagnosis. ↓ breath sounds if consolidation or pleural effusion.	↑ WBC, ↑ SGOT/SGPT, ↓ PO <sub>2</sub> , ↓ Na <sup>+</sup> , ↑ CPK, ↑ ESR, ↑ CRP, ↑ ferritin levels, microscopic hematuria, L. pneumophila antigenuria (serotypes 01–06 only) may not be positive early. Mucoid/purulent sputum with few PMNs. Sputum culture for Legionella or sputum DFA (before therapy) is diagnostic. ↑ Legionella titer ≥ 1:256 or ≥ 4-fold rise between acute and convalescent titers.	Rapidly progressive asymmetrical infiltrates clue to Legionella. Consolidation and pleural effusion not uncommon. Cavitation rare.

## Infectious Causes (cont'd)

Features (may have some, none, or all)				
Causes	History	Physical	Laboratory	Chest X-Ray
Psittacosis	Recent bird contact with psittacine birds. Severe headache.	Fever/chills, $\pm$ relative bradycardia, Horder's spots on face, epistaxis, $\pm$ splenomegaly. Signs of consolidation common.	$\uparrow$ /normal WBC, $\uparrow$ LFTs. Sputum with few PMNs. $\uparrow$ C. psittaci titers.	Dense infiltrate. Consolidation common. Pleural effusion/cavitation rare.
Q fever	Recent contact with sheep or parturient cats.	Fever/chills, $\pm$ relative bradycardia, splenomegaly, $\pm$ hepatomegaly.	$\uparrow$ /normal WBC, $\uparrow$ LFTs. Sputum with no bacteria/few PMNs (caution—biohazard). Acute Q fever with $\uparrow$ in phase II ELISA antigens.	Dense consolidation. Cavitation/pleural effusion rare.

## Noninfectious Causes

Features (may have some, none, or all)				
Causes	History	Physical	Laboratory	Chest X-Ray
Atelectasis	Ineffectual moist recurrent cough characteristic of post-operative atelectasis.	Fevers $\leq 102^\circ\text{F}$ . If large, signs of volume loss ( $\downarrow$ respiratory excursion, $\uparrow$ diaphragm, mediastinum shift toward affected side). If small, $\downarrow$ breath sounds over affected segment/lobe.	$\uparrow$ WBC (left shift), normal platelets. Other lab results related to underlying cause of atelectasis.	Segmental infiltrate. RUL/RML atelectasis obscures right heart border. In LUL atelectasis, may be triangular infiltrate extending to upper anterior mediastinum mimicking malignancy. LLL atelectasis causes $\uparrow$ density of heart shadow. No cavitation or pleural effusion.


**Noninfectious Causes** (cont'd)

Causes	Features (may have some, none, or all)			
	History	Physical	Laboratory	Chest X-Ray
Pulmonary embolus/infarct	Acute onset dyspnea/pleuritic chest pain. History of lower extremity trauma, stasis or hypercoagulable disorder.	↑ pulse/ respiratory rate.	↑ fibrin split products and D-dimers ± ↑ total bilirubin. Bloody pleural effusion. ECG with RV strain/P-pulmonale (large embolus). Positive V/Q scan and CT pulmonary angiogram.	Normal or show non-specific pleural-based infiltrates resembling atelectasis. Focal segmental/lobar hyperlucency (Westermark's sign) in some. "Hampton's hump" with infarct. Resolving infarcts ↓ in size but maintain shape/density ("melting ice cube").
Lymphoma	Fever, ↓ appetite with weight loss, night sweats, fatigue.	Adenopathy ± splenomegaly.	Normal WBC, ↑ basophils, ↑ eosinophilia, ↓ lymphocytes, ↑ platelets, ↑ ESR, ↑ alkaline phosphatase, ↑ $\alpha_{1,2}$ globulins on SPEP.	Unilateral or asymmetrical bilateral hilar adenopathy. Lung infiltrate may appear contiguous with hilar adenopathy. No clear channel between mediastinum and hilar nodes. Small pleural effusions rare.
Alveolar cell carcinoma	Fever, ↑ appetite with weight loss, night sweats.	± dullness over lobe with large lower lobe lesions.	Positive cytology by BAL/lung biopsy.	Well/ill-defined circumscribed peripheral infiltrates ± air bronchograms. May be multifocal/multilobar. Hilar adenopathy present. Stranding to the hilum ("pleural tail" sign). ± pleural effusion if lower lobe infiltrate. No cavitation.

**Noninfectious Causes** (cont'd)

Causes	Features (may have some, none, or all)			
	History	Physical	Laboratory	Chest X-Ray
Aspiration pneumonia	Swallowing disorder 2° to CNS/GI disorder; impaired consciousness; recent aspiration 2° to dental, upper GI, or pulmonary procedure.	Unremarkable.	↑ WBC, ↑ ESR.	Infiltrate usually involves superior segments of lower lobes (or posterior segments of upper lobes if aspiration occurred supine). Focal infiltrate initially, which may be followed in 7 days by cavitation/lung abscess.

## UNILATERAL FOCAL SEGMENTAL/LOBAR INFILTRATE WITH EFFUSION



### Infectious Causes

Causes	Features (may have some, none, or all)			
	History	Physical	Laboratory	Chest X-Ray
Klebsiella pneumoniae	Nosocomial pneumonia or history of alcoholism in patient with community-acquired pneumonia.	Fever, chills, signs of alcoholic cirrhosis, signs of consolidation over involved lobe.	↑ WBC, ↓ platelets, SGOT/SGPT (2° to alcoholism). "Red currant jelly" sputum with PMNs and plump gram-negative encapsulated bacilli.	"Bulging fissure" sign secondary to expanded lobar volume. Empyema rather than pleural effusion. Usually cavitation in 5–7 days (thick walled).
H. influenzae	Recent contact with H. influenzae URI.	Fever, chills.	↑ WBC. Sputum/pleural effusion with gram-negative pleomorphic bacilli. Blood cultures often positive.	Usually RLL with small/moderate effusion. No empyema. No cavitation.

## Infectious Causes (cont'd)

Causes	Features (may have some, none, or all)			
	History	Physical	Laboratory	Chest X-Ray
TB (primary)	Recent TB contact.	Unilateral lower lobe dullness related to size of pleural effusion.	PPD (-)/anergic. Exudative pleural effusion (pleural fluid with ↑ lymphocytes, ↑ glucose, ± RBCs).	Lower lobe infiltrate with small/moderate pleural effusion. No cavitation or apical infiltrates. Hilar adenopathy asymmetrical when present.
Coccidiomycosis (chronic)	Previous exposure in endemic coccidiomycosis areas. Asymptomatic.	± <i>E. nodosum</i> .	Normal WBC, no eosinophilia in chronic phase. Complement fixation IgG titer ≥ 1:32 indicates active disease.	Thick/thin-walled cavities < 3 cm ± calcifications. Air fluid level rare unless secondarily infected. Cavities usually in anterior segments of upper lobes (vs. posterior segments with TB). No surrounding tissue reaction. Pleural effusion common.
Tularemia	History of recent deer fly, rabbit, or tick exposure. Tularemia pneumonia may complicate any of the clinical presentations of tularemia.	Fever, chills, no relative bradycardia. Chest findings related to extent of infiltrate/consolidation and pleural effusion.	Normal/↑ WBC, normal LFTs. Sputum/pleural effusion with gram-negative coccobacilli (caution-biohazard). Bloody pleural effusion. Tularemia serology with ↑ microagglutination titer ≥ 1:160 acutely and ≥ 4-fold rise between acute and convalescent titers.	Pleural effusion ± hilar adenopathy.

## Infectious Causes (cont'd)

Causes	Features (may have some, none, or all)			
	History	Physical	Laboratory	Chest X-Ray
Adenovirus	Recent URI.	Fever, chills, myalgias. Sore throat and conjunctivitis not always present with adenoviral pneumonia. Chest exam $\pm$ signs of consolidation.	$\uparrow$ adenoviral titers and positive adenoviral cultures of respiratory secretions. Respiratory viral FA panel + for adenovirus.	Ill-defined infiltrate(s) without cavitation $\pm$ pleural effusion.
Legionella	Recent contact with Legionella containing water. Usually elderly. May have watery diarrhea, abdominal pain, mental confusion.	Fever/chills, relative bradycardia. Hepatic/splenic enlargement goes against the diagnosis. $\downarrow$ breath sounds if consolidation or pleural effusion.	$\uparrow$ WBC, $\uparrow$ SGOT/SGPT, $\downarrow$ $PO_4^-$ , $\downarrow$ $Na^+$ , $\uparrow$ CPK, $\uparrow$ ESR, $\uparrow$ CRP, $\uparrow$ ferritin levels, microscopic hematuria, L. pneumophila antigenuria (serotypes 01–06 only) may not be positive early. Mucoïd/purulent sputum with few PMNs. Sputum culture for Legionella or sputum DFA (before therapy) is diagnostic. $\uparrow$ Legionella titer $\geq$ 1:256 or $\geq$ 4-fold rise between acute/convalescent titers.	Rapidly progressive asymmetrical infiltrates clue to Legionella. Consolidation and pleural effusion not uncommon. Cavitation rare.
Group A streptococci	Recent exposure or recent blunt chest trauma $\pm$ chest pain.	Fever/chills. Physical signs related to size of pleural effusion.	$\uparrow$ WBC. Pleural fluid is serosanguineous. Sputum/pleural fluid with gram-positive cocci in pairs/chains. Positive pleural fluid/blood cultures.	Unilateral infiltrate may be obscured by large pleural effusion. No empyema. No cavitation.



**Infectious Causes** (cont'd)

Causes	Features (may have some, none, or all)			
	History	Physical	Laboratory	Chest X-Ray
Rhodococcus equi	Insidious onset of fever, dyspnea, chest pain, $\pm$ hemoptysis. Immunosuppressed patients with $\downarrow$ cell-mediated immunity or exposure to cattle, horses, pigs.	Unremarkable.	Normal/ $\uparrow$ WBC. Sputum/pleural fluid with gram-positive pleomorphic weakly acid-fast bacilli. Sputum, pleural fluid, blood cultures positive for R. equi.	Segmental infiltrate with upper lobe predominance $\pm$ cavitation. Air-fluid levels and pleural effusion common.


**Noninfectious Causes**

Causes	Features (may have some, none, or all)			
	History	Physical	Laboratory	Chest X-Ray
Pulmonary embolus/infarct	Acute onset dyspnea/pleuritic chest pain. History of lower extremity trauma, stasis or hypercoagulable disorder.	$\uparrow$ pulse/respiratory rate.	$\uparrow$ fibrin split products and D-dimers $\pm$ $\uparrow$ total bilirubin. Bloody pleural effusion. ECG with RV strain/P-pulmonale (large embolus). Positive V/Q scan and CT pulmonary angiogram.	Normal or show non-specific pleural-based infiltrates resembling atelectasis. Focal segmental/lobar hyperlucency (Westermark's sign) in some. "Hampton's hump" with infarct. Resolving infarcts $\downarrow$ in size but maintain shape/density ("melting ice cube").

**Noninfectious Causes** (cont'd)

Causes	Features (may have some, none, or all)			
	History	Physical	Laboratory	Chest X-Ray
Lymphoma	Fever, ↓ appetite with weight loss, night sweats, fatigue.	Adenopathy ± splenomegaly.	Normal WBC, ↑ basophils, ↑ eosinophilia, ↓ lymphocytes, ↑ platelets, ↑ ESR, ↑ alkaline phosphatase, ↑ $\alpha_{1,2}$ globulins on SPEP.	Unilateral or asymmetrical bilateral hilar adenopathy. Lung infiltrate may appear contiguous with hilar adenopathy. No clear channel between mediastinum and hilar nodes. Small pleural effusions rare.
Alveolar cell carcinoma	Fever, ↑ appetite with weight loss, night sweats.	± dullness over lobe with large lower lobe lesions.	Positive cytology by BAL/lung biopsy.	Well/ill-defined circumscribed peripheral infiltrates ± air bronchograms. May be multifocal/multilobar. Hilar adenopathy present. Stranding to the hilum ("pleural tail" sign). ± pleural effusion if lower lobe infiltrate. No cavitation.
Radiation pneumonitis	History of mantle radiation for lymphoma, lung cancer, or breast cancer.	Nonspecific.	Nonspecific.	Symmetrical infiltrates in the distribution of radiation therapy after 1 month. Infiltrates have "straight edges" ± air bronchograms. Fibrosis common over radiation field after 9–12 months. Usually no pleural effusions (small, if present).

## UNILATERAL ILL-DEFINED INFILTRATES WITHOUT EFFUSION



### Infectious Causes

Causes	Features (may have some, none, or all)			
	History	Physical	Laboratory	Chest X-Ray
Mycoplasma pneumoniae	Prolonged dry/nonproductive cough. No laryngitis. Mild sore throat/ear. Watery diarrhea.	Usually fevers $\leq 102^{\circ}\text{F}$ without relative bradycardia. Myalgias, bullous myringitis or otitis, non-exudative pharyngitis, E. multiforme. Chest exam with rales and no signs of consolidation or effusion.	Normal/ $\uparrow$ WBC; normal platelets, LFTs, $\text{PO}_4$ , CPK. $\uparrow$ cold agglutinins (early). $\uparrow$ IgM (not IgG) Mycoplasma pneumoniae titers. Respiratory secretions culture positive for Mycoplasma pneumoniae.	Ill-defined usually lower lobe indistinct infiltrates. No consolidation or air bronchograms. Small/no pleural effusion.

## Infectious Causes (cont'd)

Causes	Features (may have some, none, or all)			
	History	Physical	Laboratory	Chest X-Ray
Chlamydophilia (Chlamydia) pneumoniae	Prolonged "mycoplasma-like" illness with laryngitis.	Low-grade fevers, myalgias, non-exudate pharyngitis, laryngitis. No relative bradycardia, ear findings, or rash. Chest exam without signs of consolidation or pleural effusion.	↑ WBC, normal platelets, normal LFTs. No cold agglutinins. ↑ IgM (not IgG) C. pneumoniae titers. Respiratory secretions culture positive for C. pneumoniae.	Ill-defined, usually lower lobe indistinct infiltrate(s). May be "funnel shaped." No consolidation, cavitation, or pleural effusion.
Adenovirus	Recent URI.	Fever, chills, myalgias. Sore throat and conjunctivitis not always present with adenoviral pneumonia. Chest exam ± signs of consolidation.	↑ adenoviral titers and positive adenoviral cultures of respiratory secretions. Respiratory viral FA panel + for adenovirus.	Ill-defined infiltrate(s) without cavitation ± pleural effusion. Only viral cause of focal lobar pneumonia.
Legionella	Recent contact with Legionella containing water. Usually elderly. May have watery diarrhea, abdominal pain, mental confusion.	Fever/chills, relative bradycardia. Hepatic/splenic enlargement goes against the diagnosis. ↓ breath sounds if consolidation or pleural effusion.	↑ WBC, ↑ SGOT/SGPT, ↓ PO <sub>4</sub> <sup>-3</sup> , ↓ Na <sup>+</sup> , ↑ CPK, ↑ ESR, ↑ CRP, ↑ ferritin levels, microscopic hematuria, L. pneumophila antigenuria (serotypes 01–06 only) may not be positive early. Mucoid/purulent sputum with few PMNs. Sputum culture for Legionella or sputum DFA (before therapy) is diagnostic. ↑ Legionella titer ≥ 1:256 or ≥ 4-fold rise between acute/convalescent titers.	Rapidly progressive asymmetrical infiltrates clue to Legionella. Consolidation and pleural effusion not uncommon. Cavitation rare.

## Infectious Causes (cont'd)

Causes	Features (may have some, none, or all)			
	History	Physical	Laboratory	Chest X-Ray
Psittacosis	Recent bird contact with psittacine birds. Severe headache.	Fever/chills, ± relative bradycardia, ± Horder's spots on face, epistaxis, ± splenomegaly. Signs of consolidation common.	↑/normal WBC, ↑ LFTs. Sputum with few PMNs. ↑ C. psittaci titers.	Dense infiltrate. Consolidation common. Pleural effusion/cavitation rare.
Q fever	Recent contact with sheep or parturient cats.	Fever/chills, ± relative bradycardia, splenomegaly, ± hepatomegaly.	↑/normal WBC, ↑ LFTs. Sputum with no bacteria/few PMNs (caution—biohazard). Acute Q fever with ↑ in phase II ELISA antigens.	Dense consolidation. Cavitation/pleural effusion rare.
Nocardia	Fevers, night sweats, fatigue, ↓ cell-mediated immunity (e.g., HIV, organ transplants, PAP, immunosuppressive therapy).	Unremarkable.	Normal/↑ WBC, ↑ ESR. Sputum with gram-positive AFB.	Dense large infiltrates. May cavitate and mimic TB, lymphoma, or squamous cell carcinoma. No calcification ± pleural effusion.
Actinomycosis	Recent dental work.	± chest wall sinus tracts.	Normal/↑ WBC, ↑ ESR. Sputum with gram-positive filamentous anaerobic bacilli.	Dense infiltrates extending to chest wall. No hilar adenopathy. Cavitation rare. ± pleural effusion rare.

**Infectious Causes** (cont'd)

Causes	Features (may have some, none, or all)			
	History	Physical	Laboratory	Chest X-Ray
Cryptococcus neoformans	Exposure to air conditioner or pigeons.	Unremarkable.	Normal WBC. Cryptococcal serology with ↑ <i>C. neoformans</i> antigen levels.	Dense lower nodular mass lesions. No calcification or cavitation. ± pleural effusion.
Aspiration pneumonia	Swallowing disorder 2° to CNS/GI disorder; impaired consciousness; recent aspiration 2° to dental, upper GI, or pulmonary procedure.	Unremarkable.	↑ WBC, ↑ ESR.	Infiltrate usually involves superior segments of lower lobes (or posterior segments of upper lobes if aspiration occurred supine). Focal infiltrate initially, which may be followed in 7 days by cavitation/lung abscess.

## Noninfectious Causes

Causes	Features (may have some, none, or all)			
	History	Physical	Laboratory	Chest X-Ray
Bronchogenic carcinoma	Fever, ↓ appetite with weight loss, cough ± hemoptysis. Cough with copious clear/mucoid sputum in large cell anaplastic carcinoma. Increased risk in smokers, aluminum/uranium miners, cavitary lung disease (adenocarcinoma), previous radiation therapy from lymphoma/breast cancer.	Paraneoplastic syndromes especially with small (oat) cell/squamous cell carcinoma ± hypertrophic pulmonary osteoarthropathy.	Normal/↑ WBC, ↑ platelets, ↑ ESR, findings related to underlying malignancy, ± clubbing.	Small/squamous cell carcinomas present as central lesions/hilar masses. Adenocarcinoma/large cell anaplastic carcinomas are usually peripheral initially. "Tumor tendrils" extending into surrounding lung tissue is characteristic. No calcifications (may be present on chest CT). Cavitation with squamous cell carcinoma. ± pleural effusions.
Lymphangitic metastases	History of breast, thyroid, pancreas, cervical, prostate, or lung carcinoma.	Findings related to underlying malignancy.	Normal/↑ WBC, ↑ ESR.	Interstitial indistinct pulmonary infiltrates (may be reticulonodular) with lower lobe predominance. Usually unilateral but may be bilateral. No consolidation or cavitation ± pleural effusions.
Lung contusion	Recent closed chest trauma, chest pain.	Chest wall contusion over infiltrate.	↑ WBC (left shift).	Patchy ill-defined infiltrate(s) ± rib fractures/pneumothorax in area of infiltrate. Infiltrate clears within 1 week.

## Noninfectious Causes (cont'd)


Causes	Features (may have some, none, or all)			
	History	Physical	Laboratory	Chest X-Ray
Congestive heart failure	Coronary heart disease, valvular heart disease, cardiomyopathy.	No/low-grade fevers ↑ pulse and respiratory rate, positive jugular venous distension and hepatojugular reflex, cardiomegaly, $S_3$ , ascites, hepatomegaly, pedal edema.	↑ WBC (left shift), normal platelets, mildly ↑ SGOT/SGPT.	Cardiomegaly, pleural effusion ( $R > [R + L] > L$ ). Kerley B lines with vascular redistribution to upper lobes. Typically bilateral rather than unilateral.
Alveolar cell carcinoma	Fever, ↑ appetite with weight loss, night sweats.	± dullness over lobe with large lower lobe lesions.	Positive cytology by BAL/lung biopsy.	Well/ill-defined circumscribed peripheral infiltrates ± air bronchograms. May be multifocal/multilobar. Hilar adenopathy present. Stranding to the hilum ("pleural tail" sign). ± pleural effusion if lower lobe infiltrate. No cavitation.
Lymphoma	Fever, ↓ appetite with weight loss, night sweats, fatigue.	Adenopathy ± splenomegaly.	Normal WBC, ↑ basophils, ↑ eosinophilia, ↓ lymphocytes, ↑ platelets, ↑ ESR, ↑ alkaline phosphatase, ↑ $\alpha_{1,2}$ globulins on SPEP.	Unilateral or asymmetrical bilateral hilar adenopathy. Lung infiltrate may appear contiguous with hilar adenopathy. No clear channel between mediastinum and hilar nodes. Small pleural effusions rare.



**Noninfectious Causes** (cont'd)

Causes	Features (may have some, none, or all)			
	History	Physical	Laboratory	Chest X-Ray
Pulmonary hemorrhage	History of closed chest trauma or hemorrhagic disorder.	↑ WBC (left shift), ↑ pulse rate, respiratory rate. Signs of closed chest trauma.	Anemia plus findings secondary to underlying hemorrhagic disorder.	Localized or diffuse fluffy alveolar infiltrate(s). No cavitation, consolidation, or effusion.
Systemic lupus erythematosus (SLE)	Fatigue, chest pain. History of SLE.	Fever/myalgias, alopecia, malar rash, "cytoid bodies" in retina, painless oral ulcers, synovitis, splenomegaly, generalized adenopathy, Raynaud's phenomenon.	↑ ANA, ↑ DS-DNA, ↓ C <sub>3</sub> , polyclonal gammopathy on SPEP, ↑ ferritin. Pleural fluids with ↑ ANA, ↓ C <sub>3</sub> .	Migratory ill-defined non-segmental infiltrates ± small pleural effusions. No consolidation or cavitation.
Aspiration pneumonia	Swallowing disorder 2° to CNS/GI disorder; impaired consciousness; recent aspiration 2° to dental, upper GI, or pulmonary procedure.	Unremarkable.	↑ WBC, ↑ ESR.	Infiltrate usually involves superior segments of lower lobes (or posterior segments of upper lobes if aspiration occurred supine). Focal infiltrate initially, which may be followed in 7 days by cavitation/lung abscess.

## UNILATERAL ILL-DEFINED INFILTRATES WITH EFFUSION



### Infectious Causes

Causes	Features (may have some, none, or all)			
	History	Physical	Laboratory	Chest X-Ray
TB (primary)	Recent TB contact.	Unilateral lower lobe dullness related to size of pleural effusion.	PPD anergic. Exudative pleural effusion (pleural fluid with ↑ lymphocytes, ↑ glucose, ± RBCs).	Lower lobe infiltrate with small/moderate pleural effusion. No cavitation or apical infiltrates. Hilar adenopathy asymmetrical when present.
Nocardia	Fevers, night sweats, fatigue, ↓ cell-mediated immunity (e.g., HIV, organ transplant, PAP, immuno-suppressive therapy).	Unremarkable.	Normal/↑ WBC, ↑ ESR. Sputum with gram-positive AFB.	Dense large infiltrates. May cavitate and mimic TB, lymphoma, or squamous cell carcinoma. No calcification ± pleural effusion.

## Infectious Causes (cont'd)

Causes	Features (may have some, none, or all)			
	History	Physical	Laboratory	Chest X-Ray
Legionella	Recent contact with Legionella containing water. Usually elderly. May have watery diarrhea, abdominal pain, mental confusion.	Fever/chills, relative bradycardia. Hepatic/splenic enlargement goes against the diagnosis. ↓ breath sounds if consolidation or pleural effusion.	↑ WBC, ↑ SGOT/SGPT, ↓ PO <sub>4</sub> <sup>=</sup> , ↓ Na <sup>+</sup> , ↑ CPK, ↑ ESR, ↑ CRP, ↑ ferritin levels, microscopic hematuria, L. pneumophila antigenuria (serotypes 01–06 only) may not be positive early. Mucoid/purulent sputum with few PMNs. Sputum culture for Legionella or sputum DFA (before therapy) is diagnostic. ↑ Legionella titer ≥ 1:256 or ≥ 4-fold rise between acute/convalescent titers.	Rapidly progressive asymmetrical infiltrates clue to Legionella. Consolidation and pleural effusion not uncommon. Cavitation rare.

## Noninfectious Causes

Causes	Features (may have some, none, or all)			
	History	Physical	Laboratory	Chest X-Ray
Lymphangitic metastases	History of breast, thyroid, pancreas, cervical, prostate, or lung carcinoma.	Findings related to underlying malignancy.	Normal/↑ WBC, ↑ ESR.	Interstitial indistinct pulmonary infiltrates (may be reticulonodular) with lower lobe predominance. Usually unilateral but may be bilateral. No consolidation or cavitation ± pleural effusions.

**Noninfectious Causes** (cont'd)

Causes	Features (may have some, none, or all)			
	History	Physical	Laboratory	Chest X-Ray
Pulmonary embolus/infarct	Acute onset dyspnea/pleuritic chest pain. History of lower extremity trauma, stasis or hypercoagulable disorder.	↑ pulse/ respiratory rate.	↑ fibrin split products and D-dimers ± ↑ total bilirubin. Bloody pleural effusion. ECG with RV strain/P-pulmonale (large embolus). Positive V/Q scan and CT pulmonary angiogram.	Normal or show non-specific pleural-based infiltrates resembling atelectasis. Focal segmental/lobar hyperlucency (Westermark's sign) in some. "Hampton's hump" with infarct. Resolving infarcts ↓ in size but maintain shape/density ("melting ice cube").
Lymphoma	Fever, ↓ appetite with weight loss, night sweats, fatigue.	Adenopathy ± splenomegaly.	Normal WBC, ↑ basophils, ↑ eosinophilia, ↓ lymphocytes, ↑ platelets, ↑ ESR, ↑ alkaline phosphatase, ↑ $\alpha_{1,2}$ globulins on SPEP.	Unilateral or asymmetrical bilateral hilar adenopathy. Lung infiltrate may appear contiguous with hilar adenopathy. No clear channel between mediastinum and hilar nodes. Small pleural effusions rare.


## Noninfectious Causes (cont'd)

Causes	Features (may have some, none, or all)			
	History	Physical	Laboratory	Chest X-Ray
Bronchogenic carcinoma	Fever, ↓ appetite with weight loss ± hemoptysis. Cough with copious clear/mucoid sputum in large cell anaplastic carcinoma. Increased risk in smokers, aluminum/uranium miners, cavitory lung disease (adenocarcinoma), previous radiation therapy from lymphoma/breast cancer.	Paraneoplastic syndromes especially with small (oat) cell/squamous cell carcinoma ± hypertrophic pulmonary osteoarthropathy.	Normal/↑ WBC, ↑ platelets, ↑ ESR, findings related to underlying malignancy, ± clubbing.	Small/squamous cell carcinomas present as central lesions/hilar masses. Adenocarcinoma/large cell anaplastic carcinomas are usually peripheral initially. "Tumor tendrils" extending into surrounding lung tissue is characteristic. No calcifications (may be present on chest CT). Cavitation with squamous cell carcinoma. ± pleural effusions.
Alveolar cell carcinoma	Fever, ↑ appetite with weight loss, night sweats.	± dullness over lobe with large lower lobe lesions.	Positive cytology by BAL/lung biopsy.	Well/ill-defined circumscribed peripheral infiltrates ± air bronchograms. May be multifocal/multilobar. Hilar adenopathy present. Stranding to the hilum ("pleural tail" sign). ± pleural effusion if lower lobe infiltrate. No cavitation.

**Noninfectious Causes** (cont'd)

<b>Causes</b>	<b>Features (may have some, none, or all)</b>			
	<b>History</b>	<b>Physical</b>	<b>Laboratory</b>	<b>Chest X-Ray</b>
Metastatic carcinoma	History of breast, thyroid, renal cell, colon, pancreatic cancer or osteogenic sarcoma.	Findings related to underlying malignancy and, when present, to bone, hepatic, CNS metastases.	Secondary to effects of primary neoplasm, metastases, paraneoplastic syndrome.	Nodular lesions that vary in size. Metastatic lesions are usually well circumscribed with lower lobe predominance. Usually no bronchial obstruction (obstruction suggests colon, renal, or melanoma metastases). Usually no cavitation (except for squamous cell metastases). Calcification usually suggests osteosarcoma (rarely adenocarcinoma). Pleural effusion rare (except for breast cancer).

## BILATERAL INFILTRATES WITHOUT EFFUSION



### Infectious Causes

Causes	Features (may have some, none, or all)			
	History	Physical	Laboratory	Chest X-Ray
Influenza (human, avian, swine)	Acute onset of fever, myalgias, headache, fatigue, sore throat, rhinorrhea, dry cough, ± pleuritic chest pain.	↑ respiratory rate, cyanosis in severe cases.	↓ WBC, ↓ platelets, few/no atypical lymphocytes, ↑ A-a gradient. Influenza in respiratory secretions by culture/viral FA panel.	Early normal/near normal appearance. (< 48 hours) Later (> 48 hours) diffuse patchy bilateral interstitial infiltrates. No focal/segmental infiltrates unless secondary bacterial pneumonia present. ± small pleural effusions.

## Infectious Causes (cont'd)

Causes	Features (may have some, none, or all)			
	History	Physical	Laboratory	Chest X-Ray
MERS	Acute onset of fever/ chills, headache, myalgias, dry cough $\pm$ N/V/D  Malaise; Hemoptysis	Early fulminant respiratory failure.	(N) WBC lymphopenia/ lymphocytosis, $\uparrow$ LDH $\uparrow$ LFTs MERS-Cov in lower respiratory secretions.	Unilateral/bilateral interstitial, nodule dense infiltrate.  Consolidation common/ $\pm$ pleural effusions.
SARS	Acute onset of fever/ chills, headache myalgias, dry cough $\pm$ N/V/D.  Biphasic illness	$\uparrow$ respiratory rate, SOB in severe cases.	N/ $\downarrow$ WBC, lymphopenia, $\downarrow$ platelets, $\uparrow$ LDH, $\uparrow$ LFTs, $\downarrow$ pO <sub>2</sub> , $\uparrow$ A-a gradient. SARS – CoV in respiratory secretions.	Bilateral symmetrical patchy infiltrates. Consolidation uncommon. No cavitation or effusion.
HSV-1	Fever. Often presents in normal hosts as “failure to wean” from ventilator.	Unremarkable.	$\uparrow$ WBC, $\downarrow$ pO <sub>2</sub> , $\uparrow$ A-a gradient. Cytology + for cytopathic effects (CPE) of HSV.	Minimal bilateral diffuse infiltrates without cavitation or effusion.
RSV	Recent URI contact, dry cough, wheezing.	Mild lower respiratory tract infection in normal host. Moderate/severe pneumonia in organ transplants.	Normal WBC; $\downarrow$ pO <sub>2</sub> / $\uparrow$ A-a gradient in severe RSV. RSV in respiratory secretions by culture/viral FA panel.	Near normal chest x-ray or bilateral symmetrical patchy infiltrates. Consolidation uncommon. No cavitation or effusion.
VZV	VZV pneumonia occurs 2–3 days after rash. $\uparrow$ risk with pregnancy, smoking.	Healing vesicles, dry cough. Mild pneumonia in normal hosts. Moderate/ severe pneumonia in organ transplants.	$\uparrow$ WBC, $\uparrow$ platelets, $\uparrow$ basophils; $\downarrow$ pO <sub>2</sub> / $\uparrow$ A-a gradient in severe chickenpox pneumonia. $\uparrow$ VZV titers.	Minimal diffuse fluffy interstitial infiltrates. Diffuse small calcifications may develop years later. No calcification of hilar nodes (in contrast to TB/ histoplasmosis).



## Infectious Causes (cont'd)

Causes	Features (may have some, none, or all)			
	History	Physical	Laboratory	Chest X-Ray
CMV	↓ cell-mediated immunity (HIV, organ transplants, immunosuppressive therapy). Increasing dyspnea over 1 week.	Fever $\leq 102^{\circ}\text{F}$ .	↓ $\text{pO}_2$ , ↑ A-a gradient, ↑ LDH (PCP). PCP cysts or CMV "Cowdry Owl eye" inclusion bodies in respiratory secretions or transbronchial/open lung biopsy.	Most HIV patients with PCP also have underlying CMV. Organ transplants with CMV usually do not have underlying PCP.
P (carinii) jiroveci (PCP)	↓ cell-mediated immunity (e.g., HIV, immuno-suppressive therapy). Increasing dyspnea over 1 week. Chest pain with shortness of breath suggests pneumothorax.	↓ breath sounds bilaterally. Other findings depend on size/location of pneumothorax (if present).	Normal/↑ WBC (left shift), ↑ lymphocytes, ↑ LDH. ↓ $\text{pO}_2$ , ↓ $\text{DL}_{\text{CO}}$ , ↑ A-a gradient. PCP cysts in sputum/respiratory secretions.	Bilateral perihilar symmetrical patchy infiltrates ± pneumothorax. No calcification, cavitation, or pleural effusion.
TB (reactivation)	Fevers, night sweats, normal appetite with weight loss, cough ± hemoptysis.	± bilateral apical dullness.	Normal WBC, ↑ platelets, ↑ ESR ( $\leq 70$ mm/h). Positive PPD. AFB in sputum smear/culture.	Slowly progressive bilateral infiltrates. No pleural effusion. Usually in apical segment of lower lobes or apical/posterior segments of upper lobes. Calcifications common.

## Infectious Causes (cont'd)

Causes	Features (may have some, none, or all)			
	History	Physical	Laboratory	Chest X-Ray
Legionella	Recent contact with Legionella containing water. Usually elderly. May have watery diarrhea, abdominal pain, mental confusion.	Fever/chills, relative bradycardia. Hepatic/splenic enlargement goes against the diagnosis. ↓ breath sounds if consolidation or pleural effusion.	↑ WBC, ↑ SGOT/SGPT, ↑ pO <sub>4</sub> , ↓ Na <sup>+</sup> , ↑ CPK, ↑ ESR, ↑ CRP, ↑ ferritin levels, microscopic hematuria, L. pneumophila antigenuria (serotypes 01–06 only) may not be positive early. Mucoid/purulent sputum with few PMNs. Sputum culture for Legionella or sputum DFA (before therapy) is diagnostic. ↑ Legionella titer ≥ 1:256 or ≥ 4-fold rise between acute/convalescent titers.	Rapidly progressive asymmetrical infiltrates clue to Legionella. Consolidation and pleural effusion not uncommon. Cavitation rare.
Psittacosis	Recent bird contact with psittacine birds. Severe headache.	Fever/chills, ± relative bradycardia, ± Horder's spots on face, epistaxis, ± splenomegaly. Signs of consolidation common.	↑/normal WBC, ↑ LFTs. Sputum with few PMNs. Positive C. psittaci serology.	Dense infiltrate. Consolidation common. Pleural effusion/cavitation rare.

## Infectious Causes (cont'd)

Causes	Features (may have some, none, or all)			
	History	Physical	Laboratory	Chest X-Ray
Q fever	Recent contact with sheep or parturient cats.	Fever/chills, ± relative bradycardia, splenomegaly, ± hepatomegaly.	↑/normal WBC, ↑ LFTs. Sputum with no bacteria/few PMNs (caution–biohazard). Acute Q fever with ↑ in phase II ELISA antigens.	Dense consolidation. Cavitation/pleural effusion rare.
Nosocomial pneumonia (hema-togenous)	Fever/pulmonary symptoms ≥ 7 days in hospital. Increased risk with antecedent heart failure in previous 1–2 weeks.	Bilateral rales ± purulent respiratory secretions (tracheo-bronchitis).	↑ WBC (left shift). Blood cultures positive for pulmonary pathogens. Respiratory secretions with WBCs ± cultures positive for <i>P. aeruginosa</i> , <i>Acinetobacter</i> , <i>Enterobacter</i> , <i>Klebsiella</i> , or <i>Serratia</i> (represents colonization, <i>not</i> lower respiratory tract/lung pathogens). Definitive diagnosis by lung biopsy/culture.	Bilateral symmetrical diffuse infiltrates. May be focal/ segmental in aspiration nosocomial pneumonia. ↑ lung volumes (vs. ARDS). <i>Klebsiella</i> cavitation in 3–5 days; <i>P. aeruginosa</i> cavitation in 72 hours. No pleural effusion.

## Noninfectious Causes

Causes	Features (may have some, none, or all)			
	History	Physical	Laboratory	Chest X-Ray
Adult respiratory distress syndrome (ARDS)	Intubated on ventilator, multi-organ system failure.	± rales.	↑ WBC (left shift), normal ESR, ↓ pO <sub>2</sub> , ↓ D <sub>L</sub> CO, ↑ A-a gradient.	Bilateral fluffy infiltrates appearing ≥ 12 hours after profound hypoxemia. No cardiomegaly or pleural effusion. Reduced lung volumes (vs. nosocomial pneumonia or CHF). Bilateral consolidation ≥ 48 hours after appearance of infiltrates.
Goodpasture's syndrome	Often preceded by a URI. Most common in 20–30 year old adults. Fever, weight loss, fatigue, cough, hemoptysis, hematuria.	Findings secondary to iron deficiency anemia.	↑ WBC, anemia, ↑ creatinine, urine with RBCs/RBC casts. Positive pANCA. Linear IgG pattern on alveolar/glomerular basement membrane.	Bilateral fine reticulonodular infiltrates predominantly in lower lobes. No cavitation.
Wegener's granulomatosis	Most common in middle-aged adults. Cough, fever, fatigue.	Findings of chronic sinusitis, bloody nasal discharge.	↑ WBC, anemia, ↑ platelets, ↑ ESR, ↑ RF, negative ANA, proteinuria, hematuria. Positive cANCA.	Bilateral asymmetrical nodular infiltrates of varying size with irregular margins. Cavitation common. Inner lining of cavities irregular. Air-fluid levels rare. ± pleural effusions. No calcifications.
Pulmonary hemorrhage	History of closed chest trauma or hemorrhagic disorder.	↑ WBC (left shift), ↑ pulse rate, ↑ respiratory rate. Signs of closed chest trauma.	Anemia plus findings secondary to underlying hemorrhagic disorder.	Localized or diffuse fluffy alveolar infiltrates. No cavitation, consolidation, or effusion.

## Noninfectious Causes (cont'd)

Causes	Features (may have some, none, or all)			
	History	Physical	Laboratory	Chest X-Ray
Chronic renal failure	Chronic renal failure on dialysis.	Findings related to uremia.	Normal/↑ WBC (left shift) plus findings related to renal failure.	Bilateral symmetrical fluffy perihilar infiltrates (butterfly pattern) ± pleural effusions. No cardiomegaly (unlike CHF), but large pericardial effusion can mimic cardiomegaly.
Lung contusion	Recent closed chest trauma, chest pain.	Chest wall contusion over infiltrate.	↑ WBC (left shift).	Patchy ill-defined infiltrate(s) ± rib fractures/pneumothorax in area of infiltrate. Infiltrate clears within 1 week.
Fat emboli	1–2 days post-long bone fracture/ trauma.	↑ respiratory rate.	Urinalysis with “Maltese crosses.”	Bilateral predominantly peripheral lower lobe infiltrates. Usually clears within 1 week.
Loeffler’s syndrome	Drug or parasitic exposure.	Unremarkable.	Normal WBC, ↑ eosinophilia, ↑ ESR.	Characteristic “reversed bat-wing” pattern (i.e., peripheral infiltrates). Upper lobe predominance.
Sarcoidosis (Stage III)	Dyspnea, fatigue, nasal stuffiness.	Waxy/yellowish papules on face/ upper trunk. Funduscopic exam with “candle wax drippings.”	↑ ESR, normal LFTs, ↑ creatinine (if renal involvement), ↑ ACE levels, hypercalciuria, hypercalcemia, polyclonal gammopathy on SPEP. Anergic.	Bilateral nodular infiltrates of variable size without hilar adenopathy. Cavitation/ pleural effusion rare.

**Noninfectious Causes** (cont'd)

Causes	Features (may have some, none, or all)			
	History	Physical	Laboratory	Chest X-Ray
Alveolar cell carcinoma	Fever, ↑ appetite with weight loss, night sweats.	± dullness over lobe with large lower lobe lesions.	Positive cytology by BAL/lung biopsy.	Well/ill-defined circumscribed peripheral infiltrates ± air bronchograms. May be multifocal/multilobar. Hilar adenopathy present. Stranding to the hilum ("pleural tail" sign). ± pleural effusion if lower lobe infiltrate. No cavitation.
Metastatic carcinoma	History of breast, thyroid, renal cell, colon, pancreatic cancer or osteogenic sarcoma.	Findings related to underlying malignancy and, when present, to bone, hepatic, CNS metastases.	Secondary to effects of primary neoplasm, metastases, paraneoplastic syndrome.	Nodular lesions that vary in size. Metastatic lesions are usually well circumscribed with lower lobe predominance. Usually no bronchial obstruction (obstruction suggests colon, renal, or melanoma metastases). Usually no cavitation (except for squamous cell metastases). Calcification usually suggests osteosarcoma (rarely adenocarcinoma). Pleural effusion rare (except for breast cancer).
Lymphoma	Fever, ↓ appetite with weight loss, night sweats, fatigue.	Adenopathy ± splenomegaly.	Normal WBC, ↑ basophils, ↑ eosinophilia, ↓ lymphocytes, ↑ platelets, ↑ ESR, ↑ alkaline phosphatase, ↑ $\alpha_{1,2}$ globulins on SPEP.	Unilateral or asymmetrical bilateral hilar adenopathy. Lung infiltrate may appear contiguous with hilar adenopathy. No clear channel between mediastinum and hilar nodes. Small pleural effusions rare.

## Noninfectious Causes (cont'd)


Causes	Features (may have some, none, or all)			
	History	Physical	Laboratory	Chest X-Ray
Leukostasis (AML)	Untreated acute myelogenous leukemia (AML).	Fever, sternal tenderness, petechiae, ecchymosis.	↑ WBC ( $\geq 100 \text{ K/mm}^3$ ) with blasts in peripheral smear/bone marrow, ↓ platelets.	Diffuse symmetrical fluffy infiltrates without pleural effusion.
Drug-induced	Exposure to chemotherapeutic agents (e.g., BCNU, busulfan, methotrexate, cyclophosphamide, bleomycin) or other drugs (e.g., nitro-furantoin, sulfasalazine, amiodarone, opiates, cocaine).	Unremarkable.	Normal WBC $\pm$ ↑ eosinophilia, normal/↑ ESR/LFTs. Eosinophils in pleural effusion.	Bilateral coarse symmetrical patchy infiltrates/fibrosis $\pm$ pleural effusions. Hilar adenopathy only with drug-induced pseudolymphoma (secondary to dilantin). No cavitation.
Idiopathic pulmonary hemosiderosis (IPH)	Hemoptysis $\pm$ cough.	Findings of iron deficiency anemia.	Iron deficiency anemia. Hemosiderin in alveolar macrophages and urine.	Diffuse, bilateral ill-defined opacities or multiple "stellate" shaped infiltrates that clear between attacks. Recent hemorrhage may be superimposed on a fine reticular pattern that occurs after repeated bleeds.

**Noninfectious Causes** (cont'd)

Causes	Features (may have some, none, or all)			
	History	Physical	Laboratory	Chest X-Ray
Bronchiolitis obliterans with organizing pneumonia (BOOP)	Fever, dyspnea, cough.	Unremarkable.	↑ WBC (left shift), ↑ LDH, ↓ pO <sub>2</sub> , ↑ A-a gradient.	Classically bilateral patchy peripheral infiltrates. Often lower lobe predominance. No cavitation or pleural effusion.
Pulmonary alveolar proteinosis (PAP)	Asymptomatic if not infected with <i>Nocardia</i> .	Unremarkable.	Normal/↑ WBC, ↑ LDH.	Bilateral granular or peripheral infiltrates in butterfly pattern. No hilar adenopathy, cardiomegaly, or pleural effusion.



## BILATERAL INFILTRATES WITH EFFUSION



### Infectious Causes

Features (may have some, none, or all)				
Causes	History	Physical	Laboratory	Chest X-Ray
Legionella	Recent contact with Legionella containing water. Usually elderly. May have watery diarrhea, abdominal pain, mental confusion.	Fever/chills, relative bradycardia. Hepatic/splenic enlargement goes against the diagnosis. ↓ breath sounds if consolidation or pleural effusion.	↑ WBC, ↑ SGOT/SGPT, ↑ PO <sub>4</sub> <sup>=</sup> , ↓ Na <sup>+</sup> , ↑ CPK, ↑ ESR, ↑ CRP, ↑ ferritin levels, microscopic hematuria, L. pneumophila antigenuria (serotypes 01–06 only) may not be positive early. Mucoid/purulent sputum with few PMNs. Sputum culture for Legionella or sputum DFA (before therapy) is diagnostic. ↑ Legionella titer ≥ 1:256 or ≥ 4-fold rise between acute/convalescent titers.	Rapidly progressive asymmetrical infiltrates clue to Legionella. Consolidation and pleural effusion not uncommon. Cavitation rare.

## Infectious Causes (cont'd)

Causes	Features (may have some, none, or all)			
	History	Physical	Laboratory	Chest X-Ray
Hantavirus (HPS)	Subacute onset, shortness of breath, substernal chest discomfort. Improvement followed by rapid deterioration.	↑ respiratory rate, cyanosis in severe cases.	↓ WBC, ↓ platelets ↓ pO <sub>2</sub> , ↑ A-a gradient. ↑ hantavirus titers.	Large pleural effusions.
MERS	Recent contact with MERS case or camel contact.  Acute onset of fever/chills, headache, myalgias, dry cough ± N/V/D, Malaise, ± hemoptysis	Early fulminant respiratory failure.	(N) WBC, relative lymphopenia, ↑ LDH ↑ LFTs MERS-CoV in lower respiratory secretions.	Unilateral (early) or bilateral (late) interstitial, nodular dense infiltrates.  Recent contact with MERS case or camel contact.  Consolidation common ± pleural effusions.
Measles	Recent airborne exposure.	Measles rash, Koplik's spots.	Normal/↑ WBC (left shift), normal/↓ platelets, ↑ LFTs, ↑ CPK, normal/↓ pO <sub>2</sub> . If ↓ pO <sub>2</sub> , then ↑ A-a gradient. ↑ IgM measles titer. Warthin-Finkeldey cells in respiratory secretions.	Bilateral diffuse fine reticulonodular infiltrates ± hilar adenopathy. Lower lobe predominance. Consolidation/pleural effusion uncommon.
Strongyloides	Strongyloides exposure. 1/3 asymptomatic; 2/3 with fever, dyspnea, cough. Hyperinfection syndrome with abdominal pain, diarrhea ± GI bleed.	With hyperinfection syndrome, fever, ↓ BP, abdominal tenderness ± rebound, ± meningitis.	↑ WBC (left shift), ↑ eosinophilia, ± anemia. Blood/CSF cultures positive for enteric gram-negative bacilli. Rhabditiform larvae in sputum/stool.	Diffuse hilar patchy infiltrates without consolidation or cavitation. Eosinophilic pleural effusion common.

## Noninfectious Causes

Causes	Features (may have some, none, or all)			
	History	Physical	Laboratory	Chest X-Ray
Congestive heart failure	Coronary heart disease, valvular heart disease, cardiomyopathy.	No/low grade fevers, ↑ pulse/respiratory rate, positive jugular venous distension and hepatojugular reflex, cardiomegaly, S <sub>3</sub> , ascites, hepatomegaly, pedal edema.	↑ WBC (left shift), normal platelets, mildly ↑ SGOT/SGPT.	Cardiomegaly, pleural effusion (R > [R + L] > L). Kerley B lines with vascular redistribution to upper lobes. Typically bilateral rather than unilateral.
Chronic renal failure	Chronic renal failure on dialysis.	Findings related to uremia.	Normal/↑ WBC (left shift) plus findings related to renal failure.	Bilateral symmetrical fluffy perihilar infiltrates (butterfly pattern) ± pleural effusions. No cardiomegaly (unlike CHF), but large pericardial effusion can mimic cardiomegaly.
SLE	Fatigue, chest pain. History of SLE.	Fever/myalgias, alopecia, malar rash, "cytoid bodies" in retina, painless oral ulcers, synovitis, splenomegaly, generalized adenopathy, Raynaud's phenomenon.	↑ ANA, ↑ DS-DNA, ↓ C <sub>3</sub> , polyclonal gammopathy on SPEP, ↑ ferritin. Pleural fluids with ↑ ANA, ↓ C <sub>3</sub> .	Migratory ill-defined non-segmental infiltrates. No signs of consolidation or cavitation. ± small pleural effusions.

**Noninfectious Causes** (cont'd)

Causes	Features (may have some, none, or all)			
	History	Physical	Laboratory	Chest X-Ray
Goodpasture's Syndrome	Often preceded by a URI. Most common in 20–30 year old adults. Fever, weight loss, fatigue, cough, hemoptysis, hematuria.	Findings secondary to iron deficiency anemia.	↑ WBC, anemia, ↑ creatinine, urine with RBCs/RBC casts. Positive pANCA. Linear IgG pattern on alveolar/glomerular basement membrane.	Bilateral fine reticulonodular infiltrates predominantly in lower lobes. No cavitation.
Wegener's granulomatosis	Most common in middle-aged adults. Cough, fever, fatigue.	Findings of chronic sinusitis, bloody nasal discharge.	↑ WBC, anemia, ↑ platelets, ↑ ESR, ↑ RF, negative ANA, proteinuria, hematuria. Positive cANCA.	Bilateral asymmetrical nodular infiltrates of varying size with irregular margins. Cavitation common. Inner lining of cavities irregular. Air-fluid levels rare. ± pleural effusions. No calcifications.
Sarcoidosis (Stage III)	Dyspnea, fatigue, nasal stuffiness.	Waxy/yellowish papules on face/upper trunk. Funduscopic exam with "candle wax drippings."	↑ ESR, normal LFTs, ↑ creatinine (if renal involvement), ↑ ACE levels, hypercalciuria, hypercalcemia, polyclonal gammopathy on SPEP. Anergic.	Bilateral nodular infiltrates of variable size without hilar adenopathy. Cavitation/pleural effusion rare.

## Noninfectious Causes (cont'd)

Causes	Features (may have some, none, or all)			
	History	Physical	Laboratory	Chest X-Ray
Lymphoma	Fever, ↓ appetite with weight loss, night sweats, fatigue.	Adenopathy ± splenomegaly.	Normal WBC, ↑ basophils, ↑ eosinophilia, ↓ lymphocytes, ↑ platelets, ↑ ESR, ↑ alkaline phosphatase, ↑ $\alpha_{1,2}$ globulins on SPEP.	Unilateral or asymmetrical bilateral hilar adenopathy. Lung infiltrate may appear contiguous with hilar adenopathy. No clear channel between mediastinum and hilar nodes. Small pleural effusions rare.
Lymphangitic metastases	History of breast, thyroid, pancreas, cervical, prostate, or lung carcinoma.	Findings related to underlying malignancy.	Normal/ ↑ WBC, ↑ ESR.	Interstitial indistinct pulmonary infiltrates (may be reticulonodular) with lower lobe predominance. Usually unilateral but may be bilateral. No consolidation or cavitation ± pleural effusions.
Drug-induced	Exposure to chemotherapeutic agents (e.g., BCNU, busulfan, methotrexate, cyclophosphamide, bleomycin) or other drugs (e.g., nitrofurantoin, sulfasalazine, amiodarone, opiates, cocaine).	Unremarkable.	Normal WBC ± ↑ eosinophilia, normal/↑ ESR/ LFTs. Eosinophils in pleural effusion.	Bilateral coarse symmetrical patchy infiltrates/fibrosis ± pleural effusions. Hilar adenopathy only with drug-induced pseudolymphoma (secondary to dilantin). No cavitation.

## CAVITARY INFILTRATES (THICK-WALLED)



### Infectious Causes Based on Speed of Cavitation

Speed of Cavitation	Causes
Very rapid cavitation (3 days)	<i>S. aureus</i> , <i>P. aeruginosa</i>
Rapid cavitation (5–7 days)	<i>K. pneumoniae</i>
Slow cavitation (> 7 days)	Pyogenic lung abscess, septic pulmonary emboli
Chronic cavitation	TB (reactivation), histoplasmosis (reactivation), sporotrichosis, melioidosis, nocardia, actinomycosis, <i>Rhodococcus equi</i> , amebic abscess, alveolar echinococcosis (hydatid cysts)

## Infectious Causes

Causes	Features (may have some, none, or all)			
	History	Physical	Laboratory	Chest X-Ray
<i>S. aureus</i>	Fever, cough, dyspnea. Recent/concurrent influenza pneumonia.	↓ breath sounds ± cyanosis.	↑ WBC (left shift), ↓ pO <sub>2</sub> , ↑ A-a gradient. Sputum positive for <i>S. aureus</i> . ↑ IgM influenza titers.	Multiple thick-walled cavitory lesions superimposed on normal looking lung fields or early minimal infiltrates of influenza.
<i>P. aeruginosa</i>	Nosocomial pneumonia usually on ventilator. Nearly always rapidly fatal.	Unremarkable.	↑ WBC (left shift). Respiratory secretions culture ± for <i>P. aeruginosa</i> . Blood cultures positive for <i>P. aeruginosa</i> (hematogenous nosocomial pneumonia).	Bilateral diffuse infiltrates with rapid cavitation (≤ 72 hours).
<i>Klebsiella pneumoniae</i>	Nosocomial pneumonia or history of alcoholism in patient with community-acquired pneumonia.	Fever, chills, signs of alcoholic cirrhosis, signs of consolidation over involved lobe.	↑ WBC, ↓ platelets, ↑ SGOT/SGPT (2° to alcoholism). "Red currant jelly" sputum with PMNs and plump gram-negative encapsulated bacilli.	"Bulging fissure" sign secondary to expanded lobar volume. Empyema rather than pleural effusion. Usually cavitation in 5–7 days (thick walled).
Pyogenic lung abscess	Recent aspiration. Fevers, chills, weight loss. Swallowing disorder secondary to CNS/GI disorder.	Foul (putrid lung abscess) breath.	↑ WBC, ↑ ESR. Sputum with normal oropharyngeal anaerobic flora in putrid lung abscess.	Thick-walled cavity in portion of lung dependent during aspiration (usually basilar segment of lower lobes if aspiration occurred supine). Cavitation occurs > 7 days.

**Infectious Causes** (cont'd)

Causes	Features (may have some, none, or all)			
	History	Physical	Laboratory	Chest X-Ray
Septic pulmonary emboli	Usually IV drug abuser with fever/chills. Tricuspid regurgitation murmur or recent OB/GYN surgical procedure.	Fever > 102°F. Tricuspid valve regurgitant murmur with cannon A waves in neck.	Blood cultures positive for acute bacterial endocarditis pathogens.	Multiple peripheral nodules of varying size. Lower lobe predominance. Cavitation > 7 days characteristic of septic pulmonary emboli.
TB (reactivation)	Fevers, night sweats, normal appetite with weight loss, cough ± hemoptysis.	± bilateral apical dullness.	Normal WBC, ↑ platelets, ↑ ESR (≤ 70 mm/h). Positive PPD. AFB in sputum smear/culture.	Slowly progressive bilateral infiltrates. No pleural effusion. Usually in apical segment of lower lobes or apical/posterior segments of upper lobes. Calcifications common.
Histoplasmosis (reactivation)	Fever, night sweats, cough, weight loss, histoplasmosis exposure (river valleys of Central/Eastern United States).	± E. nodosum; otherwise unremarkable.	Normal WBC, normal/↑ eosinophilia, anemia, ↑ platelets, PPD negative. Immunodiffusion test with positive H precipitin band (diagnostic of active/chronic histoplasmosis).	Unilateral/bilateral multiple patchy infiltrates with upper lobe predilection. Bilateral hilar adenopathy uncommon. Calcifications common. No pleural effusion. Chest x-ray resembles reactivation TB.



## Infectious Causes (cont'd)

Causes	Features (may have some, none, or all)			
	History	Physical	Laboratory	Chest X-Ray
Sporotrichosis	Fever, cough, weight loss, $\pm$ hemoptysis. Antecedent lymphocutaneous or skeletal sporotrichosis, rare.	Only if secondary to residual of lymphocutaneous sporotrichosis, $\pm$ E. nodosum.	Normal WBC. No eosinophilia.	Unilateral > bilateral upper lobe nodular densities. Thick or thin-walled cavities. No hilar adenopathy. No pleural effusion.
Melioidosis	Past travel to Asia (usually > 10 years). Fever, cough, hemoptysis. $\uparrow$ risk of reactivation/dissemination with DM, alcoholics, renal failure, leukemias/lymphomas, steroids, immunosuppressives. May occur in normal hosts.	Unremarkable.	$\uparrow$ WBC (left shift), anemia, $\uparrow$ ESR, PPD negative. GNBs with bipolar "safety pin" staining in sputum. Sputum/ blood cultures positive for B. (Pseudomonas) pseudomallei.	Resembles reactivation TB, but infiltrates nodular/diffuse (septicemic) or cavitory (chronic) predominantly in middle/lower lung fields, $\pm$ air fluid level. No pleural effusion.
Nocardia	Fevers, night sweats, fatigue, $\downarrow$ cell-mediated immunity (e.g., HIV, organ transplants, PAP, immunosuppressive therapy).	Unremarkable.	Normal/ $\uparrow$ WBC, $\uparrow$ ESR. Sputum with gram-positive AFB.	Dense large infiltrates. May cavitate and mimic TB, lymphoma, or squamous cell carcinoma. No calcification $\pm$ pleural effusion.
Actinomycosis	Recent dental work.	$\pm$ chest wall sinus tracts.	Normal/ $\uparrow$ WBC, $\uparrow$ ESR. Sputum with Gram-positive filamentous anaerobic bacilli.	Dense infiltrates extending to chest wall. No hilar adenopathy. Cavitation $\pm$ pleural effusion rare.

## Infectious Causes (cont'd)

Causes	Features (may have some, none, or all)			
	History	Physical	Laboratory	Chest X-Ray
Rhodococcus equi	Insidious onset of fever, dyspnea, chest pain, $\pm$ hemoptysis. Immunosuppressed patients with $\downarrow$ cell-mediated immunity or exposure to cattle, horses, pigs.	Unremarkable.	Normal/ $\uparrow$ WBC. Sputum/pleural fluid with gram-positive pleomorphic weakly acid-fast bacilli. Sputum, pleural fluid, blood cultures positive for <i>R. equi</i> .	Segmental infiltrate with upper lobe predominance $\pm$ cavitation. Air-fluid levels and pleural effusion common.
Amebic cysts	Hepatic amebic abscess. Remote history of usually mild amebic dysentery.	$\pm$ hepatomegaly.	Normal WBC and ESR. $\uparrow$ E. histolytica HI titers.	Well-circumscribed cavitory lesions adjacent to right diaphragm. $\pm$ calcifications. Sympathetic pleural effusion above hepatic amebic abscess.
Alveolar echinococcosis (hydatid cysts)	Symptoms related to cyst size/ location: 1/3 asymptomatic; 2/3 with fever, malaise, chest pain $\pm$ hemoptysis. RUQ abdominal pain may occur.	Hepatomegaly common.	Normal/ $\uparrow$ WBC, no eosinophilia, $\uparrow$ alkaline phosphate/SGPT with hepatic cysts. Abdominal ultrasound/CT with calcified hepatic irregularly shaped cysts ("Swiss cheese" calcification characteristic). $\uparrow$ E. multilocularis IHA titers.	RLL usual location (hepatic cysts penetrate diaphragm into RLL). Nodules are 70% solitary, 30% multiple. Pleural effusion rare. Endocyst membrane on surface of cyst fluid ("water lily" sign) is characteristic.


## Noninfectious Causes

Causes	Features (may have some, none, or all)			
	History	Physical	Laboratory	Chest X-Ray
Wegener's granulomatosis	Most common in middle-aged adults. Cough, fever, fatigue.	Findings of chronic sinusitis, bloody nasal discharge.	↑ WBC, anemia, ↑ platelets, ↑ ESR, ↑ RF, negative ANA, proteinuria, hematuria. Positive cANCA.	Bilateral asymmetrical nodular infiltrates of varying size with irregular margins. Cavitation common. Inner lining of cavities irregular. Air-fluid levels rare. ± pleural effusions. No calcifications.
Squamous cell carcinoma	Long-term smoking history.	Clubbing, hypertrophic pulmonary osteoarthropathy ± findings 2° to superior vena caval syndrome and CNS/bone metastases.	↑ Ca <sup>++</sup> (without bone metastases).	Unilateral perihilar mass lesion. Cavitation common. No pleural effusion.
Lymphoma	Fever, ↓ appetite with weight loss, night sweats, fatigue.	Adenopathy ± splenomegaly.	Normal WBC, ↑ basophils, ↑ eosinophilia, ↓ lymphocytes, ↑ platelets, ↑ ESR ↑ alkaline phosphatase, ↑ α <sub>1,2</sub> globulins on SPEP.	Unilateral or asymmetrical bilateral hilar adenopathy. Lung infiltrate may appear contiguous with hilar adenopathy. No clear channel between mediastinum and hilar nodes. Small pleural effusions rare.

**Noninfectious Causes** (cont'd)

Causes	Features (may have some, none, or all)			
	History	Physical	Laboratory	Chest X-Ray
Metastatic carcinoma	History of breast, thyroid, renal cell, colon, pancreatic cancer or osteogenic sarcoma.	Findings related to underlying malignancy and, when present, to bone, hepatic, CNS metastases.	Secondary to effects of primary neoplasm, metastases, paraneoplastic syndrome.	Nodular lesions that vary in size. Metastatic lesions are usually well-circumscribed with lower lobe predominance. Usually no bronchial obstruction (obstruction suggests colon, renal, or melanoma metastases). Usually no cavitation (except for squamous cell metastases). Calcification usually suggests osteosarcoma (rarely adenocarcinoma). Pleural effusion rare (except for breast cancer).
Rheumatoid nodules	Usually in severe rheumatoid arthritis (RA); ± history of silicosis.	Findings secondary to RA. Rheumatoid nodules on exterior surfaces of arms.	Normal WBC, ↑ ESR, ↑ ANA, ↑ RF (high titer). Pleural fluid with ↓ glucose.	Lung nodules are round and well-circumscribed, predominantly in lower lobes and typically superimposed on interstitial lung disease ("rheumatoid lung"). Cavitation is common. ± pulmonary fibrosis, pleural effusion. Silicosis + RA nodules = Caplan's syndrome.

## CAVITARY INFILTRATES (THIN-WALLED)



### Infectious Causes

Causes	Features (may have some, none, or all)			
	History	Physical	Laboratory	Chest X-Ray
Atypical TB	Often occurs in setting of previous lung disease.	Unremarkable.	Normal WBC, ↑ ESR. Weakly positive PPD. Positive sputum AFB/culture.	Multiple cavitary lesions ± calcifications usually involving both lungs. Resembles reactivation TB except that cavities are thin-walled. No pleural effusion.
Coccidiomycosis (reactivation)	Previous exposure in endemic coccidiomycosis areas (e.g., Southwest USA). Asymptomatic.	± <i>E. nodosum</i> .	Normal WBC. Eosinophilia acutely but not in chronic phase, and eosinophils in pleural fluid. CF IgG titer ≥ 1:32 indicates active disease.	Thick/thin walled cavities (< 3 cm) usually in anterior segments of lower lobes ± calcifications/bilateral hilar adenopathy. Air-fluid levels rare unless secondarily infected. Pleural effusion rare.

**Infectious Causes** (cont'd)

Causes	Features (may have some, none, or all)			
	History	Physical	Laboratory	Chest X-Ray
Paragonimiasis	Ingestion of fresh-water crabs/crayfish. Acute symptoms (< 6 months): fevers, abdominal pain, diarrhea followed by episodes of pleuritic chest pain. Chronic symptoms (> 6 months): fevers, night sweats, cough $\pm$ hemoptysis. Asymptomatic in some.	Wheezing, $\pm$ urticaria (acutely).	$\uparrow$ WBC, eosinophilia. Eosinophils in pleural fluid. Sputum with Charot-Leyden crystals. Sputum/feces with operculated <i>P. Westermani</i> eggs.	Cavitary patchy or well-defined infiltrates predominantly in mid-lung fields. Hydropneumothorax common. Calcifications and pleural effusion common.
Sporotrichosis	Fever, cough, weight loss. $\pm$ hemoptysis. Antecedent lymphocutaneous or skeletal sporotrichosis, rare.	Only if secondary to residual of lymphocutaneous sporotrichosis, $\pm$ <i>E. nodosum</i> .	Normal WBC. No eosinophilia.	Unilateral > bilateral upper lobe nodular densities. Thick or thin-walled cavities. No hilar adenopathy. No pleural effusion.
Pneumatocoeles	Common in <i>S. aureus</i> pneumonia in children. Fever, cough, dyspnea. $\pm$ antecedent influenza.	Unremarkable unless pneumatocele ruptures, then signs of pneumothorax.	$\uparrow$ WBC (left shift), $\downarrow$ pO <sub>2</sub> (secondary to influenza).	Multiple thin-walled cavities in areas of <i>S. aureus</i> pneumonia. Common in children; rare in adults.

## Noninfectious Causes

Causes	Features (may have some, none, or all)			
	History	Physical	Laboratory	Chest X-Ray
Emphysema (blebs/cyst)	Long history of smoking. Rupture of apical bleb common in males > 30 years. Pneumonia rare in severe emphysema (vs. chronic bronchitis) but may occur early in unaffected areas of lung.	Asthenic "pink puffers." Barrel chest. Diaphragmatic excursions < 2 cm.	Normal WBC, ↓ pO <sub>2</sub> .	↓ lung markings ("vanishing lung") ± blebs. Hyperlucent lungs with upper lobe predominance. Flattened diaphragms, vertically elongated cardiac silhouette, ↑ retrocardiac and retrosternal airspaces. No infiltrates or pleural effusions. (In upper lobe emphysema, no vascular redistribution to upper lobes with CHF).
Bronchogenic cyst	Congenital anomaly. Usually asymptomatic. Cough if symptomatic.	Unremarkable unless secondarily infected, then signs 2° to mediastinal abscess.	Normal WBC/ESR.	Circumscribed cystic lesion originates in lung but appears high in mediastinum. If filled with fluid, appears as solitary tumor. If near the trachea, may rupture into bronchus/trachea and cyst may contain air. If communicates openly with bronchus, appears as thin-walled cavitary nodule. If infected, presents as mediastinal abscess.
Cystic bronchiectasis	Recurrent pulmonary infections with purulent sputum ± hemoptysis.	Unremarkable unless dextrocardia with sinusitis (Kartagener's syndrome).	↑ WBC, normal ESR.	Bilateral large cystic lucencies at lung bases. Upper lobes relatively spared (unless secondary chronic aspiration). Thickened bronchial markings at bases. Bronchiectasis of cystic fibrosis predominantly involves upper lobes.

## Noninfectious Causes (cont'd)

Features (may have some, none, or all)				
Causes	History	Physical	Laboratory	Chest X-Ray
Sequestered lung	Usually asymptomatic. Productive cough $\pm$ hemoptysis if communicates with bronchus or if infected.	Unremarkable.	Normal WBC, $\uparrow$ ESR.	Solid nodule unless communicates with bronchus, then thin-walled cavity $\pm$ air fluid levels. Usually posterior based segment of lower lobes (LLL > RLL). If > 3 cm, presents as mass lesion.
Histiocytosis X (eosinophilic granuloma, Langerhan's cell histiocytosis)	Patients usually 20–40 years. Usually asymptomatic. Fever, cough, dyspnea in some. Diabetes insipidus rare.	Hepato-splenomegaly, skin lesions, hemoptysis (rare).	Normal WBC, eosinophilia.	Pneumothorax superimposed on diffuse pulmonary fibrosis. Cystic bone lesions. Usually mid/upper lung fields with nodules/thin-walled cysts or infiltrates. No hilar adenopathy or pleural effusion.

## REFERENCES AND SUGGESTED READINGS

- Burgener FA, Korman M. *Differential Diagnosis in Conventional Radiology* (2<sup>nd</sup> Ed), Stuttgart, Georg Thieme Verlag, 1991.
- Chapman S, Nakielnny R. *Aids to Radiological Differential Diagnosis* (2<sup>nd</sup> Ed), London, Bailliere Tindall, 1990.
- Conn RB, Borer WZ, Snyder JW. *Current Diagnosis* 9. Philadelphia, W. B. Saunders Company, 1997.
- Crapo JD, Glassroth J, Karlinsky JB, King TE. *Baum's Textbook of Pulmonary Diseases* (7<sup>th</sup> Ed), Philadelphia, Lippincott Williams & Wilkins, 2004.
- Cunha BA. *Pneumonia Essentials* (3<sup>rd</sup> Ed), Jones & Bartlett, Sudbury MA, 2010.
- Eisenberg RL. *Clinical Imaging. An Atlas of Differential Diagnosis* (2<sup>nd</sup> Ed), Gaithersburg, Aspen Publishers, Inc., 1992.
- Gorbach SL, Bartlett JG, Blacklow NR. *Infectious Diseases* (3<sup>rd</sup> Ed), Philadelphia, Lippincott Williams & Wilkins, 2004.
- Karetzky M, Cunha BA, Brandstetter RD. *The Pneumonias*. New York, Springer-Verlag, 1993.
- Kasper L, Braunwald E, Fauci AS, Hauser SL, Longo DL, Jameson JL. *Harrison's Principles of Internal Medicine* (17<sup>th</sup> Ed), New York, The McGraw Hill Companies, 2008.
- Levison ME. *The Pneumonias: Clinical Approaches to Infectious Diseases of the Lower Respiratory Tract*. Boston, John Wright, PSG Inc., 1984.
- Lillington GA, Jamplis RW. *A Diagnostic Approach to Chest Diseases* (2<sup>nd</sup> Ed), Baltimore, The Williams & Wilkins Company, 1977.
- Mandell GL, Bennet JE, Blaser MJ (eds). *Mandell, Douglas, and Bennett's Principles and Practice of Infectious Diseases* (8<sup>th</sup> Ed), Elsevier, 2015.
- Murray JF, Nadel JA. *Textbook of Respiratory Medicine* (3<sup>rd</sup> Ed), Philadelphia, W.B. Saunders Company, 2000.
- Raof S, Feigin D, Sung A, et al. *Interpretation of Plain Chest Roentgenogram*. *Chest* 141:545–558, 2012.
- Teplick JE, Haskin ME. *Roentgenologic Diagnosis* (3<sup>rd</sup> Ed), Philadelphia, W. B. Saunders Company, 1976.
- Wright FW. *Radiology of the Chest and Related Conditions*. London, Taylor & Francis, 2002.



## Chapter 9

**Infectious Disease Differential Diagnosis****Cheston B. Cunha, MD**

Table 9.1. Fever Curves .....	475
Table 9.2. General Appearance Abnormalities .....	476
Table 9.3. Head Abnormalities .....	477
Table 9.4. Eye Abnormalities .....	478
Table 9.5. Ear Abnormalities .....	481
Table 9.6. Nasal Abnormalities .....	481
Table 9.7. Mouth Abnormalities .....	482
Table 9.8. Neck Abnormalities .....	483
Table 9.9. Chest Abnormalities .....	484
Table 9.10. Back Abnormalities .....	484
Table 9.11. Heart Abnormalities .....	485
Table 9.12. Abdominal Abnormalities .....	485
Table 9.13. Lymph Node Abnormalities .....	488
Table 9.14. Genitourinary Abnormalities .....	489
Table 9.15. Extremity Abnormalities .....	490
Table 9.16. Neurological Abnormalities .....	493
Table 9.17. WBC Abnormalities .....	495
Table 9.18. RBC Abnormalities .....	497
Table 9.19. Platelet Abnormalities .....	498
Table 9.20. Pancytopenia .....	498
Table 9.21. Serum Test Abnormalities .....	498
Table 9.22. Liver Test Abnormalities .....	501
Table 9.23. Rheumatic Test Abnormalities .....	502
Table 9.24. Urinary Abnormalities .....	502
Table 9.25. Pleural Fluid Abnormalities .....	504
Table 9.26. CSF Abnormalities .....	506
Table 9.27. CSF Diagnostic Criteria for EVD Associated ABM .....	506

**Table 9.1. Fever Curves**

Finding	Causes
Morning (am) temperature spikes	<u>Infectious:</u> TB, typhoid fever, Whipple's disease <u>Noninfectious:</u> PAN
Relative tachycardia	<u>Infectious:</u> gas gangrene, myocarditis (Coxsackie, RMSF, typhus, diphtheria, trichinosis, VLM, influenza) <u>Noninfectious:</u> Hyperthyroidism, hypoxia, cardiac arrhythmias, anxiety, pulmonary emboli
Relative bradycardia	<u>Infectious:</u> typhoid fever, typhus, leptospirosis, Legionnaire's disease, Q fever, psittacosis, RMSF, babesiosis, ehrlichiosis/anaplasmosis, yellow fever, dengue fever, arboviral hemorrhagic fevers <u>Noninfectious:</u> drug fever, CNS lesions, $\beta$ -blockers, verapamil, diltiazem, lymphoma, factitious fever
Single fever spikes	<u>Infectious:</u> Transient bacteremia <u>Noninfectious:</u> Blood/blood product transfusions
Double quotidian fevers	<u>Infectious:</u> kala-azar, right sided GC ABE, miliary TB, mixed malarial infections <u>Noninfectious:</u> adult Still's disease, antipyretics
Camelback fever pattern	<u>Infectious:</u> Colorado tick fever, dengue fever, leptospirosis, brucellosis, LCM, yellow fever, poliomyelitis, smallpox, rat-bite fever ( <i>Spirillum minus</i> ), chikungunya fever, rift valley fever, african hemorrhagic fevers (Marburg, Ebola, Lassa, etc), echovirus (echo 9), ehrlichiosis/ anaplasmosis <u>Noninfectious:</u> antipyretics

**Fever with chills/rigors**

Classically, chills/rigors are associated with some infections, diseases, e.g., influenza, bacteremia, malaria, but rare/mild with typhoid fever. Prominent/sustained rigors argues against the diagnosis of typhoid fever.

**Fever with night sweats**

Night sweats may be present with some chronic infectious diseases, e.g., abscesses, TB, SBE and various malignancies, e.g., lymphomas, leukemia. Other causes include certain medications, e.g., monoclonal antibody therapy (rituximab), HIV drugs. Night sweats are not a feature of rheumatic/inflammatory disorders.

**Table 9.2. General Appearance Abnormalities**

Finding	Causes
Hypotension/ shock	<p><u>Infectious</u>: anthrax, plague, typhus, <b>GI sources</b>: colon (perforation, colitis, abscess); <b>GU sources</b>: cystitis (only in SLE, DM, alcoholism, multiple myeloma, Vibrio vulnificus, CLL, steroids); obstruction: relative (BPH), unilateral (partial), or bilateral/ureteral (partial/total); kidney (acute pyelonephritis, renal abscess, perinephric abscess, renal calculi); prostate (acute prostatitis, prostatic abscess). <b>Other sources</b>: CVC infection, TSS (<i>S. aureus</i>, group A streptococci, <i>C. sordellii</i>), dengue fever, SARS, HPS, influenza (human, avian, swine), CAP with decreased/absent splenic function, or severe cardiopulmonary disease, influenza/ILI with CA-MRSA (PVL+)</p> <p><u>Noninfectious</u>: myocardial infarction, pulmonary embolism, acute pancreatitis, relative adrenal insufficiency, GI bleed, overzealous diuresis, inadequate volume replacement or hypotonic fluid replacement, aortic dissection, abdominal aneurysm rupture, rectus sheath hematoma</p>
Soft tissue crepitation	<p><u>Infectious</u>: mixed aerobic/anaerobic infection*</p> <p><u>Noninfectious</u>: subcutaneous emphysema, recent surgery in area of crepitation</p>
Bullae (vesicles)	<p><u>Infectious</u>: gas gangrene (hemorrhagic), <i>S. aureus</i> &gt; group A streptococci, <i>V. vulnificus</i> (hemorrhagic), necrotizing fasciitis, bullous impetigo, HSV, VZV, smallpox</p> <p><u>Noninfectious</u>: pemphigus, pemphigoid, dermatitis herpetiformis, TEN, porphyria, contact dermatitis, <i>E. multiforme</i>, DM, barbiturates</p>
Generalized edema/nephrotic syndrome	<p><u>Infectious</u>: malaria (<i>P. malariae</i>), trichinosis, VZV, yellow fever, influenza A, HBV, HIV, EBV, CMV, 2<sup>o</sup> syphilis, leprosy</p> <p><u>Noninfectious</u>: drugs, lymphomas, thiamine deficiency (wet beri-beri), kwashiorkor, post-streptococcal glomerulonephritis, DM, PAN, Henoch-Schönlein purpura, SLE, constrictive pericarditis, tricuspid regurgitation, IVC obstruction, myxedema, sickle cell disease, amyloidosis, <math>\alpha</math>-1 antitrypsin deficiency, nail-patella syndrome, Wegener's granulomatosis, sarcoidosis, renal vein thrombosis</p>
Erythroderma	<p><u>Infectious</u>: human T-cell leukemia virus 1 (HTLV-1), TSS, scarlet fever, dengue fever, chikungunya fever</p> <p><u>Noninfectious</u>: psoriasis, lymphoma, atopic dermatitis, contact dermatitis, exfoliative dermatitis, drugs, ichthyosis, erythroleukemia, pityriasis, dermatomyositis, Kawasaki's disease, Hodgkin's lymphoma</p>
Hyper- pigmentation	<p><u>Infectious</u>: kala-azar, Whipple's disease, histoplasmosis</p> <p><u>Noninfectious</u>: hemochromatosis, PBC, Addison's disease, drugs – brown (bleomycin), blue/gray (busulfan, fluorouracil, chlorpromazine), purple/gray (amiodarone), blue/gray (minocycline)</p>

\* Versus gas gangrene (minimal/no gas on muscle auscultation).

**Table 9.2. General Appearance Abnormalities (cont'd)**

Finding	Causes
Jaundice	<u>Infectious</u> : yellow fever, EBV, leptospirosis, ascariasis (in bile/pancreatic duct), viral hemorrhagic fevers, viral hepatitis  <u>Noninfectious</u> : hepatobiliary malignancy, benign biliary obstruction, PBC, cirrhosis, hemolytic anemias, alcoholic hepatitis, drugs, Gilbert's syndrome, pancreatic carcinoma
Vitiligo	<u>Infectious</u> : leprosy, yaws, pinta, HIV  <u>Noninfectious</u> : DM, hyperthyroidism, hypothyroidism, Hashimoto's thyroiditis
Lizard/elephant skin	<u>Infectious</u> : onchocerciasis  <u>Noninfectious</u> : ichthyosis
Urticaria	<u>Infectious</u> : strongyloides, dracunculiasis, HBV, F. hepatica, schistosomal dermatitis  <u>Noninfectious</u> : insect bites/stings, drugs, malignancies, cholinergic urticaria, serum sickness, urticarial vasculitis, cryoglobulinemia, systemic mastocytosis, Schnitzler's syndrome
Rash (only below waist)	<u>Infectious</u> : Parvovirus B19 (stocking/glove)  <u>Noninfectious</u> : Henoch-Schönlein purpura (HSP)

**Table 9.3. Head Abnormalities**

Finding	Causes
Prematurely gray hair	<u>Infectious</u> : HIV  <u>Noninfectious</u> : smoking, pernicious anemia
Temporal muscle wasting	<u>Infectious</u> : HIV  <u>Noninfectious</u> : myotonic dystrophy
Facial swelling (edema)	<u>Infectious</u> : arboviral hemorrhagic fevers, trichinosis, leprosy, EEE, onchocerciasis  <u>Noninfectious</u> : angioneurotic edema, nephrotic syndrome, acute glomerulonephritis, leukemic infiltrates, amyloidosis
Severe seborrheic dermatitis	<u>Infectious</u> : HIV  <u>Noninfectious</u> : CGD
Facial red spots	<u>Infectious</u> : psittacosis (Horder's spots)  <u>Noninfectious</u> : Campbell de Morgan spots

**Table 9.3. Head Abnormalities** (cont'd)

Finding	Causes
Localized alopecia	<u>Infectious</u> : leprosy, Tinea capitis, smallpox, blastomycosis, VZV, TB (cutaneous) <u>Noninfectious</u> : SLE, DM, scleroderma, common variable immune deficiency (CVID)
Generalized alopecia	<u>Infectious</u> : post-malaria, post-typhoid fever, post-kala-azar, post-yellow fever, 2° syphilis, HIV <u>Noninfectious</u> : nutritional deficiencies, drugs, hyperthyroidism
Total/partial eyebrow loss	<u>Infectious</u> : leprosy, syphilis <u>Noninfectious</u> : iatrogenic, hereditary, hypothyroidism, hypopituitarism
Long eyelashes	<u>Infectious</u> : kala-azar, trypanosomiasis <u>Noninfectious</u> : hereditary, drugs, HIV drugs
Lacrimal gland enlargement <i>unilateral</i> <i>bilateral</i>	<u>Infectious</u> : Chagas' disease <u>Noninfectious</u> : malignancies  <u>Infectious</u> : TB <u>Noninfectious</u> : Sjogren's syndrome, RA, SLE, sarcoidosis
Facial erythema	<u>Infectious</u> : facial cellulitis, erysipelas, TB (lupus vulgaris), arboviral hemorrhagic fevers <u>Noninfectious</u> : dermatomyositis, drugs, rosacea, carcinoid syndrome, SLE
Parotid enlargement	<u>Infectious</u> : mumps, Chagas' disease, rat bite fever ( <i>Streptobacillus moniliformis</i> ), melioidosis, CMV, EBV, LCM, adenovirus, Coxsackie A <u>Noninfectious</u> : Sjogren's syndrome, sarcoidosis, cirrhosis
Scalp nodules	<u>Infectious</u> : myiasis, onchocerciasis <u>Noninfectious</u> : bony exostoses, malignancies, benign cysts, pyogenic granuloma, Kimura's disease

**Table 9.4. Eye Abnormalities**

Finding	Causes
Bilateral upper lid edema	<u>Infectious</u> : EBV (Hoagland's sign) <u>Noninfectious</u> : bilateral eye irritation
Bilateral lid edema	<u>Infectious</u> : trichinosis, meningococcemia, adenovirus <u>Noninfectious</u> : Wegener's granulomatosis

**Table 9.4. Eye Abnormalities** (cont'd)

<b>Finding</b>	<b>Causes</b>
Heliotrope eyelid discoloration	<u>Infectious</u> : cholera (early), influenza A (severe) <u>Noninfectious</u> : dermatomyositis
Periorbital edema <i>unilateral</i>	<u>Infectious</u> : Chagas' disease, loiasis, gnathostomiasis, sparganosis <u>Noninfectious</u> : insect bites, unilateral eye irritation
<i>bilateral</i>	<u>Infectious</u> : RMSF, trichinosis, tularemia, dengue fever <u>Noninfectious</u> : allergies, Insect bites, TNF-receptor-1-associated periodic syndrome (TRAPS), dermatomyositis
Argyll-Robertson pupils	<u>Infectious</u> : syphilis <u>Noninfectious</u> : sarcoidosis
Iritis	<u>Infectious</u> : onchocerciasis, 2 <sup>o</sup> syphilis, relapsing fever, leprosy <u>Noninfectious</u> : SLE, dermatomyositis, Behçet's syndrome, Reiter's syndrome, RA
Conjunctivitis <i>unilateral</i>	<u>Infectious</u> : TB, HSV, tularemia, adult inclusion conjunctivitis, Chagas' disease, LGV, CSD, loiasis, ocular myiasis, diphtheria, adenovirus (types 8, 19) <u>Noninfectious</u> : SLE, eye irritation
<i>bilateral</i>	<u>Infectious</u> : TSS, measles, rubella, meningococemia, gonorrhea, adenovirus (type 3), plague, RMSF, sparganosis, LGV, listeria, relapsing fever, pertussis, influenza, microsporidia, HHV-6, arboviral hemorrhagic fevers, dengue hemorrhagic fever <u>Noninfectious</u> : Kawasaki's disease, Reiter's syndrome, Steven-Johnson syndrome, adult Still's disease, eye irritation
Hemorrhagic conjunctivitis <i>unilateral</i>	<u>Infectious</u> : adenovirus (type 8, 19) <u>Noninfectious</u> : trauma
<i>bilateral</i>	<u>Infectious</u> : trichinosis, pertussis, leptospirosis, Coxsackie A (type 24), adenovirus (type 11), enterovirus (type 70), arboviral hemorrhagic fevers <u>Noninfectious</u> : trauma
Subconjunctival hemorrhage	<u>Infectious</u> : SBE, trichinosis, meningococemia, pertussis, leptospirosis, RMSF <u>Noninfectious</u> : severe anemia, Kawasaki's disease
Conjunctival suffusion	<u>Infectious</u> : RMSF, leptospirosis, relapsing fever, ehrlichiosis, HPS, influenza (human, avian), arboviral hemorrhagic fevers, St.LE <u>Noninfectious</u> : bilateral eye irritation

**Table 9.4. Eye Abnormalities** (cont'd)

Finding	Causes
Episcleritis	<u>Infectious:</u> TB, leprosy, 2° syphilis, Lyme disease <u>Noninfectious:</u> sarcoidosis, RA, adult Still's disease, SLE, PAN, TA, RE
Scleral nodules	<u>Infectious:</u> loiasis, sparganosis, leprosy, tularemia, TB <u>Noninfectious:</u> RA, vitamin A deficiency (Bitot's spots), sarcoidosis
Dry eyes	<u>Infectious:</u> measles <u>Noninfectious:</u> vitamin A deficiency, SLE, Sjogren's syndrome, RA, sarcoidosis
Watery eyes	<u>Infectious:</u> bacterial conjunctivitis, adenovirus <u>Noninfectious:</u> PAN, allergic conjunctivitis
Uveitis	<u>Infectious:</u> TB, histoplasmosis, leprosy, syphilis, malaria, HSV, VZV, EBV, TSS, typhus, LGV, CMV, African trypanosomiasis, brucellosis, leptospirosis, RMSF, CSD <u>Noninfectious:</u> adult Still's disease, SLE, PAN, sarcoidosis, Behçet's syndrome, Reiter's syndrome, RA, relapsing polychondritis, ankylosing spondylitis, Kawasaki's disease, Wegener's granulomatosis
Corneal haziness	<u>Infectious:</u> adenovirus, leprosy, trachoma, onchocerciasis <u>Noninfectious:</u> vitamin A deficiency, cataracts
Keratitis	<u>Infectious:</u> HSV, congenital syphilis, acanthamoeba, TB, toxoplasmosis, histoplasmosis, CMV, trachoma, microsporidia, nocardia, leprosy, onchocerciasis, meliodosis <u>Noninfectious:</u> Behçet's syndrome, Reiter's syndrome, Steven-Johnson syndrome, vitamin A deficiency
Corneal ulcers	<u>Infectious:</u> HSV, listeria, acanthamoeba, tularemia, shigella, RMSF <u>Noninfectious:</u> trauma, Wegener's granulomatosis
Endophthalmitis	<u>Infectious:</u> TB, candida, aspergillus, toxocara, serratia, S. pneumoniae <u>Noninfectious:</u> retinoblastoma
Chorioretinitis	<u>Infectious:</u> toxoplasmosis, CMV, onchocerciasis, congenital syphilis, histoplasmosis, TB, WNE, coccidiomycosis, leptospirosis, chikungunya fever <u>Noninfectious:</u> sarcoidosis, SLE, PAN
Cytoid bodies (cotton wool spots)	<u>Infectious:</u> CSD, HIV, CMV, SBE <u>Noninfectious:</u> SLE, adult Still's disease, PAN, atrial myxoma, Wegener's granulomatosis, TA

**Table 9.4. Eye Abnormalities** (cont'd)

Finding	Causes
Roth spots	<u>Infectious</u> : SBE, psittacosis, RMSF, malaria <u>Noninfectious</u> : PAN, SLE, DM, severe anemia, leukemia, cholesterol emboli syndrome, atrial myxoma, TA, Takayasu's arteritis
Periphlebitis (candle wax drippings)	<u>Infectious</u> : CMV, leptospirosis (Weil's syndrome) <u>Noninfectious</u> : sarcoidosis

**Table 9.5. Ear Abnormalities**

Finding	Causes
Acute deafness	<u>Infectious</u> : ABM, mumps (aseptic meningitis), RMSF, typhus, brucellosis, Lassa fever <u>Noninfectious</u> : sound/barotrauma
External ear lesions	<u>Infectious</u> : cutaneous leishmaniasis (Chiclero ulcer), leprosy, Kaposi's sarcoma <u>Noninfectious</u> : relapsing polychondritis, eczema, carcinoma, contact dermatitis, sarcoidosis, SLE, gout (tophi), keloids, actinic keratosis

**Table 9.6. Nasal Abnormalities**

Finding	Causes
Purple nose tip/hyperpigmented	<u>Infectious</u> : Kaposi's sarcoma, TB (Bazin's erythema induratum), chikungunya fever ("Chik sign") <u>Noninfectious</u> : vasculitis, lymphoma, drugs, sarcoidosis
Nose tip gangrene	<u>Infectious</u> : Staphylococcus aureus ABE (emboli) <u>Noninfectious</u> : SLE, vasculitis
Epistaxis	<u>Infectious</u> : psittacosis, typhoid fever, nasal diphtheria, Colorado tick fever, TBRF, influenza, dengue hemorrhagic fever, arboviral hemorrhagic fevers, acute renal failure, leprosy, leptospirosis, VZV, TB, rhinosporidium, mucocutaneous leishmaniasis <u>Noninfectious</u> : Local trauma, sinus malignancies, von Willebrand's disease, polycythemia vera, Waldenstrom's macroglobulinemia, relapsing polychondritis
Nasal septal perforation	<u>Infectious</u> : leprosy, 2 <sup>o</sup> syphilis, mucocutaneous leishmaniasis, blastomycosis, pinta, yaws <u>Noninfectious</u> : cocaine, lethal midline granuloma, Wegener's granulomatosis, miasis



**Table 9.7. Mouth Abnormalities**

Finding	Causes
Trismus	<u>Infectious</u> : tetanus <u>Noninfectious</u> : temporomandibular joint dislocation/arthritis, trigeminal neuralgia
Herpes labialis	<u>Infectious</u> : HSV, pneumococcal pneumonia, meningococcal meningitis, malaria <u>Noninfectious</u> : contact dermatitis (may mimic H. labialis)
Angular cheilitis	<u>Infectious</u> : 2° syphilis, HIV <u>Noninfectious</u> : riboflavin deficiency, trauma, contact dermatitis, anemia
Gingivitis	<u>Infectious</u> : trench mouth <u>Noninfectious</u> : Wegener's granulomatosis
Tongue tenderness	<u>Infectious</u> : relapsing fever <u>Noninfectious</u> : vitamin deficiencies, pernicious anemia
Tongue ulcers	<u>Infectious</u> : histoplasmosis, HSV, syphilis <u>Noninfectious</u> : aphthous ulcers, chemotherapy, radiation therapy
Leukoplakia	<u>Infectious</u> : HIV (hairy), syphilis <u>Noninfectious</u> : lichen planus
Oral ulcers <i>solitary</i>  <i>multiple</i>	<u>Infectious</u> : syphilis, CMV, histoplasmosis, TB, Leishmania braziliensis <u>Noninfectious</u> : squamous cell carcinoma, Behçet's syndrome, Wegener's granulomatosis  <u>Infectious</u> : HSV, HFM disease, herpangina, brucellosis, Lassa fever, African tick bite fever, chickunguniya fever <u>Noninfectious</u> : SLE*, celiac disease, aphthous ulcers, squamous cell carcinoma, Behçet's syndrome, FAPA syndrome*, hyper IgE (Job's) syndrome E. multiforme, RE, cyclic neutropenia, hyper IgD syndrome, Sweet's syndrome
Frenal ulcer	<u>Infectious</u> : pertussis <u>Noninfectious</u> : trauma
Palatal petechiae	<u>Infectious</u> : Group A streptococci, EBV, CMV, HSV, VZV, toxoplasmosis, rubella, HIV, tularemia <u>Noninfectious</u> : thrombocytopenia (2° to any cause), platelet dysfunction disorders, DIC, Ehlers-Danlos syndrome, Marfan's syndrome

\* Painless ulcers

**Table 9.7. Mouth Abnormalities** (cont'd)

Finding	Causes
Palatal perforation	<u>Infectious</u> : congenital syphilis, myiasis <u>Noninfectious</u> : Post-surgical, midline granuloma, cocaine
Palatal vesicles	<u>Infectious</u> : <b>Anterior</b> : HSV, VZV, hand-foot-mouth disease (Coxsackie A). <b>Posterior</b> : herpangina (Coxsackie A) <u>Noninfectious</u> : bullous pemphigus, Steven-Johnson syndrome (drug-induced)
Uvular edema	<u>Infectious</u> : Group A streptococci <u>Noninfectious</u> : Franklin's disease, drugs, angioneurotic edema
Crimson crescents	<u>Infectious</u> : chronic fatigue syndrome (CFS), HIV <u>Noninfectious</u> : None
Tonsillar membranes	<u>Infectious</u> : diphtheria, Arcanobacterium hemolyticum <u>Noninfectious</u> : None
Tonsillar ulcers	<u>Infectious</u> : tularemia (oropharyngeal), 1° syphilis, Vincent's angina, TB <u>Noninfectious</u> : carcinoma, T-cell lymphoma, AML
Hoarseness	<u>Infectious</u> : laryngeal TB, C. pneumoniae, respiratory viruses, pertussis <u>Noninfectious</u> : recurrent laryngeal nerve paralysis, laryngeal cancer

**Table 9.8. Neck Abnormalities\***

Finding	Causes
Enlarged greater auricular nerve	<u>Infectious</u> : leprosy <u>Noninfectious</u> : none
Bull neck	<u>Infectious</u> : diphtheria, mumps, Ludwig's angina, pertussis, group A streptococcal suppurative lymphangitis <u>Noninfectious</u> : angioneurotic edema, subcutaneous emphysema
Jugular vein tenderness	<u>Infectious</u> : suppurative jugular thrombophlebitis (Lemierre's syndrome) <u>Noninfectious</u> : thrombophlebitis
Neck sinus tract	<u>Infectious</u> : actinomycosis, TB, atypical TB <u>Noninfectious</u> : branchial cleft cyst, CGD
Superior vena cava syndrome	<u>Infectious</u> : actinomycosis <u>Noninfectious</u> : lymphoma, squamous cell carcinoma

\* See Table 9.13 for cervical adenopathy (p. 488).

**Table 9.9. Chest Abnormalities**

<b>Finding</b>	<b>Causes</b>
Shoulder tenderness	<u>Infectious</u> : subdiaphragmatic abscess, septic arthritis (shoulder) <u>Noninfectious</u> : squamous cell carcinoma (Pancoast's tumor), bursitis
Sternal tenderness	<u>Infectious</u> : sternal osteomyelitis (post-open heart surgery) <u>Noninfectious</u> : metastatic carcinoma, pre-leukemia, acute leukemia, myeloproliferative disorder, trauma
Costochondral tenderness	<u>Infectious</u> : costochondritis (Coxsackie B) <u>Noninfectious</u> : trauma, plasmacytoma
Trapezius tenderness	<u>Infectious</u> : subdiaphragmatic abscess <u>Noninfectious</u> : fibromyalgia
Chest wall sinuses	<u>Infectious</u> : TB, actinomycosis, blastomycosis, abscess, M. fortuitum-chelonae (post-breast implant surgery), sternal osteomyelitis (post-sternotomy) <u>Noninfectious</u> : bronchogenic carcinoma
Spontaneous pneumothorax	<u>Infectious</u> : TB, PCP, Legionnaire's disease, lung abscess, pertussis <u>Noninfectious</u> : histiocytosis X (eosinophilic granuloma, Langerhan's cell histiocytosis), osteogenic sarcoma, emphysema, ARDS
Diffuse wheezing	<u>Infectious</u> : RSV, influenza, HMPV, HBOV, C. pneumoniae, M. pneumoniae, acute shistosomiasis (S. mansoni > S. haematobium) <u>Noninfectious</u> : pulmonary emboli, LVF, asthma, Churg-Strauss granulomatosis, carcinoid syndrome, asthmatic bronchitis, angioedema
Chest dullness <i>consolidation</i>	<u>Infectious</u> : bacterial pneumonia, psittacosis, nocardia, Q fever <u>Noninfectious</u> : large cell carcinoma
<i>pleural effusion</i>	<u>Infectious</u> : group A streptococci, tularemia, H. influenzae, 1° TB <u>Noninfectious</u> : Meig's syndrome, pancreatitis, CHF, malignancies, pulmonary embolism

**Table 9.10. Back Abnormalities**

<b>Finding</b>	<b>Causes</b>
Spinal tenderness	<u>Infectious</u> : vertebral osteomyelitis, typhoid fever, TB, brucellosis, SBE <u>Noninfectious</u> : malignancies, multiple myeloma
D'espine's sign	<u>Infectious</u> : bilateral pneumonia (consolidation), TB <u>Noninfectious</u> : sarcoidosis, large cell carcinoma, lymphoma
Unilateral CVA tenderness	<u>Infectious</u> : pyelonephritis, renal/perinephric abscess <u>Noninfectious</u> : trauma

**Table 9.11. Heart Abnormalities**

Finding	Causes
Tachycardia	<p><u>Infectious</u>: myocarditis (Coxsackie, RMSF, typhus, diphtheria, trichinosis, Toxocara canis/cati (VLM), influenza, dengue), gas gangrene</p> <p><u>Noninfectious</u>: hypovolemia, hypoxia, anxiety, hyperthyroidism, MI, pulmonary embolism, CHF, Kawasaki's disease, substance withdrawal, myocarditis (idiopathic giant cell)</p>
Heart block	<p><u>Infectious</u>: Lyme disease, ABE with paravalvular/septal abscess, diphtheria, myocarditis (Coxsackie, influenza, RMSF)</p> <p><u>Noninfectious</u>: acute (inferior) MI, drugs, myocarditis (idiopathic giant cell)</p>
Chronic	<p><u>Infectious</u>: Chagas' disease</p> <p><u>Noninfectious</u>: AV nodal ablation, sarcoidosis, Lev's/Lenegre's disease</p>
Pericardial effusion	<p><u>Infectious</u>: viral pericarditis, TB pericarditis, dengue</p> <p><u>Noninfectious</u>: SLE, uremia, malignancy</p>
Heart murmur	<p><u>Infectious</u>: SBE, 3° syphilis</p> <p><u>Noninfectious</u>: valvular heart disease, severe anemia, Takayasu's arteritis, SLE (Libman-Sacks), endomyocardial fibroelastosis, atrial myxoma, marantic endocarditis</p>

**Table 9.12. Abdominal Abnormalities**

Finding	Causes
Abdominal wall tenderness	<p><u>Infectious</u>: leptospirosis, abdominal wall cellulitis/abscess, trichinosis</p> <p><u>Noninfectious</u>: trauma, rectus sheath hematoma</p>
Abdominal wall sinus tract	<p><u>Infectious</u>: TB, abscess, ameboma, actinomycosis</p> <p><u>Noninfectious</u>: carcinomas</p>
Rose spots	<p><u>Infectious</u>: typhoid fever, shigella, trichinosis, rat bite fever, brucellosis, leptospirosis</p> <p><u>Noninfectious</u>: Campbell de Morgan spots</p>
Abdominal tenderness (2° to pancreatitis)	<p><u>Infectious</u>: Coxsackie virus, echovirus, mumps, ascariasis, EBV, CMV, HBV, HSV, VZV, HIV</p> <p><u>Noninfectious</u>: EtOH, cholelithiasis (+ common pancreatitis duct), trauma, post-ERCP, hyperlipidemia (types I, IV, V), hypertriglyceridemia, TPN, hypercalcemia, SLE, PAN, posterior penetrating duodenal ulcer, drugs (steroids, azathioprine, diuretics, furosemide, thiazides, L-asparaginase, valproic acid, pentamidine)</p>

**Table 9.12. Abdominal Abnormalities** (cont'd)

Finding	Causes
Right upper quadrant tenderness	<p><u>Infectious</u>: cholangitis, cholecystitis, pylephlebitis, splenic flexure diverticulitis, emphysematous cholecystitis (Clostridia sp.), hepatic abscess (amebic, echinococcal, bacterial), right lower lobe CAP, brucellosis, leptospirosis, typhoid fever, viral hepatitis, dengue, malaria</p> <p><u>Noninfectious</u>: burns, post-partum, ptosed right kidney, acute pancreatitis, Kawasaki's disease, cholesterol emboli syndrome, acalculous cholecystitis PAN, SLE, total parenteral nutrition, ceftriaxone (pseudocholelithiasis)</p>
Right upper quadrant tympany	<p><u>Infectious</u>: peritonitis (2° to organ perforation)</p> <p><u>Noninfectious</u>: post-abdominal surgery</p>
Right upper quadrant mass	<p><u>Infectious</u>: bacterial abscess, echinococcal cysts, amebic abscess</p> <p><u>Noninfectious</u>: ptosed right kidney, Reidel's lobe, hepatoma, malignancies, distended gallbladder (Courvoisier's sign)</p>
Right lower quadrant tenderness	<p><u>Infectious</u>: shigella, typhoid fever, typhoidal EBV, typhoidal tularemia, parvovirus B19, TB, ameboma, typhilitis, actinomycetoma, pseudoappendicitis (scarlet fever), Legionnaires' disease, yersinia, campylobacter, measles (pre-eruptive), RMSF, PID, syphilis (luetec crisis), brucellosis</p> <p><u>Noninfectious</u>: appendicitis, RE, ectopic pregnancy, diverticulitis, hyper IgE (Job's) syndrome, DM crisis, porphyria, pancreatitis, SLE</p>
Left upper abdominal quadrant tenderness (splenic tenderness)	<p><u>Infectious</u>: SBE, brucellosis, typhoid fever, malaria, splenic abscess, EBV, CMV, HHV-6</p> <p><u>Noninfectious</u>: splenic infarct</p>
Abdominal wall nodules	<p><u>Infectious</u>: leprosy</p> <p><u>Noninfectious</u>: lipomas, panniculitis, metastatic disease, sarcoidosis</p>
Peribubical purpura (thumbprint sign)	<p><u>Infectious</u>: strongyloides (hyperinfection syndrome)</p> <p><u>Noninfectious</u>: retroperitoneal hemorrhage (Cullen's sign)</p>
Hepatomegaly	<p><u>Infectious</u>: viral hepatitis, bacterial liver abscess, amebic abscess, ehrlichiosis, babesiosis, brucellosis, leptospirosis (Weil's syndrome), arboviral hemorrhagic fevers, typhus, typhoid fever, malaria, Q fever, EBV, CMV, HIV, Castleman's disease (MCD)</p>

**Table 9.12. Abdominal Abnormalities** (cont'd)

Finding	Causes
	<p><u>Noninfectious</u>: alcoholic cirrhosis, cholangiocarcinoma, carcinoma of pancreas, constrictive pericarditis, pericholangitis, veno-occlusive disease, autoimmune hepatitis, <math>\alpha</math>-1 antitrypsin deficiency, cystic fibrosis, fatty liver, hemangiomas, jejunoileal bypass, Reye's syndrome, multiple myeloma, hepatocellular carcinoma, metastatic carcinoma*, parenteral hyperalimentation (TPN)</p>
Splenomegaly	<p><u>Infectious</u>: <b>Mildly enlarged spleen</b>: malaria, kala-azar, dengue, SBE, ehrlichiosis, babesiosis, typhoid fever, typhus, leptospirosis (Weil's syndrome), viral hepatitis, EBV, CMV, relapsing fever, syphilis, toxoplasmosis, psittacosis, brucellosis, Q fever, CSD, schistosomiasis, RE, trypanosomiasis, histoplasmosis, TB, splenic abscess, HIV, hydatid cysts, colorado tick fever; Castleman's disease (MCD), <b>Moderately enlarged spleen</b>: malaria, kala-azar; Castleman's disease (MCD), <b>Massively enlarged spleen</b>: malaria, kala-azar, Castleman's disease (MCD)</p> <p><u>Noninfectious</u>: <b>Mildly enlarged spleen</b>: SLE, sarcoidosis, Felty's syndrome, hemochromatosis, Wilson's disease, Budd-Chiari syndrome, megaloblastic anemia, iron deficiency anemia, systemic mastocytosis, angioblastic lymphadenopathy, splenic cysts, splenic trauma/hemorrhage, histiocytosis X (Langerhan's eosinophilic granuloma), hyperthyroidism, serum sickness, amyloidosis, berylliosis, Kawasaki's disease</p> <p><b>Moderately enlarged spleen</b>: portal hypertension, hemolytic anemias, myeloproliferative disorders, CLL, Gaucher's disease, Niemann-Pick disease;</p> <p><b>Massively enlarged spleen</b>: CML, hairy cell leukemia, lymphoma, myelofibrosis</p>
Hepatosple-nomegaly	<p><u>Infectious</u>: malaria, typhoid fever, toxocara (visceral larval migrans), psittacosis, brucellosis, trypanosomiasis, kala-azar, 2<sup>o</sup> syphilis, EBV, CMV, acute HIV, schistosomiasis, toxoplasmosis, relapsing fever, RMSF, typhus, CSD, histoplasmosis, TB, dengue, babesiosis, ehrlichiosis, Q fever, Castleman's disease (MCD)</p> <p><u>Noninfectious</u>: hypernephroma, CGD, hyper IgD syndrome, sarcoidosis</p>
Ascites	<p><u>Infectious</u>: TB peritonitis, shistosomiasis, filariasis, spontaneous bacterial peritonitis, dengue, Whipple's disease</p> <p><u>Noninfectious</u>: malignancies, Budd-Chiari syndrome, tricuspid regurgitation, constrictive pericarditis, inferior vena cava syndrome, FMF, Henoch-Schönlein purpura, SLE, portal hypertension, pancreatic/bile ascites, CHF, Meig's syndrome</p>

\* Usually from lung, colon, pancreas, kidney, breast, stomach, or esophagus.

**Table 9.13. Lymph Node Abnormalities**

Finding	Causes
Preauricular adenopathy	<u>Infectious</u> : ipsilateral conjunctivitis, tularemia, anterior scalp infections, rat bite fever ( <i>Spirillum minus</i> ) <u>Noninfectious</u> : lymphoma
Occipital adenopathy	<u>Infectious</u> : posterior scalp infections, CSD, rubella <u>Noninfectious</u> : lymphoma
Anterior cervical adenopathy	<u>Infectious</u> : group A streptococcal pharyngitis, viral pharyngitis, mouth/dental infections, TB (scrofula), HHV-6, toxoplasmosis, CSD, brucellosis <u>Noninfectious</u> : head/neck cancer, Kawasaki's disease, lymphoma, SLE, Kikuchi's disease, Rosai-Dorfman disease
Posterior cervical adenopathy <i>unilateral</i>  <i>bilateral</i>	<u>Infectious</u> : toxoplasmosis, African trypanosomiasis (Winterbottom's sign) <u>Noninfectious</u> : posterior scalp infection, Kawasaki's disease, lymphoma, Kikuchi's disease, Rosai-Dorfman disease <u>Infectious</u> : EBV, HHV-6, CMV, Chagas' disease, HAV <u>Noninfectious</u> : lymphoma, Kawasaki's disease, Rosai-Dorfman disease, hyper IgD syndrome, sarcoidosis
Supraclavicular adenopathy	<u>Infectious</u> : TB, CSD <u>Noninfectious</u> : intra-abdominal malignancy (Virchow's sign), Kikuchi's disease
Infraclavicular adenopathy	<u>Infectious</u> : African trypanosomiasis, CSD <u>Noninfectious</u> : lymphoma
Epitrochlear adenopathy	<u>Infectious</u> : 2° syphilis, CSD, brucellosis <u>Noninfectious</u> : sarcoidosis, IVDA
Axillary adenopathy	<u>Infectious</u> : CFS (usually left), CSD, B. malayi, rat bite fever ( <i>Spirillum minus</i> ), brucellosis <u>Noninfectious</u> : lymphoma, CLL, sarcoidosis, Schnitzler's syndrome
Ulcer-node syndromes	<u>Infectious</u> : <b>Ulcer &gt; node</b> : anthrax, rickettsial fevers (except RMSF), sporotrichosis, cutaneous leishmaniasis (new world). <b>Node = ulcer</b> : 1° syphilis, chancroid, tularemia (ulceroglandular), rat bite fever ( <i>Spirillum minus</i> ). <b>Node &gt; ulcer</b> : syphilis, LGV, chancroid, HSV-2 <u>Noninfectious</u> : lymphoma
Periumbilical nodule	<u>Infectious</u> : intra-abdominal infection <u>Noninfectious</u> : malignancy (Sister Joseph's sign)

**Table 9.13. Lymph Node Abnormalities** (cont'd)

Finding	Causes
Inguinal adenopathy <i>unilateral</i>	<u>Infectious</u> : lower extremity infections, bubonic plague, rat bite fever ( <i>Spirillum minus</i> ), filariasis, leprosy, tularemia, CSD <u>Noninfectious</u> : intra-abdominal malignancy, bilateral lower extremity infection, IVDA
<i>bilateral</i>	<u>Infectious</u> : any infection causing generalized adenopathy, B. malayi, HSV-2, syphilis, LGV <u>Noninfectious</u> : any disorder causing generalized adenopathy, Schnitzler's syndrome
Generalized lymphadenopathy	<u>Infectious</u> : TB, EBV, HHV-6, CMV, rubella, measles, toxoplasmosis, CSD, LGV, brucellosis, group A streptococci, 2° syphilis, HIV, kala-azar (African), trypanosomiasis, arboviral hemorrhagic fevers, Whipple's disease, Castleman's disease (MCD) <u>Noninfectious</u> : SLE, RA, adult Still's disease, pseudolymphoma (Dilantin), Kikuchi's disease, Gaucher's disease, sarcoidosis, serum sickness, hyperthyroidism, ALL, CLL, Kimura's disease

**Table 9.14. Genitourinary Abnormalities**

Finding	Causes
Epididymoorchitis	<u>Infectious</u> : mumps, TB, blastomycosis, melioidosis, brucellosis, leptospirosis, EBV, W. bancrofti, Coxsackie A/B, S. hematobium, LCM, GC, C. trachomatis (young adults), P. aeruginosa (elderly adults), Colorado tick fever, rat bite fever ( <i>Spirillum minus</i> ), relapsing fever <u>Noninfectious</u> : lymphoma, SLE, PAN, sarcoidosis, FMF, TNF-receptor-1-associated periodic syndrome (TRAPS), trauma, torsion, malignancy
Scrotal enlargement	<u>Infectious</u> : mumps, W. bancrofti (not B. malayi), Fournier's gangrene <u>Noninfectious</u> : hydrocele, testicular torsion
Groin mass	<u>Infectious</u> : onchocerciasis (hanging groins), TB, W. bancrofti, shistosomiasis <u>Noninfectious</u> : lymphoma
Perirectal fistula	<u>Infectious</u> : peri-rectal abscess, actinomycosis, LGV <u>Noninfectious</u> : RE, malignancies
Perirectal ulcer	<u>Infectious</u> : HSV, amebiasis cutis, 1° syphilis <u>Noninfectious</u> : malignancy
Genital ulcers	<u>Infectious</u> : Syphilis (1°), HSV, Chancroid, LCV, Chikungunya fever, TB <u>Noninfectious</u> : Bìchet's syndrome, RE, malignancies
Prostate enlargement/tenderness	<u>Infectious</u> : prostatitis, prostatic abscess* <u>Noninfectious</u> : BPH

\* Acute: aerobic GNB, VSE/VRE, *Ureoplasma urealyticum*, *C. trachomatis*, *B. fragilis* (post-transrectal biopsy).  
Chronic: TB, histoplasmosis, blastomycosis, aspergillus, candida, cryptococcus, melioidosis.



**Table 9.15. Extremity Abnormalities**

Finding	Causes
Digital gangrene	<u>Infectious</u> : SBE, S. aureus bacteremia/ABE (emboli), meningococcemia, RMSF, typhus <u>Noninfectious</u> : SLE, vasculitis, peripheral vascular disease
Splinter hemorrhages	<u>Infectious</u> : SBE, ABE, trichinosis <u>Noninfectious</u> : trauma, atrial myxoma, acute leukemia, RA, scurvy, mitral stenosis, severe anemia
Clubbing	<u>Infectious</u> : SBE, lung abscess, TB <u>Noninfectious</u> : ulcerative colitis (UC), Crohn's disease (RE), cirrhosis, cyanotic congenital heart disease, bronchogenic carcinoma, PBC, celiac disease, hyperthyroidism, hyperparathyroidism, bronchiectasis, hereditary
Dactylitis	<u>Infectious</u> : kala-azar, 2° syphilis, TB <u>Noninfectious</u> : sarcoidosis, sickle cell disease, gout, psoriatic arthritis
Tender fingertips	<u>Infectious</u> : SBE, typhoid fever <u>Noninfectious</u> : SLE, vasculitis, radial artery occlusion
Lymphangitis	<u>Infectious</u> : group A streptococci, B. malayi, onchocerciasis, melioidosis <u>Noninfectious</u> : IVDA
Nodular lymphangitis	<u>Infectious</u> : sporotrichosis, atypical TB, Erysipelothrix rhusiopathiae, kala-azar, coccidiomycosis, histoplasmosis, blastomycosis, nocardia, Pseudoallescheria boydii, cryptococcus, anthrax, group A streptococci, tularemia <u>Noninfectious</u> : ganglion cyst (wrist/hand only)
Painless purple palm/sole lesions	<u>Infectious</u> : ABE (Janeway lesions) <u>Noninfectious</u> : trauma
Carpal tunnel syndrome	<u>Infectious</u> : TB, leprosy <u>Noninfectious</u> : cirrhosis, RA, scleroderma, SLE, DM, hypothyroidism, sarcoidosis, multiple myeloma, amyloidosis
Verrucous hand/arm lesions	<u>Infectious</u> : TB, leprosy, syphilis, sporotrichosis, kala-azar, bartonellosis (verruca peruana) <u>Noninfectious</u> : squamous cell carcinoma, sarcoidosis
Edema of the dorsum of hand/foot	<u>Infectious</u> : RMSF, TSS, loiasis, dengue fever, chikungunya fever <u>Noninfectious</u> : trauma, PMR, Kawasaki's disease

**Table 9.15. Extremity Abnormalities** (cont'd)

Finding	Causes
Wrist swelling	<u>Infectious</u> : loiasis, septic arthritis, parvovirus B <sub>19</sub> , rubella, chikungunya fever <u>Noninfectious</u> : RA
Arthritis (septic/reactive)	<u>Infectious</u> : septic arthritis, rat bite fever ( <i>Streptobacillus moniliformis</i> ), Lyme disease, LGV, brucellosis, GC, parvovirus B19, shigella, yersinia, salmonella, campylobacter, <i>Clostridium difficile</i> , mumps, rubella, dengue fever, HIV, HBV, Whipple's disease <u>Noninfectious</u> : FMF, RA, pseudogout, SLE, hyper IgD syndrome, lymphoma, Reiter's syndrome, post-streptococcal RF, Poncet's disease, JRA
Papular axillary lesions	<u>Infectious</u> : hydraadenitis suppurativa, 2° syphilis, yaws, blastomycosis <u>Noninfectious</u> : chronic contact dermatitis, acanthosis nigricans, Fox-Fordyce disease, seborrheic dermatitis
Tenosynovitis	<u>Infectious</u> : GC (acute), TB (chronic), atypical TB (acute/chronic) <u>Noninfectious</u> : rheumatic diseases
Thigh tenderness (bilateral, anterior)	<u>Infectious</u> : bacteremia (Louria's sign), leptospirosis, ehrlichiosis, endocarditis, parvovirus B <sub>19</sub> <u>Noninfectious</u> : myositis, vasculitis, sickle cell crisis
Calf tenderness	<u>Infectious</u> : RMSF <u>Noninfectious</u> : myositis
Tender muscles	<u>Infectious</u> : trichinosis <u>Noninfectious</u> : myositis
Thrombophlebitis	<u>Infectious</u> : psittacosis, campylobacter <u>Noninfectious</u> : Behçet's disease, malignancy
Verrucous foot/leg lesions	<u>Infectious</u> : TB, cutaneous leishmaniasis, paracoccidomycosis, sporotrichosis, leprosy, 2° syphilis, mycetoma <u>Noninfectious</u> : lichen planus, squamous cell carcinoma
Foot/leg ulcers	<u>Infectious</u> : <i>M. ulcerans</i> (Buruli ulcer may be anywhere), yaws, cutaneous diphtheria, cutaneous leishmaniasis, TB, rat bite fever ( <i>Spirillum minus</i> ) <u>Noninfectious</u> : sickle cell (medial malleolar ulcers), DM (only sole of foot/ between toes), peripheral vascular disease
Leg edema <i>unilateral</i>	<u>Infectious</u> : onchocerciasis, <i>B. malayi</i> (below knee) <u>Noninfectious</u> : malignancy, Milroy's disease

**Table 9.15. Extremity Abnormalities** (cont'd)

Finding	Causes
<i>bilateral</i>	<u>Infectious</u> : elephantiasis (chronic recurrent erysipelas), Chagas' disease <u>Noninfectious</u> : lymphatic obstruction (2° to abdominal/pelvic malignancy), congenital yellow nail syndrome
Nodular arm/leg lesions	<u>Infectious</u> : TB (Bazin's erythema induratum), filariasis, sporotrichosis, cutaneous leishmaniasis, atypical TB, leprosy, HIV <u>Noninfectious</u> : erythema nodosum, PAN, thrombophlebitis, panniculitis, nodular vasculitis, Wegener's granulomatosis, Sweet's syndrome, myiasis
Palpable purpura	<u>Infectious</u> : meningococemia, RMSF <u>Noninfectious</u> : vasculitis, cryoglobulinemia, Gardner-Diamond syndrome, Sweet's syndrome
Eschar	<u>Infectious</u> : ecthyma gangrenosum, typhus, rickettsial spotted fevers (except RMSF, rickettsial pox), anthrax, cutaneous diphtheria, African tick bite fever (multiple eschars) <u>Noninfectious</u> : burns, drugs, recluse spider bite
Scleroderma (wood hard skin lesions)	<u>Infectious</u> : actinomycosis <u>Noninfectious</u> : malignancies, DM
Hyperpigmented shins	<u>Infectious</u> : onchocerciasis <u>Noninfectious</u> : DM (dermopathy)
Cutaneous cold abscesses	<u>Infectious</u> : TB, atypical TB <u>Noninfectious</u> : hyper IgE (Job's) syndrome, CGD
Purple nodules	<u>Infectious</u> : disseminated cryptococcus, aspergillus, candida, trypanosomiasis, Kaposi's sarcoma, HIV <u>Noninfectious</u> : leukemia, lymphoma, melanoma
Painful leg nodules	<u>Infectious</u> : Kaposi's sarcoma <u>Noninfectious</u> : erythema nodosum, superficial thrombophlebitis, PAN, panniculitis, osteogenic sarcoma
Migratory rashes	<u>Infectious</u> : hookworms, dracunculiasis, loiasis, gnathostomiasis, strongyloidiasis, sparganosis <u>Noninfectious</u> : myiasis
Rash of palms/soles	<u>Infectious</u> : syphilis, RMSF, EBV, scarlet fever, echo 9, smallpox, monkeypox, chickenpox, rat bite fever (Streptobacillus moniliformis), HFM, orf, E. multiforme (M. pneumoniae, HSV) <u>Noninfectious</u> : drug rashes, E. multiforme (drugs)

**Table 9.15. Extremity Abnormalities** (cont'd)

Finding	Causes
Rash (purpuric) of hands/feet	<u>Infectious</u> : parvovirus B19, VZV, EBV, CMV, HHV-6, HHV-7, HBV, Coxsackie, rubella <u>Noninfectious</u> : trauma
Erythema nodosum	<u>Infectious</u> : TB, group A streptococci, EBV, LGV, psittacosis, coccidiomycosis, blastomycosis, histoplasmosis, CSD, yersinia, campylobacter <u>Noninfectious</u> : UC, RE, drugs, sarcoidosis, lymphoma, SLE
Erythema multiforme	<u>Infectious</u> : HSV, M. pneumoniae, Coxsackie B <u>Noninfectious</u> : drugs
Plantar hyperkeratoses	<u>Infectious</u> : 3° syphilis, yaws, pinta, HIV, tungiasis <u>Noninfectious</u> : Reiter's syndrome, arsenic
Desquamation of hands/feet	<u>Infectious</u> : erysipelas, scarlet fever, TSS, severe infections, yaws, leptospirosis, influenza, measles, arboviral hemorrhagic fevers, chikungunya fever, dengue fever (palms/soles) <u>Noninfectious</u> : Kawasaki's disease, post-edematous states, radiation therapy, vitamin A excess, pellagra, drugs

**Table 9.16. Neurological Abnormalities**

Finding	Causes
Mental confusion/encephalopathy (acute)	<u>Infectious</u> : legionnaire's disease, HSV, HHV-6, RMSF, scrub typhus (R. tsutsugamushi) listeria, mycoplasma meningoencephalitis, amebic meningitis, trichinosis, brain abscess, anthrax, brucellosis, SBE, ABE, HIV, viral encephalitis, CSD, Whipple's disease, Q fever, Colorado tick fever, chikungunya fever, melioidosis <u>Noninfectious</u> : Wernicke's encephalopathy, toxic/metabolic disorders, Behçet's syndrome, SLE, alcoholism, drugs, CHF, chronic renal failure, hepatic encephalopathy, brain tumor, CNS metastases, meningeal carcinomatosis, celiac disease, cefepime, ciprofloxacin > levofloxacin
Nuchal rigidity	<u>Infectious</u> : meningitis (bacterial, fungal, TB, viral) <u>Noninfectious</u> : meningismus, cervical arthritis
General muscle rigidity	<u>Infectious</u> : trichinosis, tetanus, rabies, viral encephalitis, brucellosis <u>Noninfectious</u> : malignancies, malignant neuroleptic syndrome, strychnine poisoning, Parkinson's disease
Transient deafness	<u>Infectious</u> : RMSF, mediterranean spotted fever, murine typhus, S. pneumoniae ABM, S. suis ABM, H. influenzae ABM, mumps, measles, VZV (Ramsey-Hunt Syndrome), EEE, congenital syphilis, brucellosis, Lassa fever <u>Noninfectious</u> : trauma, Susac's syndrome

**Table 9.16. Neurological Abnormalities** (cont'd)

Finding	Causes
Cranial nerve (CN) abnormalities <i>unilateral</i>	<u>Infectious:</u> 6 <sup>th</sup> CN palsy (TB meningitis, N. meningitidis), 7 <sup>th</sup> CN palsy (see Bell's palsy), TBRF, 8 <sup>th</sup> CN palsy (N. meningitidis) <u>Noninfectious:</u> 7 <sup>th</sup> CN palsy (neurosarcoidosis, meningeal carcinomatosis), 2 <sup>nd</sup> , 6 <sup>th</sup> CN palsy (Wegener's granulomatosis)
<i>bilateral</i>	<u>Infectious:</u> 6 <sup>th</sup> CN palsy (TB) <u>Noninfectious:</u> meningeal carcinomatosis
Optic nerve atrophy	<u>Infectious:</u> 3 <sup>o</sup> syphilis, TB, toxoplasmosis, mumps, measles, rubella <u>Noninfectious:</u> sickle cell disease, severe anemia, polycythemia vera, drugs, sarcoidosis, TA, Behçet's syndrome, SLE, PAN, MS, glaucoma
Bell's Palsy	<u>Infectious:</u> Lyme disease, HSV-1, VZV (Ramsay-Hunt syndrome), HIV, St.LE, CSD, mumps, EBV infectious mononucleosis, Mycoplasma pneumoniae, syphilis, otitis media, bacterial meningitis, leprosy <u>Noninfectious:</u> sarcoidosis, multiple sclerosis, Guillain-Barre syndrome, B-cell lymphoma, Wegener's granulomatosis, pontine lesions (infarct, tumors, demyelinate lesions), tumor compressing the facial nerve, sphenoid ridge meningiomas, acoustic neuromas, neurofibromatosis (type 2), fibrous dysplasia, osteopetrosis, Paget's disease, scleroderma, Melkersson-Rosenthal syndrome, trauma, drugs (vincristine, HBV vaccination), idiopathic
Anisocoria	<u>Infectious:</u> TB, 3 <sup>o</sup> syphilis, VZV, meningitis, encephalitis, botulism, diphtheria <u>Noninfectious:</u> DM, toxins, cavernous sinus thrombosis, glaucoma, brain tumor, intracranial aneurysm
Papilledema	<u>Infectious:</u> meningitis (bacterial, fungal, TB, viral, etc.) <u>Noninfectious:</u> pseudotumor cerebri, hypercarbia, brain tumor, DM, subarachnoid bleed, SLE, drugs, central retinal artery occlusion, cavernous sinus thrombosis, hypertensive encephalopathy
Mononeuritis multiplex	<u>Infectious:</u> leprosy, Lyme disease, HIV <u>Noninfectious:</u> DM, amyloidosis, sarcoidosis, lymphomatoid granulomatosis, vasculitis
Transverse myelitis	<u>Infectious:</u> HIV, HTLV-1, EBV, CMV, VZV, HSV, polio, rabies, TB, EV 71, epidural abscess, typhus, brucellosis, shistosomiasis, Lyme disease, syphilis, toxoplasmosis, M. pneumoniae <u>Noninfectious:</u> MS, malignancies, vaccines, SLE, sarcoidosis
Guillain-Barré syndrome	<u>Infectious:</u> influenza, Campylobacter jejuni, CMV, EBV, VZV, M. pneumoniae, EV 71, brucellosis <u>Noninfectious:</u> influenza vaccine
Flaccid paralysis (acute)	<u>Infectious:</u> polio, WNE, CMV, VZV, rabies, botulism, JE, EV 71, EV D68, Powassan encephalitis (PE), chikungunya fever, brucellosis <u>Noninfectious:</u> CVA, tick bite paralysis, hypocalcemic periodic paralysis

**Table 9.16. Neurological Abnormalities** (cont'd)

Finding	Causes
Hemiplegia/hemiparesis	<u>Infectious</u> : SBE, subdural empyema, brain abscess, JE, brucellosis, post-VZV <u>Noninfectious</u> : TIA, CVA, malignancies, CNS vasculitis, migraine, GCA/TA, SLE, neurosarcoidosis, birth control pills, protein C/S, factor V Leiden deficiency

**Table 9.17. WBC Abnormalities**

Finding	Causes
Leukocytosis	<u>Infectious</u> : most acute infections <u>Noninfectious</u> : most acute noninfectious disease disorders, any major stress, steroids, drug fever, daptomycin
Leukopenia	<u>Infectious</u> : miliary TB, typhoid/enteric fever, malaria, babesiosis dengue, chikungunya fever, tularemia, brucellosis, kala-azar, psittacosis, viral hepatitis, EBV, CMV, HHV-6, VZV, influenza, Colorado tick fever, Campylobacter histoplasmosis, relapsing fever, WNE, VEE, ehrlichiosis/anaplasmosis, SARS <u>Noninfectious</u> : drugs, pre/acute leukemias, Felty's syndrome, Gaucher's disease, splenomegaly, pernicious anemia, SLE, cyclic neutropenia, severe combined immunodeficiency disease (SCID), Chediak-Higashi syndrome, Kikuchi's disease, sarcoidosis
Relative lymphocytosis	<u>Infectious</u> : Whipple's disease, acute infection (convalescence), TB, brucellosis, pertussis, tularemia, 2° syphilis, histoplasmosis, EBV, CMV, HHV-6, mumps, viral hepatitis, rubella, VZV, kala-azar, toxoplasmosis, RMSF, chikungunya fever, typhoid/enteric fever (non-S. typhi > S. typhi), MERS <u>Noninfectious</u> : ALL, CLL, lymphomas, carcinomas, multiple myeloma, RA, Hashimoto's thyroiditis, myxedema, adrenal insufficiency, thyrotoxicosis, vasculitis, Dilantin (DPH), p-aminosalicylic acid (PAS), serum sickness
Relative lymphopenia	<u>Infectious</u> : CMV, HHV-6, HHV-8, HIV, miliary TB, Legionella, typhoid/enteric fever (S. typhi > non-S. typhi), Q fever, brucellosis, SARS, malaria, babesiosis, influenza (human, avian, swine), RMSF, histoplasmosis, dengue fever, chikungunya fever, ehrlichiosis, parvovirus B19, HPS, WNE, viral hepatitis (early), Whipple's disease, MERS <u>Noninfectious</u> : cytotoxic drugs, steroids, sarcoidosis, SLE, lymphoma, RA, radiation, Wiskott-Aldrich syndrome, severe combine immunodeficiency disease (SCID), common variable immune deficiency (CVID), Di George's syndrome, Nezelof's syndrome, intestinal lymphangiectasia, ataxia-telangiectasia, constrictive pericarditis, tricuspid regurgitation, Kawasaki's disease, idiopathic CD <sub>4</sub> cytopenia, acute/chronic renal failure, hemodialysis, myasthenia gravis, celiac disease, cirrhosis, coronary bypass, Wegener's granulomatosis, CHF, acute pancreatitis, carcinomas (terminal)

**Table 9.17. WBC Abnormalities** (cont'd)

Finding	Causes
Monocytosis	<p><u>Infectious</u>: TB, SBE, RMSF, diphtheria, histoplasmosis, brucellosis, pertussis, kala-azar, 2<sup>o</sup> syphilis, malaria, typhoid/enteric fever, babesiosis, EBV, CMV, influenza A, recovery from chronic infection</p> <p><u>Noninfectious</u>: sarcoidosis, myeloproliferative disorders (↑ pre-AML), lymphomas, Gaucher's disease, RE, UC, celiac disease, RA, SLE, PAN, TA, hyposplenism, post-splenectomy</p>
Atypical lymphocytes†	<p><u>Infectious</u>: EBV*, CMV*, HHV-6, Castleman's disease (MCD), viral hepatitis, mumps, measles, rubella, VZV, toxoplasmosis, brucellosis, HSV, arboviral hemorrhagic fevers, malaria, dengue, babesiosis, ehrlichiosis/anaplasmosis</p> <p><u>Noninfectious</u>: drug fever, Kikuchi's disease</p>
Immunoblasts	<p><u>Infectious</u>: HPS</p> <p><u>Noninfectious</u>: lymphomas</p>
Eosinophilia	<p><u>Infectious</u>: trichinosis, echinococcosis, fascioliasis, paragonimiasis, taeniasis, Strongyloides stercoralis, hookworm, filariasis, schistosomiasis, Toxocara canis/cati (VLM), histoplasmosis, coccidioidomycosis, filariasis ascariasis, gnathostomiasis, angiostrongyliasis, cysticercosis, Isospora belli, Dientamoeba fragilis, scarlet fever</p> <p><u>Noninfectious</u>: drug fever, asthma, dermatitis herpetiformis, pemphigus vulgaris, eczema, hay fever/allergic rhinitis, psoriasis, atopic dermatitis, mycosis fungoides, myeloproliferative disorders (MPDs), polycythemia vera, CML, eosinophilic leukemia, eosinophilic gastritis, acute leukemias, sickle cell anemia, lymphomas, malignancies, Churg-Strauss granulomatosis, urticaria, hyper IgE syndrome (Job's syndrome), Sweet's syndrome, bronchopulmonary aspergillosis (BPA), angioneurotic edema, serum sickness, eosinophilia-myalgia syndrome, dermatomyositis, PAN, allergic vasculitis, Loffler's syndrome, pulmonary infiltrates eosinophilia (PIE) syndrome, Loffler's endocarditis, Addison's disease, sarcoidosis, Wegener's granulomatosis, Goodpasture's syndrome, UC, RE, Wiskott-Aldrich syndrome, IgA deficiency, allergic vasculitis, Kimura's disease, peritoneal dialysis, radiation</p>
Eosmopenin	<p><u>Infectious</u>: typhoid fever</p> <p><u>Noninfectious</u>: steroids, immunosuppressive</p>
Basophilia	<p><u>Infectious</u>: smallpox, chickenpox (VZV)</p> <p><u>Noninfectious</u>: pre-leukemias, acute leukemias, lymphomas, myeloproliferative disorders (MPDs), post-splenectomy</p>
WBC inclusions	<p><u>Infectious</u>: ehrlichiosis/anaplasmosis (morula: HGA &gt; HME)</p> <p><u>Noninfectious</u>: staining artifacts</p>

\* May have &gt; 20% atypical lymphocytes.

† Not seen with malignancies. Abnormal lymphocytes (morphologically monotonous) seen with leukemias.

**Table 9.18. RBC Abnormalities**

Finding	Causes
Schistocytes (microangiopathic hemolytic anemia)	<u>Infectious</u> : meningococcemia (DIC) <u>Noninfectious</u> : DIC (due to any cause), TTP, hemolytic uremic syndrome (HUS), "Waring Blender" syndrome (prosthetic valve), malignant hypertension
Spherocytes	<u>Infectious</u> : gas gangrene, Castleman's disease (MCD) <u>Noninfectious</u> : autoimmune hemolytic anemias, hereditary spherocytosis, severe transfusion reactions, severe burns, cirrhosis
Target cells	<u>Infectious</u> : none <u>Noninfectious</u> : post-splenectomy, iron deficiency anemia, cirrhosis, hemoglobin S or C, thalassemia
RBC inclusions	<u>Infectious</u> : malaria (ring forms, RBC pigment), babesiosis (ring forms, no RBC pigment, "Maltese crosses"/tetrads) <u>Noninfectious</u> : Cabot's rings (severe hemolytic anemia, pernicious anemia), Heinz bodies (GGPD deficiency, drug induced, hereditary anemias), Pappenheimer bodies (thalassemia, sideroblastic anemias, lead poisoning), artifacts
Howell-Jolly bodies <sup>†</sup>	<u>Infectious</u> : fulminant pneumococcal sepsis <u>Noninfectious</u> : asplenia/hyposplenism*, congenital asplenia, splenectomy, splenic infarcts, splenic neoplasms, megaloblastic anemias, thalassemia "Maltese Crosses", steroids
Hemophagocytosis	<u>Infectious</u> : HIV, HSV, EBV, CMV, adenovirus, parvovirus B19, malaria, babesiosis, toxoplasmosis, kala-azar, histoplasmosis, cryptococcosis, disseminated candidiasis, typhoid/enteric fever, syphilis, listeria, SBE, Q fever, leprosy, TB, brucellosis, Castleman's disease (MCD) <u>Noninfectious</u> : histiocytosis X (eosinophilic granuloma, Langerhan's cell histiocytosis), myeloproliferative disorders (MPDs), SLE, sarcoidosis, RA, lymphomas, acute leukemias, multiple myeloma, Chediak-Higashi syndrome, Dilantin (DPH), familial hemophagocytic histiocytosis
Anemia (acute)	<u>Infectious</u> : Oroya fever, gas gangrene, malaria, babesiosis, CMV <u>Noninfectious</u> : ITP, "Waring blender" syndrome (prosthetic heart valve), hemorrhagic/necrotic pancreatitis, hemorrhage, drug induced

\* Chronic alcoholism, amyloidosis, chronic active hepatitis, IgA deficiency, intestinal lymphangectasia, myeloproliferative disorders, Waldenstrom's macroglobulinemia, NHL, celiac disease, RA, UC, thyroiditis, systemic mastocytosis, sickle cell disease, Fanconi's syndrome, Sezary's syndrome.

† Also "pitted/pocked" RBCs.



**Table 9.19. Platelet Abnormalities**

Finding	Causes
Thrombocytopenia	<p><b>Infectious:</b> acute/severe bacterial infections, measles, rubella, dengue, arboviral, hemorrhagic fevers, EBV, CMV, VZV, mumps, influenza (human, avian, swine), typhus, RMSF, WNE, ehrlichiosis, diphtheria, malaria, babesiosis, trypanosomiasis, TSS, histoplasmosis, kala-azar, HIV, miliary TB, relapsing fever, HPS, brucellosis, Campylobacter, SARS, MERS</p> <p><b>Noninfectious:</b> drugs, DIC, fat emboli syndrome, TTP, ITP, hemolytic uremic syndrome (HUS), pre/acute/leukemias, lymphomas, carcinomas, myeloproliferative disorders (MPDs), multiple myeloma, Gaucher's disease, cirrhosis, hemodialysis</p>
Thrombocytosis	<p><b>Infectious:</b> TB, chronic infections (e.g., osteomyelitis, abscess), SBE, Q fever, M. pneumoniae</p> <p><b>Noninfectious:</b> Malignancies, myeloproliferative disorders, post-splenectomy, lymphomas, Wegener's granulomatosis, vasculitis, TA, PAN, Kawasaki's disease, anemia (nemolytic, iron deficiency) fosfomycin, miconazole, ceftriaxone, oral cephalosporins, carbapenems, <math>\beta</math>-lactam/<math>\beta</math>-lactamase inhibitor combinations, iron therapy</p>

**Table 9.20. Pancytopenia**

Finding	Causes
Pancytopenia	<p><b>Infectious:</b> miliary TB, brucellosis, histoplasmosis, ehrlichiosis/anaplasmosis, parvovirus B19, HBV, CMV, EBV, HIV</p> <p><b>Noninfectious:</b> myeloproliferative disorders (MPDs), drugs, malignancies, Chediak-Higashi syndrome, megaloblastic anemias, Gaucher's disease, hypersplenism, sarcoidosis, SLE, paroxysmal nocturnal hemoglobinuria (PNH), myelophistic anemias, leukemias, lymphoma</p>

**Table 9.21. Serum Test Abnormalities**

Finding	Causes
Erythrocyte Sedimentation Rate (ESR)	<p><b>Infectious:</b> SBE, osteomyelitis, abscess, CAP (Legionnaire's disease, Q fever, S. pneumoniae), non-pulmonary TB, babesiosis</p> <p><b>Noninfectious:</b> malignancies, rheumatic disorders, vasculitis, drug fever, PMR/GCA, uremia/chronic renal failure, cirrhosis, Kawasaki's disease, Sweet's syndrome, pulmonary emboli/infarction</p>
<i>highly elevated</i> <i>(<math>\geq 100</math> mm/hr)</i> <i>low/subnormal</i>	<p><b>Infectious:</b> trichinosis, CFS, ehrlichiosis/anaplasmosis</p> <p><b>Noninfectious:</b> severe anemia, cachexia, massive hepatic necrosis, DIC, polycythemia vera, sickle cell anemia, CHF</p>
SPEP (polyclonal gammopathy)	<p><b>Infectious:</b> HIV, malaria, kala-azar, LGV, rat bite fever, Toxocara canis/cati (VLM), Q fever (chronic), Castleman's disease (MCD)</p> <p><b>Noninfectious:</b> SLE, PAN, cirrhosis, CAH, sarcoidosis, atrial myxoma, Takayasu's arteritis, Rosai-Dorfman disease</p>

**Table 9.21. Serum Test Abnormalities** (cont'd)

Finding	Causes
SPEP (monoclonal gammopathy)	<p><u>Infectious:</u> CMV, kala-azar, typhoid fever, Castleman's disease (MCD)</p> <p><u>Noninfectious:</u> multiple myeloma, Waldenström's macroglobulinemia, Schnitzler's syndrome, MGUS</p>
<p>↑ ferritin levels<sup>†</sup></p> <p><i>acute</i></p> <p><i>chronic</i></p>	<p><u>Infectious:</u> legionnaires' disease, WNE, EBV, CMV, toxoplasmosis, MSSA/MRSA ABE, malaria, babesiosis, ehrlichiosis/anaplasmosis, arboviral hemorrhagic fevers</p> <p><u>Noninfectious:</u> Kawasaki's disease, Rosai-Dorfman disease, hemophagocytic syndrome (due to any cause)</p> <hr/> <p><u>Infectious:</u> HIV, TB, filariasis</p> <p><u>Noninfectious:</u> Malignancies (preleukemias, lymphomas, multiple myeloma, hepatomas, liver/CNS metastases), myeloproliferative disorders (MPDs), RA, adult Still's disease, SLE, TA, Kawasaki's disease, chronic renal failure, hemodialysis, liver transplants, hemochromatosis, cirrhosis, <math>\alpha</math>-1 antitrypsin deficiency, CAH, (MAS), chronic HCV, cholestatic jaundice, macrophage activation syndrome sickle cell anemia, multiple blood transfusions, anemia of chronic disease</p>
↑ cold agglutinins	<p><u>Infectious:</u> Mycoplasma pneumoniae, EBV, CMV, mumps, measles, malaria, Coxsackie, Q fever, HIV, HCV, influenza, adenovirus, trypanosomiasis</p> <p><u>Noninfectious:</u> lymphomas, CLL, CML, multiple myeloma, Waldenström's macroglobulinemia, cold agglutinin disease, paroxysmal nocturnal hemoglobinuria (PNH), SLE, Rosai-Dorfman disease</p>
Monospot test (false +)	<p><u>Infectious:</u> CMV, HSV, HHV-6, malaria, dengue, babesiosis, toxoplasmosis, brucellosis, Mediterranean spotted fever (MSF), rubella, mumps, viral hepatitis, Lyme disease, HIV</p> <p><u>Noninfectious:</u> SLE, sarcoidosis, AML, drugs, idiopathic</p>
↑ Lactate dehydrogenase (LDH)	<p><u>Infectious:</u> malaria, babesiosis, ehrlichiosis/anaplasmosis, viridans streptococcal SBE, amebic liver abscess, PCP, histoplasmosis, disseminated TB, toxoplasmosis, HPS, oroya fever, gas gangrene, viral myocarditis, rubella, measles, viral hepatitis, CAP/NP, dengue, arboviral hemorrhagic fevers, trichinosis, SARS, MERS, adenovirus, CMV, EBV, influenza (human, avian, swine)</p> <p><u>Noninfectious:</u> hemolyzed blood, hemolytic anemia, malignancies, pernicious anemia, megablatic anemia, pulmonary emboli, MI, renal infarction, muscle injury, liver injury, DIC, SLE pneumonitis, adult Still's disease, "Waring blender" syndrome (prosthetic heart valve), hemorrhagic/necrotic pancreatitis</p>

**Table 9.21. Serum Test Abnormalities** (cont'd)

Finding	Causes																		
↑ Procalcitonin levels (PCT)	<p><u>Infectious</u>: bacterial pneumonias (CAP, NHAP, NP), bacteremias (gram – &gt; gram +), TB, ABM, fungal pneumonias (except PCP), viral hepatitis, toxoplasmosis, osteomyelitis, SBE*, malaria (<i>P. falciparum</i>)</p> <p><u>Noninfectious</u>: renal insufficiency, alcoholic hepatitis, lung cancer (small cell), thyroid cancer, surgery, trauma, burns, cardiogenic shock, Goodpasture's syndrome shock, GVHD, peritoneal dialysis (PD), hypotension, hemorrhagic/necrotic pancreatitis, normal variant (elderly) BMT, febrile neutropenia, drug fever, immunosuppression/steroids, OKT<sub>3</sub> therapy</p>																		
↑ Lactate	<p><u>Infections</u>: sepsis (bacterial), septic shock, malaria, cholera</p> <p><u>Noninfectious</u>: cardiogenic shock, hypovolemic shock, hypoperfusion, hypoxia, respiratory insufficiency/failure, renal insufficiency, hepatic insufficiency, hemorrhage, Ringer's lactate, metabolic acidosis (↑ anion gap), seizures, pulmonary emboli, diabetes mellitus, malignancies (leukemias, lymphomas, solid tumors), drugs (salicylates, methanol, ethylene glycol, INH, cyanide, metformin, acetaminophen, lactulose, ethanol, theophylline, niacin, cocaine, methemoglobinemia, severe anemia, carbon monoxide, short bowel syndrome (jejunoileal bypass))</p>																		
↑ Teichoic acid antibody titer (≥1:4)	<p><u>Infections</u>: MSSA/MRSA ABE, MSSA/MRSA abscesses, MSSA/MRSA osteomyelitis (chronic/acute)</p> <p><u>Noninfectious</u>: none</p>																		
Fingal serum tests	<p>Candida Aspergillus Histoplasmos Blastomycosis PCP Cryptococcus Fusaria Zygomycetes</p> <table border="0" data-bbox="83 838 948 961"> <tr> <td data-bbox="83 838 270 900">β 1, 3 D–glucan<sup>§</sup> (BG)</td> <td data-bbox="270 838 327 900">+</td> <td data-bbox="327 838 384 900">+</td> <td data-bbox="384 838 441 900">+</td> <td data-bbox="441 838 498 900">–</td> <td data-bbox="498 838 555 900">+</td> <td data-bbox="555 838 612 900">–</td> <td data-bbox="612 838 669 900">+</td> <td data-bbox="669 838 726 900">–</td> </tr> <tr> <td data-bbox="83 900 270 961">Aspergillus galactomannan (AG)</td> <td data-bbox="270 900 327 961">–</td> <td data-bbox="327 900 384 961">+</td> <td data-bbox="384 900 441 961">+</td> <td data-bbox="441 900 498 961">–</td> <td data-bbox="498 900 555 961">–</td> <td data-bbox="555 900 612 961">–</td> <td data-bbox="612 900 669 961">+</td> <td data-bbox="669 900 726 961">–</td> </tr> </table>	β 1, 3 D–glucan <sup>§</sup> (BG)	+	+	+	–	+	–	+	–	Aspergillus galactomannan (AG)	–	+	+	–	–	–	+	–
β 1, 3 D–glucan <sup>§</sup> (BG)	+	+	+	–	+	–	+	–											
Aspergillus galactomannan (AG)	–	+	+	–	–	–	+	–											

\* Viridans streptococci.

† Not an acute phase reactant when highly/persistently ↑ &gt; 2 × normal.

§ False + BG with *P. aeruginosa* bacteremia, fungal derived antibiotics (amoxicillin/clavulanate, piperacillin/tazobactam), platelet transfusions, IVIG, HD (with cellulose membranes), high serum protein levels (albumin).

**Table 9.22. Liver Test Abnormalities**

Finding	Causes
↑ Alkaline phosphatase (AP) <i>mildly elevated</i>  <i>moderately/highly elevated</i>	<p><b>Infectious:</b> legionnaires' disease, viral hepatitis, liver abscess, EBV, CMV, Q fever, syphilis (2° or tertiary), TSS, hepatic candidiasis, clonorchiasis, ehrlichiosis/anaplasmosis, diphtheria, malaria, trypanosomiasis, histoplasmosis, HIV, miliary TB, relapsing fever, HPS</p> <p><b>Noninfectious:</b> drugs, DIC, fat emboli syndrome, TTP, ITP, hemolytic uremic syndrome, pre/acute/leukemias, lymphomas, carcinomas, myeloproliferative disorders (MPDs), multiple myeloma, Gaucher's disease, cirrhosis, post-hepatic obstruction, alcoholic hepatitis, pregnancy, bone growth (children), osteomalacia, rickets, hyperthyroidism, UC, drug fever, hepatoma, liver metastases, lymphoma, Erdheim-Cheste disease (ECD), mastocytosis Schnitzler's syndrome TA, normal variant (elderly)</p> <hr/> <p><b>Infectious:</b> liver abscess</p> <p><b>Noninfectious:</b> PBC, post-hepatic obstruction, post-necrotic cirrhosis, Paget's disease, osteogenic sarcoma, hepatoma, TA, bone fractures, tigecycline</p>
↑ Serum transaminases (SGOT/AST, SGPT/ALT) <i>mildly elevated</i>  <i>highly elevated</i>	<p><b>Infectious:</b> legionnaires' disease, psittacosis, Q fever, relapsing fever, brucellosis, TSS, RMSF, babesiosis, ehrlichiosis/anaplasmosis, liver abscess, syphilis (2° or tertiary), shigellosis, clonorchiasis, EBV, CMV, HSV-1, VZV (disseminated), HHV-6, anicteric viral hepatitis, gonococcemia, malaria, gram-negative bacteremia, adenovirus, dengue</p> <p><b>Noninfectious:</b> drug fever, alcoholic cirrhosis, (SGOT/AST 2:1 SGPT/ALT) CHF, infarction (myocardial, cerebral, pulmonary), pancreatitis, intrahepatic cholestasis, sickle cell disease, amyloidosis, delirium tremens, intravascular hemolysis, UC, eosinophilia-myalgia syndrome, Kawasaki's disease, adult Still's disease</p> <hr/> <p><b>Infectious:</b> viral hepatitis, HSV hepatitis, yellow fever, arboviral hemorrhagic fevers</p> <p><b>Noninfectious:</b> shock liver, rhabdomyolysis (SGOT/AST 2:1 SGPT/ALT), exercise, drugs (statins INH, RIF)</p>
↑ Total bilirubin	<p><b>Infectious:</b> legionnaires' disease, gonococcemia, liver abscess, viral hepatitis, EBV, CMV, pneumococcal bacteremia, malaria</p> <p><b>Noninfectious:</b> hemolysis, Gilbert's syndrome, cirrhosis, alcoholic hepatitis, carcinoma (pancreatic, biliary), choledocholithiasis, amyloidosis, sickle cell disease, Rotor's syndrome, <math>\alpha</math>-1 antitrypsin deficiency, hepatic vein thrombosis, autoimmune (lupoid) hepatitis, hemochromatosis</p>
↑ GGT (GGTP)	<p><b>Infectious:</b> acute viral hepatitis, EBV</p> <p><b>Noninfectious:</b> cirrhosis, alcoholic hepatitis, PBC, fatty liver, CAH, hepatoma, liver metastases, pancreatitis (acute), myocardial infarction (acute), breast cancer, lung cancer, prostate cancer, melanoma, hypernephroma (RCC), obstructive jaundice, cholestasis, CAH</p>

**Table 9.23. Rheumatic Test Abnormalities**

Finding	Causes
↑ Rheumatoid factors (RF)	<u>Infectious:</u> SBE, TB, syphilis, kala-azar, EBV, typhoid/enteric fever, HBV, HCV, chickungunya fever, malaria <u>Noninfectious:</u> cirrhosis, CAH, ITP, silicosis, asbestosis, asthma, Wegener's granulomatosis, pulmonary fibrosis, Behçet's disease, SLE, RA, Sjögren's, dermatomyositis, sarcoidosis, PBC, B-cell malignancies, normal variant
↑ Anti-nuclear antibody titers (ANA)	<u>Infectious:</u> HIV, EBV, CMV, TB, SBE, leprosy, kala-azar, malaria <u>Noninfectious:</u> SLE, scleroderma, MCTD, CREST syndrome, RA, autoimmune (lupoid) hepatitis, CAH, dermatomyositis, subacute thyroiditis, ITP, PBC, multiple sclerosis, sarcoidosis, Wegener's granulomatosis, myasthenia gravis, ESRD on HD, normal variant (elderly)
↑ Double stranded DNA (DS-DNA)	<u>Infectious:</u> CMV, EBV <u>Noninfectious:</u> SLE, RA, CAH, PBC
↑ Angiotensin-converting enzyme levels (ACE)	<u>Infectious:</u> TB, leprosy, coccidiomycosis, viral hepatitis <u>Noninfectious:</u> sarcoidosis, allergic alveolitis, hyperparathyroidism, hyperthyroidism, PBC, Gaucher's disease, DM, ESRD, amyloidosis, multiple myeloma, lymphoma, cirrhosis, psoriasis, silicosis, asbestosis, berylliosis
↑ c-ANCA	<u>Infectious:</u> viridans streptococcal SBE, S. bovis SBE, chromomycosis, aspergillosis, amebiasis, legionnaires' disease, leptospirosis, HIV <u>Noninfectious:</u> crescentic GMN, microscopic polyangitis (MPA), Sweet's syndrome, PAN, UC, RE, SLE, Churg-Strauss granulomatosis, Wegener's granulomatosis, HSP, TA, Kawasaki's disease, sarcoidosis
↑ Anti-smooth muscle antibodies (ASM)	<u>Infectious:</u> Q fever, HCV (chronic), EBV, typhoid/enteric fever <u>Noninfectious:</u> autoimmune (lupoid) hepatitis, Wilson's disease, PBC, TA, normal variant (elderly)

**Table 9.24. Urinary Abnormalities**

Finding	Causes
Pyuria*§	<u>Infectious:</u> TB, leptospirosis, brucellosis, TSS, diphtheria, candida, GC, trichomonas, cystitis, pyelonephritis, prostatitis, acute urethral syndrome, partially treated UTI, medullary abscess, chlamydia/mycoplasma NGU, balanitis, acute appendicitis, St.LE. <u>Noninfectious:</u> interstitial nephritis, urethral irritation/inflammation, strenuous exercise, SLE, calculi, bladder tumors, chronic interstitial cystitis, RE, diverticulitis, Kawasaki's disease, sarcoidosis

\* Gross pus suggests ruptured renal abscess.

§ WBC casts suggest SLE, nephritis, or acute pyelonephritis.

**Table 9.24. Urinary Abnormalities** (cont'd)

Finding	Causes
Hematuria <i>gross</i>  <i>microscopic</i>	<u>Infectious:</u> ABE (renal septic emboli), malaria ( <i>P. falciparum</i> ), yellow fever, adenoviral cystitis (type 11), BKV (transplants)  <u>Noninfectious:</u> renal malignancy, bladder malignancy, BPH, papillary necrosis (DM), trauma  <u>Infectious:</u> SBE, renal TB, schistosomiasis ( <i>S. hematobium</i> ), <i>S. saprophyticus</i> , HPS, legionnaire's disease, Q fever, BKV (transplants), EBV  <u>Noninfectious:</u> trauma, BPH, prostatitis, malignancy, malignant hypertension, PAN, drug reactions, calculi, urethral stricture, renal vein thrombosis, hydronephrosis, polycystic kidney disease, malakoplakia, strenuous exercise, renal infarction, sarcoidosis
Proteinuria	<u>Infectious:</u> TB, HBV, syphilis, malaria (black water fever), brucellosis, SBE, chronic pyelonephritis, leprosy, schistosomiasis, any acute infection  <u>Noninfectious:</u> post-streptococcal GMN, malignant hypertension, ATN, amyloidosis, sickle cell disease, polycystic kidneys, scleroderma, sarcoidosis, PAN, Wegener's granulomatosis, Goodpasture's syndrome, multiple myeloma, lymphomas, RCC, renal trauma, strenuous exercise
Urinary pH <i>alkaline</i>  <i>acidic</i>	<u>Infectious:</u> <i>Corynebacterium urealyticum</i> , <i>Klebsiella</i> (rare), <i>Proteus</i> , <i>Providencia</i> , <i>S. saprophyticus</i> , <i>Ureaplasma urealyticum</i>  <u>Noninfectious:</u> systemic alkalosis, postprandial "alkaline tide," alkalinization therapy, old urine, vegetarian diet  <u>Infectious:</u> TB  <u>Noninfectious:</u> acidification therapy, systemic acidosis, ketosis
↓ Specific gravity (1.023 – 1.030)	<u>Infectious:</u> pyelonephritis†  <u>Noninfectious:</u> diabetes insipidus, tubo-interstitial renal diseases, sickle cell disease
Nitrite positive	<u>Infectious:</u> most uropathogens**  <u>Noninfectious:</u> none
Leukocyte esterase††	<u>Infectious:</u> acute cystitis, acute pyelonephritis  <u>Noninfectious:</u> inflammation anywhere in the upper/lower GU tract

† Cystitis is associated with a normal specific gravity. The transient decrease in specific gravity of pyelonephritis corrects to normal following effective treatment.

\*\* Negative urinary nitrite occurs with group B streptococci, group D enterococci, (*VSE/VRE*) *S. saprophyticus*, *S. aureus*, *Acinetobacter* sp., *Gardnerella* sp., *Corynebacterium urealyticum*, *Burkholderia* sp., *Pseudomonas* sp., *Candida* sp.

‡ ≥ 5 WBCs/hpf = positive.

**Table 9.24. Urinary Abnormalities** (cont'd)

Finding	Causes
Eosinophiluria <sup>†</sup>	<u>Infectious</u> : <i>Shistosoma hematobium</i> <u>Noninfectious</u> : cholesterol emboli syndrome, HSP, drug induced interstitial nephritis, renal allograft rejection
Myoglobinuria (2° to rhabdomyolysis)	<u>Infectious</u> : legionnaires' disease, gas gangrene, leptospirosis, <i>Vibrio vulnificus</i> , listeria, <i>S. aureus</i> , group A streptococci, group B streptococci, <i>S. pneumoniae</i> , tularemia, typhoid/enteric fever, echovirus, Coxsackie, influenza, adenovirus, EBV, CMV, HSV, VZV, HIV <u>Noninfectious</u> : crush injury, excessive exercise, MI, seizures, malignant hyperthermia, dermatomyositis, burns, polymyositis, SLE, diabetic ketoacidosis, McArdle's syndrome, drugs (statins, alcohol, cocaine, heroin, PCP, neuroleptics, amphetamines, ethylene glycol)
Chyluria	<u>Infectious</u> : <i>W. bancrofti</i> <u>Noninfectious</u> : abdominal or chest lymphatic obstruction

† ≥ 5 WBCs = positive.

**Table 9.25. Pleural Fluid Abnormalities**

Pleural Fluid	Causes
↓ Glucose	<u>Infectious</u> : TB, bacterial, cryptococcosis, coccidiomycosis, mycoplasma, empyema <u>Noninfectious</u> : carcinoma, rheumatoid lung, lymphoma, esophageal rupture, parapneumonic effusion
↑ Protein	<u>Infectious</u> : TB <u>Noninfectious</u> : carcinoma, lymphoma, rheumatoid lung
↑ Amylase	<u>Infectious</u> : None <u>Noninfectious</u> : pancreatitis/pseudocyst, adenocarcinoma, esophageal rupture
Extracellular debris	<u>Infectious</u> : abscesses, anaerobic empyema <u>Noninfectious</u> : rheumatoid lung
PMNs	<u>Infectious</u> : bacteria, cryptococcosis, coccidiomycosis, empyema, TB (early) <u>Noninfectious</u> : pancreatitis, subdiaphragmatic abscess (sympathetic effusion), CHF, idiopathic
Lymphocytes	<u>Infectious</u> : TB <u>Noninfectious</u> : carcinoma, lymphoma, rheumatoid lung, SLE

**Table 9.26. CSF Abnormalities**

Finding	Causes
RBCs in CSF	<p><u>Infectious:</u> listeria, leptospirosis<sup>†</sup>, TB, amebic meningoencephalitis, HSV, anthrax meningitis</p> <p><u>Noninfectious:</u> traumatic tap, CNS bleed/tumor</p>
Purulent CSF with negative Gram stain	<p><u>Infectious:</u> Neisseria meningitidis, Streptococcus pneumoniae</p> <p><u>Noninfectious:</u> none</p>
CSF with negative Gram stain and predominantly PMNs/decreased glucose	<p><u>Infectious:</u> PTBM, listeria, HSV, TB (early), parameningeal infection, emboli secondary to SBE, amebic meningoencephalitis, CNS syphilis (early)</p> <p><u>Noninfectious:</u> sarcoidosis, adult Still's disease, posterior-fossa syndrome/intracranial hemorrhage</p>
CSF with negative Gram stain and predominantly lymphocytes/normal glucose	<p><u>Infectious:</u> PTBM, viral meningitis, Lyme disease, HIV, leptospirosis<sup>†</sup>, RMSF, parameningeal infection, TB, fungi, parasitic meningitis</p> <p><u>Noninfectious:</u> sarcoidosis, meningeal carcinomatosis</p>
lymphocytes/decreased glucose	<p><u>Infectious:</u> PTBM, CNS TB, CNS fungi, LCM, mumps, enteroviral meningitis, listeria, leptospirosis<sup>†</sup>, syphilis</p> <p><u>Noninfectious:</u> neurosarcoidosis, meningeal carcinomatosis</p>
↑ CSF lactic acid levels	<p><u>Infectious:</u> &lt; 4 nm/L, Not acute bacterial meningitis (ABM); &lt; 4–6 nm/L, TB, RBCs, HSV, partially treated bacterial meningitis (PTBM), parameningeal infections;</p> <p>&gt; 6 nm/L, acute bacterial meningitis (ABM), cerebral malaria (severe)</p> <p><u>Noninfectious:</u> &lt; 6 nm/L, neurosarcoidosis, CNS SLE, cerebral anoxia, hepatic encephalopathy; meningeal carcinomatosis; CNS lymphomas, CNS tumor, intracranial hemorrhage, post-craniotomy with EVD</p>
Highly ↑ CSF Protein	<p><u>Infectious:</u> brain abscess, CNS TB (with subarachnoid block), viral meningitis, viral encephalitis</p> <p><u>Noninfectious:</u> brain tumor, MS, demyelinating CNS diseases</p>
↑ CSF adenosine deaminase (ADA) levels	<p><u>Infectious:</u> CNS TB, listeria, ABM, neurobrucellosis</p> <p><u>Noninfectious:</u> SAH, CNS malignancy</p>
CSF plasma cells	<p><u>Infectious:</u> CNS TB, WNE, VZV, HIV, neuroborreliosis, neurosyphilis</p> <p><u>Noninfectious:</u> CNS lymphoma, MS, CNS myeloma, ADEM</p>

<sup>†</sup> Leptospirosis is the only infectious disease with the CSF bilirubin > serum bilirubin.



**Table 9.26. CSF Abnormalities** (cont'd)

Finding	Causes
CSF eosinophils	<p><u>Infectious</u>: coccidioidomycosis, neurocysticercosis, gnathostomiasis, angiostrongyliasis, baylisascariasis, shistosomiasis, paragonamiasis, Toxocariasis canis/cati (VLM)</p> <p><u>Noninfectious</u>: CNS lymphomas, CNS leukemias, V-A/VP shunts, myelography (contrast material), CNS vasculitis, drugs (NSAIDs, TMP-SMX), intrathecal drugs</p>

**Table 9.27. CSF Diagnostic Criteria for EVD associated ABM****EVD Associated ABM**

- Highly elevated CSF lactic acid level (> 6 nmol/L)  
*plus*
- Marked CSF pleocytosis (> 50 WBCs/hpf)  
*plus*
- Positive CSF Gram stain (with *same morphology* as the cultured neuropathogen)  
*plus*
- Positive CSF culture with (*same morphology* as the Gram stain of the neuropathogen)

**EVD Associated Pseud meningitis**

- < 4 CSF Diagnostic Criteria for EVD Associated ABM  
Noninfections: CNS malignancy, ICH, SAH, post-craniotomy, brain trauma

LA = CSF lactic acid

ABM = acute bacterial meningitis

EVD = external-ventricular drain

## Chapter 10

**Antibiotic Pearls & Pitfalls****Burke A. Cunha, MD**

Penicillin .....	508
Ampicillin .....	508
Amoxicillin/Clavulanic Acid .....	508
Oral Anti-Staphylococcal Penicillins .....	509
Oral Anti-Pseudomonal Penicillins .....	509
Parenteral First-Generation Cephalosporins .....	509
Oral First-Generation Cephalosporins .....	509
Parenteral Second-Generation Cephalosporins .....	510
Oral Second-Generation Cephalosporins .....	510
Parenteral Third-Generation Cephalosporins .....	510
Oral Third-Generation Cephalosporins .....	511
Parenteral Fourth-Generation Cephalosporin (Cefepime) .....	511
Parenteral Anti-MRSA Cephalosporin (Ceftaroline) .....	511
Monobactams (Aztreonam) .....	511
Carbapenems (Imipenem, Meropenem, Ertapenem, Doripenem) .....	511
Beta-Lactamase Inhibitor Combinations .....	512
Tetracyclines (Doxycycline, Minocycline) .....	512
Chloramphenicol .....	512
Clindamycin .....	513
Aminoglycosides (Gentamicin, Tobramycin, Amikacin) .....	513
TMP-SMX .....	515
Quinolones (Ciprofloxacin, Ofloxacin, Levofloxacin, Moxifloxacin) .....	515
Nitrofurantoin .....	516
Vancomycin .....	516
Oxazolidinones (Linezolid, Tedizolid) .....	517
Quinupristin/Dalfopristin .....	517
Lipopeptide (Daptomycin) .....	517
Tigecycline .....	518
Macrolides (Erythromycin, Azithromycin, Clarithromycin) .....	518
Metronidazole .....	518
Lipoglycopeptides (Televancin, Oritavancin, Dalbavancin) .....	519
Fosfomycin .....	519
Colistin and Polymyxin B .....	520
Nitazoxanide .....	520

---

**PENICILLIN**

---

- *Penicillin is expensive* because it is made by a single manufacturer.
- There are very few uses for parenteral penicillin therapy today.
- PRSP has been associated with TMP-SMX and macrolides, *not penicillin*.

---

**AMPICILLIN**

---

- Unless treating serious systemic infections due to *E. faecalis* (VSE), *avoid using ampicillin*. *Ampicillin use results in increased E. coli resistance*.
- Remember ampicillin is the preferred drug to treat serious systemic infections caused by VSE but is *ineffective against nearly all E. faecium* (VRE).
- Because susceptibility is in part “concentration dependent,” *do not assume that susceptibilities of ampicillin or amoxicillin are the same*. On a same-dose basis, amoxicillin achieves twice the concentrations of ampicillin in body fluids (e.g., middle ear fluid, sinus fluid, bronchial fluid, urine).
- Remember that *unlike ampicillin, amoxicillin is infrequently associated with oral thrush or irritative diarrhea*.
- *Amoxicillin 1 gm (PO) q8h can be used in ampicillin IV-to-PO switch programs since this dose provides levels comparable to parenteral (IM) ampicillin*.

---

**AMOXICILLIN/CLAVULANIC ACID**

---

- Clavulanic acid is a beta-lactamase inhibitor which, when added to amoxicillin, restores its activity against beta-lactamase—producing strains of *H. influenzae*.
- The *newer preparations of amoxicillin/clavulanic acid result in less gastrointestinal symptoms and diarrhea* compared to older preparations with more clavulanate.
- *Ineffective against PRSP due to alterations in PBPs which are not  $\beta$ -lactamase mediated*.

---

**ORAL ANTI-STAPHYLOCOCCAL PENICILLINS**

---

- *Do not rely on oral anti-staphylococcal penicillins* such as dicloxacillin for methicillin-sensitive *S. aureus* (MSSA) infections since they are erratically/poorly absorbed and not consistently effective. When treating MSSA infections orally, a first generation cephalosporin (e.g., cephalexin) is preferable.
- *Dicloxacillin is poorly tolerated due to its metallic taste and belching.*

---

**ORAL ANTI-PSEUDOMONAL PENICILLINS**

---

- *Avoid using indanyl carbenicillin for *P. aeruginosa* UTIs due to the rapid development of resistance with *P. aeruginosa*.* Use other oral anti-*P. aeruginosa* agents for *P. aeruginosa* lower UTIs, e.g., doxycycline, levofloxacin, or fosfomycin.

---

**PARENTERAL FIRST-GENERATION CEPHALOSPORINS**

---

- *Cefazolin has limited activity against *H. influenzae*.*
- *Cefazolin remains preferred therapy for skin infections due to group A streptococci or MSSA.*
- *For treatment of biliary tract infections, cefazolin is active against *E. coli* and *Klebsiella pneumoniae* but is not active against *E. faecalis* (VSE).*
- *Cefazolin remains useful prophylaxis for cardiothoracic procedures in hospitals with low prevalence of MRSA.*

---

**ORAL FIRST-GENERATION CEPHALOSPORINS**

---

- *Cephalexin suboptimal for respiratory tract infections* i.e., *H. influenzae* a likely pathogen (otitis media, sinusitis, AECB, community-acquired pneumonia) since 1<sup>st</sup> GCs have limited *H. influenzae* activity.
- *For IV-to-PO switch programs and cephalexin 1gm (PO) q6h approximates the serum concentrations of parenteral (IM) cefazolin.*
- *Oral 3<sup>rd</sup> GCs do not have the same degree of anti-*S. aureus* MSSA activity as cephalexin.*

---

**PARENTERAL SECOND-GENERATION CEPHALOSPORINS**

---

- Cefixitin and cefotetan are useful for intra-abdominal infections.
- *Cefuroxime* offers no advantage for bacterial URIs over doxycycline, respiratory quinolones, or 3<sup>rd</sup> GCs and *does not prevent CNS "seeding" by S. pneumoniae or H. influenzae bacteremia secondary to CAP.*

---

**ORAL SECOND-GENERATION CEPHALOSPORINS**

---

- *Oral second generation cephalosporins include cefaclor, cefprozil, and cefuroxime are used primarily to treat URIs.*
- *Cefaclor has limited ability to penetrate into respiratory secretions limiting its efficacy in URIs.*
- *Among the oral second generation cephalosporins, cefprozil has the greatest degree of penetration into respiratory secretions.*

---

**PARENTERAL THIRD-GENERATION CEPHALOSPORINS**

---

- *Except for cefoperazone, 1<sup>st</sup>, 2<sup>nd</sup>, 3<sup>rd</sup> generation cephalosporins have no anti-E. faecalis (VSE) activity.*
- *Except for ceftriaxone, third generation cephalosporins predispose to C. difficile diarrhea. Ceftriaxone is associated with non-C. difficile diarrhea related to changes in colonic flora.*
- *Be aware that ceftriaxone is associated with pseudo-biliary lithiasis. Patients developing right upper quadrant pain on ceftriaxone should be suspected as having drug-induced pseudo-biliary lithiasis.*
- *For CNS infections give third generation cephalosporins in "meningeal doses"*
- *The only third-generation cephalosporin associated with resistance is ceftazidime e.g., P. aeruginosa.*
- *Ceftazidime predisposes to MDR P. aeruginosa, and increases MRSA prevalence.*
- *Ceftazidime associated with ESBL K. pneumoniae, E. coli, or Enterobacter agglomerans.*
- *Ceftazidime little anti-MSSA activity.*
- *3<sup>rd</sup> GCs without anti-B. fragilis activity are ceftazidime and ceftriaxone.*
- *3<sup>rd</sup> GCs without significant anti-P. aeruginosa activity are cefotaxime, ceftizoxime, and ceftriaxone.*

---

**ORAL THIRD-GENERATION CEPHALOSPORINS**

---

- Cefdinir, cefditoren, cefixime, cefpodoxime and ceftibuten are used for their aerobic anti-GNB activity.
  - *The only 3<sup>rd</sup> GC with good anti-MSSA activity is cefpodoxime.*
- 

**PARENTERAL FOURTH-GENERATION CEPHALOSPORIN (CEFEPIME)**

---

- Cefepime (4<sup>th</sup> GC) is an anti-Pseudomonal cephalosporin that is often active against ceftazidime resistant *P. aeruginosa* strains.
  - For *P. aeruginosa* and MDRO GNBs use high dose cefepime, i.e., 2 gm (IV) q8h.
- 

**PARENTERAL ANTI-MRSA CEPHALOSPORIN (CEFTAROLINE)**

---

- Ceftaroline is the *only cephalosporin active in vivo against both MSSA and MRSA.*
  - Ceftaroline is useful for cSSIs and MSSA/MRSA bacteremias
- 

**MONOBACTAMS (AZTHREONAM)**

---

- *Aztreonam has no gram-positive activity.*
  - Although structurally similar to the beta-lactams, *aztreonam is safe in penicillin-allergic patients.*
- 

**CARBAPENEMS (IMIPENEM, MEROPENEM, ERTAPENEM, DORIPENEM)**

---

- *Avoid imipenem in patients with seizures/CNS disorders since imipenem may cause seizures. Also, renal insufficiency increases imipenem's seizure potential.*
- Meropenem doesn't cause seizures in those with or without seizure disorders or with normal or decreased renal function.
- *All carbapenems have anti-pseudomonal activity except for ertapenem.*
- *Ertapenem has no anti-enterococcal activity, important in biliary/UTIs.*
- *Avoid IV bolus injections/rapid infusions of doripenem, ertapenem or imipenem. Only meropenem may be administered as an IV bolus, important in septic shock.*

---

**BETA-LACTAMASE INHIBITOR COMBINATIONS**

---

- *Some strains of MDR Acinetobacter baumannii are susceptible only to sulbactam/ampicillin.*
- *Tazobactam does not enhance the anti-pseudomonal activity of piperacillin.*
- *Ceftazidime/avibactam and ceftolozane/tazobactam are effective for MDR GNB UTIs and urosepsis. Active against KPCs, but not metallo-beta-lactamases.*
- *Ceftazidime/avibactam and ceftolozone/tazobactam have some B. fragilis activity, but for cIAls, should be used together with metronidazole.*

---

**TETRACYCLINES (DOXYCYCLINE, MINOCYCLINE)**

---

- *While MRSA may be susceptible to doxycycline in vitro, in vivo it frequently fails or the clinical response is delayed/incomplete. Minocycline IV/PO for MSSA/MRSA is more effective than doxycycline.*
- *Doxycycline has good CSF penetration and is often used for oral therapy of neuroborreliosis.*
- *At urinary concentrations, doxycycline is effective against P. aeruginosa CAB/lower UTIs, i.e., high urine levels > 300 mcg/mL. However, in vitro testing will report P. aeruginosa resistant to doxycycline.*
- *For nearly all non-viral zoonotic infections, doxycycline is effective except for babesiosis.*
- *Oral minocycline is useful to treat serious systemic infections due to MSSA/MRSA i.e., ABE, osteomyelitis, meningitis, etc.*

---

**CHLORAMPHENICOL**

---

- *Chloramphenicol is one of few drugs that can be given orally to treat acute bacterial meningitis due to susceptible organisms.*
- *Chloramphenicol is the only antibiotic that when given orally results in higher serum concentrations than when given at the same dose intravenously.*
- *Avoid chloramphenicol for the treatment of hepatobiliary infections and UTIs since chloramphenicol is excreted into the bile as an inactivate metabolite, and urinary levels are low.*
- *Remember, aplastic anemia is a rare idiosyncratic side effect associated with oral/ topical therapy but not IV therapy.*

- Chloramphenicol-induced aplastic anemia is an idiosyncratic reaction and *serial CBCs are unhelpful in predicting/avoiding aplastic anemia.*
  - *Serial CBCs may be obtained to monitor dose-related side effects (anemia) not idiosyncratic side effects (aplastic anemia).*
  - *Prolonged chloramphenicol therapy may cause sequential, dose-related suppression of bone marrow elements. Suppressed blood elements return in reverse sequence when chloramphenicol is discontinued.*
  - *Vacuolization on bone marrow biopsy specimens in patients receiving chloramphenicol is a manifestation of chloramphenicol effect, not aplastic anemia.*
  - *Chloramphenicol is effective for E. faecium (VRE) systemic infections.*
- 

### CLINDAMYCIN

---

- *Clindamycin misses approximately 15% of coagulase-negative S. epidermidis (CONS), the most common pathogen in foreign body/implant-related infections.*
  - *Clindamycin is one of the few antibiotics able to penetrate/dissolve staphylococcal biofilms. Clindamycin may be useful adjunctively in treating foreign body associated infections when the prosthetic device cannot be removed.*
  - *The incidence of C. difficile diarrhea is greater with PO > IV clindamycin.*
  - *Clindamycin is not active against group D enterococci (VSE/VRE).*
  - *With CA-MRSA, inducible clindamycin resistance (MLS(B)) should be suspected if erythromycin is resistant and clindamycin is susceptible. The "D test" will confirm clindamycin resistance.*
  - *Clindamycin anti-toxin properties may be useful in severe GAS infections, e.g., necrotizing fasciitis.*
- 

### AMINOGLYCOSIDES (GENTAMICIN, TOBRAMYCIN, AMIKACIN)

---

- *Among aminoglycosides, gentamicin has the most anti-gram-positive coccal activity.*
- *Amikacin has the highest degree of P. aeruginosa activity.*
- *Gentamicin and tobramycin have 6 loci that may be inactivated by acetylating, phosphorylating and phosphorylating enzymes. Amikacin has only 1 locus that may be attacked by these enzymes, making P. aeruginosa resistance less likely.*



- *Avoid aminoglycoside monotherapy in the treatment of nosocomial pneumonia since aminoglycoside activity is diminished in the presence of tissue hypoxia, WBC debris and local acidosis, which are prominent in nosocomial pneumonia.*
- *Avoid administering aminoglycosides via nebulizer since aerosolized aminoglycosides may predispose to resistance.*
- *Avoid administering aminoglycosides in split-daily doses. A single daily dose is optimal since aminoglycosides obey "concentration-dependent" killing kinetics.*
- *When aminoglycosides are given to patients with renal insufficiency, begin with the usual initial dose, then decrease the maintenance dose in proportion to the degree of renal dysfunction based on the creatinine clearance.*
- *If aminoglycosides are being used for synergy, use half the therapeutic dose (synergy dose).*
- *Aminoglycosides given on a once-daily basis optimize aminoglycoside antibacterial killing while minimizing nephrotoxic potential and eliminate the need for aminoglycoside levels.*
- *Aminoglycoside ototoxicity may occur with extremely high/prolonged peak levels. Episodic elevated peak levels are not associated with ototoxicity.*
- *When aminoglycosides are given on a once-daily basis, renal tubular cells have sufficient time between dosing intervals to decrease intracellular levels and thus avoid nephrotoxicity.*
- *Do not assess aminoglycoside nephrotoxicity based on serum creatinines. Aminoglycoside nephrotoxicity is best assessed by indicators of renal tubular damage such as urinary renal cast counts.*
- *Limiting aminoglycoside therapy to 2 weeks with split-daily dosing minimizes the risk for aminoglycoside nephrotoxicity.*
- *Aminoglycosides are suboptimal for *P. aeruginosa*. Other antibiotics (e.g., meropenem, doripenem, cefepime) are preferred.*
- *Aminoglycosides appear to be active against streptococci by in-vitro susceptibility testing, but aminoglycosides have no inherent activity against streptococci, i.e., groups A, B, C, G, D.*
- *Aminoglycosides (e.g., gentamicin) are only active against group D enterococci (*E. faecalis*, VSE) when combined with penicillin or vancomycin.*
- *Avoid, if possible, using aminoglycosides for peritoneal lavage since the peritoneum provides a very large cross-sectional area for drug absorption, increasing the risk for neuromuscular blockade/respiratory arrest.*
- *For CNS infections, aminoglycosides must be administered intrathecally (IT) since they do not cross the blood brain barrier in sufficient concentrations to be therapeutic.*

---

**TMP-SMX**

---

- Remember, *TMP-SMX* has the same spectrum as ceftriaxone (i.e., no *P. aeruginosa* coverage).
- Avoid using *TMP-SMX* to treat serious systemic *K. pneumoniae* infections. *K. pneumoniae* isolates that are sensitive to *TMP-SMX* *in-vitro* are often ineffective *in-vivo*.
- In patients with hypersensitivity reactions to *TMP-SMX*, it is always the sulfa component, not the *TMP* component, which is responsible. If continued treatment is desired with *TMP-SMX*, therapy may be completed with the *TMP* component alone.
- Although *TMP-SMX* is inactive against most streptococci, it is an excellent antibiotic against *MSSA*.
- *TMP-SMX* is active against *CA-MRSA* but is suboptimal against *HA-MRSA* and *CO-MRSA*.
- For the treatment of *hydradenitis suppurativa* due to *MSSA*, *TMP-SMX* is preferred antibiotic because of its ability to penetrate deep into infected sebaceous glands.
- For aerobic GNB bacteremias (other than *P. aeruginosa*), use *TMP-SMX* 10 mg/kg/day (IV/PO) given in 4 equally divided doses q6h.
- For CNS penetration and the treatment of unusual organisms (e.g., *PCP*), use *TMP-SMX* at dose of 20 mg/kg/day (IV/PO) given in 4 equally divided doses q6h.
- Avoid *TMP-SMX* in the treatment of respiratory tract infections since it is associated with *S. pneumoniae* resistance.

---

**QUINOLONES (CIPROFLOXACIN, OFLOXACIN, LEVOFLOXACIN, MOXIFLOXACIN)**

---

- Levofloxacin at a dose of 750 mg IV/PO is preferred to ciprofloxacin for treatment of serious systemic *P. aeruginosa* infections.
- With the exception of ciprofloxacin, quinolones do not lower the seizure threshold/cause seizures.
- Moxifloxacin is the only quinolone with a hepatobiliary mode of excretion; dose does not need to be modified and renal insufficiency but concentrations in the urine for lower UTIs may be suboptimal.
- For meningococcal prophylaxis, quinolones are equally effective (quinolones penetrate well into respiratory secretions/have a high degree of anti-meningococcal activity).

- The “respiratory quinolones”, e.g., moxifloxacin, levofloxacin have a high degree of bio-availability which makes them *ideal to treat systemic infections PO or as part of IV to PO switch therapy*. Respiratory quinolones have anti-TB activity (an option in treating MDR TB).
- *Ciprofloxacin has less activity against S. pneumoniae and S. aureus than moxifloxacin or levofloxacin.*
- Quinolones have some degree of anti-*E. faecalis* (VSE) but have no *E. faecium* (VRE) activity. *Among the quinolones, moxifloxacin has the highest degree of anti-VSE activity.*

### NITROFURANTOIN

- *Nitrofurantoin is an ideal oral agent to treat cystitis*
- *If the CrCL < 30 ml/min, nitrofurantoin may not be effective for lower UTIs/CAB.*
- *If re-emptive treatment of CAB is desired in immunocompromised hosts, e.g., diabetics, SLE, myeloma, CLL, steroids, neutropenia, nitrofurantoin is an ideal oral agent and is active against the usual CAB uropathogens, i.e., coliforms and group D enterococci (VSE/VRE).*
- *Nitrofurantoin is active against all aerobic GNB uropathogens except P. aeruginosa, Serratia marcescens, and Proteus sp.*

### VANCOMYCIN

- *If vancomycin is combined with a nephrotoxic drug and nephrotoxicity occurs, it should be ascribed to the nephrotoxic drug and not vancomycin.*
- *Vancomycin is a “concentration-dependent” drug (with MSSA/MRSA with MICs > 1 mcg/ml), but a “time dependent” drug (with MSSA/MRSA MICs < 1 mcg/ml).*
- *Other drugs preferable to vancomycin to treat cSSIs and bacteremias/ABE due to MSSA.*
- *No need for vancomycin levels for usual vancomycin dosing.* However, vancomycin levels may be useful in patients with unusually high volume of distribution ( $V_d$ ) (e.g., edema/ascites, trauma, burns).
- *Vancomycin “tolerance” is common among staphylococci and enterococci.*
- *Vancomycin is alone is inadequate for E. faecalis (VSE) bacteremia/SBE. Add gentamicin for optimal VSE activity.*
- *Vancomycin may cause cell wall thickening (“permeability-mediated” resistance) manifested by ↑ MICs.*
- *Because of its large molecular size, vancomycin does not penetrate well into synovial fluid.*
- *Vancomycin IV predisposes to VRE, but PO vancomycin does not.*

---

**OXAZOLIDINONES (LINEZOLID, TEDIZOLID)**

---

- Linezolid is highly active against the major gram-positive pathogens, including MSSA, MRSA, VSE, and VRE.
- Linezolid is *equally efficacious when administered PO or IV*.
- Unlike vancomycin, *linezolid does not increase E. faecium (VRE) prevalence*.
- Because linezolid is eliminated by hepatic mechanisms, no dosing adjustment is necessary in renal insufficiency.
- Tedizolid, like linezolid, may be given *IV/PO*.
- Once daily dosing *IV/PO* effective with tedizolid ( $t_{1/2} = 11$  hrs).
- Linezolid is *one of the few PO antibiotics that can be used to treat CNS infections*.

---

**QUINUPRISTIN/DALFOPRISTIN**

---

- Quinupristin/dalfopristin is active against *E. faecium (VRE)* but *not E. faecalis (VSE)*.
- Quinupristin/dalfopristin is highly effective against MSSA and MRSA.

---

**LIPOPEPTIDE (DAPTOMYCIN)**

---

- Daptomycin is *more active against MSSA/MRSA than VSE/VRE* (mean MSSA/MRSA MICs = 0.5 mcg/ml vs. mean VSE/VRE MICs = 1.0 mcg/ml). *If daptomycin is used to treat VSE/VRE, a higher dose is recommend (dose for MSSA/MRSA bacteremia: 6 mg/kg/day dose for VSE/VRE bacteremia: 12 mg/kg/day)*.
- Daptomycin is *inactivated by calcium in alveolar surfactant fluid* and should not be used for pneumonias, but useful for septic pulmonary emboli or lung abscesses.
- Following vancomycin therapeutic failures with MSSA/MRSA bacteremias/ABE, *daptomycin resistance may occur during therapy*.
- *An initial dose of gentamicin may increase intracellular entry/effectiveness of daptomycin when treating MSSA/MRSA infections*.

**TIGECYCLINE**

---

- Tigecycline may be given safely to patients with penicillin or sulfa drug allergy but *avoid in patients with tetracycline allergy.*
  - Tigecycline (hepatobiliary elimination) *does not need to be adjusted in renal insufficiency.*
  - With *A. baumannii* only, Etesting shows falsely high MICs with tigecycline.
  - Tigecycline is highly active against *MDR K. pneumoniae* and may be the *only antibiotic effective against such strains.*
  - Tigecycline is one of only two antibiotics effective against the nearly pan-resistant strains of NDM-1.
- 

**MACROLIDES (ERYTHROMYCIN, AZITHROMYCIN, CLARITHROMYCIN)**

---

- *Macrolides have been largely responsible for PRSP and MDRSP. Use instead doxycycline or a "respiratory quinolone."*
  - IV erythromycin lactobionate is the macrolide *most likely to be associated with QTc prolongation.* Macrolides with *lower serum levels* (e.g., azithromycin) are not associated with QTc prolongation.
  - Macrolides may cause an "irritative" diarrhea, but *not C. difficile* diarrhea.
  - Erythromycin estolate may cause cholestatic jaundice in adults, but not in children.
  - Due to widespread group A streptococci and MSSA resistance to macrolides, *avoid macrolides for the treatment of skin/soft tissue infections.*
- 

**METRONIDAZOLE**

---

- Metronidazole (IV) is *preferred therapy for C. difficile colitis.*
- Metronidazole (PO) *frequently fails and is inferior to PO vancomycin for C. difficile diarrhea.*
- A underrecognized/untoward effect of metronidazole (PO) use for *C. difficile diarrhea* is *increased VRE prevalence.*
- Metronidazole has a long serum half life ( $t_{1/2}$ ) of approximately 7 hr which permits *once daily dosing*, i.e., 1 g (IV) q24h. Except for *C. difficile colitis*, there is little rationale for dosing metronidazole 500 mg (IV) on a q6 or q8h basis.

- Metronidazole is one of the few hepatically eliminated antibiotics that requires a dosing adjustment in severe renal insufficiency ( $CrCl < 10$  ml/min).
- Don't combine metronidazole and moxifloxacin for intraabdominal infections (no rationale for double *B. fragilis* coverage).

---

### LIPOGLYCOPEPTIDES (TELEVANCIN, ORITAVANCIN, DALBAVANCIN)

---

- Lipoglycopeptides are useful for cSSIs due to MSSA/MRSA.
- Lipoglycopeptides with long  $t_{1/2}$  permits infrequent dosing, e.g., telavancin ( $t_{1/2} = 8$  hrs) q 24 hrs dosing; oritavancin ( $t_{1/2} = 245$  hrs) single dose; dalbavancin ( $t_{1/2} = 346$  hrs) q weekly dosing.
- Telavancin is not removed by HD (protein binding = 90%), but like some other antibiotics, may have decreased efficacy with  $CrCl < 50$  ml/min.
- Falsely elevated INR, PT, PTT may occur with telavancin.

---

### FOSFOMYCIN

---

- Fosfomycin is highly active against most MDR GNS uropathogens (including VSE/VRE) and may be used orally to treat lower UTIs/CAB due to susceptible organisms.
- Fosfomycin is one of the few oral antibiotics effective against most MDR GNB uropathogens. Misses some strains of MDR *Enterobacter sp.*, *Klebsiella pneumoniae*, *Acinetobacter sp.*, and *P. aeruginosa*.
- Longer fosfomycin treatment courses may be needed to eradicate some MDR GNB uropathogens.
- Achieves high urinary concentrations which minimizes the potential for GNB resistance.
- Synergistic with quinolones against MDR GNBs.
- Fosfomycin penetrates the inflamed and non-inflamed prostate in therapeutic concentrations.
- Fosfomycin is one of the few oral antibiotics useful to treat acute (inflamed prostate) prostatitis and chronic (non-inflamed) prostatitis due to MDR GNB.
- Dose of fosfomycin for chronic prostatitis due to MDR GNB may need to be higher than usual, e.g., (PO) q72h.
- Duration of fosfomycin therapy in chronic prostatitis (due to MDR GNB) may need to be prolonged (weeks – months).
- As with other antibiotics used for the treatment of chronic prostatitis, fosfomycin therapy (high dose/prolonged therapy) may fail in the setting of prostate cancer, radioactive prostate seeds, or prostatic calcifications.

- If prostatic calcifications are present, they should if possible be removed (TURP). If prostatic calcifications cannot be resected, fosfomycin (high dose/prolonged) may be effective with doxycycline is added to the regimen.
  - Fosfomycin urinary concentrations are > 1200 mcg/ml after a 3 gm (PO) dose. Urine levels > 200 mcg/ml for > 48 hours.
  - Fosfomycin is less active than GNB against VSE/VRE (MICs = 48 – 64 mcg/ml).
- 

### COLISTIN AND POLYMYXIN B

---

- *Colistin and Polymyxin B have same spectrum of activity, but differ in activity, e.g., polymyxin B has greater activity against P. aeruginosa than colistin.*
  - Colistin and Polymyxin B and colistin are active against most aerobic GNBs but no activity against *Proteus*, *Providencia*, *Morganella*, *Serratia*, or *B. cepacia*.
  - *Colistin is one of the very few antibiotics effective against the nearly pan-resistant NDM-1 strains.*
  - *Colistin an antibiotic for GNBs that may be given at each HD (q48h). HD dose: 2 mg/kg (IV) q48h (no post-HD dose needed).*
- 

### NITAZOXANIDE

---

- Nitazoxanide is active against anaerobic bacteria, e.g., *C. difficile*, *H. pylori* and *B. fragilis*.
- Oral nitazoxanide is more effective against *C. difficile* than metronidazole for either *C. difficile* diarrhea or *C. difficile* colitis.
- Nitazoxanide is effective against *B. fragilis* strains with decreased susceptibility/resistance to metronidazole.
- Nitazoxanide has no activity against aerobic Gram positive and GNB.
- Nitazoxanide has some activity against viral pathogens, e.g., norovirus, rotavirus, adenovirus.

## Chapter 11

**Antimicrobial Drug Summaries****Burke A. Cunha, MD, Damary C. Torres, PharmD****Jean E. Hage, MD, David W. Kubiak, PharmD****Nardeen Mickail, MD, Arthur Gran, MD, Muhammed Raza, MBBS****Sigridh Muñoz-Gomez, MD, Cheston B. Cunha, MD****John H. Rex, MD, Mark H. Kaplan, MD**

**This section contains prescribing information pertinent to the clinical use of antimicrobial agents in adults, as compiled from a variety of sources. Antimicrobial agents for pediatric infectious diseases are described in Chapter 7.** The information provided is not exhaustive, and the reader is referred to other drug information references and the manufacturer's product literature for further information. Clinical use of the information provided and any consequences that may arise from its use are the responsibilities of the prescribing physician. The authors, editors, and publisher do not warrant or guarantee the information contained in this section, and do not assume and expressly disclaim any liability for errors or omissions or any consequences that may occur from such. **The use of any drug should be preceded by careful review of the package insert, which provides indications and dosing approved by the U.S. Food and Drug Administration. This information can be obtained on the website provided at the end of the reference list for each drug summary.**

All orally administered antibiotics should be taken with food unless otherwise specified (included after usual oral dose)

Drugs are listed alphabetically by generic name; trade names follow in parentheses. To search by trade name, consult the index. Each drug summary contains the following information:

**Usual dose.** Represents the usual dose to treat most susceptible infections in adult patients with normal hepatic and renal function. Dosing for special situations is listed under the comments section; additional information can be found in Chapters 2, 4, 5 and the manufacturer's product literature. Loading doses for doxycycline, fluconazole, itraconazole, voriconazole, caspofungin, and other antimicrobials are described in either the usual dose or comments section. **Meningeal doses of antimicrobials used for CNS infection are described at the end of the comments section.**

**Peak serum levels.** Refers to the peak serum concentrations (mcg/ml) after the usual dose is administered. Peak serum level is useful in calculating the "kill ratios," the ratio of peak serum level to minimum inhibitory concentration (MIC) of the organism. **The higher the "kill ratio," the more effective the "dose dependent" antimicrobial is likely to be against a particular organism.**

**Bioavailability.** Refers to the percentage of the dose reaching the systemic circulation from the site of administration (PO or IM). **For PO antibiotics, bioavailability refers to the percentage of dose**



**adsorbed from the GI tract.** For IV antibiotics, “not applicable” appears next to bioavailability, since the entire dose reaches the systemic circulation. Antibiotics with high bioavailability (> 90%) are ideal for IV to PO switch therapy.

**Excreted unchanged.** Refers to the percentage of drug excreted unchanged, and **provides an indirect measure of drug concentration in the urine/feces. Antibiotics excreted unchanged in the urine in low concentration are unlikely to be useful for urinary tract infections.**

**Serum half-life (normal/ESRD).** The serum half-life ( $t_{1/2}$ ) is the time (in hours) in which serum concentration falls by 50%. **Serum half-life is useful in determining dosing interval.** If the half-life of drugs eliminated by the kidneys is prolonged in end-stage renal disease (ESRD), then the total daily dose is reduced in proportion to the degree of renal dysfunction. If the half-life in ESRD is similar to the normal half-life, then the total daily dose does not change.

**Plasma protein binding.** Expressed as the **percentage of drug reversibly bound to serum albumin. It is the unbound (free) portion of a drug that equilibrates with tissues and imparts antimicrobial activity. Plasma protein binding is not typically a factor in antimicrobial effectiveness unless binding exceeds 95%.** Decreases in serum albumin (nephrotic syndrome, liver disease) or competition for protein binding from other drugs or endogenously produced substances (uremia, hyperbilirubinemia) will increase the percentage of free drug available for antimicrobial activity, and may require a decrease in dosage. Increases in serum binding proteins (trauma, surgery, critical illness) will decrease the percentage of free drug available for antimicrobial activity, and may require an increase in dosage.

**Volume of distribution ( $V_d$ ).** Represents the apparent volume into which the drug is distributed, and is calculated as the amount of drug in the body divided by the serum concentration (in liters/kilogram).  $V_d$  is related to total body water distribution ( $V_d \text{ H}_2\text{O} = 0.7 \text{ L/kg}$ ). Hydrophilic (water soluble) drugs are restricted to extracellular fluid and have a  $V_d \leq 0.7 \text{ L/kg}$ . In contrast, **hydrophobic (highly lipid soluble) drugs penetrate most fluids/tissues of the body and have a large  $V_d$ .** Drugs that are concentrated in certain tissues (e.g., liver) can have a  $V_d$  greatly exceeding total body water.  $V_d$  is affected by organ perfusion, membrane diffusion/permeability, lipid solubility, protein binding, and state of equilibrium between body compartments. **For hydrophilic drugs, increases in  $V_d$  may occur with burns, heart failure, dialysis, sepsis, cirrhosis, or mechanical ventilation;** decreases in  $V_d$  may occur with trauma, hemorrhage, pancreatitis (early), or GI fluid losses. Increases in  $V_d$  may require an increase in total daily drug dose for antimicrobial effectiveness; decreases in  $V_d$  may require a decrease in drug dose. **In addition to drug distribution,  $V_d$  reflects binding avidity to cholesterol membranes and concentration within organ tissues (e.g., liver).**

**Mode of elimination.** Refers to the primary route of inactivation/excretion of the antibiotic, which **impacts dosing adjustments in renal/hepatic failure.**

**Dosage adjustments.** Each grid provides dosing adjustments based on renal and hepatic function. **In renal insufficiency (for renally eliminated antibiotics), the initial dose is the same as with normal renal function;** but the **maintenance dose is decreased**  $\sim \downarrow \text{CrCl}$  (see antibiotic drug summaries for specific dosing recommendations). **Antimicrobial dosing for hemodialysis (HD)/peritoneal dialysis (PD) patients (between HD/PD) is the same as a CrCl < 10 mL/min. Some antimicrobial agents require a supplemental dose immediately after hemodialysis (post-HD)/peritoneal dialysis (post-PD).** Following the post-HD or post-PD supplemental dose, antimicrobial dosing should resume as for a CrCl < 10 mL/min. “No change” indicates no change from the usual dose. “Avoid” indicates the drug should be avoided in the setting described. “None” indicates no supplemental dose is required. “No information” indicates there are insufficient data from which to make a dosing recommendation.

**Dosing recommendations are based on pharmacokinetic data, published studies, and experience.** CVVH dosing recommendations represent general guidelines, since antibiotic removal is dependent on area/type of filter, ultrafiltration rates, and sieving coefficients; replacement dosing should be individualized and guided by serum levels, if possible. **Creatinine clearance (CrCl) is used to gauge the degree of renal insufficiency, and can be estimated by the following calculation:  $\text{CrCl (mL/min)} = [(140 - \text{age}) \times \text{weight (kg)}] / [72 \times \text{serum creatinine (mg/dL)}]$ . The calculated value is multiplied by 0.85 for females.** It is important to recognize that due to age-dependent decline in renal function, elderly patients with “normal” serum creatinines may have low CrCls requiring dosage adjustments. (For example, a 70-year-old, 50-kg female with a serum creatinine of 1.2 mg/dL has an estimated CrCl of 34 mL/min.)

Slow extended daily dialysis (SLEDD) is now being used in some critical care patients. It is less likely to remove molecules > 1000 Daltons as compared to CVVH. Assuming a dialysate flow rate of 100 mL/min and a blood flow rate of 200 mL/min clearances during dialysis are about 82-98 mL/min using a Fresenius AV600 polysulfone hemodialyzer. However increasing the flow rate to 300 mL/min increases the clearance during dialysis to 170 mL/min. Therefore, doses during SLEDD may be the same as for normal renal function. **Clearances for SLEDD are comparable to CVVH except over a shorter period, so when no information is available, dosage adjustments can be made as for CVVH.** If needed, post-SLEDD doses should be given depending on the antibiotic.

“Antiretroviral Dosage Adjustment” grids indicate recommended dosage adjustments when protease inhibitors (PIs) and non-nucleoside reverse transcriptase inhibitor (NNRTIs) are combined or used in conjunction with rifampin or rifabutin.

**Drug interactions. Refers to common/important drug-drug interactions,** compiled from various sources. If a specific drug interaction is well-documented (e.g., antibiotic × with lovastatin), than other drugs from the same drug class (e.g., atorvastatin) may also be listed, based on theoretical considerations. Drug interactions may occur as a consequence of altered absorption (e.g., metal ion chelation of tetracycline), altered distribution (e.g., sulfonamide displacement of barbiturates from serum albumin), altered excretion (e.g., probenecid competition with penicillin for active transport in the kidney), altered metabolism (e.g., rifampin-induced hepatic P-450 metabolism of theophylline/warfarin; chloramphenicol inhibition of phenytoin metabolism).

**Adverse side effects.** Common/important side effects are indicated.

**Allergic potential.** Described as low or high. **Refers to the likelihood of a hypersensitivity reaction** to a particular antimicrobial.

**Safety in pregnancy.** Designated by the U.S. Food and Drug Administration’s (USFDA) use-in-pregnancy letter code (Table 11.3).

**Comments.** Includes other useful information for each antimicrobial agent.

**Cerebrospinal fluid penetration.** Expressed as a **percentage relative to serum concentration.** If an antimicrobial is used for CNS infections, the meningeal dose is indicated directly above CSF penetration. No meningeal dose is given if CSF penetration is inadequate for treatment of meningitis due to susceptible organisms.

**Biliary tract penetration.** Expressed as a percentage relative to peak serum concentrations. Percentages > 100% reflect concentrations within the biliary system. This information is useful for estimating biliary tract concentrations.

**Table 11.1 Selected Substrates, Inhibitors, and Inducers of Cytochrome P450 Isoenzymes**

	Substrates	Inhibitors	Inducers
CYP1A2	None	Erythromycin	None
CYP2C9	None	Erythromycin, INH, Metronidazole, Oritavancin	TMX-SMX, RIF
CYP3A4	Isavuconazole	Erythromycin, Isavuconazole	Oritavancin
OAT1	Oseltamivir, Cidofovir	None	None
PGP	Quinolones, Itraconazole	Isavuconazole	RIF

OAT-1 = organic anion transporter-1; TMP-SMX = trimethoprim sulfamethoxazole; RIF = rifampin; INH = isoniazid.

**Table 11.2 Antibiotics and Cytochrome P450 Isoenzymes**

Antibiotics that are <b>not</b> involved in CYP450 system	Antibiotics that are <b>involved</b> in CYP450 system either as inducer or inhibitor
Tigecycline, daptomycin, clindamycin, linezolid, vancomycin, minocycline, cephalosporins, carbapenems, nitrofurantoin, colistin, penicillins, oxacillin, aztreonam, moxifloxacin, aminoglycosides, tedizolid, telavancin, dalbavancin, ceftolozane/tazobactam, ceftazidime/avibactam	Nafcillin, quinupristin/dalfopristin, macrolides, telithromycin, fluoroquinolones (except moxifloxacin), azoles, rifampin, INH, TMP-SMX, tetracycline, doxycycline, metronidazole, chloramphenicol, oritavancin

**Table 11.3 USFDA Use-in-Pregnancy Letter Code**

Category	Interpretation
A	<b>Controlled studies show no risk.</b> Adequate, well-controlled studies in pregnant women have not shown a risk to the fetus in any trimester of pregnancy.
B	<b>No evidence of risk in humans.</b> Adequate, well-controlled studies in pregnant women have not shown increased risk of fetal abnormalities despite adverse findings in animals, or, in the absence of adequate human studies, animal studies show no fetal risk. The chance of fetal harm is remote, but remains a possibility.
C	<b>Risk cannot be ruled out.</b> Adequate, well-controlled human studies are lacking, and animal studies have shown a risk to the fetus or are lacking. There is a chance of fetal harm if the drug is administered during pregnancy, but potential benefit from use of the drug may outweigh potential risk.
D	<b>Positive evidence of risk.</b> Studies in humans or investigational or post-marketing data have demonstrated fetal risk. Nevertheless, potential benefit from use of the drug may outweigh potential risk. For example, the drug may be acceptable if needed in a life-threatening situation or serious disease for which safer drugs cannot be used or are ineffective.

**Table 11.3 USFDA Use-in-Pregnancy Letter Code** (Cont'd)

Category	Interpretation
X	<b>Contraindicated in pregnancy.</b> Studies in animals or humans or investigational or post-marketing reports have demonstrated positive evidence of fetal abnormalities or risk which clearly outweigh any possible benefit to the patient.

**Lipid-Associated Formulations of Amphotericin B.** There are 3 licensed lipid-associated formulations of amphotericin B (LFAB) (Table 11.4). Although closely related in some ways, these formulations have distinct properties and must be understood separately. The principal advantage of the LFAB over amphotericin B deoxycholate (AMBD) is greater safety. In general, the rates of both acute infection-related toxicities (fever, chills, etc.) and chronic therapy-associated toxicities (principally nephrotoxicity) are reduced with LFAB. However, the LFAB can produce all of the toxicities of AMBD (and in selected patients, LFAB have been more toxic than AMBD). Overall, (L-Amb) (AmBisome) and ABLC (Abelcet) appear to be safer than ABCD (Amphotec, Amphocil). Whichever formulation is selected for therapy, it is important to specify its name carefully when prescribing. The phrase "lipid amphotericin B" should be avoided due to its imprecision. Patients who are tolerating one formulation may develop all the standard infusion-related toxicities if switched inadvertently to a new formulation. In general (and in contrast to the usual preference for generic names), use of trade names is the clearest way to specify the choice of drug in this category. In this handbook, the phrase "lipid-associated formulation of amphotericin B" suggests use of any of the 3 formulations. The issues surrounding the selection of an LFAB for an individual patient are summarized in Table 11.4 (see comments).

**Table 11.4 Lipid-Associated Formulations of Amphotericin B**

Generic name (abbreviation)	Trade names	Licensed (IV) dosages in the United States	Comments
Amphotericin B lipid complex (ABLC)	Abelcet	5 mg/kg/d	Reliable choice; long history of use.
Liposomal amphotericin B (L-Amb)	AmBisome	3 mg/kg/d (empiric therapy) 3–5 mg/kg/d (systemic fungal infections) 6 mg/kg/d (cryptococcal meningitis in HIV patients)	Reliable choice; best studied (L-Amb); well-supported dosing recommendations by indication; probably the least nephrotoxic; good data to support increasing the dose safely.
Amphotericin B colloidal dispersion, amphotericin B cholesteryl sulfate complex (ABCD)	Amphotec, Amphocil	3–4 mg/kg/d	Infusion-related toxicities have limited its use.

## Abacavir (Ziagen) ABC

**Drug Class:** HIV NRTI (nucleoside reverse transcriptase inhibitor).

**Usual Dose:** 300 mg (PO) q12h.

**Pharmacokinetic Parameters:**

Peak serum level: 3 mcg/mL

Bioavailability: 83%

Excreted unchanged (urine): 1.2%

Serum half-life (normal/ESRD): 1.5/8 hrs

Plasma protein binding: 50%

Volume of distribution ( $V_d$ ): 0.86 L/kg

**Primary Mode of Elimination:** Hepatic

**Dosage Adjustments\***

CrCl < 10 mL/min	No change
Post-HD dose	None
Post-PD dose	None
CWH/CVHD/ CWHDF dose	No change
Mild hepatic insufficiency	200 mg (PO) q12h
Moderate—severe hepatic insufficiency	Avoid

**Drug Interactions:** Methadone (↑ methadone clearance with abacavir 600 mg bid); ethanol (↑ abacavir serum levels/half-life and may ↑ toxicity).

**Adverse Effects:** *Abacavir may cause severe hypersensitivity reactions which may be fatal (see comments); usually during the first 4–6 weeks of therapy; Patients who carry the HLA-B\*5701 allele are at high risk for experiencing a hypersensitivity reaction to abacavir. Discontinue abacavir as soon*

as a hypersensitivity reaction is suspected. Regardless of HLA-B\*5701 status, permanently discontinue abacavir if hypersensitivity cannot be ruled out, even when other diagnoses are possible. Following a hypersensitivity reaction to abacavir, never restart abacavir or any other abacavir-containing product. Report cases of hypersensitivity syndrome to Abacavir Hypersensitivity Registry at 1-800-270-0425. Drug fever/rash, abdominal pain/diarrhea, nausea, vomiting, anorexia, bad dreams/sleep disorders, weakness, headache, ↑ SGOT/SGPT, hyperglycemia, hypertriglyceridemia, lactic acidosis with hepatic steatosis (rare, but potentially life-threatening toxicity with use of NRTI's).

**Allergic Potential:** High (~5%)

**Safety in Pregnancy:** C

**Comments:** Discontinue abacavir and *do not restart in patients with signs/symptoms of hypersensitivity reaction*, which may include fever, rash, fatigue, nausea, vomiting, diarrhea, abdominal pain, anorexia, respiratory symptoms. Ethanol increases abacavir levels by 41%. When combined with didanosine rapid emergence of cross resistance via mutations to K65R, L74V, Y11F, and M184V occurs leading to failure of drug efficacy. Use with caution in patients with serious risk of coronary disease.

**Cerebrospinal Fluid Penetration:** 27–33%

### REFERENCES:

- Bedimo RJ, Westfall AO, Drechsler H, et al. Abacavir use and risk of acute myocardial infarction and cerebrovascular events in the highly active antiretroviral therapy era. *Clin Infect Dis* 53:83–94, 2011.
- Carr A, Workman C, Smith DE, et al. Abacavir substitution for nucleoside analogs in patients with HIV lipodystrophy. A randomized trial. *JAMA* 288:207–15, 2002.

\*Usual dose\* assumes normal renal/hepatic function. \*For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

Katalama C, Clotet B, Plettenberg A, et al. The role of abacavir (AVC, 1592) in antiretroviral therapy-experiences patients: results from randomized, double-blind, trial. CNA3002 European Study Team. *AIDS* 14:781–9, 2000.

Keating MR. Antiviral agents. *Mayo Clin Proc* 67:160–78, 1992.

Nishijima T, Tsukada K, Teruya K, et al. Efficacy and safety of once-daily ritonavir-based antiretroviral therapy in low-body weight treatment-naïve patients with HIV infections. *AIDS* 13:26:649–51, 2012.

Panel on Antiretroviral Guidelines for Adults and Adolescents. Department of Health and Human Services. Guidelines for the use of antiretroviral agents in HIV-infected adults and adolescents. <http://aidsinfo.nih.gov/ConsentFiles/AdultandAdolescentGL.pdf>, 2012.

Sax PE, Tierney C, Collier AC, et al. Abacavir/lamivudine versus tenofovir DF/emtricitabine as part of combination regimens for initial treatment of HIV: final results. *J Infect Dis* 204:1191–201, 2011.

Staszewski S, Keiser P, Mantaner J, et al. Abacavir-lamivudine-zidovudine vs. indinavir-lamivudine-zidovudine in antiretroviral-naïve HIV-infected adults: a randomized equivalence trial. *JAMA* 285:1155–63, 2001.

Thompson MA, Aberg JA, Cahn P, et al. Antiretroviral treatment of adult HIV infection: 2012 recommendations of the International Antiviral society — USA panel. *JAMA* 308:387–402, 2012.

Website: [www.pdr.net](http://www.pdr.net)

## Abacavir + Lamivudine (Epzicom) ABC/3TC

**Drug Class:** HIV NRTI combination.

**Usual Dose:** Epzicom tablet = abacavir 600 mg + lamivudine 300 mg. Usual dose: 1 tablet q24h.

**Pharmacokinetic Parameters:**

*Peak serum level:* 4.06/2.04 mcg/L

*Bioavailability:* 86/86%

*Excreted unchanged (urine):* 1.2/71%

*Serum half-life (normal/ESRD):* (1.5/8)/(5-7/20) hrs

*Plasma protein binding:* 50/36%

*Volume of distribution ( $V_d$ ):* 0.86/1.3 L/kg

**Primary Mode of Elimination:** Hepatic/Renal  
**Dosage Adjustments\***

CrCl < 50 mL/min	Avoid
Post-HD dose	Avoid
Post-PD dose	Avoid
CVWH/CVWHD/CVWHDf dose	Avoid
Mild hepatic insufficiency	Contraindicated
Moderate—severe hepatic insufficiency	Contraindicated

**Drug Interactions:** Methadone (↑ methadone clearance with abacavir 600 mg bid); ethanol (↑ abacavir serum levels/half-life; may ↑ toxicity); didanosine, zalcitabine (↑ risk of pancreatitis); TMP-SMX (↑ lamivudine levels); zidovudine (↑ zidovudine levels).

**Adverse Effects: Abacavir may cause severe hypersensitivity reactions that may be fatal (see comments), usually during the first 4–6 weeks of therapy;** Patients who carry the HLA-B\*5701 allele are at high risk for experiencing a hypersensitivity reaction to abacavir. Discontinue Abacavir as soon as a hypersensitivity reaction is suspected. Regardless of HLA-B\*5701 status, permanently discontinue Abacavir if hypersensitivity cannot be ruled out, even when other diagnoses are possible. Following a hypersensitivity reaction to abacavir, NEVER restart Abacavir or any other abacavir-containing product. Drug fever, rash, abdominal pain, diarrhea, nausea, vomiting, anorexia, anemia, leukopenia, photophobia, depression, insomnia, weakness, headache,

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

cough, nasal complaints, dizziness, peripheral neuropathy, myalgias, ↑ AST/ALT, hyperglycemia, hypertriglyceridemia, pancreatitis, lactic acidosis with hepatic steatosis (rare, but potentially life-threatening toxicity with the NRTI's). Immune reconstitution syndrome and redistribution/accumulation of body fat have been reported in patients treated with combination antiretroviral therapy.

**Allergic Potential:** High (~5%)/Low

**Safety in Pregnancy:** C

**Comments:** In patients with signs/symptoms of hypersensitivity reactions discontinue and do not restart, which may include fever, rash, fatigue, nausea, vomiting, diarrhea, abdominal pain, anorexia, respiratory symptoms. **Potential cross resistance with didanosine.** When combined with didanosine rapid emergence of cross resistance via mutations to K65R, L74V, Y115F, and M184V occurs leading to failure of drug efficacy. Severe acute exacerbations of hepatitis B have been reported in patients who are co-infected with hepatitis B virus (HBV) and HIV and have discontinued epivir. Hepatic function should be monitored closely for at least several months in patients who discontinue epivir.

**Cerebrospinal Fluid Penetration:** 27–33/15%

#### REFERENCES:

Panel on Antiretroviral Guidelines for Adults and Adolescents. Department of Health and Human Services. Guidelines for the use of antiretroviral agents in HIV-infected adults and adolescents. <http://aidsinfo.nih.gov/ConsentFiles/AdultandAdolescentGL.pdf>, 2012.

Thompson MA, Aberg JA, Cahn P, et al. Antiretroviral treatment of adult HIV infection: 2012 recommendations of the International Antiviral society — USA panel. *JAMA* 308:387–402, 2012.

Website: [www.pdr.net](http://www.pdr.net)

## Abacavir + Lamivudine + Zidovudine (Trizivir) ABC/3TC/ZDV

**Drug Class:** HIV NRTI combination.

**Usual Dose:** Trizivir tablet = abacavir 300 mg + lamivudine 150 mg + zidovudine 300 mg. Usual dose = 1 tablet (PO) q12h.

**Pharmacokinetic Parameters:**

*Peak serum level:* 3/1.5/1.2 mcg/mL

*Bioavailability:* 86/86/64%

*Excreted unchanged (urine):* 1.2/90/16%

*Serum half-life (normal/ESRD):* [1.5/6/1.1] / 8/20/2.2] hrs

*Plasma protein binding:* 50/36/20%

*Volume of distribution (V<sub>d</sub>):* 0.86/1.3/1.6 L/kg

**Primary Mode of Elimination:** Hepatic/Renal  
**Dosage Adjustments\***

CrCl < 50 mL/min	Avoid
Post-HD or Post-PD	Avoid
CVWH/CVWHD/ CVWHDf dose	Avoid
Moderate—severe hepatic insufficiency	Avoid

**Drug Interactions:** Amprenavir, atovaquone (↑ zidovudine levels); clarithromycin (↓ zidovudine levels); cidofovir (↑ zidovudine levels, flu-like symptoms); doxorubicin (neutropenia); stavudine (antagonistic to zidovudine; avoid combination); TMP-SMX (↑ lamivudine and zidovudine levels); zalcitabine (↓ lamivudine levels). May exacerbate neutropenia in combination with gancyclovir.

**Adverse Effects:** *Abacavir may cause severe/fatal rash/hypersensitivity reaction;* Patients who carry the HLA-B\*5701 allele are at high risk

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

for experiencing a hypersensitivity reaction to abacavir. Discontinue abacavir as soon as a hypersensitivity reaction is suspected. Regardless of HLA-B\*5701 status, permanently discontinue abacavir if hypersensitivity cannot be ruled out, even when other diagnoses are possible. Following a hypersensitivity reaction to abacavir, never restart abacavir or any other abacavir-containing product. Most common (> 5%): nausea, vomiting, diarrhea, anorexia, insomnia, fever/chills, headache, malaise/fatigue. Others (less common): peripheral neuropathy, myopathy, steatosis, pancreatitis. Lab abnormalities: anemia, leukopenia, mild hyperglycemia, ↑ LFTs, ↑ CPK, ↑ aldolase, hypertriglyceridemia. Immune reconstitution syndrome and redistribution/accumulation of body fat have been reported in patients treated with combination antiretroviral therapy.

**Allergic Potential:** High (~5%)

**Safety in Pregnancy:** C

**Comments:** Avoid in patients with CrCl < 50 mL/min. HBV hepatitis may relapse if lamivudine is discontinued.

#### REFERENCES:

McDowell JA, Lou Y, Symonds WS, et al. Multiple-dose pharmacokinetics and pharmacodynamics of abacavir alone and in combination with zidovudine in human immunodeficiency virus-infected adults. *Antimicrob Agents Chemother* 44:2061–7, 2000.

Panel on Antiretroviral Guidelines for Adults and Adolescents. Department of Health and Human Services. Guidelines for the use of antiretroviral agents in HIV-infected adults and adolescents. <http://aidsinfo.nih.gov/ConsentFiles/AdultandAdolescentGL.pdf>, 2012.

Thompson MA, Aberg JA, Cahn P, et al. Antiretroviral treatment of adult HIV infection: 2012 recommendations of the International Antiviral society — USA panel. *JAMA* 308:387–402, 2012.

Website: [www.pdr.net](http://www.pdr.net)

## Acyclovir (Zovirax)

**Drug Class:** Antiviral (see comments).

**Usual Dose:**

**HSV-1/2:** Herpes labialis: 400 mg (PO) 5x/day × 5 days. Genital herpes: *Initial therapy:* 200 mg (PO) 5x/day × 10 days. *Recurrent/intermittent therapy* (< 6 episodes/year): 200 mg (PO)

5x/day × 5 days. *Chronic suppressive therapy* (> 6 episodes/year): 400 mg (PO) q12h × 1 year.

Mucosal/genital herpes: 5 mg/kg (IV) q8h × 7 days or 400 mg (PO) 5x/day × 7 days. Nosocomial pneumonia 5 mg/kg (IV) q8h × 10 days or 400 mg (PO) 5x/day × 7 days. Meningitis/encephalitis: 10 mg/kg (IV) q8h × 10 days\*. PO not recommended. \*Severe HSV encephalitis may require 14–21 days.

**VZV:** Chickenpox: 10 mg/kg (IV) q8h × 5 days or 800 mg (PO) q6h × 5 days. VZV pneumonia: 5–10 mg/kg (IV) q8h × 10 days (normal hosts) 10 mg/kg (IV) q8h × 10 days (compromised hosts). Herpes zoster (shingles): *Dermatomal/Disseminated:* 10 mg/kg (IV) q8h × 7–10 days or 800 mg (PO) 5x/day × 7–10 days.

VZV meningitis/encephalitis: 10 mg/kg (IV) q8h × 10 days. PO not recommended.

#### Pharmacokinetic Parameters:

*Peak serum level:* 7.7 mcg/mL

*Bioavailability:* 30%

*Excreted unchanged (urine):* 70%

*Serum half-life (normal/ESRD):* 3/5 hrs

*Plasma protein binding:* 30%

*Volume of distribution (V<sub>d</sub>):* 0.7 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\* for HSV/VZV**

CrCl 10–25 mL/min	No change/ 800 mg (IV/PO) q8h
CrCl < 10 mL/min	400 mg (IV/PO) q12h/ 800 mg (IV/PO) q12h



Post-HD dose	None
Post-PD dose	None
CWVH/CVVHD/ CWVHD dose	5 mg/kg(IV) q24h/ 10 mg/kg(IV) q24h 400 mg (PO) q8h/ 800 mg (PO) q8h
Moderate—severe hepatic insufficiency	No change

**Drug Interactions:** Cimetidine, probenecid, theophylline (↑ acyclovir levels); nephrotoxic drugs (↑ nephrotoxicity); zidovudine (lethargy); theophylline (↑ levels).

**Adverse Effects:** Seizures/tremors (dose related), encephalopathy, crystalluria, ATN. HUS (only with HIV) Base dose on ideal body weight in the elderly to minimize adverse effects.

**Allergic Potential:** Low

**Safety in Pregnancy:** B

**Comments:** Na<sup>+</sup> content = 4 mEq/g. CSF levels may be increased with probenecid.

**Highly active against HSV and VZV. Some activity against CMV. No activity against EBV, RSV or adenoviruses.**

Meningeal dose = HSV or VZV encephalitis dose.

**Cerebrospinal Fluid Penetration:** 20%

#### REFERENCES:

- Gupta R, Wald A, Krantz E, et al. Valacyclovir and acyclovir for suppression of shedding of herpes simplex virus in the genital tract. *J Infect Dis* 190:1374–81, 2004.
- Hehao-Martinez AF, Weinberg A, Waldman WJ, et al. Successful treatment of acyclovir-resistant herpes simplex virus type 2 proctitis with leflunomide in an HIV-infected man. *J Clin Virol* 54:276–278, 2012.
- Keating MR. *Antiviral Agents*. *Mayo Clin Proc* 67:160–78, 1992.

Mugwanya K, Baeten JM, Mugo NR, et al. High-dose valacyclovir HSV-2 suppression results in greater reduction in plasma HIV-1 levels compared with standard dose acyclovir among HIV-1/HSV-2 coinfecting persons: a randomized, crossover trial. *J Infect Dis* 204:1912–1017, 2011.

Owens RC, Ambrose PG. Acyclovir. *Antibiotics for Clinicians* 1:85–93, 1997.

Vanpouille C, Lisco A, Introini A, et al. Exploiting the anti-HIV-1 activity of acyclovir: suppression of primary and drug-resistant HIV isolates and potentiation of the activity by ribavirin. *Antimicrob Agents Chemother* 56:2604–2611, 2012.

Vanpouille C, Lisco A, Margolis L. Acyclovir: a new use for an old drug. *Curr Opin Infect Dis* 22:583–587, 2009.

Whitley RJ, Gnann JW Jr. Acyclovir: a decade later. *N Engl J Med* 327:782–3, 1992.

Website: [www.pdr.net](http://www.pdr.net)

## Adefovir dipivoxil (Hepsera)

**Drug Class:** HBV antiviral.

**Usual Dose:** 10 mg (PO) q24h.

**Pharmacokinetic Parameters:**

*Peak serum level:* 18 ng/mL

*Bioavailability:* 59%

*Excreted unchanged (urine):* 45%

*Serum half-life (normal/ESRD):* 7.5/9 hrs

*Plasma protein binding:* 4%

*Volume of distribution (V<sub>d</sub>):* 0.4 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\***

CrCl ≥ 50 mL/min	10 mg (PO) q24h
CrCl 20–50 mL/min	10 mg (PO) q48h
CrCl 10–20 mL/min	10 mg (PO) q72h
Hemodialysis	10 mg (PO) q7days
Post-HD or PD dose	10 mg (PO)

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

CVVH/CVVHD/ CVVHDF dose	10 mg (PO) q72h
Moderate—severe hepatic insufficiency	No change

**Drug Interactions:** No significant interaction with lamivudine, TMP–SMX, acetaminophen, ibuprofen.

**Adverse Effects:** Asthenia, headache, abdominal pain, nausea, flatulence, diarrhea, dyspepsia.

**Allergic Potential:** Low

**Safety in Pregnancy:** C

**Comments:** May be taken with or without food. Does not inhibit CP450 isoenzymes. Do not discontinue abruptly to avoid exacerbation of HBV hepatitis.

**Cerebrospinal Fluid Penetration:** No data

#### REFERENCES:

- Buti M, Esteban R. Adefovir dipivoxil. *Drugs of Today* 39:127–35, 2003.
- Cundy KC, Burditch-Crovo P, Walker RE, et al. Clinical pharmacokinetics of adefovir in human HIV-1 infected patients. *Antimicrob Agents Chemother* 35:2401–2405, 1995.
- Lu H, Geng DY, Shen F, et al. Optimization of adefovir dipivoxil therapy in chronic hepatitis B according to baseline predictors and on-treatment HBV DNA: a 5-Year prospective study. *Virology* 8:444, 2011.
- Segovia MC, Chacra W, Gordon SC. Adefovir dipivoxil in chronic hepatitis B: history and current uses. *Expert Opin Pharmacother* 13:245–254, 2012.
- Sheng YJ, Liu JY, Tong SW, et al. Lamivudine plus adefovir combination therapy versus entecavir monotherapy for lamivudine-resistant chronic hepatitis B: a systematic review and meta-analysis. *Virology* 8:393, 2011.
- Yang HJ, Lee JH, Kim YJ, et al. Antiviral efficacy of combination therapy with entecavir and adefovir for entecavir/lamivudine-resistant hepatitis B virus with or without adefovir resistance. *J Med Virol* 84:424–430, 2012.
- Website: www.pdr.net

## Amantadine (Symmetrel)

**Drug Class:** Antiviral (influenza).

**Usual Dose:** 200 mg (PO) q24h.

**Pharmacokinetic Parameters:**

*Peak serum level:* 0.5 mcg/mL

*Bioavailability:* 90%

*Excreted unchanged (urine):* 90%

*Serum half-life (normal/ESRD):* 16/192 hrs

*Plasma protein binding:* 67%

*Volume of distribution ( $V_d$ ):* 6.6 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\***

CrCl > 30 mL/min	100 mg (PO) q24h
CrCl 15–30 mL/min	100 mg (PO) q48h
CrCl < 15 mL/min	200 mg (PO) q week
Post–HD	None
Post–PD	None
CVVH/CVVHD/ CVVHDF dose	100 mg (PO) q48h
Moderate—severe hepatic insufficiency	No change

**Drug Interactions:** Alcohol (↑ CNS effects); benztrapine, trihexyphenidyl (↑ interacting drug effect: dry mouth, ataxia); CNS stimulants (additive stimulation); digoxin (↑ digoxin levels); trimethoprim (↑ amantadine and trimethoprim levels); scopolamine (↑ scopolamine effect: blurred vision, slurred speech, toxic psychosis).

**Adverse Effects:** Confusion/delusions, insomnia, dysarthria, ataxia, anticholinergic effects (blurred vision, dry mouth, orthostatic hypotension, urinary retention, constipation), livedo reticularis, may ↑ QT<sub>c</sub> interval.

**Allergic Potential:** Low

**Safety in Pregnancy:** C

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

**Comments:** May precipitate heart failure. Avoid co-administration with anticholinergics, MAO inhibitors, or antihistamines. Resistant to amantadine, widespread may improve peripheral airway function/oxygenation in ventilated patients with severe influenza (human, avian, swine).

**Highly active against influenza A. No activity against other viruses.**

### Cerebrospinal Fluid Penetration:

Non-inflamed meninges = 15%

Inflamed meninges = 20%

### REFERENCES:

- Cunha BA, Amantadine may be lifesaving in severe influenza A. *Clin Infect Dis* 43:1372–3, 2006.
- Gubareva LV, Kaiser L, Hayden FG. Influenza virus neuraminidase inhibitors. *Lancet* 355:827–5, 2000.
- Jackson RJ, Cooper KL, Tappenden P, et al. Oseltamivir, zanamivir and amantadine in the prevention of influenza: A systematic review. *J Infect* 62:14–25, 2011.
- Kawai N, Ikematsu H, Iwaki N, et al. Factors influencing the effectiveness of oseltamivir and amantadine for the treatment of influenza: A multicenter study from Japan of the 2002–2003 influenza season. *Clin Infect Dis* 40:1309–16, 2005.
- Keyers LA, Karl M, Nafziger AN, et al. Comparison of central nervous system adverse effects of amantadine and rimantadine used as sequential prophylaxis of influenza A in elderly nursing home patients. *Arch Intern Med* 160:1485–8, 2000.
- Website: [www.pdr.net](http://www.pdr.net)

## Amikacin (Amikin)

**Drug Class:** Aminoglycoside.

**Usual Dose:** 1 gm or 15 mg/kg (IV) q24h (preferred to q12h dosing).

**Spectrum:** (see **Susceptibility Profiles** pp. 198–202).

**Resistance Potential:** Low

### Pharmacokinetic Parameters:

Peak serum level 65–75 mcg/mL (q24h dosing);  
20–30 mcg/mL (q12h dosing)

*Bioavailability:* Not applicable

*Excreted unchanged (urine):* 95%

*Serum half-life (normal/ESRD):* 2/50 hrs

*Plasma protein binding:* < 5%

*Volume of distribution ( $V_d$ ):* 0.25 L/kg

**Primary Mode of Elimination:** Renal  
**Dosage Adjustments\***

CrCl 50–80 mL/min	500 mg (IV) q24h or 7.5 mg/kg (IV) q24h
CrCl 10–50 mL/min	500 mg (IV) q48h or 7.5 mg/kg (IV) q48h
CrCl < 10 mL/min	250 mg (IV) q48h or 3.75 mg/kg (IV) q48h
Post–HD dose	500 mg (IV) or 7.5 mg/kg (IV)
Post–HFHD dose	500 mg (IV) or 7.5 mg/kg (IV)
Post–PD dose	250 mg (IV) or 3.75 mg/kg (IV)
CVVH/CVVHD/ CVVHDF dose	500 mg (IV) q48h or 7.5 mg/kg (IV)
Moderate—severe hepatic insufficiency	No change

**Drug Interactions:** Amphotericin B, cephalothin, cyclosporine, enflurane, methoxyflurane, NSAIDs, polymyxin B, radiographic contrast, vancomycin (↑ nephrotoxicity); cis-platinum (↑ nephrotoxicity, ↑ ototoxicity); loop diuretics (↑ ototoxicity); neuromuscular blocking agents (↑ apnea, prolonged paralysis); non-polarizing muscle relaxants (↑ apnea).

**Adverse Effects:** Neuromuscular blockade with rapid infusion/absorption. Nephrotoxicity only with prolonged/extremely high serum trough levels; may cause reversible non-oliguric

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

renal failure (ATN). Ototoxicity associated with prolonged/extremely high peak serum levels (usually irreversible): Cochlear toxicity (1/3 of ototoxicity) manifests as decreased high frequency hearing, but deafness is unusual. Vestibular toxicity (2/3 of ototoxicity) develops before ototoxicity (typically manifests as tinnitus).

**Allergic Potential:** Low

**Safety in Pregnancy:** D

**Comments: Synergy dose:** 500 mg (IV) q24h or 7.5 mg/kg (IV) q24h. **Single daily dosing virtually eliminates nephrotoxic/ototoxic potential.** Incompatible with solutions containing  $\beta$ -lactams, erythromycin, chloramphenicol, furosemide, sodium bicarbonate. IV infusion should be given slowly over 30 minutes. May be given IM. Intraperitoneal infusion  $\uparrow$  risk of neuromuscular blockade. Avoid intratracheal/aerosolized intrapulmonary instillation, which may predispose to antibiotic resistance.  $V_d$  increases with edema/ascites, trauma, burns, cystic fibrosis; may require  $\uparrow$  dose.  $V_d$  decreases with dehydration, obesity; may require  $\downarrow$  dose.

**Renal cast counts are the best indicator of aminoglycoside nephrotoxicity, not serum creatinine.** Dialysis removes ~ 50% of amikacin from serum. **CAPD dose:** 10–20 mg/L in dialysate (IP) with each exchange.

#### Therapeutic Serum Concentrations

(for therapeutic efficacy, *not toxicity*): Peak (q24h/q12h dosing): 65–75/20–30 mcg/mL  
Trough (q24h/q12h dosing): 0/4–8 mcg/mL

**Intrathecal (IT) dose:** 10–40 mg (IT) q24h.

#### Cerebrospinal Fluid Penetration:

Non-Inflamed meninges = 15%

Inflamed meninges = 20%

**Bile Penetration:** 30%

#### REFERENCES:

- Cunha BA. New uses for older antibiotics: nitrofurantoin, amikacin, colistin, polymyxin B, doxycycline, and minocycline revisited. *Med Clin North Am* 90:1089–107, 2006.
- Cunha BA. *Pseudomonas aeruginosa*: Resistance and therapy. *Semin Respir Infect* 17:231–9, 2002.
- Cunha BA. Aminoglycosides: Current role in antimicrobial therapy. *Pharmacotherapy* 8:334–50, 1988.
- Edson RS, Terrel CL. The Aminoglycosides. *Mayo Clin Proc* 74:519–28, 1999.
- Fulnecky E, Wright D, Scheld WM, et al. Amikacin and colistin for treatment of *Acinetobacter baumannii* meningitis. *Journal of Infection* 51:e249–e251, 2005.
- Layeux B, Taccone FS, Fagnoul D, et al. Amikacin monotherapy for sepsis caused by panresistant *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother* 54:4939–4941, 2010.
- Le J, McKee B, Srisupha-Olam W, Burgess DS. In vitro activity of carbapenems alone and in combination with amikacin against KPC-producing *Klebsiella pneumoniae*. *J Clin Med Res* 3:106–110, 2011.
- Pankuch GA, Seifert H, Appelbaum PC. Activity of doripenem with and without levofloxacin, amikacin, and colistin against *Pseudomonas aeruginosa* and *Acinetobacter baumannii*. *Diagn Microbiol Infect Dis* 67:191–197, 2010.
- Taccone FS, de Backer D, Laterre PF, et al. Pharmacokinetics of a loading dose of amikacin in septic patients undergoing continuous renal replacement therapy. *Int J Antimicrob Agents* 37:531–535, 2011.
- Taccone FS. Optimizing amikacin regimens in septic patients. *Int J Antimicrob Agents*. 39:264–264, 2012.
- Website: [www.pdr.net](http://www.pdr.net)

## Amoxicillin (Amoxil, A-cillin, Polymox, Trimox, Wymox)

**Drug Class:** Aminopenicillin.

**Usual Dose:** 1 gm (PO) q8h.

**Spectrum:** (see *Susceptibility Profiles* pp. 186–190).

<sup>†</sup>Usual dose<sup>†</sup> assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose  $\times$  1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

**Resistance Potential:** Low

**Pharmacokinetic Parameters:**

Peak serum level: 14 mcg/mL

Bioavailability: 90%

Excreted unchanged (urine): 60%

Serum half-life (normal/ESRD): 1.3/16 hrs

Plasma protein binding: 20%

Volume of distribution ( $V_d$ ): 0.26 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\***

CrCl 50–80 mL/min	500 mg (PO) q8h
CrCl 10–50 mL/min	500 mg (PO) q12h
CrCl < 10 mL/min	500 mg (PO) q24h
Post-HD or post-PD	500 mg
CWH/CVHD/ CWHDF dose	500 mg (PO) q24h
Moderate—severe hepatic insufficiency	No change

**Drug Interactions:** Allopurinol (↑ risk of rash).

**Adverse Effects:** Drug fever/rash, ↑ SGOT/SGPT.

**Allergic Potential:** High

**Safety in Pregnancy:** B

**Comments:** ↑ risk of rash with EBV infectious mononucleosis. No irritative diarrhea with 1 gm (PO) q8h dose due to nearly complete proximal GI absorption. **2 gm amoxicillin effective against most strains of PRSP.**

**Cerebrospinal Fluid Penetration:**

Non-inflamed/Inflamed meninges = 1%/8%

**Bile Penetration:** 3000%

(also see *Antibiotic Pearls & Pitfalls* p. 508).

**REFERENCES:**

Addo-Yobo E, Chisaka N, Hassan M, et al. Oral amoxicillin versus injectable penicillin for severe pneumonia in children age 3 to 59 months: a randomized multicentre equivalency study. *Lancet* 364:1141–48, 2004.

Camacho MT, Casal J, AGullar L, et al. Very high resistance to amoxicillin in *Streptococcus pneumoniae*: an epidemiological fact or a technical issue? *Journal of Chemotherapy* 18:303–06, 2006.

Cunha BA. Oral antibiotic therapy of serious systemic infections. *Med Clin N Am* 90:1197–1222, 2006.

Cunha BA. The aminopenicillins. *Urology* 40:186–190, 1992.

Curtin-Wirt C, Casey JR, Murray PC, et al. Efficacy of penicillin vs. amoxicillin in children with group A beta hemolytic streptococcal tonsillopharyngitis. *Clin Pediatr* 42:219–24, 2003.

Dhaon NA. Amoxicillin tablets for oral suspension in the treatment of acute otitis media: a new formulation with improved convenience. *Adv Ther* 21:87–95, 2004.

File Jr. TM. Clinical implications and treatment of multidrug-resistant *Streptococcus pneumoniae* pneumonia. *Clin Microbiol Infect* 12:31–41, 2006.

Garbutt JM, Banister C, Spitznagel E, et al. Amoxicillin for acute rhinosinusitis: a randomized controlled trial. *JAMA* 15:685–92, 2012.

Moreillon P, Wilson WR, Leclercq R, et al. Single-dose oral amoxicillin or linezolid for prophylaxis of experimental endocarditis due to vancomycin-susceptible and vancomycin-resistant *Enterococcus faecalis*. *Antimicrob Agents Chemother* 51:1661–1665, 2007.

Piglansky L, Leibovitz E, Raiz S, et al. Bacteriologic and clinical efficacy of high dose amoxicillin for therapy of acute otitis media in children. *Pediatr Infect Dis J* 22:405–13, 2003.

Ready D, Lancaster H, Qureshi F, et al. Effect of amoxicillin use on oral microbiota in young children. *Antimicrob Agents Chemother* 48:2883–7, 2004.

Website: [www.pdr.net](http://www.pdr.net)

## Amoxicillin/Clavulanic Acid (Augmentin)

**Drug Class:** Aminopenicillin/β-lactamase inhibitor.

**Usual Dose:** 500/125 mg (PO) q8h or 875/125 mg (PO) q12h for severe infections or respiratory tract infections.

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

**Spectrum:** (see **Susceptibility Profiles** pp. 186–190).

**Resistance Potential:** Low

**Pharmacokinetic Parameters:**

Peak serum level: 10.0/2.2 mcg/mL

Bioavailability: 90/60%

Excreted unchanged (urine): 80/40%

Serum half-life (normal/ESRD): [1.3/16]/[1/2] hrs

Plasma protein binding: 18/25%

Volume of distribution ( $V_d$ ): 0.26/0.3 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\*** (based on 500 mg q8h)

CrCl 10–50 mL/min	500/125 mg (PO) q12h
CrCl < 10 mL/min	250/125 mg (PO) q24h
Post-HD dose	250/125 mg (PO)
Post-PD dose	250/125 mg (PO)
CVWH/CVVD/ CVHDF dose	500/125 mg (PO) q24h
Moderate—severe hepatic insufficiency	No change

**Drug Interactions:** Allopurinol (↑ risk of rash).

**Adverse Effects:** Drug fever/rash, diarrhea, ↑ SGOT/SGPT. Rash potential same as ampicillin.

**Resistance Potential:** Low

Allergic Potential: High

**Safety in Pregnancy:** B

**Comments:** ↑ risk of rash with EBV infectious mononucleosis. **875/125 mg formulation should not be used in patients with CrCl < 30 mL/min.**

**Cerebrospinal Fluid Penetration:**

Non-Inflamed meninges = 1%

Inflamed meninges = 1%

**Bile Penetration:** 3000%

(also see **Antibiotic Pearls & Pitfalls** p. 508).

#### REFERENCES:

Cunha BA. Amoxicillin/clavulanic acid in respiratory infections: microbiologic and pharmacokinetic considerations. *Clinical Therapeutics* 14:418–25, 1992.

Donowitz GR, Mandell GL. Beta-lactam antibiotics. *N Engl J Med* 318:419–26 and 318:490–500, 1993.

Fernandez-Sabe N, Carratala J, Dorca J, et al. Efficacy and safety of sequential amoxicillin-clavulanate in the treatment of anaerobic lung infections. *Eur J Clin Microbiol Infect Dis* 22:185–7, 2003.

File TM Jr, Lode H, Kurz H, et al. Double-blind, randomized study of the efficacy and safety of oral pharmacokinetically enhanced amoxicillin-clavulanic (2,000/125 milligrams) versus those of amoxicillin-clavulanic (875/125 milligrams), both given twice daily for 7 days, in treatment of bacterial community-acquired pneumonia adults. *Antimicrob Agents Chemother* 48:3323–31, 2004.

Malingoni MA, Song J, Herrington J, et al. Randomized controlled trial of moxifloxacin compared with piperacillin-tazobactam and amoxicillin-clavulanate for the treatment of complicated intra-abdominal infections. *Ann Surg* 244:204–11, 2006.

Scaglione F, Caronzolo D, Pintucci JP, et al. Measurement of cefaclor and amoxicillin-clavulanic acid levels in middle-ear fluid in patients with acute otitis media. *Antimicrob Agents Chemother* 47:2987–9, 2003.

Vons C, Barry C, Maitre S, et al. Amoxicillin plus clavulanic acid versus appendicectomy for treatment of acute uncomplicated appendicitis: an open-label, non-inferiority, randomized controlled trial. *Lancet* 377:1573–1579, 2011.

Wright AJ, Wilkowske CJ. The penicillins. *Mayo Clin Proc* 66:1047–63, 1991.

Website: www.pdr.net

## Amoxicillin/Clavulanic Acid ES-600 (Augmentin ES-600)

**Drug Class:** Aminopenicillin/ $\beta$ -lactamase inhibitor.

**Usual Dose:** 90 mg/kg/day oral suspension in 2 divided doses (see comments).

**Spectrum:** (see **Susceptibility Profiles** pp. 186–190).

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose  $\times$  1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

**Resistance Potential:** Low

**Pharmacokinetic Parameters:**

Peak serum level: 15.7/1.7 mcg/mL

Bioavailability: 90/60%

Excreted unchanged (urine): 70/40%

Serum half-life (normal/ESRD): [1.4/16]/[1.1/2] hrs

Plasma protein binding: 18%/25%

Volume of distribution ( $V_d$ ): 0.26/0.3 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\***

CrCl < 30 mL/min	Avoid
Moderate—severe hepatic insufficiency	Use with caution

**Drug Interactions:** Allopurinol (↑ risk of rash).

**Adverse Effects:** Drug fever/rash, diarrhea, ↑ SGOT/SGPT. Rash potential same as ampicillin.

**Allergic Potential:** High

**Safety in Pregnancy:** B

**Comments:** 5 mL contains 600 mg amoxicillin and 42.9 mg clavulanatic acid. Take with meals to minimize GI upset. **Do not substitute 400 mg or 200 mg/5 mL formulation for ES-600.** Contains phenylalanine. May cause false + BG.

**Volume of ES-600 to provide 90 mg/kg/day:**

Weight	Volume (q12h)	Weight	Volume (q12h)
8 kg	3.0 mL	24 kg	9.0 mL
12 kg	4.5 mL	28 kg	10.5 mL
16 kg	6.0 mL	32 kg	12.0 mL
20 kg	7.5 mL	36 kg	13.5 mL

**Cerebrospinal Fluid Penetration:**

Non-inflamed meninges = 1%

Inflamed meninges = 1%

**Bile Penetration:** 3000%

(also see **Antibiotic Pearls & Pitfalls** p. 508).

**REFERENCES:**

- Chung KP, Huang YT, Lee LN, et al. Alarmingly decreasing rates of amoxicillin-clavulanate susceptibility among clinical isolates of Haemophilus influenzae from 2001 to 2009 in a medical center in Taiwan. J Infect 62:185–187, 2011.
- Easton J, Noble S, Perry CM. Amoxicillin/clavulanic acid: a review of its use in the management of paediatric patients with acute otitis media. Drugs 63:311–40, 2003.
- Ghaffar F, Muniz LS, Katz K, et al. Effects of large dosages of amoxicillin/clavulanate or azithromycin on nasopharyngeal carriage of Streptococcus pneumoniae, Haemophilus influenzae, nonpneumococcal alpha-hemolytic streptococci, and Staphylococcus aureus in children with acute otitis media. Clin Infect Dis 34:1301–9, 2002.
- Website: www.pdr.net

## Amoxicillin/Clavulanic Acid XR (Augmentin XR)

**Drug Class:** Aminopenicillin/β-lactamase inhibitor.

**Usual Dose:** 2000/125 mg (2 tablets) (PO) q12h (see comments).

**Spectrum:** (see **Susceptibility Profiles** pp. 186–190).

**Resistance Potential:** Low

**Pharmacokinetic Parameters:**

Peak serum level: 17/2 mcg/mL

Bioavailability: 90/60%

Excreted unchanged (urine): 70/40%

Serum half-life (normal/ESRD): [1.3/16]/[½] hrs

Plasma protein binding: 18/25%

Volume of distribution ( $V_d$ ): 0.26/0.3 L/kg

**Primary Mode of Elimination:** Renal

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

**Dosage Adjustments\***

CrCl > 30 mL/min	No change
CrCl < 30 mL/min	Avoid
Post-HD/PD dose	Avoid
CVVH/CVVHD/ CVHDF dose	Avoid
Moderate—severe hepatic insufficiency	Use with caution

**Drug Interactions:** Allopurinol (↑ risk of rash); may ↓ effectiveness of oral contraceptives.

**Adverse Effects:** Drug fever, rash, diarrhea, ↑ SGOT/SGPT, nausea, abdominal pain.

**Allergic Potential:** High

**Safety in Pregnancy:** B

**Comments:** Amoxicillin/clavulanic acid XR is a time-released formulation. Do not crush tablets. Take with food to increase absorption. XR formulation contains a different ratio of amoxicillin/clavulanic acid so other formulations cannot be interchanged. 2 tablets (1000/62.5 mg per tablet) 2000/125 mg per dose. May cause false + BG.

**Cerebrospinal Fluid Penetration:**

Non-Inflamed meninges = 1%

Inflamed meninges = 1%

**Bile Penetration:** 3000%

(also see **Antibiotic Pearls & Pitfalls** p. 508).

**REFERENCES:**

- Augmentin XR. Med Lett Drugs Ther. 45:5–6, 2003.  
 Benninger MS. Amoxicillin/clavulanate potassium extended release tablets: a new antimicrobial for the treatment of acute bacterial sinusitis and community-acquired pneumonia. Expert Opin Pharmacother. 4:1839–46, 2003.  
 File Jr TM, Jacobs M, Poole M, et al. Outcome of treatment of respiratory tract infections due to Streptococcus pneumoniae, including drug-resistant strains, with

pharmacokinetically enhanced amoxicillin/clavulanic acid. International J Antimicrobial Agents 20:235–247, 2002.

Kaye C, Allen A, Perry S, et al. The clinical pharmacokinetics of a new pharmacokinetically enhanced formulation of amoxicillin/clavulanic acid. Clin Therapeutics 23:578–584, 2001.

Website: www.pdr.net

**Amphotericin B (Fungizone)**

**Drug Class:** Antifungal.

**Usual Dose:** 0.5–0.8 mg/kg (IV) q24h.

**Pharmacokinetic Parameters:**

*Peak serum level:* 1–2 mcg/mL

*Bioavailability:* Not applicable

*Excreted unchanged (urine):* 5%

*Serum half-life (normal/ESRD):* 15/48 days

*Plasma protein binding:* 90%

*Volume of distribution (V<sub>d</sub>):* 4 L/kg

**Primary Mode of Elimination:** Metabolized

**Dosage Adjustments\***

CrCl 10–50 mL/min	No change
Post-HD or post-PD	None
Post-HFHD dose	No change
CVVH/CVVHD/CVHDF dose	No change
Moderate—severe hepatic insufficiency	No change

**Drug Interactions:** Adrenocorticoids (hypokalemia); aminoglycosides, cyclosporine, polymyxin B (↑ nephrotoxicity); digoxin (↑ digitalis toxicity due to hypokalemia); flucytosine (↑ flucytosine levels if amphotericin B produces renal dysfunction); neuromuscular blocking agents (↑ neuromuscular blockade due to hypokalemia).

**Adverse Effects:** Fevers/chills, flushing thrombophlebitis, bradycardia, seizures, hypotension, distal renal tubular acidosis (↓ K<sup>+</sup>/↓ Mg<sup>++</sup>), anemia, pancreatitis. If renal

\*Usual dose\* assumes normal renal/hepatic function. \*For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.



insufficiency is secondary to amphotericin, either ↓ daily dose by 50%, give dose every other day, or switch to an amphotericin lipid formulation.

**Allergic Potential:** Low

**Safety in Pregnancy:** B

**Comments:** Higher doses (1–1.5 mg/kg q24h) may be needed in life-threatening situations but are nephrotoxic and should only be administered under expert supervision. Do not co-administer in same IV with other drugs. Give by slow IV infusion over 2 hours initially. Aggressive hydration (1–2 liters/d) may reduce nephrotoxicity. **Test dose unnecessary.** Amphotericin B with granulocyte colony stimulating factor (G-CSF) may result in ARDS. Amphotericin B with pentamidine may cause acute tubular necrosis in HIV patients. **Fevers/chills may be reduced by meperidine, aspirin, NSAIDs, hydrocortisone or acetamino-phen, if given 30–60 minutes before infusion. Bladder irrigation dose:** 50 mg/L until cultures are negative.

**Highly active against *Candida***  
(↓ susceptibility to *C. glabrata*, *C. krusei*),  
***Cryptococcus*, *Histoplasmosis*, *Blastomycosis*,  
*Sporotrichosis*, *Penicillium marneffeii*,  
*Paracoccidiomycosis*, *Coccidiomycosis*.**  
**Some activity against *Fusarium*, *Naegleria*,  
*Leishmania*, *Malassezia* or *Mucor*.**  
**No activity against *C. lusitanae*,  
*Pseudoallescheria/Scedosporium* or  
*Trichosporon*.**

**Meningeal dose** = usual dose plus 0.5 mg 3–5x/week (IT) via Ommaya reservoir.

**Cerebrospinal Fluid Penetration:** < 10%

#### REFERENCES:

Arikan S, Lozano-Chiu M, Paetznick V, et al. In vitro synergy of caspofungin and amphotericin B against *Aspergillus* and *Fusarium* spp. *Antimicrob Agents Chemother* 46:245–7, 2002.

- Barchiesi F, Maracci M, Baldassarri I, et al. Tolerance to amphotericin B in clinical isolates of *Candida tropicalis*. *Diagn Microbiol Infect Dis* 50:179–85, 2004.
- Chocarro Martinez A, Gonzalez A, Garcia I. Caspofungin versus amphotericin B for the treatment of candidal esophagitis. *Clin Infect Dis* 35:107; discussion 107–8, 2002.
- Cruz JM, Peacock JE Jr, Loomer L, et al. Rapid intravenous infusion of amphotericin B: A pilot study. *Am J Med* 93:123–30, 1992.
- Deray G. Amphotericin B nephrotoxicity. *J Antimicrob Chemother* 49 Suppl 1:37–41, 2002.
- Ellis D. Amphotericin B: spectrum and resistance. *J Antimicrob Chemother* 49 Suppl 1:7–10, 2002.
- Gallis HA, Drew RH, Pickard WW. Amphotericin B: 30 years of clinical experience. *Rev Infect Dis* 12:308–29, 1990.
- Gubbins PO, Heldenbrand S. Clinically relevant drug interactions of current antifungal agents. *Mycoses* 53:95–113, 2010.
- Lewis, RE, Wiederhold NP, Klepser ME. In vitro pharmacodynamics amphotericin B, itraconazole, and voriconazole against *aspergillus*, *fusarium*, and *scedosporium* spp. *Antimicrob Agents and Chemotherapy* 49:945–951, 2005.
- Lyman CA, Walsh TJ. Systemically administered antifungal agents: A review of their clinical pharmacology and therapeutic applications. *Drugs* 44:9–35, 1992.
- Menzies D, Goel K, Cunha BA. Amphotericin B. *Antibiotics for Clinicians* 2:73–6, 1998.
- Miceli MH, Diaz JA, Lee SA. Emerging opportunistic yeast infections. *Lancet Infectious Diseases* 11:142–151, 2011.
- Moosa MY, Alangaden GJ, Manavathu E. Resistance to amphotericin B does not emerge during treatment for invasive aspergillosis. *J Antimicrob Chemother* 49:209–13, 2002.
- Nivoix Y, Zamfir A, Lutun P, Kara F, et al. Combination of caspofungin and an azole or an amphotericin B formulation in invasive fungal infections. *J Infect* 52:67–74, 2006.
- Pagano L, Valentini CG, Fianchi L, et al. Treatment strategies for invasive aspergillosis in neutropenic patients: voriconazole or liposomal amphotericin-B? *J Chemother* 23:5–8, 2011.

"Usual dose" assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

Pappas PG, Chetchotisakd P, Larsen RA, et al. A phase II randomized trial of amphotericin B alone or combined with fluconazole in the treatment of HIV-associated cryptococcal meningitis. *Clin Infect Dis* 48:1775–1784, 2009.

Perfect JR. "Amphoterrible": In the era of broad-spectrum azoles and candins. *Infections in Medicine*. 23:S2, 2006.

Perfect JR, Dismukes WE, Dromer F, et al. Clinical Practice guidelines for the management of cryptococcal disease: 2010 update by the Infectious Disease Society of America. *Clin Infect Dis* 50:291–322, 2010.

Safdar A, Ma J, Saliba F, et al. Drug-induced nephrotoxicity caused by amphotericin B lipid complex and liposomal amphotericin B: a review and meta-analysis. *Medicine (Baltimore)* 89:236–244, 2010.

Website: www.pdr.net

## Amphotericin B Lipid Complex (Abelcet) ABLC

**Drug Class:** Antifungal (see p. 525).

**Usual Dose:** 5 mg/kg (IV) q24h.

**Pharmacokinetic Parameters:**

*Peak serum level:* 1.7 mcg/mL

*Bioavailability:* Not applicable

*Excreted unchanged (urine):* 5%

*Serum half-life (normal/ESRD):* 173/173 hrs

*Plasma protein binding:* 90%

*Volume of distribution ( $V_d$ ):* 131 L/kg

**Primary Mode of Elimination:** Metabolized

**Dosage Adjustments\***

CrCl < 50 mL/min	No change
Post-HD/PD dose	None
Post-HFHD dose	None
CVWH/CVWHD/CVWHDF dose	No change
Moderate—severe hepatic insufficiency	No change

**Drug Interactions:** Adrenocorticoids (hypokalemia); aminoglycosides, cyclosporine, polymyxin B (↑ nephrotoxicity); digoxin (↑ digitalis toxicity due to hypokalemia); flucytosine (↑ flucytosine effect); neuromuscular blocking agents (↑ neuromuscular blockade due to hypokalemia).

**Adverse Effects:** Fevers/chills, flushing thrombophlebitis, bradycardia, seizures, hypotension, distal renal tubular acidosis (↓ K<sup>+</sup>/↓ Mg<sup>++</sup>), anemia. Fewer/less severe side effects and less nephrotoxicity than amphotericin B. Renal toxicity is dose-dependent (use caution).

**Allergic Potential:** Low

**Safety in Pregnancy:** B

**Comments:** See p. 525. Useful in patients unable to tolerate amphotericin B or in patients with amphotericin B nephrotoxicity. Infuse at 2.5 mg/kg/hr.

**Highly active against *Candida***

(↓ susceptibility to *C. glabrata*, *C. krusei*),

***Cryptococcus*, *Histoplasmosis*, *Blastomycosis*,**

***Sporotrichosis*, *Penicillum marneffe*,**

***Paracoccidiomycosis*, *Coccidiomycosis*.**

**Some activity against *Fusarium*, *Naegleria*,**

***Leishmania*, *Malassezia* or *Mucor*. No activity**

**against *C. lusitaniae*, *Pseudoallescheria*/**

***Scedosporium* or *Trichosporon*.**

**Cerebrospinal Fluid Penetration:** < 10%

### REFERENCES:

Arikan S, Rex JH. Lipid-based antifungal agents: current status. *Curr Pharm Des* 7:393–415, 2001.

Arnold TM, Dotson E, Sarosi GA, et al. Traditional and emerging antifungal therapies. *Proc Am Thorac Soc* 7:222–228, 2010.

Dupont B. Overview of the lipid formulations of amphotericin B. *J Antimicrob Chemother* 49 Suppl 1:31–6, 2002.

Hiemzen JW, Walsh TJ. Lipid formulations of amphotericin B: Recent progress and future directions. *Clin Infect Dis* 2:133–44, 1996.

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

Miceli MH, Diaz JA, Lee SA. Emerging opportunistic yeast infections. *Lancet Infectious Diseases* 11:142–151, 2011.

Mondal S, Bhattacharya P, Ali N. Current diagnosis and treatment of visceral leishmaniasis. *Expert Rev Anti Infect Ther* 8:919–944, 2010.

Website: www.pdr.net

## Amphotericin B Liposomal (AmBisome)

**Drug Class:** Antifungal (see p. 525).

**Usual Dose:** 3–6 mg/kg (IV) q24h (see comments).

**Pharmacokinetic Parameters:**

Peak serum level: 17–83 mcg/mL

Bioavailability: Not applicable

Excreted unchanged (urine): 5%

Serum half-life (normal/ESRD): 153 hrs/no data

Plasma protein binding: 90%

Volume of distribution ( $V_d$ ): 131 L/kg

**Primary Mode of Elimination:** Metabolized

**Dosage Adjustments\***

CrCl < 50 mL/min	No change
Post-HD/PD dose	None
Post-HFHD dose	None
CVWH/CVVHD/ CWHDF dose	No change
Moderate—severe hepatic insufficiency	No change

**Drug Interactions:** Adrenocorticoids (hypokalemia); aminoglycosides, cyclosporine, polymyxin B (↑ nephrotoxicity); digoxin (↑ digitalis toxicity due to hypokalemia); flucytosine (↑ flucytosine effect); neuromuscular blocking agents (↑ neuromuscular blockade due to hypokalemia).

**Adverse Effect:** Fevers/chills, flushing, thrombocytopenia, bradycardia, seizures, hypotension,

distal renal tubular acidosis (↓  $K^+$ / ↓  $Mg^{2+}$ ), anemia.

**Allergic Potential:** Low

**Safety in Pregnancy:** B

**Comments:** See p. 525. Less nephrotoxicity than amphotericin B and other amphotericin lipid preparations. **Empiric therapy of fungemia dose:** 3 mg/kg (IV) q24h can be used. **Suspected/known Aspergillus dose:** 5 mg/kg (IV) q24h. **Cryptococcal meningitis dose:** 6 mg/kg (IV) q24h. **Highly active against *Candida* (↓ susceptibility to *C. glabrata*, *C. krusei*), *Cryptococcus*, *Histoplasmosis*, *Blastomycosis*, *Sporotrichosis*, *Penicillium marneffeii*, *Paracoccidiomycosis*, *Coccidiomycosis*. **Some activity against *Fusarium*, *Naegleria*, *Leishmania*, *Malassezia* or *Mucor*. **No activity against *C. lusitanae*, *Pseudoallescheria/Scedosporium* or *Trichosporon*.******

**Cerebrospinal Fluid Penetration:** < 10%

### REFERENCES:

- Chopra R. AmBisome in the treatment of fungal infections: the UK experience. *J Antimicrob Chemother* 49 Suppl 1:43–7, 2002.
- Chu P, Sadullah S. The current role of amphotericin B lipid complex in managing systemic fungal infections. *Curr Med Res Opin* 25:3011–3020, 2009.
- Hiemenz JW, Walsh TJ. Lipid formulations of amphotericin B: Recent progress and future directions. *Clin Infect Dis* 2:133–44, 1996.
- Kuse ER, Chetchotisakd P, da Cunha CA, et al. Micafungin versus liposomal amphotericin B for candidaemia and invasive candidosis: a phase III randomised double-blind trial. *Lancet* 369:1519–1527, 2007.
- Lequaglie C. Liposomal amphotericin B (AmBisome): efficacy and safety of low-dose therapy in pulmonary fungal infections. *J Antimicrob Chemother* 49 Suppl 1:49–50, 2002.
- Miceli MH, Diaz JA, Lee SA. Emerging opportunistic yeast infections. *Lancet Infectious Diseases* 11:142–151, 2011.

\*Usual dose\* assumes normal renal/hepatic function. \*For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

Safdar A, Ma J, Saliba F, et al. Drug-induced nephrotoxicity caused by amphotericin B lipid complex and liposomal amphotericin B: a review and meta-analysis. *Medicine* (Baltimore) 89:236–244, 2010.

Website: www.pdr.net

## Amphotericin B Cholesteryl Sulfate Complex (Amphotec), ABCD (amphotericin B colloidal dispersion)

**Drug Class:** Antifungal (see p. 525).

**Usual Dose:** 3–4 mg/kg (IV) q24h.

**Pharmacokinetic Parameters:**

*Peak serum level:* 2.9 mcg/mL

*Bioavailability:* Not applicable

*Excreted unchanged (urine):* 5%

*Serum half-life (normal/ESRD):* 39/29 hrs

*Plasma protein binding:* 90%

*Volume of distribution ( $V_d$ ):* 4 L/kg

**Primary Mode of Elimination:** Metabolized

**Dosage Adjustments\***

CrCl < 50 mL/min	No change
Post-HD/PD dose	None
Post-HFHD dose	None
CVWH/CVVDH/CVWHDF dose	None
Moderate—severe hepatic insufficiency	No change

**Drug Interactions:** Adrenocorticoids (hypokalemia); aminoglycosides, cyclosporine, polymyxin B (↑ nephrotoxicity); digoxin (↑ digitalis toxicity due to hypokalemia); flucytosine (↑ flucytosine effect); neuromuscular blocking agents (↑ neuromuscular blockade due to hypokalemia).

**Adverse Effects:** Fevers/chills, flushing, thrombophlebitis, bradycardia, seizures, hypotension, distal renal tubular acidosis

(↓  $K^+$ /↓  $Mg^{++}$ ), anemia. Fewer/less severe side effects/less nephrotoxicity vs. amphotericin B.

**Allergic Potential:** Low

**Safety in Pregnancy:** B

**Comments:** See p. 525. Reconstitute in sterile water, not dextrose, saline or bacteriostatic water.

Do not co-administer with in same IV line with other drugs. Give by slow IV infusion over 2 hours (1 mg/kg/hr). Test dose unnecessary.

**Highly active against *Candida***

(↓ susceptibility to *C. glabrata* *C. krusei*),

*Cryptococcus*, *Histoplasmosis*, *Blastomycosis*,

*Sporotrichosis*, *Penicillium marneffeii*,

*Paracoccidiomycosis*, *Coccidiomycosis*.

**Some activity against *Fusarium*, *Naegleria*,**

*Leishmania*, *Malassezia* or *Mucor*.

**No activity against *C. lusitanae*,**

*Pseudoallescheria/Scedosporium* or

*Trichosporon*.

**Cerebrospinal Fluid Penetration:** < 10%

### REFERENCES:

De Marie S. Clinical use of liposomal and lipid-complexed amphotericin B. *J Antimicrob Chemother* 33:907–16, 1994.

Rapp RP, Gubbins PO, Evans ME. Amphotericin B lipid complex. *Ann Pharmacother* 31:1174–86, 1997.

Safdar A, Ma J, Saliba F, et al. Drug-induced nephrotoxicity caused by amphotericin B lipid complex and liposomal amphotericin B: a review and meta-analysis. *Medicine* (Baltimore) 89:236–244, 2010.

Website: www.pdr.net

## Ampicillin (various)

**Drug Class:** Aminopenicillin.

**Usual Dose:** 2 gm (IV) q4h, 500 mg (PO) q6h.

**Spectrum:** (see **Susceptibility Profiles** pp. 186–190.)

**Resistance Potential:** High (MSSA, aerobic GNBs)

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

**Pharmacokinetic Parameters:**

Peak serum level: 48 (IV)/5 (PO) mcg/mL

Bioavailability: 50%

Excreted unchanged (urine): 90%

Serum half-life (normal/ESRD): 0.8/10 hrs

Plasma protein binding: 20%

Volume of distribution ( $V_d$ ): 0.25 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\***

CrCl 50–80 mL/min	1 gm (IV) q4h 500 mg (PO) q6h
CrCl 10–50 mL/min	1 gm (IV) q8h 250 mg (PO) q8h
CrCl < 10 mL/min	1 gm (IV) q12h 250 mg (PO) q12h
Post-HD dose	1 gm (IV) 500 mg (PO)
Post-PD dose	1 gm (IV) 250 mg (PO)
CVWH/CVWHD/ CVWHD dose	1 gm (IV) q8h 250 mg (PO) q12h
Moderate—severe hepatic insufficiency	No change

**Drug Interactions:** Allopurinol (↑ frequency of rash); warfarin (↑ INR).

**Adverse Effects:** Drug fever/rash, nausea, GI upset, irritative diarrhea, ↑ SGOT/SGPT, ↑ incidence of rash vs. penicillin (in patients with EBV, HIV, lymphocytic leukemias, or allopurinol), C. difficile diarrhea/colitis.

**Allergic Potential:** High

**Safety in Pregnancy:** B

**Comments:** Incompatible in solutions containing amphotericin B, heparin, corticosteroids, erythromycin, aminoglycosides, or metronidazole.

Na<sup>+</sup> content = 2.9 mEq/g.

Meningeal dose = 2 gm (IV) q4h.

**Cerebrospinal Fluid Penetration:**

Non-Inflamed meninges = 1%

Inflamed meninges = 10%

**Bile Penetration:** 3000%

(also see **Antibiotic Pearls & Pitfalls** p. 508).

**REFERENCES:**

Sakoulas G, Bayer AS, Pogliano J, et al. Ampicillin enhances daptomycin- and cationic host defense peptide-mediated killing of ampicillin- and vancomycin-resistant *Enterococcus faecium*. *Antimicrob Agents Chemother.* 56:838–844, 2012.

Wright AJ. The penicillins. *Mayo Clin Proc* 74:290–307, 1999.

Website: www.pdr.net

## Ampicillin/Sulbactam (Unasyn)

**Drug Class:** Aminopenicillin/β-lactamase inhibitor.

**Usual Dose:** 1.5–3 gm (IV) q6h (see comments).

**Spectrum:** (see **Susceptibility Profiles** pp. 186–190).

**Resistance Potential:** Low (pseudoresistance with *E. coli*/Klebsiella)

**Pharmacokinetic Parameters:**

Peak serum level: 109–150/48–88 mcg/mL

Bioavailability: Not applicable

Excreted unchanged (urine): 80/80%

Serum half-life (normal/ESRD): [1/9]/[1/9] hrs

Plasma protein binding: 28/38%

Volume of distribution ( $V_d$ ): 0.25/0.38 L/kg

**Primary Mode of Elimination:** Renal/hepatic

**Dosage Adjustments\*** (based on 3 gm (IV) q6h)

CrCl 30–80 mL/min	1.5 gm (IV) q6h
CrCl 15–30 mL/min	1.5 gm (IV) q12h
CrCl < 15 mL/min	1.5 gm (IV) q24h
Post-HD dose	1.5 gm (IV)

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

Post-PD dose	None
CVWH/CVVD/ CVHDF dose	1.5–3 gm (IV) q12h
Moderate—severe hepatic insufficiency	No change

**Drug Interactions:** Probenecid (↑ ampicillin/sulbactam levels); allopurinol (↑ rash).

**Adverse Effects:** Drug fever/rash, ↑ SGOT/SGPT, hemolytic anemia, *C. difficile* diarrhea/colitis.

**Allergic Potential:** High

**Safety in Pregnancy:** B

**Comments: Mild/moderate infection dose:** 1.5 gm (IV) q6h.

**Meningeal dose:** 4.5 gm (IV) q6h (for susceptible strains of MDR *Acinetobacter baumannii* meningitis).

**Cerebrospinal Fluid Penetration:** 30%

**Bile Penetration:** 900%  
(also see **Antibiotic Pearls & Pitfalls** p. 508).

#### REFERENCES:

- Itokazu GS, Danziger LH. Ampicillin-sulbactam and ticarcillin-clavulanic acid: A comparison of their in vitro activity and review of their clinical efficacy. *Pharmacotherapy* 11:382–414, 1991.
- Jain R, Danziger LH. Multidrug-resistant *Acinetobacter* infections: an emerging challenge to clinicians. *Ann Pharmacother* 38:1449–59, 2004.
- Sensakovic JW, Smith LG. Beta-lactamase inhibitor combinations. *Med Clin North Am* 79:695–704, 1995.
- Swenson JM, Killgore GE, Tenover FC. Antimicrobial susceptibility testing of *Acinetobacter* spp. by NCCLS broth microdilution and disk diffusion methods. *J Clin Microbiol* 42:5102–8, 2004.
- Wood GC, Hanes SD, Croce MA, et al. Comparison of ampicillin-sulbactam and imipenem-cilastatin for the treatment of *Acinetobacter* ventilator-associated pneumonia. *Clin Infect Dis* 34:1425–30, 2002.
- Wright AJ. The penicillins. *Mayo Clin Proc* 73:290–307, 1999.
- Website: www.pdr.net

## Amprenavir (Agenerase) APV

**Drug Class:** HIV protease inhibitor.

**Usual Adult Dose:** 1200 mg (PO) q12h (capsules) or 1400 mg (PO) q12h (solution); age 13–16 years < 50 kg or pediatrics 4–12 years: 20 mg/kg (PO) q12h (capsules) or 1.5 ml/kg (PO) q12h (15 mg/mL solution). Maximum dose 2400 mg/d (capsules), 2800 mg/d (oral solution).

#### Pharmacokinetic Parameters:

*Peak serum level:* 7.6 mcg/mL

*Bioavailability:* No data

*Excreted unchanged (urine):* 1%

*Serum half-life (normal/ESRD):* 7-10/7-10 hrs

*Plasma protein binding:* 90%

*Volume of distribution (V<sub>d</sub>):* 6.1 L/kg

**Primary Mode of Elimination:** Hepatic

#### Dosage Adjustments\*

CrCl > 10 mL/min	No change
CrCl < 10 mL/min	No change (avoid solution)
Post-HD/PD dose	No change
CVWH/CVVD/ CVHDF dose	No change
Moderate hepatic insufficiency	450 mg (PO) q12h (capsules)
Severe hepatic insufficiency	300 mg (PO) q12h (capsules)

#### Antiretroviral Dosage Adjustments

Delavirdine	Avoid
Didanosine buffered solution	Take amprenavir 1 hour before or after
Efavirenz	No data

\*Usual dose\* assumes normal renal/hepatic function. \*For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

Indinavir	No data
Lopinavir/ritonavir	May ↓ lopinavir levels
Nelfinavir	No data
Nevirapine	No data
Ritonavir	Limited data for amprenavir 600 mg q12h + ritonavir 100 mg q12h (or amprenavir 1200 mg q24h + ritonavir 200 mg q24h)
Saquinavir	No data
Rifampin	Avoid
Rifabutin	Rifabutin 150 mg q24h or 300 mg 2–3x/week

**Drug Interactions:** Antiretrovirals, rifabutin, rifampin (see dose adjustment grid, above); bepridil, cisapride, ergotamine, statins, benzodiazepines, St. John's wort, pimozide, sildenafil, methadone (avoid if possible); carbamazepine, phenobarbital, phenytoin (may ↓ amprenavir levels, monitor anticonvulsant levels); H<sub>2</sub> blockers, proton pump inhibitors (↓ amprenavir levels).

**Adverse Effects:** Rash, Stevens-Johnson Syndrome (rare), GI upset, headache, depression, taste perversion, diarrhea, perioral paresthesias, hyperglycemia (including worsening diabetes, new-onset diabetes, DKA), ↑ cholesterol/triglycerides (evaluate risk for coronary disease/pancreatitis), fat redistribution, ↑ SGOT/SGPT, possible increased bleeding in hemophilia.

**Allergic Potential:** High. Amprenavir is a sulfonamide; use with caution in sulfa allergy

**Safety in Pregnancy:** C

**Comments:** Can be taken with or without food, but avoid high fat meals (may ↓ absorption). High vitamin E content. Capsules and solution are not interchangeable on a mg per mg basis. Decrease dosage in moderate or severe liver disease; use with caution. Oral solution contains propylene glycol: avoid in pregnancy, hepatic/renal failure, patients taking disulfiram or metronidazole, or children < 4 years old. Effective antiretroviral therapy consists of three antiretrovirals (same/different classes)

GlaxoSmithKline discontinued production and sale of amprenavir in the United States in order to focus on production of the amprenavir prodrug, fosamprenavir. This action was not taken because of concerns about safety or efficacy of amprenavir. Amprenavir may continue to be available in some countries outside the United States.

#### REFERENCES:

- Adkins JC, Faulds D. Amprenavir. *Drugs* 55:837–42, 1998.
- Croteau D, Letendre S, Best BM, et al. Therapeutic amprenavir concentrations in cerebrospinal fluid. *Antimicrob Agents Chemother* 56:1985–9, 2012.
- Kappelhoff BS, Crommentuyn KM, de Maat MM, et al. Practical guidelines to interpret plasma concentrations of antiretroviral drugs. *Clin Pharmacokinet* 43:845–53, 2004.
- Panel on Antiretroviral Guidelines for Adults and Adolescents. Department of Health and Human Services. Guidelines for the use of antiretroviral agents in HIV-infected adults and adolescents.
- Thompson MA, Aberg JA, Cahn P, et al. Antiretroviral treatment of adult HIV infection: 2010. recommendations of the international AIDS society – USA panel. *JAMA* 304:321–333, 2010.
- Website: [www.pdr.net](http://www.pdr.net)

<sup>†</sup>Usual dose assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

## Anidulafungin (Eraxis, Ecalta)

**Drug Class:** Echinocandin antifungal.

**Usual Dose:** 200 mg (IV) × 1 dose, then 100 mg (IV) q24h.

**Pharmacokinetic Parameters:**

Peak serum level: 4.2 mg/L (100/50 mg dose)/7.2 mg/L (200/100 mg dose)

Bioavailability: Not applicable

Excreted unchanged (urine): < 1%

Serum half-life (normal/ESRD): 40–50 hours/  
40–50 hours

Plasma protein binding: 99%

Volume of distribution ( $V_d$ ): 30–50L

**Primary Mode of Elimination:** Fecal (30%)

**Dosage Adjustments\***

CrCl < 50 mL/min	No change
Post-HD dose	None
Post-PD dose	None
CWH/CVHD/ CWHDF dose	No change
Moderate—severe hepatic insufficiency	No change

**Drug Interactions:** No significant interactions.

**Adverse Effects:** ↑ SGOT/SGPT/GGTP/alkaline phosphatase. Histamine-related reactions (rash, urticaria, pruritus, fever, dyspnea and hypotension) if infusion > 1.1 mg/min.

**Allergic Potential:** Histamine-related symptoms

**Safety in Pregnancy:** C

**Comments:** Not an inducer, inhibitor, or substrate of CYP450 system. Rate of infusion should not exceed 1.1 mg/min. Incompatible with other drugs. Do not infuse with other medications. Use reconstituted vials within 24 hours. Infusion solution should be stored at 25°C/77°F.

**Highly active against *Candida albicans*, non-*albicans Candida* (↓ susceptibility to *C. parasilosis*).**

**Some activity against *Aspergillus*.**

**No activity against *Cryptococcus*, *Fusarium*, *Pseudoallescheria/Scedosporium*, *Trichosporon*, *Rhodotorula* or *Mucor*.**

**Cerebrospinal Fluid Penetration:** No data

### REFERENCES:

- Bennett JE, Echinocandins for candidemia in adults without neutropenia. *N Engl J Med* 355:1154–9, 2006.
- Cohen-Wolkowicz M, Benjamin Jr. DK, Steinbach WJ, Smith PB. Anidulafungin: A new Echinocandin for the Treatment of Fungal Infections. *Drugs of Today*. 42:533–44, 2006.
- de la Torre P, Reboli AC. Anidulafungin: a new echinocandin for candidal infections. *Expert Rev Anti Infect Ther* 5:45–52, 2007.
- Dowell JA, Schranz J, Baruch A, Foster G. Safety and pharmacokinetics of coadministered voriconazole and anidulafungin. *J Clin Pharmacol* 45:1373–82, 2005.
- Dowell JA, Stogniew M, Krause D, et al. Anidulafungin does not require dosage in subjects with varying degrees of hepatic or renal impairment. *J Clin Pharmacol* 47:461–470, 2007.
- Dowell JA, Stogniew M, Krause D, et al. Lack of pharmacokinetic interaction between anidulafungin and tacrolimus. *J Clin Pharmacol* 47:305–314, 2007.
- Garber G, Laverdiere M, Libman M, et al. Anidulafungin versus fluconazole for invasive candidiasis. *N Engl J Med* 356:2472–2482, 2007.
- Kofia G, Ruhnke M. Pharmacology and metabolism of anidulafungin, caspofungin, and micafungin in the treatment of invasive candidosis: review of the literature. *Eur J Med Res* 16:159–166, 2011.
- Miceli MH, Diaz JA, Lee SA. Emerging opportunistic yeast infections. *Lancet Infectious Diseases* 11:142–151, 2011.
- Morrison VA. Echinocandin antifungals: review and update. *Expert Rev Anti Infect Ther* 4:325–42, 2006.
- Nir-Paz R, Moses AE. Anidulafungin and fluconazole for candidiasis. *N Engl J Med* 357:1347–1348, 2007.

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.



- Paderu P, Garcia-Effron G, Balashov S, et al. Serum differentially alters the antifungal properties of echinocandin drugs. *Antimicrob Agents Chemother* 51:2253–2256, 2007.
- Patel PN, Charneski L. Anidulafungin (Eraxis) for Fungal Infections. *Pharmacy and Therapeutics* 31:644–53, 2006.
- Pfaller MA, Diekema DJ, Boyken L. Effectiveness of anidulafungin in eradicating candida species in invasive candidiasis. *Antimicrob Agents Chemother* 49:4795–97, 2005.
- Raasch RH. Anidulafungin: review of a new echinocandin antifungal agent. *Expert Rev Anti Infect Ther* 2:499–508, 2004.
- Reboli AC, Rotstein C, Pappas PG, et al. Anidulafungin versus fluconazole for invasive candidiasis. *N Engl J Med* 356:2472–2482, 2007.
- Theuretzbacher U. Pharmacokinetics/pharmacodynamics of echinocandins. *Eur J Clin Microbiol Infect Dis* 23:805–12, 2004.
- Turner MS, Drew RH, Perfect JR. Emerging echinocandins for treatment of invasive fungal infections. *Expert Opin Emerg Drugs* 11:231–50, 2006.
- Vazquez JA, Sobel JD. Anidulafungin: a novel echinocandin. *Clin Infect Dis* 43:215–22, 2006.
- Wiederhold NP, Lewis RE. The echinocandin antifungals: an overview of the pharmacology, spectrum and clinical efficacy. *Expert Opin Investig Drugs* 12:1313–33, 2003.
- Wilke M. Treatment and prophylaxis of invasive candidiasis with anidulafungin, caspofungin, and micafungin and its impact on use and costs: review of the literature. *Eur J Med Res* 16:180–186, 2011.
- Website: [www.pdr.net](http://www.pdr.net)

## Atazanavir (Reyataz) ATV

**Drug Class:** HIV protease inhibitor.

**Usual Dose:** 300 mg (PO) q24h with ritonavir 100 mg (PO) q24h or 400 mg (PO) q24h with food if unable to tolerate ritonavir.

**Treatment Experienced Patients:** 300 mg with ritonavir 100 mg once daily with food.

### Pharmacokinetic Parameters:

Peak serum level: 3152 ng/mL; with ritonavir: 5233 ng/mL

Bioavailability: No data

Excreted unchanged (urine) (urine/feces): 7%/20%

Serum half-life (normal/ESRD): 6.5 hrs; with ritonavir: 8.6 hrs/no data

Plasma protein binding: 86%

Volume of distribution ( $V_d$ ): No data

**Primary Mode of Elimination:** Hepatic

### Dosage Adjustments\*

CrCl < 50 mL/min	No change
Post-HD or PD dose	None
CVWH/CVWHD/ CVWHDf dose	No change
Moderate hepatic insufficiency	300 mg (PO) q24h without ritonavir
Severe hepatic insufficiency	Avoid

### Antiretroviral Dosage Adjustments

Delavirdine	No data
Didanosine	Give atazanavir 2 hrs before didanosine with food
Efavirenz	Treatment naive: atazanavir 400 mg q24h + ritonavir 100 mg q24h + efavirenz 600 mg q24h Treatment experienced: not recommended
Indinavir	Avoid combination
Nevirapine	No data
Ritonavir	Atazanavir 300 mg/d + ritonavir 100 mg/d as single daily dose with food

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose  $\times$  1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

Saquinavir	↑ saquinavir (soft-gel) levels; no information
Rifampin	Avoid
Rifabutin	150 mg q48h or 3x/week

**Drug Interactions:** Antacids or buffered medications (↓ atazanavir levels; give atazanavir 2 hours before or 1 hour after); H<sub>2</sub>-receptor blockers (↓ atazanavir levels). In treatment-naïve patients taking an H<sub>2</sub>-receptor antagonist, give either atazanavir 400 mg once daily with food at least 2 hours before and at least 10 hours after the H<sub>2</sub>-receptor antagonist, or give atazanavir 300 mg once daily with ritonavir 100 mg once daily with food, without the need for separation from the H<sub>2</sub>-receptor antagonist. In treatment-experienced patients, give atazanavir 300 mg once daily with ritonavir 100 mg once daily with food at least 2 hours before and at least 10 hours after the H<sub>2</sub>-receptor antagonist); antiarrhythmics (↑ amiodarone, systemic lidocaine, quinidine levels; prolongs PR interval; monitor antiarrhythmic levels); antidepressants (↑ tricyclic antidepressant levels; monitor levels); calcium channel blockers (↑ calcium channel blocker levels, ↑ PR interval; ↓ diltiazem dose by 50%; use with caution; consider ECG monitoring); clarithromycin (↑ clarithromycin and atazanavir levels; consider 50% dose reduction; consider alternate agent for infections not caused by MAI); cyclosporine, sirolimus, tacrolimus (↑ immunosuppressant levels; monitor levels); ethinyl estradiol, norethindrone (↑ oral contraceptive levels; use lowest effective oral contraceptive dose); lovastatin, simvastatin (↑ risk of myopathy, rhabdomyolysis; avoid combination); sildenafil (↑ sildenafil levels; do not give more than 25 mg q48h); tadalafil (max. 10 mg/72 hours); vardenafil (max. 2.5 mg/72 hours);

St. John's wort (avoid combination); warfarin (↑ warfarin levels; monitor INR); tenofovir (tenofovir reduces systemic exposure to atazanavir).

**Contraindications: atazanavir include cisapride, pimozide, rifampin, irinotecan, midazolam, triazolam, lovastatin, simvastatin, bepridil, some ergot derivatives, indinavir, PPIs, St. John's wort, rifampentine, boceprevir.**

**Adverse Effects:** Abdominal pain, nausea, vomiting, headache, diarrhea, asthenia, anorexia, and dizziness.

**Allergic Potential:** Low

**Safety in Pregnancy:** B

**Comments:** Monitor LFTs in patients with HBV, HCV. Take 400 mg (two 200-mg capsules) once daily with food. Bioavailability is enhanced with food. Monitor LFTs in patients with HBV, HCV. In moderate liver disease reduce dose to 200 mg once daily. It is not recommended in patients with severe liver impairment.

#### REFERENCES:

- Cossarini F, Salpietro S, Galli L, et al. Monotherapy with atazanavir as a simplification strategy: results from an observational study. *J Acquir Immune Defic Syndr* 60:101–3, 2012.
- Foca E, Ripamonti D, Motta D, et al. Unboosted atazanavir for treatment of HIV infection: rationale and recommendations for use. *Drugs* 72:1161–73, 2012.
- Gandhi M, Ameli N, Bacchetti P, et al. Atazanavir concentration in hair in the strongest predictor of outcomes on antiretroviral therapy. *Clin Infect Dis* 52:1267–1275, 2011.
- Havlir DV, O'Marro SD. Atazanavir: new option for treatment of HIV infection. *Clin Infect Dis* 38:1599–604, 2004.
- Hughes PJ, Cretton-Scott E, Teague A, et al. Protease inhibitors for patients with HIV-1 infection. *Pharm Ther* 36:332–344, 2011.

<sup>†</sup>Usual dose<sup>†</sup> assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

Panel on Antiretroviral Guidelines for Adults and Adolescents. Department of Health and Human Services. Guidelines for the use of antiretroviral agents in HIV-infected adults and adolescents. <http://aidsinfo.nih.gov/ConsentFiles/AdultandAdolescentGL.pdf>, 2012.

Pillero PJ. Atazanavir: a novel HIV-1 protease inhibitor. *Expert Opin Investig Drugs* 11:1295–301, 2002.

Thompson MA, Aberg JA, Cahn P, et al. Antiretroviral treatment of adult HIV infection: 2012 recommendations of the International Antiviral society — USA panel. *JAMA* 308:387–402, 2012.

Website: [www.pdr.net](http://www.pdr.net)

## Atovaquone (Mepron)

**Drug Class:** Antiprotozoal.

**Usual Dose:** PCP treatment dose: 750 mg (PO) q12h. PCP prophylaxis dose: 1500 mg (PO) q24h

**Pharmacokinetic Parameters:**

*Peak serum level:* 12–24 mcg/mL

*Bioavailability:* 30% (47% with food; food ↑ bioavailability by 2-fold)

*Excreted unchanged (feces):* 94%

*Serum half-life (normal/ESRD):* 2.9/2.9 days

*Plasma protein binding:* 99.9%

*Volume of distribution ( $V_d$ ):* 0.6 L/kg

**Primary Mode of Elimination:** Hepatic

**Dosage Adjustments\***

CrCl < 80 mL/min	No change
Post-HD/PD dose	None
CWH/CVVD/ CWHDF dose	No change
Mild-moderate hepatic insufficiency	No change
Severe hepatic insufficiency	No change

**Drug Interactions:** Rifabutin, rifampin (↓ atovaquone effect); zidovudine (↑ zidovudine

levels). Rifampin decreases atovaquone levels by 50%.

**Adverse Effects:** Rash, nausea, vomiting, diarrhea, headache, drowsiness, dizziness, fever, anemia, leukopenia, ↑ SGOT/SGPT, ↑ amylase.

**Allergic Potential:** Low

**Safety in Pregnancy:** C

**Comments:** Active against *T. gondii*, PCP, Plasmodia, and Babesia.

**Cerebrospinal Fluid Penetration:** < 1%

### REFERENCES:

- Artymowicz RJ, James VE. Atovaquone: A new anti-pneumocystis agent. *Clin Pharmacol* 12:563–70, 1993.
- Baggish AL, Hill DR. Antiparasitic agent atovaquone. *Antimicrob Agents Chemother* 46:1163–73, 2002.
- Bonoan JT, Johnson DH, Schoch PE, Cunha BA. Life threatening babesiosis treated by exchange transfusion with azithromycin and atovaquone. *Heart & Lung* 27:42–8, 1998.
- Haile LG, Flaherty JF. Atovaquone: a review. *Ann Pharmacother* 27:1488–94, 1993.
- Tommasi C, Bellagamba R, Tempestilli M, et al. Marked increase in etravirine and saquinavir plasma concentrations furring atovaquone/proguanil prophylaxis. *Malar J* 21:141, 2011.
- Wormser GP, Prasad A, Neuhaus E, et al. Emergence of resistance to azithromycin-atovaquone in immunocompromised patients with babesia microti infection. *Clin Infect Dis* 50:381–386, 2010.
- Website: [www.pdr.net](http://www.pdr.net)

## Atovaquone + Proguanil (Malarone)

**Drug Class:** Antimalarial.

**Usual Dose:** Malaria prophylaxis: 1 tablet (250 mg/100 mg) (PO) q24h for 2 days before entering endemic area, daily during exposure, and daily

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose x 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

× 1 week post-exposure. **Malaria treatment:** 4 tablets (1000 mg/400 mg) (PO) as single dose × 3 days.

**Pharmacokinetic Parameters:**

Peak serum level: 38 mcg/mL

Bioavailability: 23/90%

Excreted unchanged (feces): 94/50%

Serum half-life (normal/ESRD):

[60/60]/[21/no data] hrs

Plasma protein binding: 99/75%

Volume of distribution ( $V_d$ ): 3.5/42 L/kg

**Primary Mode of Elimination:** Metabolized

**Dosage Adjustments\***

CrCl > 30 mL/min	No change
CrCl 10–30 mL/min	Avoid (for prophylaxis)
CrCl < 10 mL/min	Avoid (for prophylaxis)
Post-HD/PD dose	Avoid
CVWH/CVVD/ CVHDF dose	Avoid
Mild-moderate hepatic insufficiency	No change
Severe hepatic insufficiency	No change

**Drug Interactions:** Chloroquine (↑ incidence of mouth ulcers); metoclopramide, rifabutin, rifampin, tetracycline (↓ atovaquone + proguanil effect); ritonavir (↑ or ↓ atovaquone + proguanil effect); typhoid vaccine (↓ typhoid vaccine effect).

**Adverse Effects:** Rash, headache, drowsiness, dizziness, nausea, vomiting, diarrhea, fever, anemia, leukopenia, ↑SGOT/↑SGPT, ↑ amylase,

**Allergic Potential:** Low

**Safety in Pregnancy:** C

**Comments:** Malarone tablet = 250 mg atovaquone + 100 mg proguanil. **Effective against chloroquine sensitive/resistant strains of *P. falciparum*, but not *P. ovale*, *P. vivax*, or *P. malariae*.** Dosage may be decreased in patients with diarrhea. Take with food/milk.

**Cerebrospinal Fluid Penetration:** < 1%

**REFERENCES:**

- [No authors listed.] Atovaquone/proguanil (Malarone) for malaria. *Med Lett Drugs Ther* 42:109–11, 2000.
- Camus D, Djossou F, Schithuis HJ, et al. Atovaquone-proguanil versus chloroquine-proguanil for malaria prophylaxis in nonimmune pediatric travelers: results of an international, randomized, open-label study. *Clin Infect Dis* 38:1716–23, 2004.
- Looaeesuwan S, Chulay JD, Canfield CJ. Malarone (atovaquone and proguanil hydrochloride): a review of its clinical development for treating malaria. Malarone clinical trials study group. *Am J Trop Med Hyg* 60:533–41, 1999.
- Marra F, Salzman JR, Ensom MH. Atovaquone-proguanil for prophylaxis and treatment of malaria. *Ann Pharmacother* 37:1266–75, 2003.
- McKeage K, Scott L. Atovaquone-proguanil: a review of its use for the prophylaxis of *Plasmodium falciparum* malaria. *Drugs* 63:597–623, 2003.
- Thapar MM, Ashton M, Lindegardh N, et al. Time-dependent pharmacokinetics and drug metabolism of atovaquone plus proguanil (Malarone) when taken as chemoprophylaxis. *Eur J Clin Pharmacol* 15:19–27, 2002.
- Website: www.pdr.net

## Azithromycin (Zithromax)

**Drug Class:** Macrolide (Azolide).

**Usual Dose:** 500 mg (IV/PO) × 1 dose, then 250 mg (IV/PO) q24h (see comments).

**Spectrum:** (see **Susceptibility Profiles** pp. 186–190).

**Resistance Potential:** High (*S. pneumoniae*, GAS, MSSA)

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

**Pharmacokinetic Parameters:**

Peak serum level: 1.1 (IV)/0.2 (PO) mcg/mL

Bioavailability: 35%

Excreted unchanged (urine): 6%

Serum half-life (normal/ESRD): 68/68 hrs

Plasma protein binding: 50%

Volume of distribution ( $V_d$ ): 31 L/kg

**Primary Mode of Elimination:** Hepatic

**Dosage Adjustments\***

CrCl 10–50 mL/min	No change
CrCl < 10 mL/min	Use with caution
Post-HD/PD dose	None
CWH/CVWHD/ CWHDF dose	No change
Moderate—severe hepatic insufficiency	No change

**Drug Interactions:** Amiodarone, carbamazepine, cisapride, clozapine, corticosteroids, midazolam, triazolam, valproic acid (not studied/not reported); cyclosporine (↑ cyclosporine levels with toxicity); digoxin (↑ digoxin levels); pimozide (may ↑ QT interval, torsade de pointes).

**Adverse Effects:** ↑ incidence cardiac/non-cardiac sudden deaths after 5 days of therapy. Nausea, GI upset, non-C. difficile diarrhea. May ↑ QT<sub>c</sub> interval.

**Allergic Potential:** Low

**Safety in Pregnancy:** B

**Comments:** Bioavailability is decreased by food. C. trachomatis urethritis dose: 1 gm (PO) × 1 dose.

**N. gonorrhoea urethritis dose:** 2 gm (PO) × 1 dose.

**MAI prophylaxis dose:** 1200 mg (PO) weekly.

**MAI therapy dose:** 600 mg (PO) q24h.

**Cerebrospinal Fluid Penetration:** < 10%

**Bile Penetration:** > 3000%

(also see **Antibiotic Pearls & Pitfalls** p. 519)

**REFERENCES:**

- Alvarez-Elcoro S,ENZLER MJ. The macrolides: Erythromycin, clarithromycin and azithromycin. *Mayo Clin Proc* 74:613–34, 1999.
- Bergman M, Huikko S, Huovinen P, et al. Macrolide and azithromycin use are linked to increased macrolide resistance in *Streptococcus pneumoniae*. *Antimicrob Agents Chemother* 50:3646–3650, 2006.
- Centers for Disease Control. *Neisseria gonorrhoeae* with reduced susceptibility to azithromycin. *Morb Mortal Wkly Rep* 60:579–581, 2011.
- Cochereau I, Goldschmidt P, Goepogui A, et al. Efficacy and safety of short duration azithromycin eye drops versus azithromycin single oral dose for the treatment of trachoma in children: a randomized, controlled, double-masked clinical trial. *Br J Ophthalmol* 91:667–672, 2007.
- Cunha BA. Macrolides, doxycycline, and fluoroquinolones in the treatment of Legionnaires' Disease. *Antibiotics for Clinicians* 2:117–8, 1998.
- Daneman N, McGeer A, Green K, et al. Macrolide resistance in bacteremic pneumococcal disease: Implications for patient management. *Clin Infect Dis* 43:432–438, 2006.
- DuPont HL. Azithromycin for the self-treatment of traveler's diarrhea. *Clin Infect Dis* 44:347–349, 2007.
- Fisher AJ. Azithromycin and bronchiolitis obliterans syndrome after lung transplantation: is prevention is better than cure? *Eur Respir J* 37:10–12, 2011.
- Frénch RW, Mansour A, Nakhla I, et al. Short-course azithromycin for the treatment of uncomplicated typhoid fever in children and adolescents. *Clin Infect Dis* 38:951–7, 2004.
- Griffin AT, Peyrani P, Wiemken T, et al. Macrolides versus quinolones in Legionella pneumonia: results from the Community-Acquired Pneumonia Organization international study. *Int J Tuberc Lung Dis* 14:495–499, 2010.

\*Usual dose assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

- Hook EW III, Behets F, Van Damme E, et al. A phase III equivalence trial of azithromycin versus benzathine penicillin for treatment of early syphilis. *J Infect Dis* 201;1:1729–1735, 2010.
- Jain R, Danzinger LH. The macrolide antibiotics: a pharmacokinetic and pharmacodynamic overview. *Curr Pharm Des* 10:3045–53, 2004.
- Jenkins SG, Farrell DJ. Increase in *Pneumococcus* Macrolide Resistance, United States. *Emerging Infect Dis* 15:1260–1264, 2009.
- Kim Y-S, Yun H-J, Shim SK, et al. A comparative trial of a single dose of azithromycin versus doxycycline for the treatment of mild scrub typhus. *Clin Infect Dis* 39:1329–35, 2004.
- Magbanua JP, Goh BT, Michel CE, et al. Chlamydia trachomatis variant not detected by plasmid based nucleic acid amplification tests: molecular characterization and failure of single dose azithromycin. *Sex Transm Infect* 83:339–343, 2007.
- Malhotra-Kumar S, Lammens C, Coenen S, et al. Effect of azithromycin and clarithromycin therapy on pharyngeal carriage of macrolide-resistant streptococci in healthy volunteers: a randomized, double-blind, placebo-controlled study. *Lancet* 369:482–490, 2007.
- Mena LA, Mroczkowski TF, Nsuami M, et al. A randomized comparison of azithromycin and doxycycline for the treatment of *Mycoplasma genitalium* – positive urethritis in men. *Clin Infect Dis* 48:1649–1654, 2009.
- Paradisi F, Corti G. Azithromycin. *Antibiotics for Clinicians* 3:1–8, 1999.
- Phillips P, Chan K, Hogg R, et al. Azithromycin prophylaxis for *Mycobacterium avium* complex during the era of highly active antiretroviral therapy: evaluation of a provincial program. *Clin Infect Dis* 37:3471–8, 2002.
- Pichichero ME, Hoeger WJ, Casey JR. Azithromycin for the treatment of pertussis. *Pediatr Infect Dis J* 22:847–9, 2003.
- Plouffe JF, Breiman RF, Fields BS, et al. Azithromycin in the treatment of *Legionella* pneumonia requiring hospitalization. *Clin Infect Dis* 37:1475–80, 2003.
- Saiman L, Marshall BC, Mayer-Hamblett N, et al. Azithromycin in patients with cystic fibrosis chronically infected with *Pseudomonas aeruginosa*: a randomized controlled trial. *JAMA*. 290:1749–5, 2003.
- Savaris RF, Teixeira LM, Torres TG, et al. Comparing ceftriaxone plus azithromycin or doxycycline for pelvic inflammatory disease: a randomized controlled trial. *Obstet Gynecol* 110:53–60, 2007.
- Schlossberg D. Azithromycin and clarithromycin. *Med Clin N Amer* 79:803–816, 1995.
- Suzuki S, Yamazaki T, Narita M. Clinical evaluation of macrolide-resistant *Mycoplasma pneumoniae*. *Antimicrobial Agents and Chemotherapy*. 50:709–12, 2006.
- Swainston Harrison T, Keam SJ. Azithromycin extended release: a review of its use in the treatment of acute bacterial sinusitis and community-acquired pneumonia in the US. *Drugs* 773–79, 2007.
- To KK, Chan KH, Fung YF, et al. Azithromycin treatment failure in macrolide-resistant *Mycoplasma pneumoniae* pneumonia. *Eur Respir J* 36:969–971, 2010.
- Toltzis P, Dul M, Blumer J. Change in pneumococcal susceptibility to azithromycin during treatment for acute otitis media. *Pediatr Infect Dis* 26:647–649, 2007.
- Topic A, Skerf V, Puntaric A, et al. Azithromycin: 1.0 or 3.0 gram dose in the treatment of patients with asymptomatic urogenital chlamydial infections. *Journal of Chemotherapy* 18:115–16, 2006.
- Videler WJ, Badia L, Harvey RJ. Lack of efficacy of long-term, low-dose azithromycin in chronic rhinosinusitis: a randomized controlled trial. *Allergy* 66:1457–1468, 2011.
- Wilson R, Wells AU. Azithromycin in bronchiectasis: when should it be used? *Lancet*. 380:627–629, 2012.
- Wormser GP, Prasad A, Neuhaus E, et al. Emergence of resistance to azithromycin-atovaquone in immunocompromised patients with *Babesia microti* infection. *Clin Infect Dis* 50:381–386, 2010.
- Zuckerman JM, Qamar F, Bono BR. Review of macrolides (azithromycin, clarithromycin), ketolids (telithromycin) and glycylyclines (tigecycline). *Med Clin North Am* 95:761–791, 2011.
- Website: [www.pdr.net](http://www.pdr.net)

"Usual dose" assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

## Aztreonam (Azactam)

**Drug Class:** Monobactam.

**Usual Dose:** 2 gm (IV) q8h (see comments).

**Spectrum:** (see *Susceptibility Profiles* pp. 198–202).

**Resistance Potential:** Low

**Pharmacokinetic Parameters:**

Peak serum level: 204 mcg/mL

Bioavailability: Not applicable

Excreted unchanged (urine): 60–70%

Serum half-life (normal/ESRD): 1.7/7 hrs

Plasma protein binding: 56%

Volume of distribution ( $V_d$ ): 0.2 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\*** (based on 2 gm IV q8h)

CrCl 10–50 mL/min	1 gm (IV) q8h
CrCl < 10 mL/min	500 mg (IV) q8h
Post-HD dose	500 mg (IV)
Post-PD dose	500 mg (IV)
CWH/CVWHD/ CWHDF dose	1–2 gm (IV) q12h
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	No change

**Drug Interactions:** None.

**Adverse Effects:** None.

**Allergic Potential:** Low

**Safety in Pregnancy:** B

**Comments:** Incompatible in solutions containing vancomycin or metronidazole. **No cross allergenicity with penicillins,  $\beta$ -lactams; safe to use in penicillin allergic patients.**

**CAPD dose:** 1 gm (IP), then 250 mg/L of dialysate (IP) with each exchange.

Meningeal dose = 2 gm (IV) q6h.

**Cerebrospinal Fluid Penetration:**

Non-Inflamed meninges = 1%

Inflamed meninges = 40%

**Bile Penetration:** 300%

(also see *Antibiotic Pearls & Pitfalls* p. 511).

### REFERENCES:

- Assael BM. Aztreonam inhalation solution for suppressive treatment of chronic *Pseudomonas aeruginosa* lung infection in cystic fibrosis. *Expert Rev Anti Infect Ther* 9:967–973, 2011.
- Brogden RN, Heal RC. Aztreonam: A review of its antibacterial activity, pharmacokinetic properties, and therapeutic use. *Drugs* 31:96–130, 1986.
- Critchley IA, Sahn DF, Kelly LJ, et al. In vitro synergy studies using aztreonam and fluoroquinolone combinations against six species of Gram-negative bacilli. *Chemotherapy* 49:44–8, 2003.
- Cunha BA. Aztreonam: A review. *Urology* 41:249–58, 1993.
- Cunha BA. Cross allergenicity of penicillin with carbapenems and monobactams. *J Crit Illness* 13:344, 1998.
- Fleming DR, Ziegler C, Baize T, et al. Cefepime versus ticarcillin and clavulanate potassium and aztreonam for febrile neutropenia therapy in high-dose chemotherapy patients. *Am J Clin Oncol* 26:285–8, 2003.
- Gasink LB, Neil OF, Nachamkin I, et al. Risk factors for the impact of infection or colonization with aztreonam-resistant *Pseudomonas aeruginosa*. *Infection Control and Hospital Epidemiology* 28:1175–1180, 2007.
- Hellinger WC, Brewer NS. Carbapenems and monobactams: Imipenem, meropenem, and aztreonam. *Mayo Clin Proc* 74:420–34, 1999.
- Johnson, Cunha BA. Aztreonam. *Med Clin North Am* 79:733–43, 1995.
- Jordan EF, Nye MB, Luque AE. Successful treatment of *Pasteurella multocida* meningitis with aztreonam. *Scand J Infect Dis* 39:72–74, 2007.
- Zeitler K, Salvias B, Stevens V, et al. Aztreonam lysine for inhalation: new formulation of an old antibiotic. *Am J Health Syst Pharm* 69:107–115, 2012.
- Website: [www.pdr.net](http://www.pdr.net)

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose  $\times$  1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

## Capreomycin (Capastat)

**Drug Class:** Anti-TB drug.

**Usual Dose:** 1 gm (IM) q24h.

**Pharmacokinetic Parameters:**

*Peak serum level:* 30 mcg/mL

*Bioavailability:* Not applicable

*Excreted unchanged (urine):* 50%

*Serum half-life (normal/ESRD):* 5/30 hrs

*Plasma protein binding:* No data

*Volume of distribution ( $V_d$ ):* 0.4 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\***

CrCl 50–80 mL/min	500 mg (IM) q24h
CrCl 10–50 mL/min	500 mg (IM) q48h
CrCl < 10 mL/min	500 mg (IM) q72h
Post-HD dose	500 mg (IM)
Post-PD dose	None
CVWH/CVVD/ CVHDF dose	500 mg (IM) q48h
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	No change

**Drug Interactions:** None.

**Adverse Effects:** Eosinophilia, leukopenia, drug fever/rash, ototoxicity (vestibular), nephrotoxicity (glomerular/tubular).

**Allergic Potential:** Moderate

**Safety in Pregnancy:** C

**Comments:** Pain/phlebitis at IM injection site. Additive toxicity with aminoglycosides/viomycin.

**Cerebrospinal Fluid Penetration:** < 10%

## REFERENCES:

- Caminero JA, Sotgiu G, Zumla A, et al. Best drug treatment for multidrug-resistant and extensively drug-resistant tuberculosis. *Lancet Infect Dis* 10:621–629, 2010.
- Davidson PT, Le HQ. Drug treatment of tuberculosis - 1992. *Drugs* 43:651–73, 1992.
- Furin J, Nardell EA. Multidrug-resistant tuberculosis: an update on the best regimens. *J Respir Dis* 27:172–82, 2006.
- Iseman MD. Treatment of multidrug resistant tuberculosis. *N Engl J Med* 329:784–91, 1993.
- Website: www.pdr.net

## Caspofungin (Cancidas)

**Drug Class:** Echinocandin antifungal.

**Usual Dose:** **Loading Dose** 70 mg (IV) × 1 dose, then **Maintenance Dose:** 50 mg (IV) q24h (see comments).

**Pharmacokinetic Parameters:**

*Peak serum level:*

*70 mg:* 12.1 (single dose)/14.83 mcg/mL (multiple dose)

*50 mg:* 7.6/8.7 mcg/mL (multiple dose)

*Bioavailability:* Not applicable

*Excreted unchanged (urine):* 1.4%

*Serum half-life (normal/ESRD):* 10/10 hrs

*Plasma protein binding:* 97%

*Volume of distribution ( $V_d$ ):* No data

**Primary Mode of Elimination:** Hepatic

**Dosage Adjustments\***

CrCl < 10 mL/min	No change
Post-HD/PD dose	None
CVWH/CVVD/ CVHDF dose	None

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.



Moderate hepatic insufficiency	35 mg (IV) q24h (maintenance dose after 70 mg (IV) loading dose)
Severe hepatic insufficiency	Use with caution

**Drug Interactions:** Carbamazepine, rifampin, dexamethasone, efavirenz, nelfinavir, nevirapine, phenytoin (↓ caspofungin levels; ↑ caspofungin maintenance dose to 70 mg/day); cyclosporine (↑ caspofungin levels, ↑ SGOT/SGPT; co-administration is discouraged unless careful monitoring can be ensured and unless the benefit of caspofungin outweighs the risk of hepatotoxicity); tacrolimus (↓ tacrolimus levels, ↑ SGOT/SGPT).

**Adverse Effects:** Drug fever/rash.

**Allergic Potential:** Low

**Safety in Pregnancy:** C

**Comments:** Administer by slow IV infusion over 1 hour; do not give IV bolus. Do not mix/co-infuse with glucose solutions.

**Highly-resistant organisms dose: 70 mg (IV) q24h. Esophageal candidiasis dose:** begin therapy with 50 mg dose.

**Highly active against *Candida albicans* and non-albicans *Candida*, *Hansenula*.**

**Some activity against *C. parapsilosis*, *Aspergillus*. No activity against *Cryptococcus*, *Fusarium*, *Pseudoallescheria/Scedosporium*, *Rhodotorula*, *Trichosporon*, or *Mucor*.**

**Cerebrospinal Fluid Penetration:** 20%

#### REFERENCES:

Arathoon EG, Gotuzzo E, Noriega LM, et al. Randomized, double-blind, multicenter study of caspofungin versus amphotericin B for treatment of oropharyngeal and esophageal candidiasis. *Antimicrob Agents Chemother* 46:451–7, 2002.

Armstrong-James D, Stebbing J, John L, et al. A trial of caspofungin salvage treatment in PCP pneumonia. *Thorax* 66:537–538, 2011.

Barchiesi F, Spreghini E, Tomassetti S, et al. Efficacy of caspofungin against invasive *Candida* or invasive *Aspergillus* infections in neutropenic patients. *Antimicrob Agents Chemother* 50:2719–27, 2006.

Bennett JE, Echinocandins for candidemia in adults without neutropenia. *N Engl J Med* 355:1154–9, 2006.

Betts RF, Nucci M, Talwar D, et al. A multicenter, double-blind trial of a high-dose caspofungin treatment regimen versus a standard caspofungin treatment regimen for adult patients with invasive candidiasis. *Clin Infect Dis* 48:1676–1684, 2009.

Betts RF, Glasmacher A, Maertens J, et al. Efficacy of caspofungin against invasive *Candida* or invasive *Aspergillus* infections in neutropenic patients. *Cancer* 106:466–73, 2006.

Colombo AL, Ngai AL, Bourque M, et al. Caspofungin use in patients with invasive candidiasis caused by common non-albicans *Candida* species: review of the caspofungin database. *Antimicrob Agents Chemother* 54:1864–1871, 2010.

Deresinski SC, Stevens DA. Caspofungin. *Clin Infect Dis* 36:1445–57, 2003.

Falagas ME, Ntziora F, Betsi GI, et al. Caspofungin for the treatment of fungal infections: a systematic review of randomized controlled trials. *Int J Antimicrob Agents* 29:136–143, 2007.

Glasmacher A, Cornely OA, Orlopp K, et al. Caspofungin treatment in severely ill, immunocompromised patients: a case-documentation study of 118 patients. *J Antimicrob Chemother* 57:127–34, 2006.

Hiemenz JW, Raad II, Maertens JA, et al. Efficacy of caspofungin as salvage therapy for invasive aspergillosis compared to standard therapy in a historical cohort. *Eur J Clin Microbiol Infect Dis* 29:1387–1394, 2010.

Hope WW, Shoham S, Walsh TJ. The pharmacology and clinical use of caspofungin. *Expert Opin Drug Metab Toxicol* 3:263–274, 2007.

Keating G, Figgitt D. Caspofungin: a review of its use in esophageal candidiasis, invasive candidiasis and invasive aspergillosis. *Drugs* 63:2235–63, 2003.

"Usual dose" assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

- Kofia G, Ruhnke M. Pharmacology and metabolism of anidulafungin, caspofungin, and micafungin in the treatment of invasive candidosis: review of the literature. *Eur J Med Res* 16:159–166, 2011.
- Lomaestro BM. Caspofungin. *Hospital Formulary* 36:527–36, 2001.
- Lortholary O, Desnos-Ollivier M, Sitbon K, et al. Recent exposure to caspofungin or fluconazole influences the epidemiology of candidemia: a prospective multicenter study involving 2,441 patients. *Antimicrob Agents Chemother* 55:532–538, 2011.
- Maertens J, Raad I, Petrikos G, et al. Efficacy and safety of caspofungin for treatment of invasive aspergillosis in patients refractory to or intolerant of conventional antifungal therapy. *Clin Infect Dis* 39:1563–71, 2004.
- Mattiuzzi GN, Alvarado G, Giles FJ, et al. Open-label, randomized comparison of itraconazole versus caspofungin for prophylaxis in patients with hematologic malignancies. *Antimicrob Agents Chemother* 50:143–7, 2006.
- Miceli MH, Diaz JA, Lee SA. Emerging opportunistic yeast infections. *Lancet Infectious Diseases* 11:142–151, 2011.
- Mora-Duarte J, Betts R, Rotstein C, et al. Comparison of caspofungin and amphotericin B for invasive candidiasis. *N Engl J Med* 347:2020–9, 2002.
- Morrison VA. Echinocandin antifungals: review and update. *Expert Rev Anti Infect Ther* 4:325–42, 2006.
- Navado J, De Alarcon A, Hernandez A. Caspofungin: a new therapeutic option for fungal endocarditis. *Clin Microbiol Infect* 11:248, 2005.
- Nguyen TH, Hoppe-Tichy T, Geiss HK, et al. Factors influencing caspofungin plasma concentrations in patients of a surgical intensive care unit. *J Antimicrob Chemother* 60:100–106, 2007.
- Nivoix Y, Zamfir A, Lutun P, Kara F, et al. Combination of caspofungin and an azole or an amphotericin B formulation in invasive fungal infections. *J Infect* 52:67–74, 2006.
- Pappas PG, Rotstein CM, Betts RF, et al. Micafungin versus caspofungin for treatment of candidemia and other forms of invasive candidiasis. *Clin Infect Dis* 45:883–893, 2007.
- Pfaller MA, Messer SA, Boyken L, et al. Caspofungin activity against clinical isolates of fluconazole-resistant *Candida*. *J Clin Microbiol* 41:5729–31, 2003.
- Sobel JD, Bradshaw SK, Lipka CJ, et al. Caspofungin in the treatment of symptomatic candiduria. *Clin Infect Dis* 44:e46–e49, 2007.
- Steinmann J, Buer J, Rath PM. Caspofungin: cross-reactivity in the *Aspergillus* antigen assay. *J Clin Microbiol* 48:2313, 2010.
- Walsh TJ, Tepler H, Donowitz GR, et al. Caspofungin versus liposomal amphotericin B for empirical antifungal therapy in patients with persistent fever and neutropenia. *N Engl J Med* 351:1391–402, 2004.
- Wiederhold NP, Najvar LK, Bocanegra RA, et al. Caspofungin dose escalation for invasive candidiasis due to resistant *Candida albicans*. *Antimicrob Agents Chemother* 55:3254–3260, 2011.
- Wilke M. Treatment and prophylaxis of invasive candidiasis with anidulafungin, caspofungin, and micafungin and its impact on use and costs: review of the literature. *Eur J Med Res* 16:180–186, 2011.
- Winkler M, Pratschke J, Schulz U, et al. Caspofungin for post solid organ transplant invasive fungal disease: results of a retrospective observational study. *Transpl Infect Dis* 12:230–237, 2010.
- Website: [www.pdr.net](http://www.pdr.net)

## Cefaclor (Ceclor)

**Drug Class:** 2<sup>nd</sup> generation oral cephalosporin.

**Usual Dose:** 500 mg (PO) q8h.

**Spectrum:** (see **Susceptibility Profiles** pp. 191–197).

**Resistance Potential:** Low

**Pharmacokinetic Parameters:**

*Peak serum level:* 13 mcg/mL

*Bioavailability:* 80%

*Excreted unchanged (urine):* 60–85%

*Serum half-life (normal/ESRD):* 0.8/3 hrs

*Plasma protein binding:* 25%

<sup>†</sup>Usual dose<sup>†</sup> assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

Volume of distribution ( $V_d$ ): 0.30 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\***

CrCl < 10 mL/min	250 mg (PO) q8h
Post-HD dose	250 mg (PO)
Post-PD dose	None
CVWH/CVHD/CVHDF dose	No change
Moderate—severe hepatic insufficiency	No change

**Drug Interactions:** None.

**Adverse Effects:** Drug fever/rash, *C. difficile* diarrhea/colitis.

**Allergic Potential:** High

**Safety in Pregnancy:** B

**Comments:** Limited penetration into respiratory secretions. **Ceclor CD 500 mg (PO) q12h is equivalent to cefaclor 250 mg (PO) q8h, not cefaclor 500 mg (PO) q8h. Ceclor CD 500 mg (PO) q12h is equivalent to 250 mg (PO) q8h of cefaclor.**

**Cerebrospinal Fluid Penetration:** < 10%

**Bile Penetration:** 60%

(also see **Antibiotic Pearls & Pitfalls** p. 510).

#### REFERENCES:

- Cazzola M, Di Perna F, Boveri B. Interrelationship between the pharmacokinetics and pharmacodynamics of cefaclor advanced formulation in patients with acute exacerbation of chronic bronchitis. *J Chemother* 12:216–22, 2000.
- Mazzei T, Novelli A, Esposito S, et al. New insight into the clinical pharmacokinetics of cefaclor: Tissue penetration. *J Chemother* 12:53–62, 2000.
- Meyers BR. Cefaclor revisited. *Clin Ther* 22:154–66, 2000.
- Website: www.pdr.net

## Cefadroxil (Duricef, Ultracef)

**Drug Class:** 1<sup>st</sup> generation oral cephalosporin.

**Usual Dose:** 1000 mg (PO) q12h.

**Spectrum:** (see **Susceptibility Profiles** pp. 191–197).

**Resistance Potential:** Low

**Pharmacokinetic Parameters:**

*Peak serum level:* 16 mcg/mL

*Bioavailability:* 99%

*Excreted unchanged (urine):* 85%

*Serum half-life (normal/ESRD):* 0.5/22 hrs

*Plasma protein binding:* 20%

*Volume of distribution ( $V_d$ ):* 0.31 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\***

CrCl 10–25 mL/min	500 mg (PO) q24h
CrCl < 10 mL/min	500 mg (PO) q36h
Post-HD dose	500 mg (PO)
Post-PD dose	250 mg (PO)
CVWH/CVHD/CVHDF dose	500 mg (PO) q24h
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	No change

**Drug Interactions:** None.

**Adverse Effects:** Drug fever/rash, *C. difficile* diarrhea/colitis.

**Allergic Potential:** High

**Safety in Pregnancy:** B

**Comments:** Penetrates oral/respiratory secretions well.

**Cerebrospinal Fluid Penetration:** < 10%

**Bile Penetration:** 20%

(also see **Antibiotic Pearls & Pitfalls** p. 509).

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose  $\times$  1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

**REFERENCES:**

- Bucko AD, Hunt BJ, Kidd SL, et al. Randomized, double-blind, multicenter comparison of oral cefditoren 200 or 400 mg BID with either cefuroxime 250 mg BID or cefadroxil 500 mg BID for the treatment of uncomplicated skin and skin-structure infections. *Clin Ther* 24:1134–47, 2002.
- Cunha BA. Antibiotics selection for the treatment of sinusitis, otitis media, and pharyngitis. *Infect Dis Pract* 7:5324–5326, 1998.
- Gustaferro CA, Steckelberg JM. Cephalosporins: Antimicrobial agents and related compounds. *Mayo Clin Proc* 66:1064–73, 1991.
- Website: [www.pdr.net](http://www.pdr.net)

**Cefamandole (Mandol)**

**Drug Class:** 2<sup>nd</sup> generation cephalosporin.

**Usual Dose:** 2 gm (IV) q6h.

**Spectrum:** (see *Susceptibility Profiles* pp.191–197)

**Resistance Potential:** High (H. influenzae, Enterobacter sp.)

**Pharmacokinetic Parameters:**

*Peak serum level:* 240 mcg/mL

*Bioavailability:* Not applicable

*Excreted unchanged (urine):* 85%

*Serum half-life (normal/ESRD):* 1/11 hrs

*Plasma protein binding:* 76%

*Volume of distribution (V<sub>d</sub>):* 0.29 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\***

CrCl 50–80 mL/min	1 gm (IV) q6h
CrCl 10–50 mL/min	1 gm (IV) q6h
CrCl < 10 mL/min	1 gm (IV) q12h
Post-HD dose	1 gm (IV)
Post-PD dose	1 gm (IV)
CVWH/CVVD/ CVHDF dose	1 gm (IV) q6h
Moderate—severe hepatic insufficiency	No change

**Drug Interactions:** Alcohol (disulfiram-like reaction); antiplatelet agents, heparin, thrombolytics, warfarin (↑ risk of bleeding).

**Adverse Effects:** Drug fever/rash, ↑ INR.

**Allergic Potential:** High

**Safety in Pregnancy:** B

**Comments:** Incompatible in solutions with Mg<sup>++</sup> or Ca<sup>++</sup>. **Contains MTT side chain, but no increase in clinical bleeding.**

**Cerebrospinal Fluid Penetration:** < 10%

**Bile Penetration:** 300%

(also see *Antibiotic Pearls & Pitfalls* p. 510).

**REFERENCES:**

- Cunha BA, Klimek JJ, Qunitiliani R. Cefamandole nafate in respiratory and urinary tract infections. *Curr Ther Res* 25:584–9, 1971.
- Gentry LO, Zeluff BJ, Cooley DA. Antibiotic prophylaxis in open-heart surgery: A comparison of cefamandole, cefuroxime, and cefazolin. *Ann Thorac Surg* 46:167–71, 1988.
- Website: [www.pdr.net](http://www.pdr.net)

**Cefazolin (Ancef, Kefzol)**

**Drug Class:** 1<sup>st</sup> generation cephalosporin.

**Usual Dose:** 1 gm (IV) q8h.

**Spectrum:** (see *Susceptibility Profiles* pp. 191–197).

**Resistance Potential:** Low

**Pharmacokinetic Parameters:**

*Peak serum level:* 185 mcg/mL

*Bioavailability:* Not applicable

*Excreted unchanged (urine):* 96%

*Serum half-life (normal/ESRD):* 1.8/40 hrs

*Plasma protein binding:* 85%

*Volume of distribution (V<sub>d</sub>):* 0.2 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\***

CrCl 35–55 mL/min	1 gm (IV) q12h
CrCl 10–35 mL/min	500 mg (IV) q12h

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

CrCl < 10 mL/min	500 mg (IV) q24h
Post-HD dose	1 gm (IV)
Post-HFHD dose	2 gm (IV)
Post-PD dose	500 mg (IV)
CVWH/CVWHD/CVWHDf dose	500 mg (IV) q12h
HD q 48–72h dose	2 gm (IV) q HD*
Moderate—severe hepatic insufficiency	No change

\*Give at end of HD.

**Drug Interactions:** None.

**Adverse Effects:** Drug fever/rash, AIHA.

**Allergic Potential:** High

**Safety in Pregnancy:** B

**Comments:** Incompatible in solutions containing erythromycin, aminoglycosides, cimetidine, theophylline.

**Cerebrospinal Fluid Penetration:** < 10%

**Bile Penetrations:** 300%

(also see **Antibiotic Pearls & Pitfalls** p. 509).

#### REFERENCES:

- Adebri C, Ristori R, Chelazzi C, et al. Cefazolin bolus and continuous administration for elective cardiac surgery: improved pharmacokinetic and pharmacodynamic parameters. *J Thorac Cardiovasc Surg* 140:471–475, 2010.
- Cunha BA, Gossling HR, Pasternak HS, et al. Penetration of cephalosporins into bone. *Infection* 12:80–4, 1984.
- Gentry LO, Zeluff BJ, Cooley DA. Antibiotic prophylaxis in open-heart surgery: A comparison of cefamandole, cefuroxime, and cefazolin. *Ann Thorac Surg* 46:167–71, 1988.
- Marshall WF, Blair JE. The cephalosporins. *Mayo Clin Proc* 74:187–95, 1999.
- Nightingale CH, Klimek JJ, Quintiliani R. Effect of protein binding on the penetration of nonmetabolized cephalosporins into atrial appendage and pericardial

fluids in open-heart surgical patients. *Antimicrob Agents Chemother* 17:595–8, 1980.

Quintiliani R, Nightingale CH. Cefazolin. *Ann Intern Med* 89:650–6, 1978.

Turnidge JD. Cefazolin and Enterobacteriaceae: rationale for revised susceptibility testing breakpoints. *Clin Infect Dis* 52:917–924, 2011.

Youngster I, Shenoy ES, Hooper D, et al. Comparative evaluation of the tolerability of cefazolin and nafcillin for treatment of methicillin-susceptible *Staphylococcus aureus* infections in the outpatient setting. *Clin Infect Dis*. 59:369-75, 2014.

Website: [www.pdr.net](http://www.pdr.net)

## Cefdinir (Omnicef)

**Drug Class:** 3<sup>rd</sup> generation oral cephalosporin.

**Usual Dose:** 600 mg (PO) q24h.

**Spectrum:** (see **Susceptibility Profiles** pp. 191–197).

**Resistance Potential:** Low

**Pharmacokinetic Parameters:**

*Peak serum level:* 2.9 mcg/mL

*Bioavailability:* 16% (tab) / 25% (suspension)

*Excreted unchanged (urine):* 11–18%

*Serum half-life (normal/ESRD):* 1.7/3 hrs

*Plasma protein binding:* 70%

*Volume of distribution (V<sub>d</sub>):* 0.35 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\***

CrCl < 30 mL/min	300 mg (PO) q24h
Post-HD dose	300 mg (PO)
Post-PD dose	None
CVWH/CVWHD/CVWHDf dose	300 mg (PO) q24h
Moderate—severe hepatic insufficiency	No change

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

**Drug Interactions:** Probenecid (↑ cefdinir levels).

**Adverse Effects:** Drug fever/rash, *C. difficile* diarrhea/colitis.

**Allergic Potential:** High

**Safety in Pregnancy:** B

**Comments:** Good activity against bacterial respiratory pathogens. Available as capsules or suspension. **CAP dose:** 300 mg (PO) q12h; **Other respiratory infections dose:** 600 mg (PO) q24h.

**Cerebrospinal Fluid Penetration:** < 10% (also see **Antibiotic Pearls & Pitfalls** p. 511).

#### REFERENCES:

- Cefdinir: A new oral cephalosporin. *Med Lett Drugs Therap* 40:85–7, 1998.
- Fogarty CM, Bettis RB, Griffin TJ, et al. Comparison of a 5 day regimen of cefdinir with a 10 day regimen of cefprozil for treatment of acute exacerbations of chronic bronchitis. *J Antimicrob Chemother* 45:851–8, 2000.
- Guay DR. Cefdinir: an advanced-generation, broad-spectrum oral cephalosporin. *Clin Ther* 24:473–89, 2002.
- Hedrick JA. Community-acquired upper respiratory tract infections and the role of third-generation oral cephalosporins. *Expert Rev Anti Infect Ther* 8:15–21, 2010.
- Perry CM, Scott LJ. Cefdinir: a review of its use in the management of mild-to-moderate bacterial infections. *Drugs* 64:1433–64, 2004.
- Website: www.pdr.net

## Cefditoren (Spectracef)

**Drug Class:** 3<sup>rd</sup> generation oral cephalosporin.

**Usual Dose:** 400 mg (PO) q12h.

**Spectrum:** (see **Susceptibility Profiles** pp. 191–197).

**Resistance Potential:** Low

**Pharmacokinetic Parameters:**

*Peak serum level:* 1.8 mcg/mL

*Bioavailability:* 16%

*Excreted unchanged (urine):* 20%

*Serum half-life (normal/ESRD):* 1.5/5 hrs

*Plasma protein binding:* 88%

*Volume of distribution (V<sub>d</sub>):* 0.13 L/kg

**Primary Mode of Elimination:** Renal  
**Dosage Adjustments\***

CrCl 50–80 mL/min	No change
CrCl 30–50 mL/min	200 mg (PO) q12h
CrCl < 30 mL/min	200 mg (PO) q24h
Post-HD dose	200 mg (PO)
Post-PD dose	None
CVVH/CVVHD/ CVVHDF dose	200 mg (PO) q24h
Moderate—severe hepatic insufficiency	No change

**Drug Interactions:** H<sub>2</sub> receptor antagonists, Al<sup>3+</sup>, Mg<sup>2+</sup> antacids (↓ absorption of cefditoren); Probenecid (↓ elimination of cefditoren).

**Adverse Effects:** Drug fever/rash, *C. difficile* diarrhea/colitis.

**Allergic Potential:** Low

**Safety in Pregnancy:** B

**Comments:** Serum concentrations increased ~ 50% if taken without food. ↑ elimination of carnitine; do not use in carnitine deficiency or hereditary carnitine metabolism disorder.

**Contains Na+ caseinate; avoid in patients with casein hypersensitivity.**

**Cerebrospinal Fluid Penetration:** < 1%

**Bile Penetration:** 200%

(also see **Antibiotic Pearls & Pitfalls** p. 511).

#### REFERENCES:

- Alvarez-Sala JL, Kardos P, et al. Martinez-Beltran, et al. Clinical and bacteriological efficacy in treatment of acute exacerbations of chronic bronchitis with

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

- cefditoren-pivoxil versus cefuroxime-axetil. *Antimicrob Agents and Chemother* 50:1762–67, 2006.
- Balbis EA. Cefditoren, a new aminothiazolyl cephalosporin. *Pharmacotherapy* 22:1278–93, 2002.
- Chow J, Russel M, Bolk S, et al. Efficacy of cefditoren pivoxil vs. amoxicillin-clavulanic in acute maxillary sinusitis. Presented at the 40<sup>th</sup> Interscience Conference on Antimicrobial Agents and Chemotherapy; Abstract 835, Toronto, ON 2000.
- Clark CL, Nagai K, Dewasse BE, et al. Activity of cefditoren against respiratory pathogens. *J Antimicrob Chemother* 50:33–41, 2002.
- Darkes MJ, Plosker GL. Cefditoren pivoxil. *Drugs* 62:319–36, 2002.
- Guay DR. Review of cefditoren, an advanced-generation, broad-spectrum oral cephalosporin. *Clin Ther* 23:1924–37, 2002.
- Tempera G, Furneri PM, Carlone NA, et al. Antibiotic susceptibility of respiratory pathogens recently isolated in Italy: focus on cefditoren. *J Chemother* 22:153–159, 2010.
- Wellington K, Curran MP. Cefditoren pivoxil: a review of its use in the treatment of bacterial infections. *Drugs* 64:2597–618, 2004.
- Website: www.pdr.net

## Cefepime (Maxipime)

- Drug Class:** 4<sup>th</sup> generation cephalosporin.
- Usual Dose:** 2 gm (IV) q12h (see comments).
- Spectrum:** (see **Susceptibility Profiles** pp. 191–197).
- Resistance Potential:** Low
- Pharmacokinetic Parameters:**  
*Peak serum level:* 163 mcg/mL  
*Bioavailability:* Not applicable  
*Excreted unchanged (urine):* 80%  
*Serum half-life (normal/ESRD):* 2.2/18 hrs  
*Plasma protein binding:* 20%  
*Volume of distribution ( $V_d$ ):* 0.29 L/kg
- Primary Mode of Elimination:** Renal
- Dosage Adjustments\***

CrCl 30–60 mL/min	2 gm (IV) q12h
CrCl 10–30 mL/min	2 gm (IV) q24h
CrCl < 10 mL/min	1 gm (IV) q24h
Post–HD dose	2 gm (IV)
Post–HFHD dose	2 gm (IV)
Post–PD dose	1 gm (IV)
CAPD	2 gm (IV) q48h
CVWH/CVWHD/ CVWHD dose	1 gm (IV) q12h
HD q 48–72h dose	2 gm (IV) q HD*
Moderate—severe hepatic insufficiency	No change

\*Give at end of HD.

**Drug Interactions:** None.

**Adverse Effects:** Drug fever/rash, *C. difficile* diarrhea/colitis. **Neurotoxicity, particularly with doses not adjusted for renal function.**

**Neurotoxic manifestations include encephalopathy, non-convulsive status epilepticus, and myoclonus.**

**Allergic Potential:** Moderate

**Safety in Pregnancy:** B

**Comments:** For ESBL + Enterobacteriaceae (MIC 4–8 mcg/mL, proven serious systemic *P. aeruginosa* infections, febrile neutropenia, or cystic fibrosis, use 2 gm (IV) q8h. For aerobic GNB bacteremias use 2 gm (IV) q12h.

Meningeal dose = 2 gm (IV) q8h.

**Cerebrospinal Fluid Penetration:**

Non-Inflamed meninges = 1%

Inflamed meninges = 15%

**Bile Penetration:** 10%

(also see **Antibiotic Pearls & Pitfalls** p. 511).

"Usual dose" assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

## REFERENCES:

- Akhabue E, Synnestvedt M, Weiner MG, et al. Cefepime-resistant *Pseudomonas aeruginosa*. *Emerg Infect Dis* 17:1037–1043, 2011.
- Al-Hassan MN, Eckel-Passow JE, Baddour LM. Cefepime effectiveness in Gram-negative bloodstream infections. *J Antimicrob Chemother* 66:1156–1160, 2011.
- Altshuler J. Treatment of extended-spectrum beta-lactamase Enterobacteriaceae with cefepime: the dose matters, too. *Clin Infect Dis* 57:915–916, 2013.
- Bazan JA, Martin SI, Kaye KM. Newer beta-lactam antibiotics: doripenem, ceftobiprole, ceftaroline fosamil, and cefepime. *Med Clin North Am* 95:743–760, 2011.
- Bow EJ, Rotstein C, Noskin GA, et al. A Randomized, open-label, multicenter comparative study of the efficacy and safety of piperacillin-tazobactam and cefepime for the empirical treatment of febrile neutropenic episodes in patients with hematologic malignancies. *Clin Infect Dis* 43:447–59, 2006.
- Chapman TM, Perry CM. Cefepime: a review of its use in the management of hospitalized patients with pneumonia. *Am J Respir Med* 2:75–107, 2003.
- Cheatham SC, Shea KM, Healy DP, et al. Steady-state pharmacokinetics and pharmacodynamics of cefepime administered by prolonged infusion in hospitalized patients. *Int J Antimicrob Agents* 37:46–50, 2011.
- Cunha BA. *Pseudomonas aeruginosa*: Resistance and therapy. *Semin Respir Infect* 17:231–9, 2002.
- Cunha BA, Gill MV. Cefepime. *Med Clin North Am* 79:721–32, 1995.
- Durand-Maugard C, Lemaire-Hurtel A, Gras-Champel V, et al. Blood and CSF monitoring of cefepime-induced neurotoxicity: nine case reports. *J Antimicrob Chemother* 1093:1297–1298, 2012.
- Fugate JE, Kalimullah EA, Hocker SE. Cefepime neurotoxicity in the intensive care unit: a cause of severe, underappreciated encephalopathy. *Crit Care* 17:R264, 2013.
- Gangireddy VG, Mitchell LC, Coleman T. Cefepime neurotoxicity despite renal adjusted dosing. *Scand J Infect Dis* 43:827–829, 2011.
- Lee SY, Kuti JL, Nicolau DP. Cefepime pharmacodynamics in patients with extended spectrum beta-lactamase (ESBL) and non-ESBL infections. *J Infect* 54:463–468, 2007.
- Montalar J, Segura A, Bosch C, et al. Cefepime monotherapy as an empirical initial treatment of patients with febrile neutropenia. *Med Oncol* 19:161–6, 2002.
- Perez KK, Hughes DW, Maxwell PR, et al. Cefepime for Gram-negative bacteremia in long-term hemodialysis: a single-center experience. *Am J Kidney Dis* 59:740–742, 2012.
- Picao RC, Jones RN, Mendes RE, et al. Klebsiella pneumoniae carbapenemase-producing enterobacteriaceae testing susceptible to cefepime by reference methods. *Clin Micro* 51:2388–2390, 2013.
- Siedne MJ, Galar A, Guzman-Suarez BB, et al. Cefepime vs other antibacterial agents for the treatment of Enterobacter species bacteremia. *Clin Infect Dis* 58:1554–1563, 2014.
- Spriet I, Meersseman W, De Troy E, et al. Meropenem–valproic acid interaction in patients with cefepime-associated status epilepticus. *Am J Health Syst Pharm* 64:54–58, 2007.
- Tamma PD, Girdwood SCT, Gopaul R, et al. The use of cefepime for treating AmpC  $\beta$ -lactamase-producing Enterobacteriaceae. *Clin Infect Dis* 57:781–788, 2013.
- Tam VH, McKinnon PS, Akins RL, et al. Pharmacokinetics and pharmacodynamics of cefepime in patients with various degrees of renal function. *Antimicrob Agents Chemother* 47:1853–61, 2003.
- Ugai T, Morisaki K, Tsuda K. Cefepime induced encephalopathy in patients with hematological malignancies: clinical features and risk factors. *Scand J Infect Dis*. 46:272–279, 2014.
- Yahave D, Paul M, Fraser A, et al. Efficacy and safety of cefepime: a systematic review and meta-analysis. *Lancet Infect Dis* 7:338–348, 2007.
- Wilson FP, Bachhuber MA, Caroff D, et al. Low cefepime concentrations during high blood and dialysate flow continuous venovenous hemodialysis. *Antimicrob Agents Chemother* 56:2178–2180, 2012.
- Website: [www.pdr.net](http://www.pdr.net)

"Usual dose" assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose  $\times$  1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.



## Cefixime (Suprax)

**Drug Class:** 3<sup>rd</sup> generation oral cephalosporin.

**Usual Dose:** 400 mg (PO) q12h.

**Spectrum:** (see **Susceptibility Profiles** pp. 191–197).

**Resistance Potential:** Low

### Pharmacokinetic Parameters:

*Peak serum level:* 3.7 mcg/mL

*Bioavailability:* 50%

*Excreted unchanged (urine):* 50%

*Serum half-life (normal/ESRD):* 3.1/11 hrs

*Plasma protein binding:* 65%

*Volume of distribution ( $V_d$ ):* 0.1 L/kg

**Primary Mode of Elimination:** Renal

### Dosage Adjustments\*

CrCl 20–50 mL/min	400 mg (PO) q24h
CrCl < 20 mL/min	200 mg (PO) q24h
Post-HD dose	400 mg (PO)
Post-PD dose	200 mg (PO)
CWVH/CWHD/ CWVDF dose	200 mg (PO) q12h
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	No change

**Drug Interactions:** Carbamazepine (↑ carbamazepine levels).

**Adverse Effects:** Drug fever/rash, *C. difficile* diarrhea/colitis.

**Allergic Potential:** High

**Safety in Pregnancy:** B

**Comments:** **PRNG dose:** 400 mg (PO) × 1 dose. Also useful for quinolone-resistant GC. **Little/no activity against *S. aureus* (MSSA).**

**Cerebrospinal Fluid Penetration:** < 10%

**Bile Penetration:** 800%

(also see **Antibiotic Pearls & Pitfalls** p. 511).

### REFERENCES:

Chisholm SA, Mouton JW, Lewis DA, et al. Cephalosporin MIC creep among gonococci: time for a pharmacodynamic rethink? *J Antimicrob Chemother* 65:2141–2148, 2010.

Quintillani R. Cefixime in the treatment of patients with lower respiratory tract infections: results of US clinical trials. *Clinical Therapeutics* 18:373–90, 1996. Update to CDC's Sexually Transmitted Diseases Treatment Guidelines, 2006: fluoroquinolones no longer recommended for treatment of gonococcal infections. *MMWR* 56:332–336, 2007.

Website: [www.pdr.net](http://www.pdr.net)

## Cefoperazone (Cefobid)

**Drug Class:** 3<sup>rd</sup> generation cephalosporin.

**Usual Dose:** 2 gm (IV) q12h.

**Spectrum:** (see **Susceptibility Profiles** pp. 191–197).

**Resistance Potential:** Low

### Pharmacokinetic Parameters:

*Peak serum level:* 240 mcg/mL

*Bioavailability:* Not applicable

*Excreted unchanged (urine):* 20%

*Serum half-life (normal/ESRD):* 2.4/2.4 hrs

*Plasma protein binding:* 90%

*Volume of distribution ( $V_d$ ):* 0.17 L/kg

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CWVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

**Primary Mode of Elimination:** Hepatic

**Dosage Adjustments\***

CrCl < 10 mL/min	No change
Post-HD dose	None
Post-PD dose	None
CVVH/CVVHD/ CVVHDF dose	No change
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	1 gm (IV) q12h

**Drug Interactions:** Alcohol (disulfiram-like reaction); antiplatelet agents, heparin, thrombolytics, warfarin ( $\uparrow$  risk of bleeding).

**Adverse Effects:** Drug fever/rash.  $\uparrow$  INR due to MTT side chain, but no increase in clinical bleeding. Prophylactic vitamin K unnecessary.

**Allergic Potential:** Low

**Safety in Pregnancy:** B

**Comments:** One of the few antibiotics to penetrate into an obstructed biliary tract. May be administered IM. **Concentration dependent serum half life.**

Meningeal dose = 2 gm (IV) q8h.

**Cerebrospinal Fluid Penetration:**

Non-Inflamed meninges = 1%

Inflamed meninges = 10%

**Bile Penetration:** 1200% (also see **Antibiotic Pearls & Pitfalls** p. 510).

**REFERENCES:**

Cunha BA. 3<sup>rd</sup> generation cephalosporins: A review. Clin Ther 14:616–52, 1992.

Klein NC, Cunha BA. Third-generation cephalosporins. Med Clin North Am 79:705–19, 1995.

Marshall WF, Blair JE. The cephalosporins. Mayo Clin Proc 74:187–95, 1999.

Oie S, Uematsu T, Sawa A, et al. In vitro effects of combinations of antipseudomonal agents against seven strains of multidrug-resistant *Pseudomonas aeruginosa*. J Antimicrob Chemother 52:911–4, 2004.

Website: www.pdr.net

## Cefotaxime (Claforan)

**Drug Class:** 3<sup>rd</sup> generation cephalosporin.

**Usual Dose:** 2 gm (IV) q6h.

**Spectrum:** (see **Susceptibility Profiles** pp. 191–197).

**Resistance Potential:** Low

**Pharmacokinetic Parameters:**

*Peak serum level:* 214 mcg/mL

*Bioavailability:* Not applicable

*Excreted unchanged (urine):* 20–36%; 15–25% excreted as active metabolite

*Serum half-life (normal/ESRD):* 1/15 hrs

*Plasma protein binding:* 37%

*Volume of distribution ( $V_d$ ):* 0.25 L/kg

**Primary Mode of Elimination:** Renal  
**Dosage Adjustments\***

CrCl 10–50 mL/min	1 gm (IV) q12h
CrCl < 10 mL/min	1 gm (IV) q24h
Post-HD dose	1 gm (IV)
Post-PD dose	1 gm (IV)
CVVH/CVVHD/ CVVHDF dose	2 gm (IV) q12h
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	No change

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose  $\times$  1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

**Drug Interactions:** None.

**Adverse Effects:** Drug fever/rash, *C. difficile* diarrhea/colitis.

**Allergic Potential:** Moderate

**Safety in Pregnancy:** B

**Comments:** Incompatible in solutions containing sodium bicarbonate, metronidazole, or aminoglycosides. **Desacetyl metabolite** ( $t_{1/2} = 1.5$  hrs) **synergistic with cefotaxime against *S. aureus/B. fragilis*.**

Meningeal dose = 3 gm (IV) q6h.

**Cerebrospinal Fluid Penetration:**

Non-Inflamed meninges = 1%

Inflamed meninges = 10%

**Bile Penetration:** 75%

(also see **Antibiotic Pearls & Pitfalls** p. 510).

#### REFERENCES:

Brogden RN, Spencer CM. Cefotaxime: A reappraisal of its antibacterial activity and pharmacokinetic properties and a review of its therapeutic efficacy when administered twice daily for the treatment of mild to moderate infections. *Drugs* 53:483–510, 1987.

File Jr. TM. Clinical implications and treatment of multiresistant *Streptococcus pneumoniae pneumonia*. *Clin Microbiol Infect* 12:31–41, 2006.

Klein NC, Cunha BA. Third-generation cephalosporins. *Med Clin North Am* 79:705–19, 1995.

Marshall WF, Blair JE. The cephalosporins. *Mayo Clin Proc* 74:187–95, 1999.

Patel KB, Nicolau DP, Nightingale CH, et al. Comparative serum bactericidal activities of ceftizoxime and cefotaxime against intermediately penicillin-resistant *Streptococcus pneumoniae*. *Antimicrob Agents Chemother* 40:2805–8, 1996.

Website: www.pdr.net

## Cefotetan (Cefotan)

**Drug Class:** 2<sup>nd</sup> generation cephalosporin (Cephamycin).

**Usual Dose:** 2 gm (IV) q12h.

**Spectrum:** (see **Susceptibility Profiles** pp. 191–197).

**Resistance Potential:** Low

**Pharmacokinetic Parameters:**

*Peak serum level:* 237 mcg/mL

*Bioavailability:* Not applicable

*Excreted unchanged (urine):* 50–80%

*Serum half-life (normal/ESRD):* 4/10 hrs

*Plasma protein binding:* 88%

*Volume of distribution ( $V_d$ ):* 0.17 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\***

CrCl 30–80 mL/min	2 gm (IV) q12h
CrCl 10–30 mL/min	2 gm (IV) q24h
CrCl < 10 mL/min	2 gm (IV) q48h
Post-HD dose	1 gm (IV)
Post-PD dose	1 gm (IV)
CVVH/CVVHD/ CVVHDF dose	750 mg (IV) q12h
Moderate—severe hepatic insufficiency	No change

**Drug Interactions:** Alcohol (disulfiram-like reaction); antiplatelet agents, heparin, thrombolytics, warfarin (↑ risk of bleeding).

**Adverse Effects:** Drug fever/rash, hemolytic anemia, *C. difficile* diarrhea/colitis. ↑ **INR due to MTT side chain, but no increase in clinical bleeding.**

**Allergic Potential:** Low

**Safety in Pregnancy:** B

**Comments:** Less effective than cefoxitin against *B. fragilis* D.O.T strains.

**Cerebrospinal Fluid Penetration:** < 10%

**Bile Penetration:** 20% (also see **Antibiotic Pearls & Pitfalls** p. 510).

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

**REFERENCES:**

- Moes GS, MacPherson BR. Cefotetan-induced hemolytic anemia: A case report and a review of the literature. *Arch Pathol Lab Med* 124:1344–6, 2000.
- Ray EK, Warkentin TE, O'Hoski PL. Delayed onset of life-threatening immune hemolysis after perioperative antimicrobial prophylaxis with cefotetan. *Can J Surg* 43:461–2, 2000.
- Stroncek D, Procter JL, Johnson J. Drug-induced hemolysis: cefotetan-dependent hemolytic anemia mimicking an acute intravascular immune transfusion reaction. *Am J Hematol* 64:67–70, 2000.
- Website: [www.pdr.net](http://www.pdr.net)

## Cefoxitin (Mefoxin)

**Drug Class:** 2<sup>nd</sup> generation cephalosporin (Cephamycin).

**Usual Dose:** 2 gm (IV) q6h.

**Spectrum:** (see **Susceptibility Profiles** pp. 191–197).

**Resistance Potential:** Low

**Pharmacokinetic Parameters:**

*Peak serum level:* 221 mcg/mL

*Bioavailability:* Not applicable

*Excreted unchanged (urine):* 85%

*Serum half-life (normal/ESRD):* 1/21 hrs

*Plasma protein binding:* 75%

*Volume of distribution (V<sub>d</sub>):* 0.12 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\***

CrCl 30–50 mL/min	2 gm (IV) q12h
CrCl 10–30 mL/min	1 gm (IV) q12h
CrCl < 10 mL/min	1 gm (IV) q24h
Post-HD dose	2 gm (IV)
Post-PD dose	1 gm (IV)

CW/H/CVHD/CWHDF dose	2 gm (IV) q12h
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	No change

**Drug Interactions:** None.

**Adverse Effects:** Drug fever/rash, *C. difficile* diarrhea/colitis.

**Allergic Potential:** Low

**Safety in Pregnancy:** B

**Comments:** Effective against *B. fragilis*, including *D.O.T.* strains (*B. distasonis*, *B. ovatus*, *B. thetaiotaomicron*).

**Cerebrospinal Fluid Penetration:** < 10%

**Bile Penetration:** 250%

(also see **Antibiotic Pearls & Pitfalls** p. 510).

**REFERENCES:**

- da Costa Darini AL, Palazzo IC. Cefoxitin does not include production of penicillin binding protein 2a in methicillin-susceptible *Staphylococcus aureus* strains. *J Clin Microbiol* 42:4412, 2004.
- Donowitz GR, Mandell GL. Beta-lactam antibiotics. *N Engl J Med* 318:419–26 and 318:490–500, 1993.
- Hansen EA, Cunha BA. Cefoxitin. *Antibiotics for Clinicians* 5:33–41, 2001.
- Marshall WF, Blair JE. The cephalosporins. *Mayo Clin Proc* 74:187–95, 1999.
- Website: [www.pdr.net](http://www.pdr.net)

## Cefpodoxime (Vantin)

**Drug Class:** 3<sup>rd</sup> generation oral cephalosporin.

**Usual Dose:** 200 mg (PO) q12h (see comments).

**Spectrum:** (see **Susceptibility Profiles** pp. 191–197).

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

**Resistance Potential:** Low

**Pharmacokinetic Parameters:**

*Peak serum level:* 2.3 mcg/mL

*Bioavailability:* 50%

*Excreted unchanged (urine):* 30%

*Serum half-life (normal/ESRD):* 2.3/9.8 hrs

*Plasma protein binding:* 21–33%

*Volume of distribution ( $V_d$ ):* 0.9 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\***

CrCl 30–80 mL/min	No change
CrCl < 30 mL/min	200 mg (PO) q24h
Post-HD dose	200 mg (PO)
Post-PD dose	200 mg (PO)
CWVH/CVWHD/ CVWHD dose	200 mg (PO) q12h
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	No change

**Drug Interactions:** None.

**Adverse Effects:** Drug fever/rash, *C. difficile* diarrhea/colitis pulmonary infiltrates with eosinophilia, ↑ SGOT/SGPT.

**Allergic Potential:** High

**Safety in Pregnancy:** B

**Comments:** Only oral 3<sup>rd</sup> generation cephalosporin active against *S. aureus* (MSSA).

**PPNG dose:** 400 mg (PO) × 1 dose.

**cSSSI dose:** 400 mg (PO) q12h.

**Cerebrospinal Fluid Penetration:** < 10%

**Bile Penetration:** 100%

(also see **Antibiotic Pearls & Pitfalls** p. 511).

**REFERENCES:**

- Adam D, Bergogne-Berezin E, Jones RN. Symposium on cefpodoxime proxetil: A new third generation oral cephalosporin. *Drugs* 42:1–66, 1991.
- Cohen R. Clinical efficacy of cefpodoxime in respiratory tract infection. *J Antimicrob Chemother* 50(Suppl):23–7, 2002.
- Liu P, Muller M, Grant M, et al. Interstitial tissue concentrations of cefpodoxime. *J Antimicrob Chemother* 50(Suppl):19–22, 2002.
- Schatz BS, Karavokiros KT, Taeubel MA, et al. Comparison of cefprozil, cefpodoxime proxetil, loracarbef, cefixime, and cefbuten. *Ann Pharmacother* 30:258–68, 1996.
- Website: www.pdr.net

## Cefprozil (Cefzil)

**Drug Class:** 2<sup>nd</sup> generation oral cephalosporin.

**Usual Dose:** 500 mg (PO) q12h.

**Spectrum:** (see **Susceptibility Profiles** pp. 191–197).

**Resistance Potential:** Low

**Pharmacokinetic Parameters:**

*Peak serum level:* 10 mcg/mL

*Bioavailability:* 95%

*Excreted unchanged (urine):* 60%

*Serum half-life (normal/ESRD):* 1.3/5.9 hrs

*Plasma protein binding:* 36%

*Volume of distribution ( $V_d$ ):* 0.23 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\***

CrCl < 30 mL/min	250 mg (PO) q12h
Post-HD dose	500 mg (PO)
Post-PD dose	250 mg (PO)
CWVH/CVWHD/ CVWHD dose	500 mg (PO) q24h

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CWVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

**Drug Interactions:** None.

**Adverse Effects:** Drug fever/rash, *C. difficile* diarrhea/colitis.

**Allergic Potential:** Low

**Safety in Pregnancy:** B

**Comments:** Penetrates oral/respiratory secretions well. No resistance.

**Cerebrospinal Fluid Penetration:** < 10% (also see **Antibiotic Pearls & Pitfalls** p. 510).

#### REFERENCES:

Cunha BA. New antibiotics for the treatment of acute exacerbations of chronic bronchitis. *Adv Ther* 13:313–23, 1996.

Gainer RB 2<sup>nd</sup>. Cefprozil: A new cephalosporin; its use in various clinical trials. *South Med J* 88:338–46, 1995.

Marshall WF Blair JE. The cephalosporins. *Mayo Clin Proc* 74:187–95, 1999.

Schatz BS, Karavokiros KT, Tæubel MA, et al. Comparison of cefprozil, cefpodoxime, proxetil, loracarbef, cefixime, and cefibuten. *Ann Pharmacother* 30:258–68, 1996. Website: www.pdr.net

## Ceftaroline fosamil (Teflaro)

**Drug class:** Anti-MRSA cephalosporin

**Usual dose:** 600 mg (IV) q12h.

**Spectrum:** (see **Susceptibility Profiles** pp. 191–197).

**Resistance Potential:** Low

**Pharmacokinetic Parameters:**

*Peak serum level:* 19 mcg/mL

*Bioavailability:* not applicable

*Excreted unchanged:* 64%

*Serum half-life (normal/ESRD):* 2.7 hours / 6.16 hours

*Plasma protein binding:* 20%

*Volume of distribution (V<sub>d</sub>):* 20.3 L/kg

**Primary Mode of Elimination:** Renal

**Drug Interactions:** None.

**Adverse Reactions:** *C. difficile*, diarrhea, nausea, vomiting, allergic reactions, rash, constipation,

CrCl <sup>a</sup> > 50	No change
CrCl > 30 to ≤ 50	400 mg (IV) (over 1 hour) q12h
CrCl ≥ 15 to ≤ 30	300 mg (IV) (over 1 hour) q12h
ESRD on HD*	200 mg (IV) (over 1 hour) q12h <sup>c</sup>
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	No change

\* should be administered after HD on hemodialysis days.

↑SGOT/↑SGPT hypokalemia, direct Coombs test, phlebitis.

**Safety in Pregnancy:** B

**Comments:** The only cephalosporin effective against MRSA *in vivo*. Not an inducer, inhibitor or substrate of CYP450 system. Infuse over 1 hour.

**Cerebrospinal Fluid Penetration:** < 10% (also see **Antibiotic Pearls & Pitfalls** p. 511).

#### REFERENCES:

Biek D, Critchley IA, Riccobene TA, et al. Ceftaroline fosamil: a novel broad-spectrum cephalosporin with expanded anti-Gram-positive activity. *J Antimicrob Chemother* 65:9–16, 2010.

Casapao AM, Steed ME, Levine DP, et al. Ceftaroline fosamil for community-acquired bacterial pneumonia and acute bacterial skin and skin structure infection. *Expert Opin Pharmacother* 13:1177–1186, 2012.

Corey GR, Wilcox M, Talbot GH, et al. Integrated analysis of CANVAS 1 and 2: phase 3, multicenter, randomized, double-blind studies to evaluate the safety and efficacy of ceftaroline versus vancomycin plus

<sup>a</sup>Usual dose<sup>a</sup> assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

- aztreonam in complicated skin and skin-structure infection. *Clin Infect Dis* 51:641–650; 2010.
- Fille TM Jr, Low DE, Edckburg PB, et al. Efficacy and safety of ceftaroline fosamil versus ceftriaxone in patients with community-acquired pneumonia. *Clin Infect Dis* 51:1395–1405, 2011.
- Ho TT, Cadena J, Childs LM, et al. Methicillin-resistant *Staphylococcus aureus* bacteremia and endocarditis treated with ceftaroline salvage therapy. *J Antimicrob Chemother* 67:1267–1270, 2012.
- Kaushik D, Rathi S, Jain A. Ceftaroline: a comprehensive update. *Int J Antimicrob Agents* 37:389–395, 2011.
- Koeth LM, Matuschek E, Kahlmeter G, et al. Development of EUCAST zone diameter breakpoints and quality control range for *Staphylococcus aureus* with ceftaroline. *Eur J Clin Microbiol Infect Dis* 33:1511–1517, 2014.
- Laudano JB. Ceftaroline fosamil: a new broad-spectrum cephalosporin. *J Antimicrob Chemother* 66:11–18, 2011.
- Lim L, Sutton E, Brown J. Ceftaroline: A new broad-spectrum cephalosporin. *Am J Health Syst Pharm* 15:491–4198, 2011.
- Mendes RE, Tsakris A, Sader HS, et al. Characterization of methicillin-resistant *Staphylococcus aureus* displaying increased MICs of ceftaroline. *J Antimicrob Chemother* 67:1231–1234, 2012.
- Nannini Ec, Stryjewskin ME, Corey GR. Ceftaroline for complicated skin and skin-structure infections. *Expert Opin Pharmacother* 11:1197–1206, 2010.
- Panagiotidis G, Backstrom T, Asker-Hagelberg C, et al. Effect of ceftaroline on normal human intestinal microflora. *Antimicrob Agents Chemother* 54:1811–1814, 2010.
- Poon H, Chang MH, Fung HB. Ceftaroline fosamil: a cephalosporin with activity against methicillin-resistant *Staphylococcus aureus*. *Clin Ther* 34:743–765, 2012.
- Saravolatz LD, Pawlak J, Johnson L. In vitro activity of ceftaroline against community-associated methicillin-resistant, vancomycin-intermediate, vancomycin-resistant, and daptomycin-nonsusceptible *Staphylococcus aureus* isolates. *Antimicrob. Agents Chemother* 54:3027–3030, 2010.
- Saravolatz LD, Stein GE, Johnson LB. Ceftaroline: A novel cephalosporin with activity against methicillin-resistant *Staphylococcus aureus*. *Clin Infect Dis* 52:1156–1163, 2011.
- Watkins RR. Successful treatment of osetomyelitis from methicillin-resistant *Staphylococcus aureus* with ceftaroline. *Infectious Disease in Clinical Practice* 21:1:48–49, 2013.
- Zainah H, Vazquez JA. Use of ceftaroline in complicated methicillin-resistant *Staphylococcus aureus* bacteremia. *Infectious Disease in Clinical Practice* 21:1:58–60, 2013.
- Website: www.pdr.net

## Ceftazidime (Fortaz, Tazicef, Tazidime)

**Drug Class:** 3<sup>rd</sup> generation cephalosporin.

**Usual Dose:** 2 gm (IV) q8h (see comments).

**Spectrum:** (see **Susceptibility Profiles** pp. 191–197).

**Resistance Potential:** High (*P. aeruginosa*, *Klebsiella pneumoniae*; ↑ prevalence of MRSA)

### Pharmacokinetic Parameters:

*Peak serum level:* 170 mcg/mL

*Bioavailability:* Not applicable

*Excreted unchanged (urine):* 80–90%

*Serum half-life (normal/ESRD):* 1.9/21 hrs

*Plasma protein binding:* 10%

*Volume of distribution ( $V_d$ ):* 0.36 L/kg

**Primary Mode of Elimination:** Renal

### Dosage Adjustments\*

CrCl 30–80 mL/min	2 gm (IV) q12h
CrCl 10–30 mL/min	2 gm (IV) q24h
CrCl < 10 mL/min	1 gm (IV) q48h
Post–HD dose	1 gm (IV)
Post–PD dose	500 mg (IV)
CVWH/CVWH/D/ CVHDF dose	2 gm (IV) q12h
HD q48–72h dose	2 gm (IV) q HD*
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	No change

\*Give at end of HD.

**Drug Interactions:** None.

**Adverse Effects:** Phlebitis, drug fever/rash, *C. difficile* diarrhea/colitis.

**Allergic Potential:** High

**Safety in Pregnancy:** B

**Comments:** Incompatible in solutions containing vancomycin or aminoglycosides.

**Use increases MRSA prevalence and *P. aeruginosa* resistance. Potent inducer of *Klebsiella/Enterobacter/E. coli* ESBLs.**

**CAPD dose:** 125 mg/L of dialysate (I.P.) with each exchange.

Meningeal dose = 2 gm (IV) q8h.

**Cerebrospinal Fluid Penetration:**

Non-Inflamed meninges = 1%

Inflamed meninges = 20%

**Bile Penetration:** 50%

(also see **Antibiotic Pearls & Pitfalls** p. 510).

#### REFERENCES:

- Briscoe-Dwyer L. Ceftazidime. *Antibiotics for Clinicians* 1:41–8, 1997.
- Fortaleza CM, Freire MP, de Carvalho Ramos M. Risk factors for recovery of imipenem- or ceftazidime-resistant *Pseudomonas aeruginosa* among patients admitted to a teaching hospital in Brazil. *Infection Control and Hosp Epidemiol* 27:901–06, 2006.
- Klein NC, Cunha BA. Third-generation cephalosporins. *Med Clin North Am* 79:705–19, 1995.
- Lu Q, Yang J, Liu Z, et al. Nebulized ceftazidime and amikacin in ventilator-associated pneumonia caused by *Pseudomonas aeruginosa*. *Am J Respir Crit Care Med*. 184:106–115, 2011.
- Nicolau DP, Nightingale CH, Banevicius MA, et al. Serum bactericidal activity of ceftazidime: Continuous infusion versus intermittent injections. *Antimicrob Agents Chemother* 40:61–4, 1996.
- Owens JC, Jr, Ambrose PG, Quintiliani R. Ceftazidime to cefepime formulary switch: Pharmacodynamic rationale. *Conn Med* 61:225–7, 1997.
- Rains CP, Bryson HM, Peters DH. Ceftazidime: An update of its antibacterial activity, pharmacokinetic properties, and therapeutic efficacy. *Drugs* 49:577–617, 1995.
- Website: [www.pdr.net](http://www.pdr.net)

## Ceftazidime/Avibactam (Avycaz)

**Drug Class:** Antipseudomonal cephalosporin/ $\beta$ -lactamase inhibitor

**Usual Dose:** 2.5 gm (IV) q8h

**Spectrum:** (see **Susceptibility Profiles** pp 191-197)

**Resistance Potential:** Low

**Pharmacokinetic Parameters:**

*Peak Serum level:* 170 mcg/ml/15 mcg/ml

*Bioavailability:* Not applicable

*Excreted unchanged:* 80–90%/85%

*Serum half-life (normal/ESRD):* [1.9h/21 h]/ [2.0h/22.82 h]

*Plasma protein binding:* 10%/8.2%

*Volume of distribution ( $V_d$ ):* 0.36 L/kg/0.32 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\***

CrCl 31–50 mL/min	1.25 gm (IV) q8h
CrCl 16–30 mL/min	0.94 gm (IV) q12h
CrCl 6–15 mL/min	0.94 gm (IV) q24h
CrCl < 5 mL/min	0.94 gm (IV) q48h
Post-HD dose	0.94 gm (IV) q48h
Post-PD dose	No data
CVVH dose	No data
Hepatic insufficiency	No change

**Drug Interactions:** None

**Adverse Effects:** Headache, dizziness, insomnia, constipation, diarrhea, abdominal pain, nausea, vomiting,  $\uparrow$  SGOT/SGPT, drug fever/rash, phlebitis, hypersensitivity reactions, *C. difficile* diarrhea/colitis.

**Allergic Potential:**

**Safety in Pregnancy:** B



**Comments:** Avibactam is a novel, non- $\beta$ -lactam,  $\beta$ -lactamase inhibitor, which acts a reversible inhibitor (compared to irreversible action of other  $\beta$ -lactamase inhibitors) and has an expanded spectrum of  $\beta$ -lactamase inhibition including: class A (includes *Klebsiella pneumoniae* carbapenemase-KPC), class C, some class D. Avibactam's spectrum does not include class B  $\beta$ -lactamase (metalloenzymes). Use of avibactam with ceftazidime restores susceptibility against ceftazidime-nonsusceptible organisms. Active against multi-drug resistant Gram negative organisms, including *P. aeruginosa*, ESBL producing pathogens, and CRE. Infuse slowly over 2 hours.

For complicated intra-abdominal infections (cIAI), ceftazidime/avibactam must be used in combination with metronidazole.

**Cerebrospinal Fluid Penetration:** < 10% (also see **Antibiotic Pearls & Pitfalls** p. 519).

#### REFERENCES:

- Ehmann DE, Jahic H, Ross PL, et al. Kinetics of avibactam inhibition against Class A, C, and D beta-lactamases. *J Biol Chem.* 288:27960-27971, 2013.
- Keepers TR, Gomez M, Celeri C, et al. Bactericidal activity, absence of serum effect, and time-kill kinetics of ceftazidime-avibactam against beta-lactamase-producing Enterobacteriaceae and *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother.* 58:5297-5305, 2014.
- Lahiri SD, Johnstone MR, Ross PL, et al. Avibactam and class C beta-lactamases: mechanism of inhibition, conservation of the binding pocket, and implications for resistance. *Antimicrob Agents Chemother.* 58:5704-5713, 2014.
- Lahiri SD, Mangani S, Durand-Reville T, et al. Structural insight into potent broad-spectrum inhibition with reversible recyclization mechanism: avibactam in complex with CTX-M-15 and *Pseudomonas aeruginosa* AmpC beta-lactamases. *Antimicrob Agents Chemother.* 57:2496-2505, 2013.
- Lucasti C, Popescu I, Ramesh MK, et al. Comparative study of the efficacy and safety of ceftazidime/avibactam plus metronidazole versus meropenem in the treatment of complicated intra-abdominal infections in hospitalized adults: results of a randomized, double-

blind, Phase II trial. *J Antimicrob Chemother.* 68:1183-1192, 2013.

Mushtaq S, Warner M, Livermore DM. In vitro activity of ceftazidime+NXL 104 against *Pseudomonas aeruginosa* and other non-fermenters. *J Antimicrob Chemother.* 65:2376-2381, 2010.

Vazquez JA, Gonzalez Patzan LD, Stricklin D, et al. Efficacy and safety of ceftazidime-avibactam versus imipenem/cilastatin in the treatment of complicated urinary tract infections, including acute pyelonephritis, in hospitalized adults: results of a prospective, investigator-blinded, randomized study. *Curr Med Res Opin.* 28:1921-1931, 2012.

Zhanell GG, Lawson CD, Adam H, et al. Ceftazidime-avibactam: a novel cephalosporin/beta-lactamase inhibitor combination. *Drugs.* 73:159-177, 2013.

Website: www.pdr.net

## Ceftibuten (Cedax)

**Drug Class:** 3<sup>rd</sup> generation oral cephalosporin.

**Usual Dose:** 400 mg (PO) q24h.

**Spectrum:** (see **Susceptibility Profiles** pp. 191–197).

**Resistance Potential:** Low

**Pharmacokinetic Parameters:**

*Peak serum level:* 18 mcg/mL

*Bioavailability:* 80%

*Excreted unchanged:* 56% (urine); 39% (fecal) *Serum half-life (normal/ESRD):* 2.4/22 hrs

*Plasma protein binding:* 65%

*Volume of distribution ( $V_d$ ):* 0.2 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\***

CrCl 30–50 mL/min	200 mg (PO) q24h
CrCl < 30 mL/min	100 mg (PO) q24h
Post-HD dose	400 mg (PO)
Post-PD dose	200 mg (PO)
CWH/CWHD/ CWHDF dose	200 mg (PO) q24h

Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	No change

**Drug Interactions:** None.

**Adverse Effects:** Drug fever/rash.

**Allergic Potential:** High

**Safety in Pregnancy:** B

**Comments:** Little/no activity against *S. aureus*/*S. pneumoniae*.

**Cerebrospinal Fluid Penetration:** < 10% (also see **Antibiotic Pearls & Pitfalls** p. 511).

#### REFERENCES:

Andrews JW, Wise R, Baldwin DR, et al. Concentrations of cefitibuten in plasma and the respiratory tract following a single 400 mg oral dose. *Int J Antimicrob Agents* 5:141–144, 1995.

Guay DR. Cefitibuten: A new expanded-spectrum oral cephalosporin. *Ann Pharmacother* 31:1022–33, 1997.

Owens RC Jr, Nightingale CH, Nicolau DP. Cefitibuten: An overview. *Pharmacother* 17:707–20, 1997.

Wiseman LR, Balfour JA. Cefitibuten: A review of its antibacterial activity, pharmacokinetic properties and clinical efficacy. *Drugs* 47:784–808, 1994.

Website: www.pdr.net

## Ceftizoxime (Cefizox)

**Drug Class:** 3<sup>rd</sup> generation cephalosporin.

Usual Dose: 2 gm (IV) q8h (see comments).

**Spectrum:** (see **Susceptibility Profiles** pp. 191–197).

**Resistance Potential:** Low

Pharmacokinetic Parameters:

*Peak serum level:* 132 mcg/mL

*Bioavailability:* Not applicable

*Excreted unchanged (urine):* 90%

*Serum half-life (normal/ESRD):* 1.7/35 hrs

*Plasma protein binding:* 30%

*Volume of distribution ( $V_d$ ):* 0.32 L/kg

**Primary Mode of Elimination:** Renal  
Dosage Adjustments\*

CrCl 50–80 mL/min	1 gm (IV) q8h
CrCl 10–50 mL/min	1 gm (IV) q12h
CrCl < 10 mL/min	1 gm (IV) q24h
Post-HD dose	1 gm (IV)
Post-PD dose	1 gm (IV)
CVWH/CVWHD/ CVWHD dose	1 gm (IV) q12h
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	No change

**Drug Interactions:** None.

**Adverse Effects:** Drug fever/rash, *C. difficile* diarrhea/colitis.

Allergic Potential: High

Safety in Pregnancy: B

**Comments:** PRNG dose: 500 mg (IM) × 1 dose.

Meningeal dose = 3 gm (IV) q6h.

**Cerebrospinal Fluid Penetration:**

Non-Inflamed meninges = 1%

Inflamed meninges = 10%

**Bile Penetration:** 50%

(also see **Antibiotic Pearls & Pitfalls** p. 510).

#### REFERENCES:

Klein NC, Cunha BA. Third-generation cephalosporins. *Med Clin North Am* 79:705–19, 1995.

Donowitz GR, Mandell GL. Beta-lactam antibiotics. *N Engl J Med* 318:419–26 and 318:490–500, 1993.

Marshall WF, Blair JE. The cephalosporins. *Mayo Clin Proc* 74:187–95, 1999.

Website: www.pdr.net

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

## Ceftolozane/Tazobactam (Zerbaxa)

**Drug Class:** Anti-pseudomonal cephalosporin/ $\beta$ -lactamase inhibitor.

Usual Dose: 1.5 gm (IV) q8h

**Spectrum:** (see *Susceptibility Profiles* pp. 194–197).

**Resistance Potential:** Low

**Pharmacokinetic Parameters:**

Peak serum level: 74.4/34 mcg/mL

Bioavailability: Not applicable

Excreted unchanged (urine): 95/80%

Serum half-life (normal/ESRD):

Plasma protein binding: 16-21%/30%

Volume of distribution ( $V_d$ ): 0.21 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments**

CrCl 30–50 mL/min	750 mg (IV) q8h
CrCl 15–29 mL/min	375 mg (IV) q8h
HD dose	LD of 750 mg (IV) $\times$ 1 dose, 150 mg (IV) q8h (on HD days, administer immediately following HD)
PD dose	No data
CVWH/CVVD/ CVHDF dose	No data
Moderate- severe hepatic insufficiency	No change

**Drug Interactions:** No significant interactions are anticipated between ceftolozane/tazobactam and substrates, inhibitors and inducers of cytochrome P450 enzymes

**Adverse Effects:** Drug fever/rash, nausea, diarrhea, headache, insomnia, constipation, *C. difficile* diarrhea/colitis, mild transient  $\uparrow$  SGOT/SGPT, hypotension,  $\uparrow$  platelets (thrombocytosis), anemia.

**Allergic Potential:**

**Safety in Pregnancy:** Category B

**Comments:** Useful for MDR Gram Negative pathogens, including *P. aeruginosa* and ESBL producing pathogens. Ineffective against CRE. For treating cIAI, must add Metronidazole 500 mg (IV) q8h.

(also see **Antibiotic Pearls & Pitfalls** p. 519).

### REFERENCES:

- Lucasti C, Hershberger E, Miller B, et al. Multicenter, double-blind, randomized, phase II trial to assess the safety and efficacy of ceftolozane/tazobactam plus metronidazole compared with meropenem in adult patients with complicated intra abdominal infections. *Antimicrob Agents and Chemotherapy*. 58:5350-5357, 2014.
- Miller B, Hershberger E, Benziger D, et al. Pharmacokinetics and safety of intravenous ceftolozane/tazobactam in healthy adult subjects following single and multiple ascending doses. *Antimicrob Agents Chemo* 56:3086-3091, 2012.
- Sorbera M, Chung E, Ho CW, et al. Ceftolozane/tazobactam: a new option in the treatment of complicated gram-negative infections. *P & T* 39:825-832, 2014.
- Zhanell GG, Chung P, Adam H, et al. Ceftolozane/tazobactam: a novel cephalosporin/beta-lactamase inhibitor combination with activity against multidrug-resistant gram-negative bacilli. *Drugs* 74:31-51, 2014.
- Website: [www.pdr.net](http://www.pdr.net)

## Ceftriaxone (Rocephin)

**Drug Class:** 3<sup>rd</sup> generation cephalosporin.

**Usual Dose:** 1–2 gm (IV) q24h (see comments).

"Usual dose" assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose  $\times$  1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

**Spectrum:** (see **Susceptibility Profiles**

pp. 191–197).

**Resistance Potential:** Low**Pharmacokinetic Parameters:***Peak serum level:* 151–257 mcg/mL*Bioavailability:* Not applicable*Excreted unchanged (urine/feces):* 33–67%*Serum half-life (normal/ESRD):* 8/16 hrs*Plasma protein binding:* 90%*Volume of distribution ( $V_d$ ):* 0.08–0.3 L/kg**Primary Mode of Elimination:** Renal/hepatic**Dosage Adjustments\***

CrCl < 10 mL/min	No change
Post-HD dose	None
Post-HFHD dose	None
Post-PD dose	None
CVVH/CVVHD/ CVVHDF dose	No change
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	No change (max dose: 2 gm/d)

**Drug Interactions:** May cause intravascular precipitation when co-administered with calcium-containing solutions/products.**Adverse Effects:** Drug fever/rash, non-C. difficile diarrhea, pseudobiliary lithiasis, hemolytic anemia, may interfere with platelet aggregation, thrombocytosis.**Allergic Potential:** High**Safety in Pregnancy:** B. **Avoid near term in 3<sup>rd</sup> trimester** (↑ incidence of kernicterus in newborns)**Cerebrospinal Fluid Penetration:**

Non-Inflamed meninges = 1%

Inflamed meninges = 10%

**Bile Penetration:** 500%(also see **Antibiotic Pearls & Pitfalls** p. 510).**REFERENCES:**

- Araz N, Okan V, Demirci M, et al. Pseudolithiasis due to ceftriaxone treatment for meningitis in children: report of 8 cases. *Tohoku J Exp Med* 211:285–290, 2007.
- Borg R, Dotevall L, Hagberg L, et al. Intravenous ceftriaxone compared with oral doxycycline for the treatment of Lyme neuroborreliosis. *Scand J Infect Dis* 37:449–454, 2005.
- Cunha BA, Klein NC. The selection and use of cephalosporins: A review. *Adv Ther* 12:83–101, 1995.
- Dietrich ES, Bieser U, Frank U, et al. Ceftriaxone versus other cephalosporins for perioperative antibiotic prophylaxis: a meta analysis of 43 randomized controlled trials. *Chemotherapy* 48:49–56, 2002.
- Faella F, Pagliano P, Fusco U, Attanasio V, Conte M. Combined treatment with ceftriaxone and linezolid of pneumococcal meningitis: a case series including penicillin-resistant strains. *Clin Microbiol Infect* 12:391–4, 2006.
- Feurle GE, Junga NS, Marth T. Efficacy of ceftriaxone or meropenem as initial therapies in Whipple's disease. *Gastroenterology* 138:478–486, 2010.
- File TM Jr. Clinical implications and treatment of multiresistant *Streptococcus pneumoniae* pneumonia. *Clin Microbiol Infect* 12:31–41, 2006.
- File TM Jr, Low DE, Edckburg PB, et al. Efficacy and safety of ceftaroline fosamil versus ceftriaxone in patients with community-acquired pneumonia. *Clin Infect Dis* 51:1395–1405, 2011.
- Freeman CD, Nightingale CH, Nicolau DP, et al. Serum bactericidal activity of ceftriaxone plus metronidazole against common intra-abdominal pathogens. *Am J Hosp Pharm* 51:1782–7, 1994.
- Grassi C. Ceftriaxone. *Antibiotics for Clinicians* 2:49–57, 1998.
- Klein NC, Cunha BA. Third generation cephalosporins. *Med Clin North Am* 79:705–19, 1995.
- Lamb HM, Ormrod D, Scott LJ, et al. Ceftriaxone: an update of its use in the management of community-acquired and nosocomial infections. *Drugs* 62:1041–89, 2002.

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVVH = continuous veno-veno hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

Marshall WF, Blair JE. The cephalosporins. *Mayo Clin Proc* 74:187–95, 1999.

Pichichero ME. Use of selected cephalosporins in penicillin-allergic patients: a paradigm shift. *Diagn Microbiol Infect Dis* 57:135–185, 2007.

Roberts JA, Boots R, Rickard CM, et al. Is continuous infusion ceftriaxone better than once-a-day dosing in intensive care? A randomized controlled pilot study. *J Antimicrob Chemother* 59:285–291, 2007.

Savaris RF, Teixeira LM, Torres TG, et al. Comparing ceftriaxone plus azithromycin or doxycycline for pelvic inflammatory disease: a randomized controlled trial. *Obstet Gynecol* 110:53–60, 2007.

Shelburne SA 3<sup>rd</sup>, Greenburg SB, Aslam S, et al. Successful ceftriaxone therapy of endocarditis due to penicillin non-susceptible viridans streptococci. *J Infect* 54:e108–e110, 2007.

Woodfield JC, Van Rij AM, Pettigrew RA, et al. A comparison of the prophylactic efficacy of ceftriaxone and cefotaxime in abdominal surgery. *Am J Surg* 185:45–9, 2003.

Website: [www.pdr.net](http://www.pdr.net)

## Cefuroxime (Kefurox, Zinacef, Ceftin)

**Drug Class:** 2<sup>nd</sup> generation IV/oral cephalosporin.

**Usual Dose:** 1.5 gm (IV) q8h; 500 mg (PO) q12h (see comments).

**Spectrum:** (see **Susceptibility Profiles** pp. 191–197).

**Resistance Potential:** Low

**Pharmacokinetic Parameters:**

*Peak serum level:* 100 (IV)/7 (PO) mcg/mL

*Bioavailability:* 52%

*Excreted unchanged (urine):* 89%

*Serum half-life (normal/ESRD):* 1.2/17 hrs

*Plasma protein binding:* 50%

*Volume of distribution ( $V_d$ ):* 0.15 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\***

CrCl 20–80 mL/min	No change
CrCl 10–20 mL/min	750 mg (IV) q12h 500 mg (PO) q12h
CrCl < 10 mL/min	750 mg (IV) q24h 500 mg (PO) q24h
Post–HD dose	750 mg (IV) 250 mg (PO)
Post–PD dose	750 mg (IV) 250 mg (PO)
CVVH/CVVHD/ CVVHDF dose	1 gm (IV) q12h 500 mg (PO) q12h
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	No change

**Drug Interactions:** None.

**Adverse Effects:** Drug fever/rash, *C. difficile* diarrhea/colitis.

**Allergic Potential:** High

**Safety in Pregnancy:** B

**Comments:** Oral preparation penetrates oral/respiratory secretions well. **PPNG dose:** 1 gm (IM) × 1 dose. **Ineffective for *H. influenzae* meningitis prophylaxis or therapy.**

**Cerebrospinal Fluid Penetration:** < 10% (also see **Antibiotic Pearls & Pitfalls** p. 510).

### REFERENCES:

Alvarez-Sala JL, Kardos P, Martinez-Beltran, et al.

Clinical and bacteriological efficacy in treatment of acute exacerbations of chronic bronchitis with cefditoren-pivoxil versus cefuroxime-axetil. *Antimicrob Agents and Chemother* 50:1762–67, 2006.

Bucko AD, Hunt BJ, Kidd SL, et al. Randomized, double-blind, multicenter comparison of oral cefditoren

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

200 or 400 mg BID with either cefuroxime 250 mg BID or cefadroxil 500 mg BID for the treatment of uncomplicated skin and skin-structure infections. Clin Ther 24:1134–47, 2002.

Gentry LO, Zeluff BJ, Cooley DA. Antibiotic prophylaxis in open-heart surgery: A comparison of cefamandole, cefuroxime, and cefazolin. Ann Thorac Surg 46:167–71, 1988.

Marshall WF, Blair JE. The cephalosporins. Mayo Clin Proc 74:187–95, 1999.

Perry Cm, Brogden RN. Cefuroxime axetil. A review of its antibacterial activity, pharmacokinetic properties, and therapeutic efficacy. Drugs 52:125–58, 1996.

Website: www.pdr.net

## Cephalexin (Keflex)

**Drug Class:** 1<sup>st</sup> generation oral cephalosporin.

**Usual Dose:** 500 mg (PO) q6h.

**Spectrum:** (see *Susceptibility Profiles* pp. 191–197).

**Resistance Potential:** Low

**Pharmacokinetic Parameters:**

*Peak serum level:* 18 mcg/mL

*Bioavailability:* 99%

*Excreted unchanged (urine):* > 90%

*Serum half-life (normal/ESRD):* 0.7/16 hrs

*Plasma protein binding:* 10%

*Volume of distribution ( $V_d$ ):* 0.35 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\***

CrCl 10–50 mL/min	500 mg (PO) q12h
CrCl < 10 mL/min	500 mg (PO) q24h
Post-HD dose	500 mg (PO)
Post-PD dose	250 mg (PO)

CVVH/CVVHD/ CVVHDF dose	250 mg (PO) q8h
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	No change

**Drug Interactions:** None.

**Adverse Effects:** Drug fever/rash, *C. difficile* diarrhea/colitis.

**Allergic Potential:** High

**Safety in Pregnancy:** B

**Comments:** Highly active against *S. aureus* (MSSA) and Group A streptococci. **Limited activity against *H. influenzae*.**

**Cerebrospinal Fluid Penetration:** < 10%

**Bile Penetration:** 200%

(also see *Antibiotic Pearls & Pitfalls* p. 509).

### REFERENCES:

Chow M, Quintilliani R, Cunha BA, et al. Pharmacokinetics of high dose oral cephalosporins. J Pharmacol 19:185–194, 1979.

Cunha BA. Cephalexin remains the preferred antibiotic for uncomplicated cellulitis: revisited. J Chemo. 26:65–66, 2014.

Cunha BA. Cephalexin remains preferred oral antibiotic therapy for uncomplicated cellulitis. Am J Med 121:e11–15, 2008.

Cunha BA. Oral antibiotic therapy of serious systemic infections. Med Clin N Am 90:1197–1222, 2006.

Marshall WF Blair JE. The cephalosporins. Mayo Clin Proc 74:187–95, 1999.

Smith GH. Oral cephalosporins in perspective. DICP 24:45–51, 1990.

Website: www.pdr.net

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

## Chloramphenicol (Chloromycetin)

**Drug Class:** No specific class.

**Usual Dose:** 500 mg (IV/PO) q6h.

**Spectrum:** (see *Susceptibility Profiles* pp. 186–190).

**Resistance Potential:** Low

**Pharmacokinetic Parameters:**

Peak serum level: 9 mcg/mL

Bioavailability: 90%

Excreted unchanged (urine): 10%

Serum half-life (normal/ESRD): 2.5/3 hrs

Plasma protein binding: 50%

Volume of distribution ( $V_d$ ): 1 L/kg

**Primary Mode of Elimination:** Hepatic

**Dosage Adjustments\***

CrCl < 50 mL/min	No change
Post-HD dose	500 mg (IV/PO)
Post-PD dose	None
CVWH/CVWHD/ CVWHDf dose	No change
Moderate—severe hepatic insufficiency	No change

**Drug Interactions:** Barbiturates (↑ barbiturate effect, ↓ chloramphenicol effect); cyclophosphamide (↑ cyclophosphamide toxicity); cyanocobalamin, iron (↓ response to interacting drug); warfarin (↑ INR); phenytoin (↑ phenytoin toxicity); rifabutin, rifampin (↓ chloramphenicol levels); sulfonyleureas (↑ sulfonyleurea effect, hypoglycemia).

**Adverse Effects:**

Dose-related bone marrow suppression  
Reversible. Bone marrow aspirate shows vacuolated WBCs (“chloramphenicol effect,” not toxicity). Does not precede aplastic anemia

Idiosyncratic bone marrow toxicity

Irreversible aplastic anemia. May occur after only one dose; monitoring with serial CBCs is useless. Very rare. Usually associated with IM, intraocular, or oral administration. Rarely, if ever, with IV chloramphenicol.

**Allergic Potential:** Low

**Safety in Pregnancy:** A

**Comments:** Incompatible in solutions containing diphenylhydantoin, methylprednisone, aminophylline, ampicillin, gentamicin, erythromycin, vancomycin. **Dose-related marrow suppression is common, and reversible.** Hepatic toxicity related to prolonged high doses (> 4 gm/d). **Oral administration results in higher serum levels than IV administration.** Do not administer IM. Chloramphenicol is inactivated in bile. **Urinary concentrations are therapeutically ineffective.**

Meningeal dose = usual dose.

**Cerebrospinal Fluid Penetration:**

Non-Inflamed meninges = 90%

Inflamed meninges = 90%

**Bile Penetration:** 0%

(also see *Antibiotic Pearls & Pitfalls* p. 513).

### REFERENCES:

- Ariyal A, Basnyal B, Koirala S, et al. Gatifloxacin versus chloramphenicol for uncomplicated enteric fever: an open-label, randomized, controlled trial. *Lancet Infect Dis* 11:445–454, 2011.
- Cunha BA. New uses of older antibiotics. *Postgrad Med* 100:68–88, 1997.
- Cunha BA. Oral antibiotic therapy of serious systemic infections. *Med Clin N Am* 90:1197–1222, 2006.
- Feder HM Jr, Osier C, Maderazo EG. Chloramphenicol: A review of its use in clinical practice. *Rev Infect Dis* 3:479–91, 1981.

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

Kasten MJ. Clindamycin, metronidazole, and chloramphenicol. *Mayo Clin Proc* 74:825–33, 1999.

Safdar A, Bryan CS, Stinson S, et al. Prosthetic valve endocarditis due to vancomycin-resistant *Enterococcus faecium*: treatment with chloramphenicol plus minocycline. *Clin Infect Dis* 34:61–3, 2002.

Smilack JD, Wilson WE, Cockeril FR 3<sup>rd</sup>. Tetracycline, chloramphenicol, erythromycin, clindamycin, and metronidazole. *Mayo Clin Proc* 66:1270–80, 1991.

Wareham DW, Wilson P. Chloramphenicol in the 21<sup>st</sup> century. *Hosp Med* 63:157–61, 2002.

Website: [www.pdr.net](http://www.pdr.net)

## Cidofovir (Vistide)

**Drug class:** HSV, CMV (see comments).

**Usual dose:** 5 mg/kg (IV) weekly for 2 weeks, then 5 mg/kg every 2 weeks (for CMV retinitis in HIV patients, patients must have all of the following: SrCr < 1.5 mg/dL, CrCl > 55 mL/min, urine protein < 100 mg/dL).

**Pharmacokinetic Parameters:**

*Peak serum level:* 19.6 mcg/mL

*Bioavailability:* not applicable

*Excreted unchanged:* 1%

*Serum half-life (normal/ESRD):* 2.2 h/no data

*Plasma protein binding:* 6%

*Volume of distribution (V<sub>d</sub>):* 0.54 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\***

Cr increase of 0.3–0.4 mg/dL above baseline	3 mg/kg (IV) (see usual dose for duration)
Cr increase of ≥ 0.5 mg/dL or + 3 proteinuria	Discontinue cidofovir
Moderate—severe hepatic insufficiency	No change

**Drug Interactions:** Amphotericin B, aminoglycosides, foscarnet, pentamidine, tenofovir, zidovudine (↑ nephrotoxicity).

**Adverse Effects:** Nephrotoxicity; neutropenia; nausea, vomiting associated with probenecid, take with food; metabolic acidosis; decreased intraocular pressure; anterior uveitis and iritis.

**Contraindications:** if CrCl < 55 mL/min.

**Allergic Potential:** Low for cidofovir, high for probenecid.

**Safety in Pregnancy:** C

**Comments:** All patients should receive probenecid (2 g (PO) 3 hours before infusion start, then 1 g at 2 and again at 8 hours after completion of cidofovir infusion, take with food to minimize nausea and vomiting) and hydration with at least 1 L of normal saline with each dose of cidofovir; infuse over 1 hour; monitor neutrophils, growth factor support may be needed; contraindicated in severe hypersensitivity to probenecid or other sulfa drugs; direct ocular injection is contraindicated; effective contraception required for male and female patients. **Highly active against CMV and HSV. Some activity against BK virus, JC virus, VZV, and adenoviruses. No activity against HHV-6, EBV, RSV, or influenza viruses.**

**Cerebrospinal Fluid Penetration:** < 1%

**REFERENCES:**

Ganguly N, Clough LA, Dubois LK, et al. Low-dose cidofovir in the treatment of symptomatic BK virus infection in patients undergoing allogeneic hematopoietic stem cell transplantation: a retrospective analysis of an algorithmic approach. *Transpl Infect Dis* 12:406–411, 2010.

Hua DK, Howard K, Craig JC, et al. Cost-effectiveness of cidofovir treatment of polyomavirus nephropathy in kidney transplant recipients. *Transplantation* 27:188–94, 2012.

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.



Lurain NS, Chou S. Antiviral drug resistance of human cytomegalovirus, *Clin Microbiol Rev* 23:689–712, 2010.  
 Pallet N, Burgard M, Quamouss O, et al. Cidofovir may be deleterious in BK virus-associated nephropathy. *Transplantation* 89:1542–1544, 2010.  
 Website: www.pdr.net

## Ciprofloxacin (Cipro)

**Drug Class:** Fluoroquinolone.

**Usual Dose:** 400 mg (IV) q12h; 500–750 mg (PO) q12h (see comments).

**Spectrum:** (see **Susceptibility Profiles** pp. 198–202).

**Resistance Potential:** High (*S. pneumoniae*, *E. coli*, *P. aeruginosa*; ↑ prevalence of MRSA)

**Pharmacokinetic Parameters:**

*Peak serum level:* 4.6 (IV)/2.9 (PO) mcg/mL

*Bioavailability:* 70%

*Excreted unchanged (urine):* 70%

*Serum half-life (normal/ESRD):* 4/8 hrs

*Plasma protein binding:* 20–40%

*Volume of distribution ( $V_d$ ):* 2.5 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\***

CrCl 30–50 mL/min	400 mg (IV) q12h 500 mg (PO) q12h
CrCl < 30 mL/min	400 mg (IV) q24h 500 mg (PO) q24h
Post-HD dose	200–400 mg (IV) 250–500 mg (PO)
Post-PD dose	200–400 mg (IV) 250–500 mg (PO)
CVWH/CVHD/CVHDF dose	400 mg (IV) q12h 250 mg (PO) q12h
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	No change

**Drug Interactions:** Al<sup>+</sup>, Ca<sup>+</sup>, Fe<sup>+</sup>, Mg<sup>+</sup>, Zn<sup>+</sup> antacids, citrate/citric acid, dairy products (↓ absorption of ciprofloxacin only if taken together); caffeine, cyclosporine, theophylline (↑ interacting drug levels); cimetidine (↑ ciprofloxacin levels); foscarnet (↑ risk of seizures); oral hypoglycemics (slight ↑ or ↓ in blood glucose); NSAIDs (may ↑ risk of seizures/CNS stimulation); phenytoin (↑ or ↓ phenytoin levels); probenecid (↑ ciprofloxacin levels); warfarin (↑ INR).

**Adverse Effects:** Drug fever/rash, *C. difficile* diarrhea/colitis, headache, dizziness, insomnia, malaise, delirium, encephalopathy, seizures. FQ use has an increased risk of tendinitis/tendon rupture. Risk in highest in the elderly, those on steroids, and those with heart, lung or renal transplants. Peripheral neuropathy may occur early and may be permanent. May ↑ QT<sub>c</sub>. Vitamins C and E (↓ ciprofloxacin levels).

**Allergic Potential:** Low

**Safety in Pregnancy:** C

**Comments:** Enteral feeding decreases ciprofloxacin absorption ≥ 30%. Use with caution in patients with severe renal insufficiency or seizure disorder. Administer 2 hours before or after H<sub>2</sub> antagonists, omeprazole, sucralfate, calcium, iron, zinc, multivitamins, or aluminum/magnesium containing medications. Administer ciprofloxacin (IV) as an intravenous infusion over 1 hour.

**Nosocomial pneumonia dose:**

400 mg (IV) q8h.

↑ **incidence of *C. difficile* diarrhea/colitis with PPIs (for patients on PPIs during FQ therapy, D/C PPIs or switch to H<sub>2</sub> blocker for duration of FQ therapy).**

**Cerebrospinal Fluid Penetration:**

Non-Inflamed meninges = 10%

Inflamed meninges = 26%

**Bile Penetration:** 3000%

(also see **Antibiotic Pearls & Pitfalls** p. 516).

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

## REFERENCES:

- Chou HW, Wang JL, Chang CH, et al. Risk of severe dysglycemia among diabetic patients receiving levofloxacin, ciprofloxacin, or moxifloxacin in Taiwan. *Clin Infect Dis* 57:971–980, 2013.
- Didier JP, Villet R, Huggler E, et al. Impact of ciprofloxacin exposure on *Staphylococcus aureus* genomic alterations linked with emergence of rifampin resistance. *Antimicrob Agents Chemother* 55:1946–1952, 2011.
- Fantin B, Duval X, Massias L, et al. Ciprofloxacin dosage and emergence of resistance in human commensal bacteria. *Journal of Infect Dis* 200: 390–398, 2009.
- Howell MD, Novack V, Grgurich P, et al. Iatrogenic gastric acid suppression and the risk of nosocomial *Clostridium difficile* infection. *Arch Intern Med* 170:784–790, 2010.
- Lewis GJ, Fang Z, Gooch M, et al. Decreased resistance of *Pseudomonas aeruginosa* with restriction of ciprofloxacin in a large teaching hospitals intensive care and intermediate care units. *Infect Control Hosp Epidemiol* 33:368–73, 2012.
- Linsky A, Gupta K, Lawler EV, et al. Proton pump inhibitors and risk for recurrent *Clostridium difficile* infection. *Arch Intern Med* 170:772–778, 2010.
- Masadeh MM, Mhaidat NM, Alzoubi KH, et al. Ciprofloxacin-induced antibacterial activity is reversed by vitamin E and vitamin C. *Curr Microbiol* 64:457–62, 2012.
- Miliani K, L'Heriteau, F, Lacave L, et al. Imipenem and ciprofloxacin consumption as factors associated with high incidence rates of resistance *Pseudomonas aeruginosa* in hospitals in northern France. *J Hosp Infect* 77:343–347, 2011.
- Sanders CC. Ciprofloxacin: In vitro activity, mechanism of action, resistance. *Rev Infect Dis* 10:516–27, 1998.
- Sandberg T, Skoog G, Hermansson AB, et al. Ciprofloxacin for 7 days versus 14 days in women with acute pyelonephritis: a randomized, open-label and double-blind, placebo-controlled, non-inferiority trial. *Lancet* 4:484–90, 2012.
- Website: [www.pdr.net](http://www.pdr.net)

## Ciprofloxacin Extended-Release (Cipro XR)

**Drug Class:** Fluoroquinolone.

**Usual Dose:** 500 mg or 1000 mg (PO) q24h (see comments).

**Spectrum:** (see **Susceptibility Profiles** pp. 198–202).

**Resistance Potential:** High (*S. pneumoniae*, *E. coli*, *P. aeruginosa*; ↑ prevalence of MRSA)

**Pharmacokinetic Parameters (500/1000 mg):**

*Peak serum level:* 1.59/3.11 mcg/mL

*Bioavailability:* 70%

*Excreted unchanged (urine):* 50–70%; 22% excreted as active metabolite

*Serum half-life:* 6.6/6.3 hrs

*Plasma protein binding:* 20–40%

*Volume of distribution ( $V_d$ ):* 2.5 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\***

CrCl < 30 mL/min	No change for 500 mg dose; reduce 1000 mg dose to 500 mg (PO) q24h
Post-HD/PD dose	500 mg (PO)
CVWH/CVWHD/ CVWHDf dose	same as for CrCl < 30 mL/min
Moderate—severe hepatic insufficiency	No change

**Drug Interactions:** Al<sup>++</sup>, Ca<sup>++</sup>, Fe<sup>++</sup>, Mg<sup>++</sup>, Zn<sup>++</sup> antacids, citrate/citric acid, dairy products, didanosine (↓ absorption of ciprofloxacin only if taken together); caffeine, cyclosporine,

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

theophylline (↑ interacting drug levels); cimetidine (↑ ciprofloxacin levels); foscarnet (↑ risk of seizures); insulin, oral hypoglycemics (slight ↑ or ↓ in blood glucose); NSAIDs (may ↑ risk of seizures/CNS stimulation); phenytoin (↑ or ↓ phenytoin levels); probenecid (↑ ciprofloxacin levels); warfarin (↑ INR).

**Adverse Effects:** Drug fever/rash, *C. difficile* diarrhea/colitis, seizures. FQ use has an increased risk of tendinitis/tendon rupture. Risk in highest in the elderly, those on steroids, and those with heart, lung or renal transplants.

**Allergic Potential:** Low

**Safety in Pregnancy:** C

**Comments: Uncomplicated UTI (acute cystitis) dose:** 500 mg (PO) q24h; **complicated UTI/acute uncomplicated pyelonephritis dose:** 1000 mg (PO) q24h. May be administered with or without food. Administer at least 2 hours before or 6 hours after H<sub>2</sub> antagonists, omeprazole, sucralfate, calcium, iron, zinc, multivitamins, or aluminum/magnesium containing medications.

↑ **incidence of *C. difficile* diarrhea/colitis with PPIs (for patients on PPIs during FQ therapy, D/C PPIs or switch to H<sub>2</sub> blocker for duration of FQ therapy).**

**Cerebrospinal Fluid Penetration:**

Non-Inflamed meninges = 10%

Inflamed meninges = 26%

**Bile Penetration:** 3000%

(also see **Antibiotic Pearls & Pitfalls** p. 516).

#### REFERENCES:

Davis R, Markham A, Balfour JA. Ciprofloxacin: An updated review of its pharmacology, therapeutic efficacy, and tolerability. *Drugs* 51:1019–74, 1996.

Website: www.pdr.net

## Clarithromycin (Biaxin)

**Drug Class:** Macrolide.

**Usual Dose:** 500 mg (PO) q12h.

**Spectrum:** (see **Susceptibility Profiles** pp. 186–190).

**Resistance Potential:** High (*S. pneumoniae*, GAS, MSSA)

**Pharmacokinetic Parameters:**

*Peak serum level:* 1–4 mcg/mL

*Bioavailability:* 50%

*Excreted unchanged (urine):* 20%

*Serum half-life (normal/ESRD):* 3–7/4 hrs

*Plasma protein binding:* 70%

*Volume of distribution (V<sub>d</sub>):* 3 L/kg

**Primary Mode of Elimination:** Hepatic

**Dosage Adjustments\***

CrCl < 30 mL/min	250 mg (PO) q12h
Post-HD dose	500 mg (PO)
Post-PD dose	None
CVWH/CVWHD/ CVWHDf dose	No change
Moderate—severe hepatic insufficiency	No change

**Drug Interactions:** Amiodarone, procainamide, sotalol, astemizole, terfenadine, cisapride, pimozide (may ↑ QT interval, torsade de pointes); carbamazepine (↑ carbamazepine levels, nystagmus, nausea, vomiting, diarrhea); cimetidine, digoxin, ergot alkaloids, midazolam, triazolam, phenytoin, ritonavir, tacrolimus, valproic acid (↑ interacting drug levels); clozapine, corticosteroids (not studied); cyclosporine (↑ cyclosporine levels with toxicity); efavirenz (↓ clarithromycin levels); rifabutin, rifampin (↓ clarithromycin levels, ↑ interacting drug levels);

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

statins (↑ risk of rhabdomyolysis); theophylline (↑ theophylline levels, nausea, vomiting, seizures, apnea); warfarin (↑ INR); zidovudine (↓ zidovudine levels).

**Adverse Effects:** Nausea, vomiting, GI upset,

**Non-C. difficile diarrhea,** abdominal pain.

May ↑ QT<sub>c</sub>; avoid with other medications that prolong the QT<sub>c</sub> interval and in patients with cardiac arrhythmias/heart block.

**Allergic Potential:** Low

**Safety in Pregnancy:** C

**Comments:** Peculiar “aluminum sand” taste sensation on swallowing.

**Cerebrospinal Fluid Penetration:** < 10%

**Bile Penetration:** 7000%

(also see **Antibiotic Pearls & Pitfalls** p. 519).

#### REFERENCES:

Alcaide F, Calatayud L, Santin M, et al. Comparative in vitro activities of linezolid, telithromycin, clarithromycin, levofloxacin, moxifloxacin and four conventional antimycobacterial drugs against mycobacterium kansasii. *Antimicrob Agents Chemother* 48:4562–5, 2004.

Alvarez-Elcoro S, Enzler MJ. The macrolides: Erythromycin, clarithromycin and azithromycin. *Mayo Clin Proc* 46:13–34, 1999.

Benson CA, Williams PL, Cohn DL, and the ACTG 196/CPCRA 009 Study Team. Clarithromycin or rifabutin alone or in combination for primary prophylaxis of *Mycobacterium avium* complex disease in patients with AIDS: A randomized, double-blinded, placebo-controlled trial. *J Infect Dis* 181:1289–97, 2000.

Berg HF, Tjhie JH, Scheffer GJ, et al. Emergence and persistence of macrolide resistance in oropharyngeal flora and elimination of nasal carriage of *Staphylococcus aureus* after therapy with slow-release clarithromycin: a randomized, double-blind, placebo-controlled study. *Antimicrob Agents Chemother* 48:4138–8, 2004.

Bergman M, Huikko S, Huovinen P, et al. Macrolide and azithromycin use are linked to increased macrolide resistance in *Streptococcus pneumoniae*. *Antimicrob Agents Chemother* 50:3646–3650, 2006.

Daneman N, McGeer A, Green K, et al. Macrolide resistance in bacteremic pneumococcal disease: Implications for patient management. *Clin Infect Dis* 43:432–438, 2006.

Dean NC, Sperry P, Wikler M, et al. Comparing clarithromycin in pneumonia symptom resolution and process of care. *Antimicrob Agents Chemother* 50:1164–69, 2006.

Jain R, Danzinger LH. The macrolide antibiotics: a pharmacokinetic and pharmacodynamic overview. *Curr Pharm Des* 10:3045–53, 2004.

Jenkins SG, Farrell DJ. Increase in pneumococcus macrolide resistance, United States. *Emerging Infect Dis* 15:1260–1264, 2009.

Kafetzis DA, Chantzi F, Tigaani G, et al. Safety and tolerability of clarithromycin administered to children at higher-than-recommended doses. *Eur J Clin Microbiol Infect Dis* 26:99–103, 2007.

Kraft M, Cassell AGH, Jpak J, et al. *Mycoplasma pneumoniae* and *Chlamydia pneumoniae* in asthma: effect of clarithromycin. *Chest* 121:1782–8, 2002.

Malfertheiner P, Bazzoli F, Delchier JC, et al. *Helicobacter pylori* eradication with a capsule containing bismuth subcitrate potassium, metronidazole, and tetracycline given with omeprazole versus clarithromycin-based triple therapy: a randomized open-label, non-inferiority, phase 3 trial. *Lancet* 33:905–913, 2011.

McConnell SA, Amsden GW. Review and comparison of advanced-generation macrolides clarithromycin and dirithromycin. *Pharmacotherapy* 19:404–15, 1999.

Ozsoylar G, Sayin A, Bolay H. Clarithromycin monotherapy-induced delirium. *J Antimicrob Chemother* 59:331, 2007.

Periti P, Mazzei T. Clarithromycin: Pharmacokinetic and pharmacodynamic interrelationships and dosage regimen. *J Chemother* 11:11–27, 1999.

Portier H, Filipecki J, Weber P, et al. Five day clarithromycin modified release versus 10 day penicillin V for group A streptococcal pharyngitis: a multi-centre, open-label, randomized study. *J Antimicrob Chemother* 49:337–44, 2002.

Rodvold KA. Clinical pharmacokinetics of clarithromycin. *Clin Pharmacokinetics* 37:385–98, 1999.

Schlossberg D. Azithromycin and clarithromycin. *Med Clin North Am* 79:803–16, 1995.

“Usual dose” assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

Stein GE, Schooley S. Serum bactericidal activity of extended-release clarithromycin against macrolide-resistant strains of *Streptococcus pneumoniae*. *Pharmacotherapy* 22:593–6, 2002.

Svensson M, Strom M, Nilsson M et al. Pharmacodynamic effects of nitroimidazoles alone and in combination with clarithromycin on *Helicobacter pylori*. *Antimicrob Agents Chemother* 46:2244–8, 2002.

Tartaglione TA. Therapeutic options for the management and prevention of *Mycobacterium avium* complex infection in patients with acquired immunodeficiency syndrome. *Pharmacotherapy* 16:171–82, 1996.

Website: www.pdr.net

## Clarithromycin XL (Biaxin XL)

**Drug Class:** Macrolide.

**Usual Dose:** 1 gm (PO) q24h.

**Spectrum:** (see **Susceptibility Profiles**

pp. 186–190).

**Resistance Potential:** High (*S. pneumoniae*, GAS, MSSA)

**Pharmacokinetic Parameters:**

*Peak serum level:* 3 mcg/mL

*Bioavailability:* 50%

*Excreted unchanged (urine):* 20%; 10–15%

*excreted as active metabolite*

*Serum half-life (normal/ESRD):* 4/4 hrs

*Plasma protein binding:* 70%

*Volume of distribution ( $V_d$ ):* 3 L/kg

**Primary Mode of Elimination:** Hepatic

**Dosage Adjustments\***

CrCl < 30 mL/min	500 mg (PO) q24h
Post-HD dose	None
Post-PD dose	None
CVWH/CVVD/ CVHDF dose	No change

Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	No change

**Drug Interactions:** Amiodarone, procainamide, sotalol, astemizole, terfenadine, cisapride, pimozide (may ↑ QT interval, torsade de pointes); carbamazepine (↑ carbamazepine levels, nystagmus, nausea, vomiting, diarrhea); cimetidine, digoxin, ergot alkaloids, midazolam, triazolam, phenytoin, ritonavir, tacrolimus, valproic acid (↑ interacting drug levels); clozapine, corticosteroids (not studied); cyclosporine (↑ cyclosporine levels with toxicity); efavirenz (↓ clarithromycin levels); rifabutin, rifampin (↓ clarithromycin levels, ↑ interacting drug levels); statins (↑ risk of rhabdomyolysis); theophylline (↑ theophylline levels, nausea, vomiting, seizures, apnea); warfarin (↑ INR); zidovudine (↓ zidovudine levels).

**Adverse Effects:** Nausea, vomiting **Non-C. difficile diarrhea.** May ↑ QT<sub>c</sub>; avoid with other medications that prolong the QT<sub>c</sub> interval and in patients with cardiac arrhythmias/heart block.

**Allergic Potential:** Low

**Safety in Pregnancy:** C

**Comments:** Two 500 mg tablets of XL preparation permits once daily dosing and decreases GI intolerance.

**Cerebrospinal Fluid Penetration:** < 10%

**Bile Penetration:** 7000%

(also see **Antibiotic Pearls & Pitfalls** p. 519).

### REFERENCES:

Adler JL, Jannetti W, Schneider D, et al. Phase III, randomized, double-blind study of clarithromycin extended-release and immediate-release formulations in the treatment of patients with acute exacerbation of chronic bronchitis. *Clin Ther* 22:1410–20, 2000.

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

- Anzueto A, Fisher CL Jr, Busman T. Comparison of the efficacy of extended-release clarithromycin tablets and amoxicillin/clavulanate tablets in the treatment of acute exacerbation of chronic bronchitis. *Clin Ther* 23:72–86, 2000.
- Bergman N, Huikko S, Huovinen P, et al. Macrolide and azithromycin use are linked to increased macrolide resistance in *Streptococcus pneumoniae*. *Antimicrob Agents Chemother* 50:3646–3650, 2006.
- Daneman N, McGeer A, Green K, et al. Macrolide resistance in bacteremic pneumococcal disease: Implications for patient management. *Clin Infect Dis* 43:432–438, 2006.
- Gotfried HM. Clarithromycin (Biaxin) extended-release tablet: a therapeutic review. *Expert Rev Anti Infect Ther* 1:9–20, 2004.
- Jenkins SG, Farrell DJ. Increase in pneumococcus macrolide resistance, United States. *Emerging Infect Dis* 15:1260–1264, 2009.
- Laine L, Estrada R, Trujillo M, et al. Once-daily therapy for *H. pylori* infection: Randomized comparison of four regimens. *Am J Gastroenterol* 94:962–6, 1999.
- Mandell LA, Wunderink RG, Anzueto A, et al. Infectious Diseases Society of America/American Thoracic Society consensus guidelines on the management of community-acquired pneumonia in adults. *Clin Infect Dis* 44:S27–72, 2007.
- Nalepa P, Dobryniowska M, Busman T, et al. Short-course therapy of acute bacterial exacerbation of chronic bronchitis: a double-blind, randomized, multicenter comparison of extended-release versus immediate-release clarithromycin. *Curr Med Res Opin* 19:411–20, 2003.
- Website: www.pdr.net

## Clindamycin (Cleocin)

**Drug Class:** Lincosamide.

**Usual Dose:** 600 mg (IV) q8h;  
150–300 mg (PO) q6h.

**Spectrum:** (see **Susceptibility Profiles** pp. 186–190).

**Resistance Potential:** Low

**Pharmacokinetic Parameters:**

*Peak serum level:* 2.5–10 mcg/mL

*Bioavailability:* 90%

*Excreted unchanged (urine):* 10%; 3.6% excreted as active metabolite

*Serum half-life (normal/ESRD):* 2.4/4 hrs

*Plasma protein binding:* 90%

*Volume of distribution ( $V_d$ ):* 1 L/kg

**Primary Mode of Elimination:** Hepatic

**Dosage Adjustments\***

CrCl < 10 mL/min	No change
Post-HD/PD dose	None
CVVH/CVVHD/ CVVHDF dose	No change
Moderate—severe hepatic insufficiency	No change

**Drug Interactions:** Muscle relaxants, neuromuscular blockers (↑ apnea, respiratory paralysis); kaolin (↓ clindamycin absorption); theophylline (↑ theophylline levels, seizures).

**Adverse Effects:** C. difficile diarrhea/colitis, neuromuscular blockade.

**Allergic Potential:** Low

**Safety in Pregnancy:** B

**Comments:** **Anti-spasmodics contraindicated in C. difficile diarrhea.** C. difficile diarrhea more common with PO vs. IV clindamycin. **Clindamycin misses some Peptostreptococci, Fusobacteria, non-perfringens Clostridia, and Prevotella.**

**Cerebrospinal Fluid Penetration:** < 10%

**Bile Penetration:** 300%

(also see **Antibiotic Pearls & Pitfalls** p. 513).

### REFERENCES:

- Back EE, O'Grady EJ, Back JD. High rates of perinatal group B *Streptococcus* clindamycin and erythromycin resistance in an upstate New York Hospital. *Antimicrob Agents Chemother* 56:739–742, 2012.

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

- Coyle EA, Cha R, Rybak MJ. Influences of linezolid, penicillin, and clindamycin, alone and in combination, on streptococcal pyrogenic exotoxin A release. *Antimicrob Agents Chemother* 47:1752–5, 2003.
- Cunha BA. Oral Antibiotic Therapy of Serious Systemic Infections. *Med Clin N Am* 90:1197–1222, 2006.
- Czekaj J, Dinh A, Moldovan A, et al. Efficacy of a combined oral clindamycin-rifampicin regimen for therapy of staphylococcal osteoarticular infections. *Scand J Infect Dis* 43:962–967, 2011.
- Falagas ME, Gorbach SL. Clindamycin and metronidazole. *Med Clin North Am* 79:845–67, 1995.
- Kasten MJ. Clindamycin, metronidazole, and chloramphenicol. *Mayo Clin Proc* 74:825–33, 1999.
- Klepser ME, Nicolau DP, Quintillani R, et al. Bactericidal activity of low-dose clindamycin administered at 8- and 12-hour intervals against *Streptococcus pneumoniae* and *Bacteroides fragilis*. *Antimicrob Agents Chemother* 41:630–5, 1997.
- Lell B, Dremsner PG. Clindamycin as an antimalarial drug: review of clinical trials. *Antimicrob Agents Chemother* 46:2315–20, 2002.
- Levin TP, Suh B, Axelrod P, et al. Potential clindamycin resistance in clindamycin-susceptible, erythromycin-resistant staphylococcus aureus: Report of a clinical failure. *Antimicrob Agents Chemother* 49:1222–24, 2005.
- Patel M, Waites KB, Moser SA, et al. Prevalence of inducible clindamycin resistance among community- and hospital-associated *Staphylococcus aureus* isolates. *Journal of Clinical Microbiology* 44:2481–84, 2006.
- Website: www.pdr.net

## Colistin (Coly-Mycin M)

**Drug Class:** Cell membrane altering antibiotic.

**Usual Dose:** 5 mg/kg (IV) q8h.

**Spectrum:** (see **Susceptibility Profiles** pp. 198–202).

**Resistance Potential:** Low

**Pharmacokinetic Parameters:**

*Peak serum level:* 5 mcg/mL

*Bioavailability:* Not applicable

*Excreted unchanged (urine):* 90%

*Serum half-life (normal/ESRD):* 9/13 hrs

*Plasma protein binding:* 50%

*Volume of distribution ( $V_d$ ):* 0.09 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\***

CrCl 30–49 mL/min	3.5 mg/kg (IV) q24h
CrCl 10–29 mL/min	2.5 mg/kg (IV) q24h
CrCl < 10 mL/min	1.5 mg/kg (IV) q24h
Post-HD dose	None
Post-PD dose	None
CVWH/CVWH/ CVHDF dose	2.5 mg/kg (IV) q48h
Mild hepatic insufficiency	No change
Moderate—severe hepatic insufficiency	No change

**Drug Interactions:** Neuromuscular blocking agents (↑ neuromuscular blockade); nephrotoxic drugs (↑ nephrotoxic potential).

**Adverse Effects:** Dose dependent/reversible nephrotoxic potential (acute tubular necrosis). Paresthesias, vertigo, dizziness, slurred speech, blurry vision, respiratory arrest.

**Allergic Potential:** Low

**Safety in Pregnancy:** C

**Comments:** 1 mg = 12,500 U. Colistin (Polymyxin E)

has **less nephrotoxic potential than previously thought. Useful for MDR *P. aeruginosa* and *Acinetobacter* species.**

For *P. aeruginosa* or *Acinetobacter* meningitis *also give amikacin* 10–40 mg (IT) q24h or *colistin* 10 mg (IT) q24h. **Intrathecal colistin dose:** 10 mg (IT) Q24h. **Nebulizer dose:** 80 mg in saline q8h; for recurrent infection use 160 mg q8h (freshly prepare solution and use within 24 hours). If possible, avoid aerosolized colistin, which may result in pulmonary/systemic toxicity

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

due to the polymyxin E1 metabolite. **Continuous infusion dose:** give ½ of daily dose (IV) over 5–10 min, then give remaining ½ dose (in D5W or NS) 1 hour after initial dose over next 24 hours.

**Cerebrospinal Fluid Penetration:** < 25% (also see **Antibiotic Pearls & Pitfalls** p. 520).

#### REFERENCES:

- Cai Y, Chai D, Wang R, et al. Colistin resistance of *Acinetobacter baumannii*: clinical reports, mechanisms and antimicrobial strategies. *J Antimicrob Chemother* 67:1607–15, 2012.
- Cascio A, Conti A, Sinardi L, et al. Post-neurosurgical multidrug-resistant *Acinetobacter baumannii* meningitis successfully treated with intrathecal colistin. A new case and a systematic review of the literature. *Int J Infect Dis* 14:e572–9, 2010.
- Cheng CY, Sheng WH, Wang JT, et al. Safety and efficacy of intravenous colistin (colistin methanesulphonate) for severe multidrug-resistant Gram-negative bacterial infections. *Int J Antimicrob Agents* 35:297–300, 2010.
- Couet W, Gregoire N, Marchan S, et al. Colistin pharmacokinetics: the fog is lifting. *Clin Microbiol Infect* 18: 30–39, 2012.
- Cunha BA. New uses for older antibiotics: nitrofurantoin, amikacin, colistin, polymyxin B, doxycycline, and minocycline revisited. *Med Clin North Am* 90:1089–1107, 2006.
- Daikos GL, Skiada A, Pavleas J, et al. Serum bactericidal activity of three different dosing regimens of colistin with implications for optimum clinical use. *J Chemother* 22:175–178, 2010.
- Deryke CA, Crawford AJ, Uddin N, et al. Colistin dosing and nephrotoxicity in a large community teaching hospital. *Antimicrob Agents Chemother* 54:4503–4505, 2010.
- Durante-Mangoni E, Signoriello G, Andini R, et al. Colistin and rifampicin compared with colistin alone for the treatment of serious infections due to extensively drug-resistant *Acinetobacter baumannii*. *Clin Infect Dis* 57: 349–358, 2013.
- Falagas ME, Rafailidis PL. Colistin in ventilator-associated pneumonia. *Clin Infect Dis* 54:681–3, 2012.
- Falagas ME, Rafailidis PI, Kasiakou SK, et al. Effectiveness and nephrotoxicity of colistin monotherapy vs colistin-meropenem combination therapy for multidrug-resistant Gram-negative bacterial infections. *Clin Microbiol Infect* 12:1227–30, 2006.
- Falagas ME, Rafailidis PI. Nephrotoxicity of colistin: new insight into an old antibiotic. *Clin Infect Dis* 48:1729–1731, 2009.
- Falagas ME, Rafailidis PI, Ioannidou E, et al. Colistin therapy for microbiologically documented multi drug-resistant Gram-negative bacterial infections: a retrospective cohort study of 258 patients. *Int J Antimicrob Agents* 35:194–199, 2010.
- Gauthier TP. Rifampicin plus colistin in the era of extensively drug-resistant *Acinetobacter baumannii* infections. *Clin Infect Dis* 57:359–362, 2013.
- Hartzell JD, Neff R, Ake J, et al. Nephrotoxicity associated with intravenous colistin (colistimethate sodium) treatment at a tertiary care medical center. *Clin Infect Dis* 48:1724–1728, 2009.
- Hindler JA, Humphries RM. Colistin MIC Variability by method for contemporary clinical isolates of multidrug-resistant gram-negative bacilli. *J Clin Micro* 51:1678–1684, 2013.
- Horton J, Pankey GA. Polymyxin B, colistin, and sodium colistimethate. *Med Clin North Am* 66:135–42, 1982.
- Imberti R. Intravenous colistimethate administration and colistin lung tissue concentrations. *Intensive Care Med* 36:1795, 2010.
- Kempf M, Djouhri-Bouktab L, Brunel JM, et al. Synergistic activity of sulbactam combined with colistin against colistin-resistant *Acinetobacter baumannii*. *Int J Antimicrob Agents* 39:180–1, 2012.
- Kofteridis DP, Alexopoulou C, Valachis A, et al. Aerosolized plus intravenous colistin versus intravenous colistin alone for the treatment of ventilator-associated pneumonia: a matched case-control study. *Clin Infect Dis* 51: 1238–1244, 2010.
- Kwa AL, Falagas ME, Michalopoulos A, et al. Benefits of aerosolized colistin for ventilator-associated pneumonia: absence of proof versus proof of absence? *Clin Infect Dis* 52:1278–1279, 2011.
- Levin AS, Baron AA, Penco J, et al. Intravenous colistin as therapy for nosocomial infections caused by multidrug resistant *Pseudomonas aeruginosa* and *Acinetobacter baumannii*. *Clin Infect Dis* 28:1008–11, 1999.

<sup>†</sup>Usual dose<sup>†</sup> assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.



- Liang W, Liu XR, Huang, J, et al. Activities of colistin- and minocycline-based combinations against extensive drug resistant *Acinetobacter baumannii* isolates from intensive care unit patients. *BMC Infect Dis* 11:109, 2011.
- Linden PK, Kusne S, Coley K, et al. Use of parenteral colistin for the treatment of serious infection due to antimicrobial-resistant *Pseudomonas aeruginosa*. *Clin Infect Dis* 37:154–60, 2003.
- Marchand S, Frat JP, Petitpas F, et al. Removal of colistin during intermittent haemodialysis in two critically ill patients. *J Antimicrob Chemother* 65:1836–1837, 2010.
- Markantonis SL, Markou N, Fousteri M, et al. Penetration of colistin into cerebrospinal fluid. *Antimicrob Agents Chemother* 53:4907–4910, 2009.
- Oliveira MS, Prado GV, Costa SF, et al. Polymyxin B and colistimethate are comparable as to efficacy and renal toxicity. *Diagn Microbiol Infect Dis* 65:431–434, 2009.
- Ozbek B, Otuk G. In vitro activities of tigecycline alone and in combination with colistin su fate or sulbactam against carbapenem-susceptible and -resistant *Acinetobacter baumannii* strains isolated from Intensive Care Units. *Int J Antimicrob Agents* 36:191–192, 2010.
- Paul M, Bishara J, Levcovich A, et al. Effectiveness and safety of colistin: prospective comparative cohort study. *J Antimicrob Chemother* 65:1019–1027, 2010.
- Papagelopoulos PJ, Mavrogenis AF, Giannitsioti E, et al. Management of a multidrug-resistant *Pseudomonas aeruginosa* infected knee arthroplasty using colistin. A care report and review of the literature. *J Arthroplasty* 22:457–463, 2007.
- Plachouras D, Giamerlos-Bourboulis EJ, Kentepozidis N, et al. In vitro postantibiotic effect of colistin on multidrug-resistant *Acinetobacter baumannii*. *Diagn Microbiol Infect Dis* 57:419–422, 2007.
- Rattanaumpawan P, Ungprasert P, Thamilkitkul V. Risk factors for colistin-associated nephrotoxicity. *J Infect* 62:187–190, 2011.
- Reed MD, Stern RC, O'Riordan MA, et al. The pharmacokinetics of colistin in patients with cystic fibrosis. *J Clin Pharmacol* 41:645–54, 2001.
- Rynn C, Wooton M, Bowker KE, et al. In vitro assessment of colistin's antipseudomonal antimicrobial interactions with other antibiotics. *Clin Microbiol Infect* 5:32–6, 1999.
- Santamaría C, Mykietiak A, Temporiti E, et al. Nephrotoxicity associated with the use of intravenous colistin. *Scand J Infect Dis* 41:767–769, 2009.
- Stein A, Raoult D. Colistin: an antimicrobial for the 21<sup>st</sup> century? *Clin Infect Dis* 35:901–2, 2002.
- Tan TY, Ng SY. Comparison of Etest, Viteck and agar dilution for susceptibility testing of colistin. *Clin Microbiol Infect* 13:541–544, 2007.
- Tarchini G. Nephrotoxicity associated with intravenous colistin. *Clin Infect Dis* 49:1773, 2009.
- Van Leeuwen DH. A closer look at aerosolized colistin. *Clin Infect Dis* 52:1472–1473, 2011.
- Volkow-Fernandez P, Rodriguez CF, Cornejo-Juarez P. Intravesical colistin irrigation to treat multidrug-resistant *Acinetobacter baumannii* urinary tract infection. *J of Med Case Reports* 6:426–427, 2012.
- Wahby K, Chopra T, Chandrasekar P. Intravenous and Inhalational colistin-induced respiratory failure. *Clin Infect Dis* 50:e38–40, 2010.
- Yahav D, Farbmam L, Leibovici L, et al. Colistin: New lessons on an old antibiotic. *Clin Microbiol Infect* 18:18–29, 2012.
- Website: [www.pdr.net](http://www.pdr.net)

## Cycloserine (Seromycin)

**Drug Class:** TB drug.

**Usual Dose:** 250 mg (PO) q12h.

**Pharmacokinetic Parameters:**

*Peak serum level:* 20 mcg/mL

*Bioavailability:* 90%

*Excreted unchanged (urine):* 65%

*Serum half-life (normal/ESRD):* 10–25 hrs/no data

*Plasma protein binding:* No data

*Volume of distribution ( $V_d$ ):* 0.2 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\***

\*Usual dose\* assumes normal renal/hepatic function. \*For renal insufficiency, give usual dose  $\times$  1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

CrCl 10–50 mL/min	250 mg (PO) q12–24h
CrCl < 10 mL/min	250 mg (PO) q24h
Post-HD dose	None
Post-PD dose	250 mg (PO)
CVVH/CVVHD/ CVVHDF dose	No change
Moderate—severe hepatic insufficiency	No change

**Drug Interactions:** Alcohol (seizures); ethambutol, ethionamide (drowsiness, dizziness); phenytoin (↑ phenytoin levels).

**Adverse Effects:** Peripheral neuropathy, seizures (dose related), psychosis/delirium.

Allergic Potential: Low

Safety in Pregnancy: C

**Comments:** Avoid in patients with seizures.

Ethambutol, ethionamide, or ethanol may increase CNS toxicity.

Meningeal dose = usual dose.

#### Cerebrospinal Fluid Penetration:

Non-Inflamed meninges = 90%

Inflamed meninges = 90%

#### REFERENCES:

Davidson PT, Le HQ. Drug treatment of tuberculosis 1992.

Drugs 43:651–73, 1992.

Furin J, Nardell EA. Multidrug-resistant tuberculosis: An update on the best regimens. *J Respir Dis* 27:172–82, 2006.

Iseman MD. Treatment of multidrug resistant tuberculosis.

*N Engl J Med* 329:784–91, 1993.

Website: www.pdr.net

## Dalbavancin (Dalvance)

**Drug Class:** Lipoglycopeptide

**Usual Dose:** 1 gm (IV) × 1 dose, then 500 mg (IV) × 1 dose 7 days later.

**Spectrum:** (see *Susceptibility Profiles* pp. 198-202.

**Resistance Potential:** Low

#### Pharmacokinetic Parameters:

*Peak serum level:* 287 mcg/mL

*Bioavailability not applicable*

*Excreted unchanged (urine/feces):* 33%/20%

*Serum half-life (normal/ESRD):* 346 h/376 h

*Plasma protein binding:* 93%

*Volume of distribution (V<sub>d</sub>):* 0.1–0.19 L/kg

**Primary Mode of Elimination:** Renal

#### Dosage Adjustments\*

CrCl < 30 mL/min	750 mg (IV) then 325 mg (IV) 7 days later
Post-HD dose	No change
Post-PD dose	No change
CVVH dose	No data
Hepatic insufficiency	No change

**Drug Interactions:** None

**Adverse Effects:** Headache, nausea, vomiting, diarrhea, rash, pruritus, increased AST/ALT, C. difficile.

**Allergic Potential:** Moderate

**Safety in Pregnancy:** C

**Comments:** Infuse slowly over 30 minutes to avoid “Red Man Syndrome”; hypersensitivity reactions (use caution if patient had previous hypersensitivity reaction to other glycopeptides, e.g. vancomycin).

(also see *Antibiotic Pearls & Pitfalls* p. 519).

#### REFERENCES:

Bailey J, Summers KM. Dalbavancin: a new lipoglycopeptide antibiotic. *Am J Health Syst Pharm.* 65:599-610, 2008.

Boucher HW, Wilcox M, Talbot GH, et al. Once-weekly dalbavancin versus daily conventional therapy for skin infection. *N Engl J Med.* 370:2169-2179, 2014.

\*Usual dose\* assumes normal renal/hepatic function. \*For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

Citron OM, Tyrrell KL, Goldstein EJ. Comparative in vitro activities of dalbavancin and seven comparator agents against 41 *Staphylococcus* species cultured from osteomyelitis infections and 18 VISA and hVISA strains. *Diagn Microbiol Infect Dis*. 79:438-440, pp. 2014.

Dowell JA, Goldstein BP, Buckwalter M, et al. Pharmacokinetic-pharmacodynamic modeling of dalbavancin, a novel glycopeptide antibiotic. *J Clin Pharmacol*. 48:1063-1068, 2008.

Holmes NE, Howden BP. What's new in the treatment of serious MRSA infection? *Curr Opin Infect Dis*. 27:471-478, 2014.

Jaregui LE, Babazadeh S, Seltzer E, et al. Randomized, double-blind comparison of once-weekly dalbavancin versus twice-daily linezolid therapy for the treatment of complicated skin and skin structure infections. *Clin Infect Dis*. 41:1407-1415, 2005.

Marbury T, Dowell JA, Seltzer E, et al. Pharmacokinetics of dalbavancin in patients with renal or hepatic impairment. *J Clin Pharmacol*. 49:465-476, 2009.

Seltzer E, Dorr MB, Goldstein BP, et al. Once-weekly dalbavancin versus standard-of-care antimicrobial regimens for treatment of skin and soft-tissue infections. *Clin Infect Dis*. 37:1298-1303, 2003.

Vilay AM, Shah KH, Churchwell MD, et al. Modeled dalbavancin transmembrane clearance during intermittent and continuous renal replacement therapies. *Blood Purif*. 30:37-43, 2010.

Website: [www.pdr.net](http://www.pdr.net)

## Dapsone

**Drug Class:** Antiparasitic, anti-leprosy, anti-PCP drug.

**Usual Dose:** 100 mg (PO) q24h (see comments).

**Pharmacokinetic Parameters:**

*Peak serum level:* 1.8 mcg/mL

*Bioavailability:* 85%

*Excreted unchanged (urine):* 10%

*Serum half-life (normal/ESRD):* 25/30 hrs

*Plasma protein binding:* 80%

*Volume of distribution ( $V_d$ ):* 1.2 L/kg

**Primary Mode of Elimination:** Hepatic/renal

**Dosage Adjustments\***

CrCl < 10 mL/min	No change
Post-HD dose	None
Post-PD dose	None
CVWH/CVHD/ CVHDF dose	No change
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	No data

**Drug Interactions:** Didanosine (↓ dapsone absorption); oral contraceptives (↓ oral contraceptive effect); pyrimethamine, zidovudine (↑ bone marrow suppression); rifabutin, rifampin (↓ dapsone levels); trimethoprim (↑ dapsone and trimethoprim levels, methemoglobinemia).

**Adverse Effects:** Drug fever/rash, nausea, vomiting, hemolytic anemia in G6PD deficiency, methemoglobinemia.

**Allergic Potential:** High

**Safety in Pregnancy:** C

**Comments:** Useful in sulfa (SMX) allergic patients. Avoid, if possible, in G6PD deficiency or hemoglobin M deficiency.

**PCP prophylaxis dose:** 100 mg (PO) q24h. **PCP therapy dose:** 100 mg (PO) q24h.

**Cerebrospinal Fluid Penetration:** < 50%

**REFERENCES:**

- East J, Blanton LS. Symptomatic hyperbilirubinemia secondary to dapsone-induced hemolysis and atazanavir therapy. *Antimicrob Chemother* 67:465-8, 2012.
- El-Sadr WM, Murphy RL, Yurik TM, et al. Atovaquone compared with dapsone to the prevention of *Pneumocystis carinii* in patients with HIV infection who cannot tolerate trimethoprim, sulfonamides, or both. *N Engl J Med* 339:1889-95, 1998.
- Medina I, Mills J, Leoung G, et al. Oral therapy for *Pneumocystis carinii* pneumonia in the acquired immunodeficiency syndrome. A controlled trial of trimethoprim-sulfamethoxazole versus

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

trimethoprim-dapsone. *N Engl J Med* 323:776–82, 1990.

Podzmaczer D, Salazar A, Jimenez J, et al. Intermittent trimethoprim-sulfamethoxazole compared with dapsone-pyrimethamine for the simultaneous primary prophylaxis of *Pneumocystis pneumonia* and toxoplasmosis in patients infected with HIV. *Ann Intern Med* 122:755–61, 1995.

Website: [www.pdr.net](http://www.pdr.net)

## Daptomycin (Cubicin)

**Drug Class:** Lipopeptide.

**Usual Dose:** Complicated skin/skin structure infections: 4 mg/kg (IV) q24h. Bacteremia, endocarditis: 6 mg/kg (IV) q24h. Bacteremia, endocarditis unresponsive to 6 mg/kg (IV) q24h × 3 days: 12 mg/kg (IV) q24h.

**Spectrum:** (see **Susceptibility Profiles** pp. 198–202).

**Resistance Potential:** Low

**Pharmacokinetic Parameters:**

*Peak serum level:*

4 mg/kg = 58 mcg/mL; 6 mg/kg = 99 mcg/mL

*Bioavailability:* Not applicable

*Excreted unchanged (urine):* 80%

*Serum half-life (normal/ESRD):* 8.1/29.8 hrs

*Plasma protein binding:* 92%

*Volume of distribution ( $V_d$ ):* 0.096 L/kg

**Primary Mode of Elimination:** Renal  
**Dosage Adjustments\***

CrCl > 30 mL/min	No change
CrCl < 30 mL/min	4 mg/kg (IV) q48h or 6 mg/kg (IV) q48h or 12 mg/kg (IV) q48h
Post-HD dose	None
Post-PD dose	None
CVVH/CVVHD/ CVVHDF dose	8 mg/kg (IV) q48h
HD q 48–72h dose	10 mg/kg (IV) q HD*
Moderate—severe hepatic insufficiency	No change

\*Usual dose\* assumes normal renal/hepatic function. \*For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

\*Give at end of HD.

**Drug Interactions:** Warfarin/statin (no significant interaction in small number of volunteers; consider temporary suspension of statins during daptomycin use).

**Adverse Effects:** Constipation, nausea, headache, peripheral neuropathy. Rarely, ↑ CPK; in clinical trials, same or less (0.2%) than comparators. Eosinophilic pneumonia (rare and reversible when drug discontinued). False-positive ↑ INR.

**Allergic Potential:** No data

**Safety in Pregnancy:** B

**Comments:** Concentration-dependent killing may be enhanced with an initial dose of gentamicin. Antecedent vancomycin (↑ cell wall thickness of staphylococci) predisposes to daptomycin resistance (penetration mediated resistance). ↑ daptomycin MICs with strains of *S. aureus* may be ↓ by adding a β-lactam (to ↑ cell wall permeability) for staphylococci add nafcillin to daptomycin; for enterococci add ampicillin to daptomycin. **High-dose daptomycin 12 mg/kg (IV) q24h has been used for MSSA/MRSA bacteremias/ABE unresponsive to other antistaphylococcal drugs without toxicity.** Post-antibiotic effect (PAE) up to 6 hours. For rapid effect or need to limit infusion volume, daptomycin **may be given IV push over 5 minutes** (500 mg dose given in 10 mL; 1 g dose given in 20 mL). Cannot be given IM.

**Cerebrospinal Fluid Penetration:** ~ 5% (also see **Antibiotic Pearls & Pitfalls** p. 518).

### REFERENCES:

Alder JD. Daptomycin: a new drug class for the treatment of gram-positive infections. *Drugs Today* 41:81–90, 2005.  
Bassetti M, Nicco E, Ginocchio F, et al. High-dose daptomycin in documented *Staphylococcus aureus* infections. *Int J Antimicrob Agents* 36:459–461, 2010.

- Bhavani SM, Prakhya A, Hammel JP, et al. Cost-Effectiveness of daptomycin versus vancomycin and gentamicin for patients with methicillin-resistant *Staphylococcus aureus* bacteremia and/or endocarditis. *Clin Infect Dis* 49:691–698, 2009.
- Benevuto M, Benziger DP, Yankelev S, Vigniani G. Pharmacokinetics and tolerability of daptomycin at doses up to 12 milligrams per kilogram of body weight once daily in healthy volunteers. *Antimicrobial Agents and Chemotherapy* 50: 3245–49, 2006.
- Burkhardt O, Kielstein JT. A simplified three-times weekly daptomycin dosing regimen for chronic hemodialysis patients. *Expert Rev Anti Infect Ther* 8:11–14, 2010.
- Cantón R, Ruiz-Garbajosa P, Chaves RL, Johnson AP. A potential role for daptomycin in enterococcal infections: what is the evidence? *J Antimicrob Chemother* 65:1126–1136, 2010.
- Cervera C, Casteneda X, Pericas JM, et al. Clinical utility of daptomycin in infective endocarditis caused by Gram-positive cocci. *Int J Antimicrob Agents* 38:365–370, 2011.
- Crompton JA, North DS, McConnell SA, et al. Safety and efficacy of daptomycin in the treatment of osteomyelitis: results from the CORE Registry. *J Chemother* 21:414–420; 2009.
- Crompton JA, North DS, Yoon M, et al. Outcomes with daptomycin in the treatment of *Staphylococcus aureus* infections with a range of vancomycin MICs. *J Antimicrob Chemother* 65:1784–1791, 2010.
- Cunha BA, Mickail N, Eisenstein L. E. faecalis vancomycin sensitive enterococci (VSE) bacteremia unresponsive to vancomycin successfully treated with high dose daptomycin. *Heart & Lung* 36: 456–461, 2007.
- Cunha BA, Eisenstein LE, Hamid NS. Pacemaker-induced *Staphylococcus aureus* mitral valve acute bacterial endocarditis complicated by persistent bacteremia from a coronary stent: Cure with prolonged/high-dose daptomycin without toxicity. *Heart & Lung* 35:207–11, 2006.
- Cunha BA. Persistent *S. aureus* bacteremia: clinical pathway for diagnosis and treatment. *Antibiotics for Clinicians* 10:539–46, 2006b.
- Cunha BA, Hamid N, Kessler H, Panchuri S. Daptomycin cure after ceftazolin treatment failure of methicillin-sensitive *Staphylococcus aureus* (MSSA) tricuspid valve acute bacterial endocarditis from a peripherally inserted central catheter (PICC) line. *Heart & Lung* 34:442–7, 2005.
- De Rosa FG, Mollaretti O, Cometto C, et al. Early experience with high-dosage daptomycin for prosthetic infections. *Clin Infect Dis* 49:1772–1773, 2009.
- Dhand A, Bayer AS, Pogliano J, et al. Use of antistaphylococcal beta-lactams to increase daptomycin activity in eradicating persistent bacteremia due to methicillin-resistant *Staphylococcus aureus*: role of enhanced daptomycin binding. *Clin Infect Dis* 53:158–163, 2011.
- Doan TL, DePetrillo J, Singer C. Daptomycin-induced eosinophilia without pulmonary involvement. *Am J Health Syst Pharm* 67:2107–2109, 2010.
- Dretler R, Branch T. MRSA knee infection treated successfully with daptomycin after two failed prolonged high-dose courses of vancomycin. *Infect Dis Clin Pract* 13:327–329, 2005.
- Falagas ME, Giannopoulou KP, Ntziora F, et al. Daptomycin for endocarditis and/or bacteremia: a systematic review of the experimental and clinical evidence. *J Antimicrob Chemother* 60:7–19, 2007.
- Falcone M, Russo A, Venditti M, et al. Considerations for higher doses of daptomycin in critically ill patients with methicillin-resistant *Staphylococcus aureus* bacteremia. *Clin Infect Dis* 57:1568–1576, 2013.
- Figuerola DA, Mangini E, Amodio-Groton M, et al. Safety of high-dose intravenous daptomycin treatment: three-year cumulative experience in a clinical program. *Clin Infect Dis* 15:49:177–80, 2009.
- Gallagher JC, Perez ME, Marino EA, et al. Daptomycin therapy for vancomycin-resistant enterococcal bacteremia: a retrospective case series of 30 patients. *Pharmacotherapy* 29:792–799, 2009.
- Hair PI, Keam SJ. Daptomycin: a review of its use in the management of complicated skin and soft-tissue infections and *Staphylococcus aureus* bacteremia. *Drugs* 67:1483–1512, 2007.
- Hayes D Jr, Anstead MI, Kuhn RJ. Eosinophilic pneumonia induced by daptomycin. *J Infect* 54:e211–e213, 2007.
- Kanafani ZA, Boucher H, Fowler V, et al. Daptomycin compared to standard therapy for the treatment of

"Usual dose" assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

- native valve endocarditis. *Enferm Infecc Microbiol Clin* 28:498–503, 2010.
- King EA, McCoy D, Desai S, et al. Vancomycin-resistant enterococcal bacteraemia and daptomycin: *J Antimicrob Chemother* 66:2112–2118, 2011.
- Kosmidis C, Levine DP. Daptomycin: pharmacology and clinical use. *Expert Opin Pharmacother* 11:615–625, 2010.
- Kullar R, Davis SL, Levine DP, et al. High-dose daptomycin for treatment of complicated gram-positive infections: a large, multicenter, retrospective study. *Pharmacotherapy* 31:527–536, 2011.
- Lichterfeld M, Ferraro MJ, Davis BT. High-dose daptomycin for the treatment of endocarditis caused by *Staphylococcus aureus* with intermediate susceptibility to glycopeptides. *Int J Antimicrob Agents* 35:96, 2010.
- Miller BA, Gray A, Leblanc TW, et al. Acute eosinophilic pneumonia secondary to daptomycin: a report of three cases. *Clin Infect Dis* 50:e63–68, 2010.
- Mohan S, McDermott BP, Cunha BA. Methicillin-resistant *Staphylococcus aureus* (MRSA) prosthetic valve endocarditis (PVE) with paravalvular abscess treated with daptomycin. *Heart & Lung* 34:69–71, 2005.
- Nguyen MH, Eels SJ, Tan J, et al. Pharmacokinetics of daptomycin during cardiopulmonary bypass surgery: A prospective, open-label investigation. *Antimicrob Agents Chemother* 55:2499–2505, 2011.
- Palavecino EI, Burnell JM. False daptomycin-nonsusceptible MIC results by microscan panel relative to Etest results for *Staphylococcus aureus* and enterococci. *J Clin Micro*. 51:281–283, 2013.
- Patel JB, Jevitt LA, Hageman J, et al. An association between reduced susceptibility to daptomycin and reduced susceptibility to vancomycin in *Staphylococcus aureus*. *Clin Infect Dis* 42:1652–3, 2006.
- Patel N, Cardone K, Grabe DW, et al. Use of pharmacokinetic and pharmacodynamic principles to determine optimal administration of daptomycin in patients receiving standardized thrice-weekly hemodialysis. *Antimicrob Agents Chemother* 55:1677–1683, 2011.
- Rangaraj G, Cleveland KO, Gelfand MS. Comparative analysis of daptomycin and vancomycin in the treatment of vertebral osteomyelitis. *Infect Dis Clinical Practice* 22:2019–222, 2014.
- Ritchie ND, Lovering AM, Seaton RA. Daptomycin in synovial fluid during treatment of methicillin-resistant *Staphylococcus aureus* septic arthritis. *J Antimicrob Chemother* 65:1314–1315, 2010.
- Schutt AC, Bohm NM. Multidrug-resistant *Enterococcus faecium* endocarditis treated with combination tigecycline and high-dose daptomycin. *Ann Pharmacother* 43:2108–2112, 2009.
- Stewart PS, Davison WM, Steenbergen JN. Daptomycin rapidly penetrates a *Staphylococcus epidermidis* biofilm. *Antimicrob Agents Chemother* 53:3505–3507, 2009.
- Tascini C, Congiorni C, Cori A, et al. Cardiovascular implantable electronic device endocarditis treated with daptomycin with or without transvenous removal. *Heart & Lung* 41:24–30, 2012.
- Traunmüller F, Schintler MV, Metzler J, et al. Soft tissue and bone penetration abilities of daptomycin in diabetic patients with bacterial foot infections. *J Antimicrob Chemother* 65:1252–1257, 2010.
- Van Hai SJ, Paterson DL, Gosbell IB. Emergence of daptomycin resistance following vancomycin-unresponsive *Staphylococcus aureus* bacteraemia in daptomycin-native patient—a review of the literature. *Eur J Clin Microbiol Infect Dis* 30:603–610, 2011.
- Vilay AM, Grio M, Depestel DD, et al. Daptomycin pharmacokinetics in critically ill patients receiving continuous venovenous hemodialysis. *Crit Care Med* 39:19–25, 2011.
- Wu G, Abraham T, Rapp J, et al. Daptomycin: evaluation of a high-dose treatment strategy. *Int J Antimicrob Agents* 38:192–196, 2011.
- Zhang N, Huang L, Liu S, et al. Neutropenia associated with prolonged daptomycin use. *Clin Infect Dis* 56:1353–1355, 2013.
- Website: [www.pdr.net](http://www.pdr.net)

## Darunavir (Prezista) DRV

**Drug Class:** HIV protease inhibitor.

**Usual Dose:** Treatment naive / treatment experienced with no darunavir resistance-associated substitutions: 800 mg (PO) q24h with ritonavir 100 mg (PO) q24h. Treatment

"Usual dose" assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

experienced / genotyping not available: 600 mg (PO) q12h with ritonavir 100 mg (PO) q12h.

### Pharmacokinetic Parameters:

Peak serum level: 3578 ng/mL

Bioavailability: 37% (alone) 82% (with ritonavir)

Excreted unchanged: 41.2% (feces), 7.7% (urine)

Serum half-life (normal/ESRD): 15/15 hrs

Plasma protein binding: 95%

Volume of distribution ( $V_d$ ): not studied

**Primary Mode of Elimination:** Fecal/Renal

### Dosage Adjustments\*

CrCl 10–50 mL/min	No change
CrCl < 10 mL/min	No data
Post-HD dose	No change
Post-PD dose	No change
CVWH/CVWHD/CVWHDF dose	No change
Mild—moderate hepatic insufficiency	No change
Severe hepatic insufficiency	Use with caution

### Antiretroviral Dosage Adjustments

Efavirenz	No data
Nevirapine	No change
Didanosine	1 hour before or 1 hour after darunavir
Tenofovir	No change
Fosamprenavir	No change
Indinavir	No data
Lopinavir/ritonavir	Avoid
Saquinavir	Avoid
Rifabutin	150 mg (PO) qod

**Drug Interactions:** Indinavir, ketoconazole, nevirapine, tenofovir (↑ darunavir levels); lopinavir/ritonavir, saquinavir, efavirenz (↓ darunavir levels); concomitant administration of darunavir/ritonavir with agents highly-dependent on CYP3A for clearance, astemizole, cisapride, dihydroergotamine, ergonovine, ergotamine, methylegonovine, midazolam, pimoziide, terfenadine, midazolam, triazolam (may ↓ darunavir levels and ↓ effectiveness); sildenafil, vardenafil, tadalafil (↑ PDE-5 inhibitors; sildenafil do not exceed 25 mg in 48 hrs, vardenafil do not exceed 2.5 mg in 72 hrs, or tadalafil do not exceed 10 mg in 72 hrs); clarithromycin (↑ QT<sub>c</sub>).

**Adverse Effects:** Rash, diarrhea, nausea, headache, nasopharyngitis drug induced hepatitis. Use with caution in patients with preexisting liver disease. New-onset or exacerbations of pre-existing diabetes mellitus and hyperglycemia, and increased bleeding in hemophiliacs (class effect). Immune reconstitution syndrome.

**Contraindicated:** lovastatin, simvastatin, atorvastatin (> 20 mg/day), rifampentine, boceprevir, telaprevir.

**Allergic Potential:** High (see comments)

**Safety in Pregnancy:** C

**Comments:** Always take with food (increases AUC, C<sub>max</sub> by approximately 30%). Must be given with ritonavir to boost bioavailability. Darunavir contains a sulfonamide moiety (as do fosamprenavir and tipranavir); use with caution in patients with sulfonamide allergies. astemizole, terfenadine, ergot derivatives dihydroergotamine, ergonovine, ergotamine, methylegonovine, cisapride, pimoziide, midazolam, triazolam, are contraindicated with darunavir. Phenobarbital, phenytoin, carbamazepine, or products containing

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

St. John's wort, rifampin. Lovostatin and simvastatin may be taken with caution.

**Cerebrospinal Fluid Penetration:** No data

#### REFERENCES:

- De Meyer S, Azijn H, Surleraux D, et al. TMC114, a novel human immunodeficiency virus type 1 protease inhibitor active against protease inhibitor-resistant viruses, including a broad range of clinical isolates. *Antimicrob Agents Chemother* 49:2314–2321, 2005.
- Dominique L. N. G, Surleraux T, Abdellah Tahri T, et al. Discovery and selection of TMC114, a next generation HIV-1 protease inhibitor. *J Med Chem* 48:1813–1822, 2005.
- Okoli C, Siccardi M, Thomas-William S, et al. Once Daily maraviroc 300 mg or 150 mg in combination with ritonavir-boosted darunavir 800/100 mg. *J Antimicrob Chemother* 67:671–4, 2012.
- Patterson K, Jennings S, Falcon R, et al. Darunavir, ritonavir, etravirine pharmacokinetics in the cervicovaginal fluid and blood plasma of HIV-infected women. *Antimicrob Agents Chemother* 55:1120–2, 2011.
- Sorbera LA, Castaner J, Bayes M. Darunavir: Anti-HIV agent HIV protease inhibitor. *Drugs of the Future* 30:441–449, 2005.
- Thompson MA, Aberg JA, Cahn P, et al. Antiretroviral treatment of adult HIV infection: 2012 recommendations of the International Antiviral Society — USA panel. *JAMA* 308:387–402, 2012.
- Website: [www.pdr.net](http://www.pdr.net)

## Delavirdine (Rescriptor) DLV

**Drug Class:** HIV NNRTI (non-nucleoside reverse transcriptase inhibitor).

**Usual Dose:** 400 mg (PO) q8h.

**Pharmacokinetic Parameters:**

*Peak serum level:* 35 mcg/ml

*Bioavailability:* 85%

*Excreted unchanged (urine):* 5%

*Serum half-life (normal/ESRD):* 5.8 hrs/no data

*Plasma protein binding:* 98%

*Volume of distribution ( $V_d$ ):* 0.5 L/kg

**Primary Mode of Elimination:** Hepatic

**Dosage Adjustments\***

CrCl 50–80 mL/min	No change
CrCl 10–50 mL/min	No change
CrCl < 10 mL/min	No change
Post-HD dose	None
Post-PD dose	None
CVH/CVHD/CVHDF dose	No change
Moderate hepatic insufficiency	Use with caution
Severe hepatic insufficiency	Use with caution

#### Antiretroviral Dosage Adjustments

Amprenavir	Increased agenerase levels
Efavirenz	No data
Indinavir	Indinavir 600 mg q8h
Lopinavir/ritonavir	No data
Nelfinavir	No information (monitor for neutropenia)
Nevirapine	No data
Ritonavir	Delavirdine: no change; ritonavir: No data
Saquinavir soft-gel	Saquinavir soft-gel 800 mg q8h (monitor transaminases)
Rifampin, rifabutin	Avoid
Statins	Avoid

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose  $\times$  1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.



**Drug Interactions:** With antiretrovirals adjust according to table above. Avoid in combination with rifabutin/rifampin. Contraindicated with astemizole, terfenadine, Dihydroergotamine, ergonovine, ergotamine, methylergonovine, Cisapride, Pimozide, Alprazolam, midazolam, triazolam. Phenytoin may decrease delaviridine levels. Clarithromycin, dapsone, nifedipine, warfarin cause increase in interacting drug levels. Sildenafil (do not exceed 25 mg in 48 hrs), tadalafil (max 10mg/72 hrs); vardenafil (max 2.5 mg/72hrs). St. John's wort substantially reduces delaviridine levels.

**Adverse Effects:** Drug fever/rash, Stevens–Johnson Syndrome (rare), headache, nausea/vomiting, diarrhea, ↑ SGOT/SGPT. May cause immune reconstitution syndrome.

**Allergic Potential:** High

**Safety in Pregnancy:** C

**Comments:** May be taken with or without food, but food decreases absorption by 20%. May disperse four 100-mg tablets in > 3 oz. water to produce slurry; 200-mg tablets should be taken as intact tablets and not used to make an oral solution. Separate dosing with ddl or antacids by 1 hour.

**Cerebrospinal Fluid Penetration:** 0.4%

#### REFERENCES:

Conway B. Initial therapy with protease inhibitor-sparing regimens: Evaluation of nevirapine and delaviridine. *Clin Infect Dis* 2:130–4, 2000.

Demeter LM, Shafer RW, Meehan PM, et al. Delaviridine susceptibilities and associated reverse transcriptase mutations in human immunodeficiency virus type 1 isolates from patients in a phase I/II trial of delaviridine monotherapy (ACTG260). *Antimicrob Agents Chemother* 44:794–7, 2000.

Justesen US, Klitgaard NA, Brosen K, et al. Dose-dependent pharmacokinetics of delaviridine in combination

with amprenavir in healthy volunteers. *J Antimicrob Chemother* 54:206–10, 2004.

Thompson MA, Aberg JA, Cahn P, et al. Antiretroviral treatment of adult HIV infection: 2010 recommendations of the international AIDS society – USA panel. *JAMA* 304:321–333, 2010.

Website: [www.pdr.net](http://www.pdr.net)

## Didanosine EC (Videx) ddl

**Drug Class:** HIV NRTI nucleoside reverse transcriptase inhibitor.

**Usual Dose:** 400 mg (PO) q24h for > 60 kg; 250 (PO) q24h for < 60 kg.

**Pharmacokinetic Parameters:**

*Peak serum level:* 29 mcg/mL

*Bioavailability:* 42%

*Excreted unchanged (urine):* 60%

*Serum half-life (normal/ESRD):* 1.6/4.1 hrs

*Plasma protein binding:* ≤ 5%

*Volume of distribution (V<sub>d</sub>):* 1.1 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\***

CrCl 10–59 mL/min	125 mg (PO) q24h
CrCl < 10 mL/min	125 mg (PO) q24h: > 60 kg Avoid: < 60 kg
Post–HD dose	None
Post–PD dose	None
CVWH/CVWHD/ CVWHD dose	150 mg (PO) q24h [100 mg (PO) q24h]
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	No change

**Drug Interactions:** Alcohol, lamivudine, pentamidine and valproic acid increase the risk of

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

pancreatitis. Patients on tenofovir should have didanosine dose reduced to 250 mg/24 hrs. Concomitant use with tenofovir increases risk of virologic failure. The appropriate dose of didanosine when taken with tenofovir has not been determined in patients with creatinine clearance of  $\leq 60$  ml/min. Peripheral neuropathy is enhanced by stavudine, INH, metronidazole, nitrofurantoin, vincristine and zalcitabine. Dapsone levels are dramatically reduced by powder form especially and increase risk of PCP. Decreases levels of ciprofloxacin delavirdine, indinavir. Methadone will decrease didanosine levels. Coadministration of didanosine with ribavirin is not recommended.

**Adverse Effects:** Fatal and non-fatal pancreatitis have occurred during therapy with didanosine used alone or in combination regimens. Tenofovir can increase this risk as well. Lactic acidosis and severe hepatomegaly with steatosis can also occur especially in pregnant women. Combination with stavudine potentiates this risk and should be avoided especially in pregnancy and in patients with concomitant HCV/HBV. Other effects include diarrhea, peripheral neuropathy, rash and abdominal pain, anemia, leukopenia, alopecia. Depigmentation and optic neuritis have occurred. Patients with underlying liver disease are at increased risk of hepatic toxicity.

**Allergic Potential:** Low

**Safety in Pregnancy:** B

Avoid in pregnancy as it may cause fatal pancreatitis

**Comments:** Available as buffered powder for oral solution and enteric-coated extended-release capsules (Videx EC 400 mg PO q24h). Take 30 minutes before or 2 hours after meal (food decreases serum concentrations by 49%). Avoid in patients with alcoholic cirrhosis/history of pancreatitis. Use with caution in combination with tenofovir, which can potentiate risk of pancreatitis and require a significant dose reduction of ddl. Avoid use with ribavirin.

**Cerebrospinal Fluid Penetration:** 20%

#### REFERENCES:

- Barreiro P, Corbaton A, Nunez M, et al. Tolerance of didanosine as enteric-coated capsules versus buffered tablets. *AIDS Patient Care STDS* 18:329–31, 2004.
- Negredo E, Molto J, Munoz-Moreno JA, et al. Safety and efficacy of once-daily didanosine, tenofovir and nevirapine as a simplification antiretroviral approach. *Antivir Ther* 9:335–42, 2004.
- Perry CM, Balfour JA. Didanosine: An update on its antiviral activity, pharmacokinetic properties, and therapeutic efficacy in the management of HIV disease. *Drugs* 52:928–62, 1996.
- Rathbun RC, Martin ES 3<sup>rd</sup>. Didanosine therapy in patients intolerant of or failing zidovudine therapy. *Ann Pharmacother* 26:1347–51, 1992.
- Thompson MA, Aberg JA, Cahn P, et al. Antiretroviral treatment of adult HIV infection: 2012 recommendations of the International Antiviral society — USA panel. *JAMA* 308:387–402, 2012.
- Website: [www.pdr.net](http://www.pdr.net)

## Dolutegravir (Tivicay) DTG

**Drug class:** HIV integrase strand transfer inhibitor (INSTI).

#### Usual dose:

Treatment-naïve or treatment experienced INSTI-naïve patients: 50 mg (PO) of q24h

Treatment-naïve or treatment experienced INSTI-naïve patients also on efavirenz,

fosamprenavir/ritonavir, tipranavir/ritonavir, or rifampin: 50 mg (PO) of q12h

INSTI-experienced with certain INSTI-associated resistance substitutions or clinically suspected

INSTI resistance: 50 mg (PO) q12h

#### Pharmacokinetic Parameters:

*Peak serum level:* 3.6–4.1 mcg/L

*Bioavailability:* Not applicable

<sup>†</sup>Usual dose<sup>†</sup> assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose  $\times 1$  followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl  $< 10$  mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

Excreted unchanged: 53% feces and 31% urine

Serum half-life (normal/ESRD): 14h/14h

Plasma protein binding: 98.9%

Volume of distribution ( $V_d$ ): 17.4 L/kg

**Primary Mode of Elimination:** Hepatic

**Dosage Adjustments\***

CrCl <10 mL/min	No change
Post-HD dose	No data
Post-PO dose	No data
CWVH dose	No data
Moderate to severe hepatic insufficiency	Avoid (Child-Pugh C)

**Drug Interactions:** Should be taken 2 hours before or 6 hours after taking cation-containing antacids or laxatives, sucralfate, oral iron supplements, oral calcium supplements, or buffered medications; metabolized through UGT1A1 and to a lesser degree through CYP3A and also a substrate of UGT1A3, UGT1A9, BCRP and P-gp. Drugs that induce or inhibit these enzymes or transporters may affect dolutegravir levels; avoid concomitant oxcarbazepine, phenytoin, phenobarbital, carbamazepine, rifampin, St. John's wort; should not be used with etravirine without coadministration of atazanavir/ritonavir, darunavir/ritonavir, or lopinavir/ritonavir; coadministration with nevirapine should be avoided because there are insufficient data to make dosing recommendations; combinations of antiretrovirals that do not include metabolic inducers should be considered where possible for INSTI-experienced patients with certain INSTI-associated resistance substitutions or clinically suspected INSTI resistance.

**Adverse Effects:** Insomnia, headache, diarrhea, increased SGOT/SGPT, lipase, bilirubin hyperglycemia, increased CPK

**Allergic Potential:** High.

**Safety in Pregnancy:** B

**Comments:** Coadministration with dufetilide in contraindicated; hypersensitivity reactions characterized by rash, constitutional findings, and sometimes organ dysfunction, including liver injury, discontinue

dolutegravir if these occur; patients with underlying hepatitis B or C may be at increased risk for worsening or development of transaminase elevations; Safety not established in patients < 40 kg and < 12 yo

#### REFERENCES:

Cottrell ML, Hadzic T, Kashuba AD. Clinical pharmacokinetic, pharmacodynamic and drug-interaction profile of the integrase inhibitor dolutegravir. Clin Pharmacokinet 52:981–994, 2013.

Raffi F, Jaeger H, Quiros-Roldan E, et al. Once-daily dolutegravir versus twice-daily raltegravir in antiretroviral-naïve adults with HIV-1 infection (SPRING-2 study): 96 week results from a randomized, double-blind, non-inferiority trial. Lancet Infect Dis 13:927–935, 2013.

Walmsely SL, Antela A, Clumeck N, et al. Dolutegravir plus abacavir-lamivudine for the treatment of HIV-1 infection. N Engl J Med 369:1807–1818, 2013.

Website: www.pdr.net

## Doripenem (Doribax)

**Drug Class:** Carbapenem.

**Usual Dose:** 1 gm (IV) q8h.

**Spectrum:** (see **Susceptibility Profiles** pp. 198–202).

**Resistance potential:** Low

**Pharmacokinetic Parameters:**

*Peak serum level:* 23 mcg/mL

*Bioavailability:* Not applicable

*Excreted unchanged (urine):* 40%

*Serum half-life (normal/ESRD):* 1/6.20 hrs

*Plasma protein binding:* 8.1%

*Volume of distribution ( $V_d$ ):* 16.8 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\***

CrCl 30–50 mL/min	1 gm (IV) q12h
CrCl 10–30 mL/min	500 mg (IV) q12h
CrCl <10 mL/min	500 mg (IV) q24h
Post-HD dose	250 mg (IV)

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CWVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

Post-HFHD dose	500 mg (IV)
Post-PD dose	500 mg (IV)
CVVH/CVVHD/ CVVHDf dose	500 mg (IV) q12h
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	No change

**Drug Interactions:** Valproic acid (↓ valproic acid levels).

**Adverse Effects:** Headache, nausea, diarrhea, rash, phlebitis, pneumonitis if administered by inhalation (do not give by inhalation).

**Allergic Potential:** Anaphylaxis (rare)

**Safety in Pregnancy:** B

**Comments:** cSSSIs, cIAls, or UTIs dose:

500 mg (IV) q8h. Infuse over 1 hour. **MDR P.**

**aeruginosa, MDR Acinetobacter baumannii/nosocomial pneumonia dose:** 1 gm (IV) q8h.

**For NP/VAP, infuse IV over 4 hours.**

**Unlike meropenem, doripenem should not be used for meningitis.**

**Cerebrospinal Fluid Penetration:** < 1%

**Bile Penetration:** 117%

(also see **Antibiotic Pearls & Pitfalls** p. 511).

#### REFERENCES:

- Anderson DL. Doripenem. *Drugs for Today* 42:399–404, 2006.
- Betriu C, Gómez M, López-Fabal F, et al. Activity of doripenem against extended-spectrum beta-lactamase-producing Enterobacteriaceae and *Pseudomonas aeruginosa* isolates. *Eur J Clin Microbiol Infect Dis* 29:1179–1181, 2010.
- Castanheira M, RN Jones, DM Livermore. Antimicrobial activities of doripenem and other carbapenems against *Pseudomonas aeruginosa*, other nonfermentative bacilli, and *Aeromonas* spp. *Diagn Microbiol Infect Dis* 63:426–433, 2012.

Hellwig TR, Onisk ML, Chapman BA. Potential interaction between valproic acid and doripenem. *Curr Drug Saf* 6:54–58, 2011.

Lister PD. Carbapenems in the USA: focus on doripenem. *Expert Rev Anti Infect Ther* 5:793–809, 2007.

Pankuch GA, Seifert H, Appelbaum PC. Activity of doripenem with and without levofloxacin, amikacin, and colistin against *Pseudomonas aeruginosa* and *Acinetobacter baumannii*. *Diagn Microbiol Infect Dis* 67:191–197, 2010.

Paterson DL, DePestel DD. Doripenem. *Clin Infect Dis* 49:291–298, 2009.

Wall GC, Nayima VA, Neumeister KM. Assessment of hypersensitivity reactions in patients receiving carbapenem antibiotics who report antibiotics a history of penicillin. *J Chemo*. 26:150-153, 2014.

Website: [www.pdr.net](http://www.pdr.net)

## Doxycycline (Vibramycin, Vibra-tabs)

**Drug Class:** 2<sup>nd</sup> generation IV/PO tetracycline.

**Usual Dose:** 100 mg (IV/PO) q12h or 200 mg (IV/PO) q24h (PI dose). For **serious systemic infections**, begin with a **Loading Regimen:**

200 mg (IV/PO) q12h × 3 days (**PK dose**), then continue at same dose **or** decrease to

**Maintenance dose:** 100 mg (IV/PO) q12h to complete therapy.

**Spectrum:** (see **Susceptibility Profiles** pp. 186–190).

**Resistance Potential:** Low

**Pharmacokinetic Parameters:**

*Peak serum level:* 100/200 mg = 4/8 mcg/mL

*Bioavailability:* 93%

*Excreted unchanged (urine):* 40%

*Serum half-life (normal/ESRD):* 18–22 hrs/18–22 hrs

*Plasma protein binding:* 82%

*Volume of distribution (V<sub>d</sub>):* 0.75 L/kg

**Primary Mode of Elimination:** Hepatic

**Dosage Adjustments\***

PI = package insert

PK = pharmacokinetic dose

CrCl < 10 mL/min	No change
Post-HD or PD dose	None
CVH/CVHD/ CWHDF dose	No change
Moderate—severe hepatic insufficiency	No change

**Drug Interactions:** Antacids, Al<sup>++</sup>, Ca<sup>++</sup>, Fe<sup>++</sup>, Mg<sup>++</sup>, Zn<sup>++</sup>, multivitamins, sucralfate (↓ doxycycline absorption); barbiturates, carbamazepine, phenytoin (↓ doxycycline half-life); bicarbonate (↓ doxycycline absorption, ↑ doxycycline clearance); warfarin (↑ INR).

**Adverse Effects:** Nausea if not taken with food. Phlebitis if given IV in inadequate volume. Avoid in pregnancy. **Does not ↑ SGOT/SGPT.** Photosensitivity rare.

**Allergic Potential:** Low

**Safety in Pregnancy:** D

**Comments:** Tetracycline susceptibilities do not predict doxycycline effectiveness against *S. pneumoniae*, MSSA, MRSA. Due to high urinary concentrations doxycycline IV/PO effective in treating lower UTIs due to GNBS, e.g., *P. aeruginosa* even those resistant by susceptibility testing (related to achievable serum concentrations).

Protective against *C. difficile*. Absorption minimally effected by iron, bismuth, milk, or antacids containing Ca<sup>++</sup>, Mg<sup>++</sup>, or Al<sup>++</sup>. Serum half life increases with multiple doses. **For MSSA, MRSA use minocycline instead of doxycycline**

Meningeal dose = 200 mg (IV/PO) q12h.

**Cerebrospinal Fluid Penetration:**

Non-Inflamed meninges = 25%

Inflamed meninges = 25%

**Bile Penetration:** 3000%

(also see **Antibiotic Pearls & Pitfalls** p. 512).

#### REFERENCES:

- Bernadino ALF, Kaushal D, Philipp MT. The antibiotics doxycycline and minocycline inhibit the inflammatory responses to the Lyme disease spirochete *Borrelia burgdorferi*. *Journal of Infectious Diseases* 199;1379–1388, 2009.
- Borg R, Dotevall L, Hagberg L, et al. Intravenous ceftriaxone compared with oral doxycycline for the treatment of Lyme neuroborreliosis. *Scand J Infect Dis* 37:449–454, 2005.
- Cunha BA. Minocycline versus doxycycline for methicillin-resistant *Staphylococcus aureus* (MRSA): in vitro susceptibility versus in vivo effectiveness. *Int J Antimicrob Agents* 35:517–8, 2012.
- Cunha BA. Oral doxycycline for non-systemic urinary tract infections (UTIs) due to *P. aeruginosa* and other Gram negative uropathogens. *Eur J Clin Microbiol Infect Dis* 31:2865–8, 2012.
- Cunha BA. Doxycycline for nursing home acquired pneumonia (NHAP). *Scandinavian Journal of Infectious Disease* 41:77–78, 2009.
- Cunha BA. New uses for older antibiotics: nitrofurantoin, amikacin, colistin, polymyxin B, doxycycline, and minocycline revisited. *Med Clin North Am* 90:1089–1107, 2006.
- Cunha BA. Oral antibiotic therapy of serious systemic infections. *Med Clin N Am* 90:1197–1222, 2006.
- Cunha BA. Doxycycline for community-acquired pneumonia. *Clin Infect Dis* 37:870, 2003.
- Cunha BA, Domenico PD, Cunha CB. Pharmacodynamics of doxycycline. *Clin Micro Infect Dis* 6:270–3, 2000.
- Cunha BA. Doxycycline re-revisited. *Arch Intern Med* 159:1006–7, 1999.
- Cunha BA. Doxycycline re-revisited. *Arch Intern Med* 159:1006–7, 1999.
- Cunha BA. Doxycycline. *Antibiotics for Clinicians* 3:21–33, 1999.
- Doernberg SB, Winston LG, Deck LH, et al. Does Doxycycline protect against development of *Clostridium difficile* infection? *Clin Infect Dis* 55:615–20, 2012.
- Haggerty CL, Ness RB. Newest approaches to treatment of pelvic inflammatory disease: a review of recent randomized clinical trials. *Clin Infect Dis* 44:953–960, 2007.

Pl = package insert

PK = pharmacokinetic dose

- Hasin T, Davidovitch N, Cohen R, et al. Postexposure treatment with doxycycline for the prevention of tick-borne relapsing fever. *N Engl J Med* 355:147–55, 2006.
- Johnson JR. Doxycycline for treatment of community-acquired pneumonia. *Clin Infect Dis* 35:632, 2002.
- Jones RN, Sader HS, Fritsche TR. Doxycycline use for community-acquired pneumonia: contemporary in vitro spectrum of activity against *Streptococcus pneumoniae*. *Diagn Microbiol Infect Dis* 49:147–9, 2004.
- Karkkonen K, Stiernstedt SH, Karisson M. Follow-up of patients treated with oral doxycycline for Lyme neuroborreliosis. *Scand J Infect Dis* 33:259–262, 2001.
- Kong FYS, Tabrizi SN, Law M, et al. Azithromycin Versus Doxycycline for the Treatment of Genital Chlamydia Infection: A Meta-analysis of Randomized Controlled Trials. *Clin Infect Dis* 59:193–205, 2014.
- Somas KC, Brun M, Causse A, et al. Efficacy of a ceftriaxone and doxycycline in the treatment of early syphilis. *Magn Reson Imaging* 30:271–82, 2012.
- Ranjbar M, Keramat F, Mamani M, et al. Comparison between doxycycline-rifampin-amikacin and doxycycline regimens in the treatment of brucellosis. *Int J Infect Dis* 11:152–156, 2007.
- Raoult D. Doxycycline for *Mansonella perstans* Infection. *N Engl J Med* 362:272–273, 2010.
- Roushan MRH, Mohraz M, Hajiahmadi M, et al. Efficacy of gentamycin plus doxycycline versus streptomycin plus doxycycline in the treatment of brucellosis in humans. *Clin Infect Dis* 42:1075–80, 2006.
- Schwartz BS, Graber CJ, Diep BA, et al. Doxycycline, not minocycline, induces its own resistance in multidrug-resistant community-associated methicillin-resistant *Staphylococcus aureus* clone USA300. *Clin Infect Dis* 48:1483–1484, 2009.
- Shea KW, Ueno Y, Abumustafa F, Cunha BA. Doxycycline activity against *Streptococcus pneumoniae*. *Chest* 107:1775–6, 1995.
- Tan KR, Magill AJ, Parise ME, et al. Doxycycline for malaria chemoprophylaxis and treatment: report from the CDC expert meeting on malaria chemoprophylaxis. *Am J Trop Med Hyg* 84:517–531, 2011.
- Teh B, Grayson ML, Johnson PD, et al. Doxycycline vs. macrolides in combination therapy for treatment of

community-acquired pneumonia. *Clin Microbiol Infect* 18:71–3, 2012.

Turner JD, Mand S, Debrah AY, et al. A randomized, double-blind clinical trial of a 3-week course of doxycycline plus albendazole and ivermectin for the treatment of *Wuchereria bancrofti* infection. *Clin Infect Dis* 42:1081–9, 2006.

Volovitz B, Shkap R, Amir J, et al. Absence of tooth staining with doxycycline treatment in young children. *Clin Pediatr* 46:121–126, 2007.

Website: [www.pdr.net](http://www.pdr.net)

## Efavirenz (Sustiva) EFV

**Drug Class:** HIV NNRTI (non-nucleoside reverse transcriptase inhibitor).

**Usual Dose:** 600 mg (PO) q24h.

**Pharmacokinetic Parameters:**

*Peak serum level:* 12.9 mcg/mL

*Bioavailability:* Increased with food

*Excreted unchanged (urine):* 14–34%

*Serum half-life (normal/ESRD):* 40–55 hrs/no data

*Plasma protein binding:* 99%

*Volume of distribution ( $V_d$ ):* No data

**Primary Mode of Elimination:** Hepatic

**Dosage Adjustments\***

CrCl < 60 mL/min	No change
Post-HD or PD dose	None
CVVH/CVVHD/CVVHDF dose	No change
Moderate—severe hepatic insufficiency	No change

## Antiretroviral Dosage Adjustments

Delavirdine	No data
Indinavir	Indinavir 1000 mg q8h
Lopinavir/ritonavir (l/r)	Consider l/r 533/133 mg q12h in PI-experienced patients
Nelfinavir	No change

\*Usual dose\* assumes normal renal/hepatic function. \*For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

Nevirapine	No data
Ritonavir	Ritonavir 600 mg q12h (500 mg q12h for intolerance)
Saquinavir	Avoid use as sole PI
Rifampin	No change
Rifabutin	Rifabutin 450–600 mg q24h or 600 mg 2–3x/week if not on protease inhibitor
Telaprevir increase telaprevir 1125 mg q7–9h	
Voriconazole: give efavirenz 300 mg q24h, voriconazole 400 mg q12h	

**Drug Interactions:** Antiretrovirals, rifabutin, rifampin (see dose adjustment grid, above); astemizole, terfenadine, cisapride, ergotamine, midazolam, carbamazepine, phenobarbital, phenytoin (monitor anticonvulsant levels; use with caution); verapamil, diltiazem (↓ calcium channel blocker levels) caspofungin, itraconazole, posaconazole (↓ anti fungal levels); methadone, clarithromycin (↓ interacting drug levels; titrate methadone dose to effect; consider using azithromycin instead of clarithromycin).

**Contraindications:** Triazolam, rifampentine, boceprevir.

**Adverse Effects:** Drug fever/rash, CNS symptoms (nightmares, dizziness, neuropsychiatric symptoms, difficulty concentrating, somnolence), ↑ SGOT/SGPT, E. multiforme/Steven's-Johnson Syndrome (rare), false positive cannabinoid test.

**Allergic Potential:** High

**Safety in Pregnancy:** D

**Comments:** Rash/CNS symptoms usually resolve spontaneously over 2–4 weeks. Take at bedtime. Avoid taking after high fat meals (levels ↑ 50%). 600-mg dose available as single tablet.

**Cerebrospinal Fluid Penetration:** 1%

#### REFERENCES:

- Haas DW, Fessel WJ, Delapenha RA, et al. Therapy with efavirenz plus indinavir in patients with extensive prior nucleoside reverse-transcriptase inhibitor experience: A randomized, double-blind, placebo-controlled trial. *J Infect Dis* 183:392–400, 2001.
- la Porte CJ, de Graaff-Teulen MJ, Colbers EP, et al. Effect of efavirenz treatment on the pharmacokinetics of nelfinavir boosted by ritonavir in healthy volunteers. *Br J Clin Pharmacol* 58:632–40, 2004.
- Marzolini C, Telenti A, Decosterd LA, et al. Efavirenz plasma levels can predict treatment failure and central nervous system side effects in HIV-1-infected patients. *AIDS* 15:71–5, 2001.
- McComsey GA, Kitch D, Sax PE, et al. Peripheral and central fat changes in subjects randomized to abacavir-lamivudine or tenofovir-emtricitabine with atazanavir-ritonavir or efavirenz: ACTG Study A5224s. *Clin Infect Dis* 53:185–96, 2011.
- Thompson MA, Aberg JA, Cahn P, et al. Antiretroviral treatment of adult HIV infection: 2012 recommendations of the International Antiviral Society — USA panel. *JAMA* 308:387–402, 2012.
- Young J, Schafer J, Fux CA, et al. Renal function in patients with HIV starting therapy with tenofovir and either efavirenz, lopinavir or atazanavir. *AIDS* 26:567–75, 2012. Website: [www.pdr.net](http://www.pdr.net)

## Efavirenz + Emtricitabine + Tenofovir disoproxil fumarate (Atripla) EFV/FTC/TDF

**Drug Class:** HIV

**Usual Dose:** 1 tablet (efavirenz 600 mg/ emtricitabine 200 mg/tenofovir 300 mg) (PO) q24h on an empty stomach.

**Pharmacokinetic Parameters:**

*Peak serum level:* 4.0/1.8 mcg/mL/296 ng/mL

*Bioavailability:* NR/93%/25%

"Usual dose" assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

Excreted unchanged: < 1% unchanged and 14–30% as metabolites/86%/32%

Serum half-life (normal/ESRD): (40–55 hrs/~10 hrs on hemodialysis)/(10 hrs/extended)/(17 hrs/no data)

Plasma protein binding: 99/< 4/< 0.7%

Volume of distribution ( $V_d$ ): NR/NR/1.2 L/kg

**Primary Mode of Elimination:** Hepatic/Renal/  
Renal

### Dosage Adjustments\*

CrCl 50–80 mL/min	No change
CrCl 10–50 mL/min	Avoid
CrCl < 10 mL/min	Avoid
Post-HD dose	Avoid
Post-PD dose	Avoid
CWVH/CVVHD/ CWHDF dose	Avoid
Mild hepatic insufficiency	No data
Moderate—severe hepatic insufficiency	No data

### Antiretroviral Dosage Adjustments

Fosamprenavir/ ritonavir	An additional 100 mg/day (300 mg total) of ritonavir is recommended when ATRIPLA is administered with fosamprenavir/ritonavir q24h. No change in ritonavir dose when ATRIPLA is administered with fosamprenavir/ritonavir q12h
Atazanavir	Atazanavir 300 mg q24h Ritonavir 100 mg q12h
Indinavir	Indinavir 1000 mg q8h

Lopinavir/ ritonavir	Increase lopinavir/ritonavir to 600/150 mg (3 tablets) q12h
Ritonavir	No data
Saquinavir	Avoid
Didanosine	Avoid
Rifabutin	Rifabutin 450–600 mg q24h or 600 mg 2–3x/week if not on protease inhibitor
Rifampin	No data

**Drug Interactions:** Antiretrovirals, rifabutin (see dose adjustment grid above); astemizole, cisapride, ergotamine, methylergonovine, midazolam, triazolam, St John's Wort (↓ efavirenz levels), voriconazole (↓ voriconazole levels; avoid); caspofungin (↓ caspofungin levels); carbamazepine, phenytoin, phenobarbital (monitor anticonvulsant levels; use with caution; potential for ↓ efavirenz levels); statins (may ↓ statin levels); methadone, (↓ methadone levels); clarithromycin (may ↓ clarithromycin effectiveness, consider using azithromycin). Should not be administered concurrently with astemizole, cisapride, midazolam, triazolam, or ergot derivatives. Significantly reduces voriconazole drug levels.

**Adverse Effects:** Dizziness, insomnia, impaired concentration, somnolence, abnormal dreaming, euphoria, confusion, agitation, amnesia, hallucinations, stupor, abnormal thinking, and depersonalization. Psychiatric symptoms of depression suicidal ideation, nonfatal suicide attempts, aggressive behavior, paranoid reactions, and manic reactions. Mild to moderate skin rash, Steven's Johnson syndrome. Pancreatitis can occur. Increase in lipids and

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.



cholesterol. Immune reconstitution Syndrome and redistribution/accumulation of body fat have been reported in patients treated with combination antiretroviral therapy.

**Allergic Potential:** High

**Safety in Pregnancy:** D

**Comments:** Rash/CNS effects usually resolve in a few weeks. Take at bedtime on empty stomach. High fat meals can ↑ efavirenz by 50%. Use with caution in patients with history of seizures (↑ risk of convulsions). May cause false positive tests for cannabinoids.

**Cerebrospinal Fluid Penetration:** 1%

#### REFERENCES:

Cooper DA, Heera J, Goodrich J, et al. Maraviroc versus efavirenz, both in combination with zidovudine-lamivudine, for the treatment of antiretroviral-naïve subjects with CCR5-tropic HIV-1 infection. *J Infect Dis* 201;803–813, 2010.

Gallant JE, DeJesus E, Arribas JR, et al: Tenofovir DF, emtricitabine, and efavirenz vs. zidovudine, lamivudine, and efavirenz for HIV. *N Engl J Med* 354:251–260, 2006.

Thompson MA, Aberg JA, Cahn P, et al. Antiretroviral treatment of adult HIV infection: 2012 recommendations of the International Antiviral Society — USA panel. *JAMA* 308:387–402, 2012.

Website: [www.pdr.net](http://www.pdr.net)

## Elvitegravir + Cobicistat + Emtricitabine + Tenofovir disoproxil fumarate (Stribild) EVG/COBI/FTC/TDF

**Drug class Antiretroviral agent:** HIV Integrase strand transfer inhibitor (elvitegravir 150 mg), pharmacokinetic enhancer (cobicistat 150 mg), and 2 nucleos(t)ide analog HIV-1 reverse transcriptase inhibitors (emtricitabine 200 mg and tenofovir disoproxil fumarate 300 mg)

**Usual dose:** 1 tablet once daily with food

#### Pharmacokinetic Parameters:

*Peak serum level:* 1.7/1.1/1.9/0.45 mcg/mL

*Bioavailability:* Increased with food

*Excreted unchanged:* minimal for elvitegravir and cobicistat, ~ 100% for emtricitabine and tenofovir disoproxil fumarate

*Serum half-life (normal/ESRD):* 9/3/12/11 (not recommended in renal failure)

*Plasma protein binding:* 98/98/4/0.7%

*Volume of distribution (V<sub>d</sub>):* No data

**Primary Mode of Elimination:** Hepatic

(elvitegravir and cobicistat); Renal (emtricitabine and tenofovir disoproxil fumarate)

#### Dosage Adjustments\*

CrCl < 50 mL/min	Avoid
Post-HD dose	Avoid
Post-PD dose	Avoid
CVVH dose	Avoid
Severe hepatic insufficiency	Avoid

**Drug Interactions:** Inhibitor of CYP3A, CYP2D6 and CYP2C9; and p-glycoprotein; metabolized by CYP3A and CYP2D6; antacids (may ↓ elvitegravir absorption).

**Contraindications:** nephrotoxic drugs

**Adverse Effects:** nausea, diarrhea, new onset/worsening of renal failure, decreased bone mineral density redistribution/accumulation of body fat, immune reconstitution syndrome, lactic acidosis and hepatomegaly

**Allergic Potential:** not available

**Safety in Pregnancy:** B

**Comments:** not approved for HIV patients co-infected with HBV

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

**REFERENCES:**

DeJesus E, Rockstroh JK, Henry K, et al. Co-formulated elvitegravir, cobicistat, emtricitabine, and tenofovir disoproxil fumarate versus ritonavirboosted atazanavir plus co-formulated emtricitabine and tenofovir disoproxil fumarate for initial treatment of HIV-1 infection: a randomised, double-blind, phase 3, non-inferiority trial. *Lancet* 379:2429–38, 2012.

Desimmmie BA, Schrijvers R, Debyser Z. Elvitegravir: a once daily alternative to raltegravir. *Lancet Infect Dis* 12:3–5, 2012.

Olin JL, Spooner LM, Klibanov OM. Elvitegravir/Cobicistat/Emtricitabine/Tenofovir Disoproxil Fumarate Single Tablet for HIV-1 Infection Treatment. *Ann Pharmacotherapy* 46: epub November 7, 2012.

Sax PE, DeJesus E, Mills A, et al. Co-formulated elvitegravir, cobicistat, emtricitabine, and tenofovir versus co-formulated efavirenz, emtricitabine, and tenofovir for initial treatment of HIV-1 infection: a randomised, double-blind, phase 3 trial, analysis of results after 48 weeks. *Lancet* 379:2439–48, 2012.

Website: www.pdr.net

**Emtricitabine (Emtriva) FTC**

**Drug Class:** HIV NRTI (nucleoside reverse transcriptase inhibitor).

**Usual Dose:** 200 mg (PO) q24h.

**Pharmacokinetic Parameters:**

*Peak serum level:* 1.8 mcg/mL

*Bioavailability:* 93%

*Excreted unchanged (urine):* 86%

*Serum half-life (normal/ESRD):* 10 hrs/extended

*Plasma protein binding:* 4%

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\***

CrCl $\geq$ 50 mL/min	200 mg (PO) q24h
CrCl 30–49 mL/min	200 mg (PO) q48h

CrCl 15–29 mL/min	200 mg (PO) q72h
CrCl < 15 mL/min	200 mg (PO) q96h
Post–HD dose	200 mg (PO)
Post–PD dose	200 mg (PO)
CWVH/CWVHD/ CWVHDF dose	200 mg (PO) q72h
Moderate—severe hepatic insufficiency	No change

**Drug Interactions:** No significant interactions with indinavir, stavudine, zidovudine, famciclovir, tenofovir.

**Adverse Effects: Lactic acidosis hepatomegaly with steatosis, post-treatment exacerbation of hepatitis B.**

Headache, diarrhea, nausea and rash as well as abdominal pain, fatigue, dizziness abnormal dreams. Rare cases of skin hyperpigmentation on palms and soles. Immune reconstitution syndrome may occur.

**Allergic Potential:** Low

**Safety in Pregnancy:** B

**Comments:** May be taken with or without food. Does not inhibit CYP450 enzymes. Mean intracellular half-life of 39 hours. Potential cross-resistance to lamivudine and zalcitabine. Low affinity for DNA polymerase- $\gamma$ . Should not be coadministered with drugs containing lamivudine.

**Cerebrospinal Fluid Penetration:** No data

**REFERENCES:**

Dando TM, Wagstaff AJ. Emtricitabine/tenofovir disoproxil fumarate. *Drugs* 64:2075–82, 2004.

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose  $\times$  1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CWVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

- Gallant JE, DeJesus D, Arribas JR, et al. Tenofovir DF, emtricitabine, and efavirenz vs. zidovudine, lamivudine, and efavirenz for HIV. *N Engl J Med* 354:251–60, 2006.
- Lim SG, Ng TN, Kung N, et al. A double-blind placebo-controlled study of emtricitabine in chronic hepatitis B. *Arch Intern Med* 166:49–56, 2006.
- Saag MS. Emtricitabine, a new antiretroviral agent with activity against HIV and hepatitis B virus. *Clin Infect Dis* 42:128–31, 2006.
- Sax PE, Tierney C, Collier AC, et al. Abacavir/lamivudine versus tenofovir DF/emtricitabine as part of combination regimens for initial treatment of HIV: final results. *J Infect Dis* 204:1191–201, 2011.
- Thompson MA, Aberg JA, Cahn P, et al. Antiretroviral treatment of adult HIV infection: 2012 recommendations of the International Antiviral Society — USA panel. *JAMA* 308:387–402, 2012.
- Website: [www.pdr.net](http://www.pdr.net)

## Emtricitabine + Tenofovir disoproxil fumarate (Truvada) FTC/TDF

**Drug Class:** HIV NRTI (nucleoside reverse transcriptase inhibitor) + nucleotide analogue.

**Usual Dose:** One tablet (PO) q24h (each tablet contains 200 mg of emtricitabine + 300 mg of tenofovir).

**Pharmacokinetic Parameters:**

*Peak serum level:* 1.8/0.3 mcg/L

*Bioavailability:* 93%/27% if fasting (39% with high fat meal)

*Excreted unchanged (urine):* 86/32%

*Serum half-life (normal/ESRD):* (10 hrs/extended)/(17 hrs/no data)

*Plasma protein binding:* 4/0.7–7.2%

*Volume of distribution ( $V_d$ ):* no data/1.3 L/kg

**Primary Mode of Elimination:** Renal/Renal  
**Dosage Adjustments\***

CrCl $\geq$ 50 mL/min	No change
CrCl 30–49 mL/min	One capsule (PO) q48h
CrCl 15–29 mL/min	Avoid
CrCl < 15 mL/min	Avoid
Post–HD dose	Avoid
Post–PD dose	Avoid
CVH/CVHD/ CVHDF dose	Avoid
Moderate—severe hepatic insufficiency	No change

**Drug Interactions:** No significant interactions with indinavir, stavudine, zidovudine, famciclovir, lamivudine, lopinavir/ritonavir, efavirenz, methadone, oral contraceptives. Tenofovir  $\uparrow$  didanosine levels which may result in severe pancreatitis. Tenofovir reduces systemic exposure to atazanavir; whenever the two are co-administered, the recommended dose of atazanavir is 300 mg once daily with ritonavir 100 mg once daily. Atazanavir may increase tenofovir levels. Avoid coadministration with didanosine.

**Adverse Effects:** Headache, diarrhea, nausea, vomiting, GI upset, rash, lactic acidosis with hepatic steatosis (rare but potentially life-threatening with NRTI's). Decreases bone mineral density. May result in flare of hepatitis after stopping medication in patients with chronic hepatitis B infection. Cases of osteomalacia and renal tubular acidosis have occurred. Renal dysfunction is more common in diabetic patients. Immune reconstitution syndrome and redistribution/accumulation of body fat have been reported in patients treated with combination antiretroviral therapy.

**Allergic Potential:** Low  
**Safety in Pregnancy:** B

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose  $\times$  1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

**Comments:** May be taken with or without food. Does not inhibit CYP450 enzymes. Mean intracellular half-life with emtricitabine is 39 hours. Potential cross-resistance to lamivudine, zalcitabine, abacavir, didanosine. Low affinity for DNA polymerase-gamma.

**Cerebrospinal Fluid Penetration:** No data

#### REFERENCES:

Dando TM, Wagstaff AJ. Emtricitabine/tenofovir disoproxil fumarate. *Drugs* 64:2075–82, 2004.

Thompson MA, Aberg JA, Cahn P, et al. Antiretroviral treatment of adult HIV infection: 2012 recommendations of the International Antiviral Society — USA panel. *JAMA* 308:387–402, 2012.

Website: [www.pdr.net](http://www.pdr.net)

## Emtricitabine/Rilpivirine/ Tenofovir (Complera) FTC/RPV/TDF

**Drug Class:** HIV 2 NRTI/1 NNRTI.

**Usual Dose:** 1 tablet (PO) q24h (contains 200 mg emtricitabine, 25 mg rilpivirine, 300 mg tenofovir).

**Pharmacokinetic Parameters:**

*Peak serum level:* 1.8/1/0.3 mcg/L

*Bioavailability:* 93/No data/25%

*Excreted unchanged:* 86/1/75%

*Serum half-life (normal):* 10/50/17 hrs

*Plasma protein binding:* unknown/99/1%

*Volume of distribution ( $V_d$ ):* no data/no data/no data

**Primary Mode of Elimination:** Renal/hepatic/renal

**Dosage Adjustments\***

CrCl < 50 mL/min	Avoid
Post-HD dose	Avoid
Post-PD dose	Avoid
CVVH/CVVHD/ CVVHDF dose	Avoid
Mild—moderate hepatic insufficiency	No change
Severe hepatic insufficiency	No data

**Drug Interactions:** Do not administer with other NNRTIs. Antacid and H<sub>2</sub> blockers should be separated. Any drug that induces or inhibits CYP3A has the potential to increase or decrease rilpivirine levels and should only be used if benefit outweighs risk, clarithromycin (↓ clarithromycin levels).

**Contraindications:** carbamazepine, oxycarbazepine, phenobarbital, phenytoin, rifabutin, rifampin, rifapentine, proton pump inhibitors, dexamethasone (except single dose), St. John's Wort.

**Adverse Effects:** Nausea, diarrhea, fatigue, headache, dizziness, depression, insomnia, abnormal dreams, rash, **lactic acidosis/severe hepatomegaly with steatosis and post-treatment acute exacerbation of hepatitis B**, immune reconstitution syndrome, fat redistribution syndrome.

**Allergic Potential:** Low

**Safety in Pregnancy:** B

**Comments:** Monitor BMD in patients at risk for pathological fractures. *Not approved for patients co-infected with HBV and HIV.*

**Cerebrospinal Fluid Penetration:** No data

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

**REFERENCES:**

Molina JM, Cahn P, Ginzstein B, et al. Rilipivirine versus efavirenz with tenofovir and emtricitabine in treatment-naïve adults infected with HIV-1 (ECHO): a phase 3 randomised double-blind active-controlled trial. *Lancet Inf Dis* 378:238–246, 2011.

Thompson MA, Aberg JA, Cahn P, et al. Antiretroviral treatment of adult HIV infection: 2012 recommendations of the International Antiviral Society — USA panel. *JAMA* 308:387–402, 2012.

Website: [www.pdr.net](http://www.pdr.net)

**Enfuvirtide (Fuzeon) ENF, T20**

**Drug Class:** HIV fusion inhibitor.

**Usual Dose:** 90 mg (SC) q12h.

**Pharmacokinetic Parameters:**

*Peak serum level:* 4.9 mcg/mL

*Bioavailability:* 84.3%

*Serum half-life (normal/ESRD):* 3.8 hrs/no data

*Plasma protein binding:* 92%

*Volume of distribution ( $V_d$ ):* 5.5 L/kg

**Primary Mode of Elimination:** Metabolized

**Dosage Adjustments\***

CrCl > 35 mL/min	No change
CrCl < 35 mL/min	No change
Post-HD dose	No change
Post-PD dose	No change
CWH/CVHD/ CWHDF dose	No change
Moderate—severe hepatic insufficiency	No change

**Drug Interactions:** No clinically significant interactions with other antiretrovirals. Does not inhibit CYP450 enzymes.

**Adverse Effects:** Local injection site reactions are common. Diarrhea, nausea, fatigue may occur. Laboratory abnormalities include mild/transient

eosinophilia. Pneumonia may occur, but cause is unclear and may not be due to drug therapy. Pancreatitis, myalgia, conjunctivitis (rare).

**Allergic Potential:** Hypersensitivity reactions may occur, including fever, chills, hypotension, rash, ↑ serum transaminases. Do not rechallenge following a hypersensitivity reaction

**Safety in Pregnancy:** B

**Comments:** Enfuvirtide interferes with entry of HIV-1 into cells by blocking fusion of HIV-1 and CD<sub>4</sub> cellular membranes by binding to HR1 in the gp41 subunit of the HIV-1 envelope glycoprotein. Additive/synergistic with NRTI's, NNRTI's, and PI's, and no cross resistance to other antiretrovirals in cell culture. Reconstitute in 1.1 mL of sterile water. SC injection should be given into upper arm, anterior thigh, or abdomen. Rotate injection sites; do not inject into moles, scars, bruises. After reconstitution, use immediately or refrigerate and use within 24 hours (no preservatives added).

**REFERENCES:**

Coleman CI, Musial BL, Ross J. Enfuvirtide: the first fusion inhibitor for the treatment of patients with HIV-1 infection. *Formulary* 38:204–222, 2003.

Kilby JM, Lalezari JP, Eron JJ, et al. The safety, plasma pharmacokinetics, and antiviral activity of subcutaneous enfuvirtide (T-20), a peptide inhibitor of gp41-mediated virus fusion, in HIV-infected adults. *AIDS Res Hum Retroviruses* 18:685–93, 2002.

Lalezari JP, Henry K, O'Hearn M, et al. TORO 1 Study Group. Enfuvirtide, an HIV-1 fusion inhibitor, for drug-resistant HIV infection in North and South America. *N Engl J Med* 348:2175–85, 2003.

Leen C, Wat C, Nieforth K. Pharmacokinetics of enfuvirtide in a patient with impaired renal function. *Clin Infect Dis* 4:339–55, 2004.

Website: [www.pdr.net](http://www.pdr.net)

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

## Entecavir (Baraclude)

**Drug Class:** HBV nucleoside analogue.

**Usual Dose:** 0.5 mg (PO) q24h; 1 mg (PO) q24h for lamivudine-refractory patients or with decompensated liver disease. (Take without food)

**Pharmacokinetic Parameters:**

*Peak serum level:* 4.2 ng/mL (for 0.5 mg dose)

*Bioavailability:* Similar for tablet and oral solution

*Excreted unchanged (urine):* 62–73%

*Serum half-life (normal/ESRD):* 128–139 hrs

*Plasma protein binding:* No information

*Volume of distribution ( $V_d$ ):* No information

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\*** (based on 0.5 mg dose)

CrCl > 50 mL/min	No change
CrCl 30–50 mL/min	0.25 mg (PO) q24h
CrCl 10–30 mL/min	0.15 mg (PO) q24h
CrCl < 10 mL/min	0.05 mg (PO) q24h
Post-HD	0.05 mg (PO or 0.5 mg (PO) q week)
Post-PD	0.05 mg (PO)
Moderate—severe hepatic insufficiency	No change

**Drug Interactions:** Cyclosporin, tacrolimus (may ↑ entecavir levels).

**Adverse Effects:** Fatigue, headache, upper respiratory tract infection, upper abdominal pain, ↑ ALT, cough, nausea.

**Allergic Potential:** No information

**Safety in Pregnancy:** C

**Comments:** Used to treat chronic HBV. Effective in lamivudine-resistant patients. Available as tablets and oral solution.

### REFERENCES:

- Honkoop P, De Man RA. Entecavir: a potent new antiviral drug for hepatitis B. *Expert Opin Investig Drugs* 12:683–8, 2003.
- Keating GM. Entecavir: a review of its use in the treatment of chronic hepatitis B in patients with decompensated liver disease. *Drugs* 24:2511–29, 2011.
- Marzella N. Entecavir (Baraclude) for the treatment of chronic hepatitis B virus (HBV) infection. *Pharmacy & Therapeutics* 31:313–20, 2006.
- [No authors listed]. Entecavir: A new nucleoside analogue for the treatment of chronic Hepatitis B. *Drugs for Today* 43:201–220, 2007.
- Petersen J, Ratzju, Buti M, et al. Entecavir plus tenofovir combination as recur therapy in pre-treated chronic hepatitis B patients: an international multicenter cohort study. *J Hepatol* 56:520–6, 2012.
- Pol S, Lampertico, P. First-line treatment of chronic hepatitis B with entecavir or tenofovir in 'real-life' setting; from clinical trials to clinical practice. *J viral Hepat* 19:377–86, 2012.
- Rivkin A. A review of entecavir in the treatment of chronic hepatitis B infection. *Curr Med Res Opin* 21:1845–56, 2005.
- Scotto G, D'Addiego G, Giammario A, et al. Tenofovir plus entecavir as rescue therapy for multidrug-resistant chronic hepatitis B. *Liver Int* 32:171–2, 2012.
- Sherman M, Yurdaydin C, et al. Entecavir therapy for lamivudine-refractory chronic hepatitis B: improved virologic, biochemical, and serology outcomes through 96 weeks. *Hepatology* 48:99–108, 2008.
- Tsai Mc, Lee CM, Chiu KW, et al. A comparison of telbivudine and entecavir for chronic hepatitis B in real-world clinical practice. *J Antimicrob Chemother* 67:696–9, 2012
- Yang HJ, Lee JH, Kim YJ, et al. Antiviral efficacy of combination therapy with entecavir and adefovir for entecavir/lamivudine-resistant hepatitis B virus with or without adefovir resistance. *J Med Virol* 84:424–430, 2012.
- Website: [www.pdr.net](http://www.pdr.net)

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

## Ertapenem (Invanz)

**Drug Class:** Carbapenem.

**Usual Dose:** 1 gm (IV/IM) q24h.

**Spectrum:** (see *Susceptibility Profiles* pp. 198–202).

**Resistance Potential:** Low

**Pharmacokinetic Parameters:**

Peak serum level: 150 mcg/mL

Bioavailability: 90% (IM)

Excreted unchanged (urine): 40%; 40% excreted as active metabolite

Serum half-life (normal/ESRD): 4/14 hrs

Plasma protein binding: 95%

Volume of distribution ( $V_d$ ): 0.12 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\***

CrCl 30–80 mL/min	No change
CrCl < 30 mL/min	500 mg (IV) q24h
Post-HD dose	If dosed < 6 h prior to HD, give 150 mg (IV)
Post-PD dose	No information
CVWH/CVHD/ CWVHDF dose	500 mg (IV) q24h
Moderate – Severe hepatic insufficiency	No change

\* If dosed > 6 h prior to HD, no Post-HD needed.

**Drug Interactions:** Not a substrate/inhibitor of cytochrome P-450 enzymes; probenecid (↓ clearance of ertapenem). Valproic acid (↓ seizure threshold).

**Adverse Effects:** Mild headache, infrequent nausea or diarrhea. No seizures. Probenecid (↓ clearance of ertapenem).

**Allergic Potential:** Low

**Safety in Pregnancy:** B

**Comments:** Concentration-dependent protein binding. Compared to imipenem or meropenem, ertapenem has little activity against VSE, Acinetobacter or P. aeruginosa. For deep muscle (IM) injection, mix 1 gm with 3.2 mL of 1% lidocaine. **Probably safe to use in penicillin allergic patients.**

**Cerebrospinal Fluid Penetration:** 5–20%

**Bile Penetration:** 10%

(also see *Antibiotic Pearls & Pitfalls* p. 511).

### REFERENCES:

- Beovic B, Krefl S, Seme K, et al. Does ertapenem alter the susceptibility of Pseudomonas aeruginosa to carbapenems? J Chemother 23:216–220, 2011.
- Boselli E, Breilh D, Djabarouti S, et al. Diffusion of ertapenem into bone and synovial tissues. J Antimicrob Chemother 60:893–896, 2007.
- Cardone KE, Grabe DW, Kulawy RW, et al. Ertapenem pharmacokinetics and pharmacodynamics during continuous ambulatory peritoneal dialysis. Antimicrob Agents Chemother 56:725–30, 2012.
- Collins VL, Marchaim D, Pogue JM, et al. Efficacy of ertapenem for treatment of bloodstream infections caused by extended-spectrum-B-lactamase-producing Enterobacteriaceae. Antimicrob Agents Chemother 56:2173–7, 2012.
- Congeni BL. Ertapenem. Expert Opin Pharmacother 11:669–672, 2010.
- Cunha BA, Jose A, Hage J. Ertapenem: Lack of allergic reactions in hospitalized adults reporting a history of penicillin allergy. Int J Antimicrob Agents 42:585–586, 2013.
- Cunha BA, Lenopoli S, Hage JE. Klebsiella pneumoniae septic wrist arthritis successfully treated with ertapenem and levofloxacin. J Chemother 23:376–7, 2011.
- Cunha BA. Ertapenem. A review of its microbiologic, pharmacokinetic and clinical aspects. Drugs Today (Barc). 38:195–213, 2002.

"Usual dose" assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

- Fong JJ, Rose L, Radigan EA. Clinical outcomes with ertapenem as a first-line treatment option of infections caused by extended-spectrum  $\beta$ -lactamase producing gram-negative bacteria. *Ann Pharmacother* 46:347–52, 2012.
- Graham DR, Lucasti C, Malafaia O, et al. Ertapenem once daily versus piperacillin-tazobactam 4 times per day for treatment of complicated skin and skin-structure infections in adults: results of a prospective, randomized, double-blind multicenter study. *Clin Infect Dis* 34:1460–8, 2002.
- Hyle EP, Ferraro MJ, Silver M, et al. Ertapenem-resistant enterobacteriaceae: risk factors for acquisition and outcomes. *Infect Control Hosp Epidemiol* 31:1242–1249, 2010.
- Itani KMF, Wilson SE, Awad SS, et al. Ertapenem versus cefotetan prophylaxis in elective colorectal surgery. *N Engl J Med* 355:2640–51, 2006.
- Liao FF, Huang YB, Chen CY. Decrease in serum valproic acid levels during treatment with ertapenem. *Am J Health Syst Pharm* 67:1260–1264, 2010.
- Lauf L, Ozsvar Z, Mitha I, et al. Phase 3 study comparing tigecycline and ertapenem in patients with diabetic foot infections with and without osteomyelitis. *Diagn Microbiol Infect Dis*. 78:469–480, 2014.
- Lipsky BA, Armstrong DG, Citron DM, et al. Ertapenem versus piperacillin/tazobactam for diabetic foot infections (SIDESTEP): prospective, randomized, controlled, double-blinded, multicentre trial. *Lancet* 366:1695–703, 2005.
- Namias N, Solomkin JS, Jensen EH, et al. Randomized, multicenter, double-blind study of efficacy, safety, and tolerability of intravenous ertapenem versus piperacillin/tazobactam in treatment of complicated intra-abdominal infections in hospitalized adults. *Surg Infect* 8:15–28, 2007.
- Orsi GB, Garcia-Fernandez A, Giordano V, et al. Risk factors and clinical significance of ertapenem-resistant *Klebsiella pneumoniae* in hospitalized patients. *J Hosp Infect* 78:54–58, 2011.
- Virdier MC, Seguin P, Le Touvet B, et al. Ertapenem in plasma and peritoneal fluid from patients with severe intra-abdominal infections. *J Antimicrob Chemother* 66:1934–1936, 2011.
- Yakovlev SV, Stratchounskin LS, Woods GL, et al. Ertapenem versus cefepime for initial empirical treatment of pneumonia acquired in skilled-care facilities or in hospitals outside the intensive care unit. *Eur J Clin Micro Inf Dis* 25:633–641, 2006.
- Wall GC, Nayima VA, Neumeister KM. Assessment of hypersensitivity reactions in patients receiving carbapenem antibiotics who report antibiotics a history of penicillin. *J Chem*. 26:150–153, 2014.
- Website: [www.pdr.net](http://www.pdr.net)

## Erythromycin lactobionate, base (various)

**Drug Class:** Macrolide.

**Usual Dose:** 1 gm (IV) q6h; 500 mg (PO) q6h (Take without food).

**Spectrum:** (see **Susceptibility Profiles** pp. 186–190).

**Resistance Potential:** High (*S. pneumoniae*, GAS, MSSA)

### Pharmacokinetic Parameters:

*Peak serum level:* 12 (IV); 1.2 (PO) mcg/mL

*Bioavailability:* 50%

*Excreted unchanged (urine):* 5%; 5% excreted as active metabolite

*Serum half-life (normal/ESRD):* 1.4/5.4 hrs

*Plasma protein binding:* 80%

*Volume of distribution ( $V_d$ ):* 0.5 L/kg

**Primary Mode of Elimination:** Hepatic  
**Dosage Adjustments\***

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose  $\times$  1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.



CrCl < 10 mL/min	No change
Post-HD/PD dose	None
CVWH/CVVD/ CVHDF dose	No change
Moderate—severe hepatic insufficiency	No change

**Drug Interactions:** Amiodarone, procainamide, sotalol, astemizole, terfenadine, cisapride, pimozide ( $\uparrow$  QTc); carbamazepine ( $\uparrow$  carbamazepine levels, nystagmus, nausea, vomiting, diarrhea; avoid combination); cimetidine, digoxin, ergot alkaloids, felodipine, midazolam, triazolam, phenytoin, ritonavir, tacrolimus, valproic acid ( $\uparrow$  interacting drug levels); clozapine ( $\uparrow$  clozapine levels; CNS toxicity); corticosteroids ( $\uparrow$  corticosteroid effect); cyclosporine ( $\uparrow$  cyclosporine levels with toxicity); efavirenz ( $\downarrow$  erythromycin levels); rifabutin, rifampin ( $\downarrow$  erythromycin levels,  $\uparrow$  interacting drug levels); statins ( $\uparrow$  risk of rhabdomyolysis); theophylline ( $\uparrow$  theophylline levels, nausea, vomiting, seizures, apnea); warfarin ( $\uparrow$  INR); zidovudine ( $\downarrow$  zidovudine levels).

**Adverse Effects:** Nausea, vomiting, GI upset, irritative diarrhea, abdominal pain, phlebitis. May  $\uparrow$  QT<sub>c</sub>; avoid with other medications that prolong the QT<sub>c</sub> interval and in patients with cardiac arrhythmias/heart block.

**Allergic Potential:** Low

**Safety in Pregnancy:** B

**Comments:** Do not mix erythromycin with B/C vitamins, glucose solutions, cephalothin, tetracycline, chloramphenicol, heparin, or warfarin. Increases GI motility and QT<sub>c</sub> interval.

**Limited anti-Legionella activity.**

**S. pneumoniae resistance prevalent.**

**Cerebrospinal Fluid Penetration:** < 10% (also see **Antibiotic Pearls & Pitfalls** p. 519).

#### REFERENCES:

- Alvarez-Elcoro S, Enzler MJ. The macrolides: Erythromycin, clarithromycin and azithromycin. *Mayo Clin Proc* 74:613–34, 1999.
- Amsden GW. Erythromycin, clarithromycin, and azithromycin: Are the differences real? *Clinical Therapeutics* 18:572, 1996.
- Bergman M, Huikko S, Huovinen P, et al. Macrolide and azithromycin use are linked to increased macrolide resistance in Streptococcus pneumoniae. *Antimicrob Agents Chemother* 50:3646–3650, 2006.
- Cunha BA. The virtues of doxycycline and the evils of erythromycin. *Adv Ther* 14:172–80, 1997.
- Daneman N, McGeer A, Green K, et al. Macrolide resistance in bacteremic pneumococcal disease: Implications for patient management. *Clin Infect Dis* 43:432–438, 2006.
- Gagliotti C, Nobilio L, Milandri M, et al. Macrolide Prescriptions and Erythromycin Resistance or Streptococcus pyogenes. *Clin Infect Dis* 42:1153–6, 2006.
- Jain R, Danzinger LH. The macrolide antibiotics: a pharmacokinetic and pharmacodynamic overview. *Curr Pharm Des* 10:3045–53, 2004.
- Jenkins SG, Farrell DJ. Increase in pneumococcus macrolide resistance, United States. *Emerging Infect Dis* 15:1260–1264, 2009.
- Mandell LA, Wunderink RG, Anzueto A, et al. Infectious Diseases Society of America/American Thoracic Society consensus guidelines on the management of community-acquired pneumonia in adults. *Clin Infect Dis* 44:527–72, 2007.
- Smilack JD, Wilson WE, Cockerill FR 3<sup>rd</sup>. Tetracycline, chloramphenicol, erythromycin, clindamycin, and metronidazole. *Mayo Clin Proc* 66:1270–80, 1991.
- Suzuki S, Yamazaki T, Narita M. Clinical evaluation of macrolide-resistant Mycoplasma pneumoniae. *Antimicrobial Agents and Chemotherapy* 50:709–12, 2006.
- Website: [www.pdr.net](http://www.pdr.net)

"Usual dose" assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose  $\times$  1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

## Ethambutol (Myambutol) EMB

**Drug Class:** TB drug.

**Usual Dose:** 15 mg/kg (PO) q24h (see comments).

**Pharmacokinetic Parameters:**

*Peak serum level:* 2–5 mcg/mL

*Bioavailability:* 80%

*Excreted unchanged (urine):* 50%

*Serum half-life (normal/ESRD):* 4/10 hrs

*Plasma protein binding:* 20%

*Volume of distribution ( $V_d$ ):* 2 L/kg

**Primary Mode of Elimination:** Renal/hepatic

**Dosage Adjustments\***

CrCl < 30 mL/min	15 mg/kg (PO) 3x/ week
CrCl 15–30 mL/min	15 mg/kg (PO) q36h
Post-HD/PD dose	None
CVWH/CVWHD/ CVWHDf dose	15 mg/kg (PO) q36h
Moderate—severe hepatic insufficiency	No change

**Drug Interactions:** Aluminum salts, didanosine buffer (↓ ethambutol and interacting drug absorption).

**Adverse Effects:** Drug fever/rash, ↓ visual acuity, central scotomata, color blindness (red-green), metallic taste, mental confusion, peripheral neuropathy, ↑ uric acid.

**Allergic Potential:** Low

**Safety in Pregnancy:** C

**Comments:** Optic neuritis may occur with high doses (≥ 15 mg/kg/day). **TB D.O.T.**

**dose:** 4 gm (PO) 2 x /week or 3 gm (PO) 3 x / week. **MAI dose:** 15 mg/kg (PO) q24h (with azithromycin 500 mg [PO] q24h).

Meningeal dose = 25 mg/kg (PO) q24h.

**Cerebrospinal Fluid Penetration:**

Non-Inflamed meninges = 1%

Inflamed meninges = 40%

**REFERENCES:**

- Chaisson RE, Keiser P, Pierce M, et al. Clarithromycin and ethambutol with or without clofazimine for the treatment of bacteremic Mycobacterium avium complex disease in patients with HIV infection. *AIDS* 11:311–317, 1997.
- Davidson PT, Le HQ. Drug treatment of tuberculosis 1992. *Drugs* 43:651–73, 1992.
- [No authors listed]. *Drugs for tuberculosis*. *Med Lett Drugs Ther* 35:99–101, 1993.
- McIllerson H, Wash P, Burger A, et al. Determinants of rifampin, isoniazid, pyrazinamide and ethambutol pharmacokinetics in a cohort of tuberculosis patients. *Antimicrob Agents Chemother* 50:1170–77, 2006.
- Perlman DC, Segal Y, Rosenkranz S, et al. The clinical pharmacokinetics of rifampin and ethambutol in HIV-infected persons with tuberculosis. *Clinical Infectious Disease* 41:1638–47, 2005.
- Schlossberg D. Treatment of multi-drug resistant tuberculosis. *Antibiotics for Clinicians* 9:317–21, 2005.
- Van Scoy RE, Wilkowske CJ. Antituberculous agents. *Mayo Clin Proc* 67:179–87, 1992.
- Website: [www.pdr.net](http://www.pdr.net)

## Ethionamide (Trecator)

**Drug Class:** TB drug.

**Usual Dose:** 500 mg (PO) q12h.

**Pharmacokinetic Parameters:**

*Peak serum level:* 5 mcg/mL

*Bioavailability:* 99%

*Excreted unchanged (urine):* 1%

*Serum half-life (normal/ESRD):* 2/9 hrs

*Plasma protein binding:* 30%

*Volume of distribution ( $V_d$ ):* No data

**Primary Mode of Elimination:** Renal/hepatic

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

**Dosage Adjustments\***

CrCl < 40 mL/min	No change
Post-HD/PD dose	No information
CWH/CVWH/ CWHDF dose	No information
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	500 mg (PO) q24h

**Drug Interactions:** Cycloserine (↑ neurologic toxicity); ethambutol (↑ GI distress, neuritis, hepatotoxicity); INH (peripheral neuritis, hepatotoxicity); pyrazinamide, rifampin (hepatotoxicity).

**Adverse Effects:** ↑ SGOT/SGPT, headache, nausea/vomiting, abdominal pain, tremor, olfactory abnormalities, alopecia, gynecomastia, hypoglycemia, impotence, neurotoxicity (central/peripheral neuropathy).

**Allergic Potential:** Low

**Safety in Pregnancy:** C

**Comments:** Additive toxicity with thiacetazone

Meningeal dose = usual dose.

**Cerebrospinal Fluid Penetration:** 100%

**REFERENCES:**

- Davidson PT, Le HQ. Drug treatment of tuberculosis 1992. *Drugs* 43:651–73, 1992.  
 [No authors listed.] Drugs for tuberculosis. *Med Lett Drugs Ther* 35:99–101, 1993.  
 Furin J, Nardell EA. Multidrug-resistant tuberculosis: An update on the best regimens. *J Respir Dis* 27:172–82, 2006.  
 Iseman MD. Treatment of multidrug resistant tuberculosis. *N Engl J Med* 329:784–91, 1993.  
 Schlossberg D. Treatment of multi-drug resistant tuberculosis. *Antibiotics for Clinicians* 9:317–321, 2005.  
 Website: www.pdr.net

**Etravirine (Intelence) ETR**

**Drug Class:** HIV NNRTI.

**Usual Dose:** 200 mg (PO) q12h (take with meals).

**Pharmacokinetic Parameters:**

*Peak serum level:* 2.97 mcg/mL

*Bioavailability:* no data

*Excreted unchanged:* 0%

*Serum half-life (normal/ESRD):* 41/41 hrs

*Plasma protein binding:* 99%

*Volume of distribution (V<sub>d</sub>):* No data

**Primary Mode of Elimination:** Hepatic

**Dosage Adjustments\***

CrCl < 30 mL/min	No change
Post-HD dose	None
Post-PD dose	None
CVWH/CVWH/ CVHDF dose	No change
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	use with caution

**Drug Interactions:** Do not administer with drugs that are inducers, inhibitors or substrates of CYP 3A4, 2C9 and/or 2C19 (tipranavir/ritonavir, atazanavir/ritonavir, protease inhibitors without ritonavir and/or NNRTIs), clarithromycin (↓ clarithromycin levels), statins (↓ statin levels), atovaquone/proguanil (↓ atovaquone/proguanil levels).

**Contraindications:** carbamazepine, phenytoin, phenobarbital, St. John's wort, rifapentine, atazanavir, efavirenz, fosamprenavir, nevirapine, rilpivirine, tipranavir.

**Adverse Effects:** Rash, peripheral neuropathy, immune reconstitution syndrome, fat redistribution.

**Allergic Potential:** Low

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

**Safety in Pregnancy:** B

**Comments:** Severe, life-threatening and fatal skin reactions (Steven-Johnson Syndrome, toxic epidermal necrolysis, hypersensitivity reaction and erythema multiforme). Doesn't ↓ methadone levels. If etravirine given with ritonavir–boosted protease inhibitor, rifabutin is contraindicated. Otherwise rifabutin can be given at 300 mg q24h.

**REFERENCES:**

Patterson K, Jennings S, Falcon R, et al. Darunavir, ritonavir, etravirine pharmacokinetics in the cervicovaginal fluid and blood plasma of HIV-infected women. *Antimicrob Agents Chemother* 55:1120–2, 2011.

Thompson MA, Aberg JA, Cahn P, et al. Antiretroviral treatment of adult HIV infection: 2012 recommendations of the International Antiviral Society — USA panel. *JAMA* 308:387–402, 2012.

Website: www.pdr.net

**Famciclovir (Famvir)**

**Drug Class:** HSV, VZV, HHV-6 Antiviral

(see comments).

**Usual Dose:**

**HSV-1/2:** Herpes labialis: 500 mg (PO) q12h × 7 days. Genital herpes: *Initial therapy*: 1 g (PO) q12h × 1 day. *Recurrent/intermittent therapy* (< 6 episodes/year): normal host: 125 mg (PO) q12h × 5 days; HIV-positive: 500 mg (PO) q12h × 7 days. *Chronic suppressive therapy* (> 6 episodes/year): 250 mg (PO) q12h × 1 year.

Meningitis/encephalitis: 500 mg (PO) q8h × 10 days.

**VZV:** Chickenpox: 500 mg (PO) q8h × 5 days. Herpes zoster (shingles) (dermatomal/disseminated): 500 mg (PO) q8h × 7–10 days.

VZV Pneumonia: 500 mg (PO) q8h × 7–10 days.

**Pharmacokinetic Parameters:**

*Peak serum level*: 3.3 mcg/mL

*Bioavailability*: 77%

*Excreted unchanged (urine)*: 60%

*Serum half-life (normal/ESRD)*: 2.5/13 hrs

*Plasma protein binding*: 20%

*Volume of distribution (V<sub>d</sub>)*: 1.1 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments for HSV/VZV\*** (based on 250 mg [PO] q8h/500 mg [PO] q8h)

CrCl 40–60 mL/min	No change/500 mg (PO) q12h
CrCl 20–40 mL/min	125 mg (PO) q24h/ 500 mg (PO) q24h
CrCl < 20 mL/min	125 mg (PO) q24h/ 250 mg (PO) q24h
Post-HD dose	125 mg (PO)/ 250 mg (PO)
Post-PD dose	None
CVWH/CVHHD/ CVHDF dose	125 mg (PO) q24h/ 500 mg (PO) q24h
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	Use with caution

**Drug Interactions:** Digoxin (↑ digoxin levels).

**Adverse Effects:** Headache, seizures/tremors (dose related), nausea, diarrhea, renal insufficiency (dose related).

**Allergic Potential:** Low

**Safety in Pregnancy:** B

**Comments:** 99% converted to penciclovir in liver/GI tract. **Highly active against HSV, VZV, and HHV-6. Some activity against CMV. No activity against EBV, RSV or adenoviruses.**

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

Meningeal dose = VZV dose.

**Cerebrospinal Fluid Penetration:** 50%

**REFERENCES:**

Luber AD, Flaherty JF Jr. Famciclovir for treatment of herpesvirus infections. *Ann Pharmacother* 30:978–85, 1996.

Website: www.pdr.net

## Fluconazole (Diflucan)

**Drug Class:** Antifungal.

**Usual Dose: Loading Dose (LD) = twice the maintenance dose × 1 dose** followed by the **Maintenance Dose (MD)**.

Candidemia:

LD = 800 mg (IV/PO) × 1 then  
MD = 400 mg (IV/PO) q24h

Mucocutaneous Candidiasis:

LD = 400 mg (IV/PO) × 1 then  
MD = 200 mg (IV/PO) q24h

Candida esophagitis:

LD = 200 mg (IV/PO) × 1 then  
MD = 100 mg (IV/PO) q24h

Candiduria:

LD = 200 mg (IV/PO) × 1 then  
MD = 100 mg (IV/PO) q24h

**Pharmacokinetic Parameters:**

Peak serum level: 6.7 mcg/mL

Bioavailability: 90%

Excreted unchanged (urine): 80%; 11% excreted as active metabolite

Serum half-life (normal/ESRD): 27/100 hrs

Plasma protein binding: 12%

Volume of distribution ( $V_d$ ): 0.7 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\***

CrCl < 50 mL/min	100 mg (IV/PO) q24h
Post-HD dose	200 mg (IV/PO)
Post-PD dose	200 mg (IV/PO)
CVVH/CVVHD/ CVVHDF dose	200–400 mg (IV/PO) q24h
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	No change

**Drug Interactions:** Astemizole, cisapride, terfenadine, amiodarone (may ↑ QT interval, torsades de pointes); cyclosporine, oral hypoglycemics, tacrolimus, theophylline, zidovudine (↑ interacting drug levels with possible toxicity); hydrochlorothiazide (↑ fluconazole levels); phenytoin, rifabutin, rifampin (↓ fluconazole levels, ↑ interacting drug levels); warfarin (↑ INR).

**Adverse Effects:** Not hepatotoxic. **Does not ↑ SGOT/SGPT even in high dose.** Hypokalemia.

**Allergic Potential:** Low

**Safety in Pregnancy:** C

**Comments:** Not effective against most non-albicans *Candida*.

**Highly active against *C. albicans*, *C. tropicalis*, *C. parasilosis*, and *C. lusitaniae*. Moderately active against *Cryptococcus*, *Histoplasmosis*, *Blastomycosis*, *Sporotrichosis*, *Penicillium marneffeii*, *Paracoccidiomycosis* and *Coccidiomycosis*. Some (dose dependent) activity against**

***C. glabrata*. No activity against *C. krusei*,**

***C. guilliermondii*, *C. rugosa*, *Mucor*,**

***Fusarium*, *Pseudoallescheria/Scedosporium*,**

***Trichosporon*, *Malassezia*, *Hansenula*,**

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

**Geotrichum, Rhodotorula or Aspergillus.**

Meningeal dose = 400 mg (IV/PO) q24h.

**Cerebrospinal Fluid Penetration:**

Non-Inflamed meninges = 50–90%

Inflamed meninges = 50–90%

**REFERENCES:**

- Cha R, Sobel JD. Fluconazole for the treatment of candidiasis: 15 years experience. *Expert Rev Anti Infect Ther* 2:357–66, 2004.
- Clancy CJ, Yu VL, Morris AJ, et al. Fluconazole MIC and fluconazole dose/MIC ratio correlate with therapeutic response among patients with candidemia. *Antimicrobial Agents and Chemotherapy* 49:3171–3177, 2005.
- Hamza OJM, Matee MIN, Bruggemann RJM, et al. Single-dose fluconazole versus standard 2-week therapy for oropharyngeal candidiasis in HIV-infected patients: a randomized, double-blind, double-dummy trial. *Clin Infect Dis* 47:1270–1276, 2008.
- Goa KL, Barradell LB. Fluconazole: An update of its pharmacodynamics and pharmacokinetic properties and therapeutic use in major superficial and systemic mycoses in immunocompromised patients. *Drugs* 50:658–90, 1995.
- Kowalsky SF, Dixon DM. Fluconazole: A new antifungal agent. *Clin Pharmacol* 10:179–94, 1991.
- Miceli MH, Diaz JA, Lee SA. Emerging opportunistic yeast infections. *Lancet Infectious Diseases* 11: 142–151, 2011.
- Nussbaum JC, Jackson A, Namarika D, et al. Combination flucytosine and high-dose fluconazole compared with fluconazole monotherapy for the treatment of Cryptococcal meningitis: a randomized trial in Malawi. *Clin Infect Dis* 50:338–344, 2010.
- Pappas PG, Chetchotisakd P, Larsen RA, et al. A phase II randomized trial of amphotericin B alone or combined with fluconazole in the treatment of HIV-associated cryptococcal meningitis. *Clin Infect Dis* 48:1775–1784, 2009.
- Perfect JR, Dismukes WE, Dromer F, et al. Clinical practice guidelines for the management of cryptococcal disease: 2010 update by the Infectious

Disease Society of America. *Clin Infect Dis* 50:291–322, 2010.

- Rex JH, Pappas PG, Karchmer AW, et al. National Institute of Allergy and Infectious Diseases Mycoses Study Group. A randomized and blinded multicenter trial of high-dose fluconazole plus placebo versus fluconazole plus amphotericin B as therapy for candidemia and its consequences in nonneutropenic subjects. *Clin Infect Dis* 36:1221–8, 2003.
- Sabatelli F, Patel R, Mann PA, et al. In vitro activities of posaconazole, fluconazole, itraconazole, voriconazole, and amphotericin B against a large collection of clinically important molds and yeasts. *Antimicrob Agents Chemother* 50:2009–15, 2006.
- Smego RA Jr, Ahmad H. The role of fluconazole in the treatment of Candida endocarditis: a meta-analysis. *Medicine* 90:237–249, 2011.
- Sousa AQ, Frutuoso MS, Moraes EA, et al. High-dose oral fluconazole therapy effective for cutaneous leishmaniasis due to *Leishmania (Vianna) braziliensis*. *Clin Infect Dis* 53:693–695, 2011.
- Terrell CL. Antifungal agents: Part II. The azoles. *Mayo Clin Proc* 74:78–100, 1999.
- Zhai B, Zhou H, Yang L, et al. Polymyxin B, in combination with fluconazole, exerts a potent fungicidal effect. *J Antimicrob Chemother* 65:931–938, 2010.
- Website: [www.pdr.net](http://www.pdr.net)

**Flucytosine (Ancobon) 5-FC**

**Drug Class:** Antifungal.

**Usual Dose:** 25 mg/kg (PO) q6h.

**Pharmacokinetic Parameters:**

*Peak serum level:* 3.5 mcg/ml

*Bioavailability:* 80%

*Excreted unchanged (urine):* 90%

*Serum half-life (normal/ESRD):* 4/85 hrs

*Plasma protein binding:* 4%

*Volume of distribution (V<sub>d</sub>):* 0.6 L/kg

**Primary Mode of Elimination:** Renal

**Dose Adjustments\***

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

CrCl 20–40 mL/min	37.5 mg/kg (PO) q12h
CrCl 10–20 mL/min	37.5 mg/kg (PO) q18h
CrCl < 10 mL/min	37.5 mg/kg (PO) q24h
Post-HD dose	37.5 mg/kg (PO)
Post-PD dose	1 gm (PO)
CVH/CVHD/ CWVHDF dose	37.5 mg/kg (PO) q18h
Moderate—severe insufficiency	No change

**Drug Interactions:** Cytarabine (↓ flucytosine effect); zidovudine (neutropenia).

**Adverse Effects:** Leukopenia, anemia, thrombocytopenia, nausea, vomiting, abdominal pain, ↑ SGOT/SGPT, drug fever/rashes.

**Allergic Potential:** High

**Safety in Pregnancy:** C

**Comments:** Always use in combination with amphotericin B for cryptococcal meningitis.

**Highly active against *Cryptococcus neoformans*. Some activity against *C. albicans*. No activity against other yeasts/fungi.**

Meningeal dose = usual dose.

**Cerebrospinal Fluid Penetration:**

Non-Inflamed meninges = 100%

Inflamed meninges = 100%

#### REFERENCES:

Lyman CA, Walsh TJ. Systemically administered antifungal agents. A review of their clinical pharmacology and therapeutic applications: Part I. Amphotericin B preparations and flucytosine. *Mayo Clin Proc* 73:1205–25, 1998.

Mosquera J, Shardt A, Moore CB, et al. In vitro interaction of terbinafine with itraconazole, fluconazole, amphotericin

B and 5-flucytosine against *Aspergillus* spp. *J Antimicrob Chemother* 50:189–94, 2002.

Nussbaum JC, Jackson A, Namarika D, et al. Combination Flucytosine and High-Dose Fluconazole Compared with Fluconazole Monotherapy for the Treatment of Cryptococcal Meningitis: A Randomized Trial in Malawi. *Clin Infect Dis* 50:338–344, 2010.

Pfaller MA, Messer SA, Boyken L, et al. In vitro activities of 5-fluorocytosine against 8,803 clinical isolates of *Candida* spp.: Global assessment of primary resistance using National Committee for Clinical Laboratory Standards susceptibility testing methods. *Antimicrob Agents Chemother* 46:3518–3521, 2002.

Perfect JR, Dismukes WE, Dromer F, et al. Clinical practice guidelines for the management of cryptococcal disease: 2010 update by the Infectious Disease Society of America. *Clin Infect Dis* 50:291–322, 2010.

Te Dorsthorst DT, Verweij PE, Meletidis J, et al. In vitro interaction of flucytosine combined with amphotericin B or fluconazole against thirty-five yeast isolates determined by both the fractional inhibitory concentration index and the response surface approach. *Antimicrob Agents Chemother* 46:2982–9, 2002.

Wintermeyer SM, Mahata MC. Stability of flucytosine in an extemporaneously compounded oral liquid. *Am J Health Syst Pharm* 53:407–9, 1996.

Website: www.pdr.net

## Fosamprenavir (Lexiva) FPV

**Drug Class:** HIV protease inhibitor.

**Usual Dose:** 1400 mg (PO) q24h plus ritonavir 200 mg (PO) q24h or 700 mg (PO) q12h plus ritonavir 100 mg (PO) q12h. For PI-experienced patients: 700 mg (PO) q12h plus ritonavir 100 mg (PO) q12h.

**Pharmacokinetic Parameters:**

*Peak serum level:* 4.8 mcg/mL

*Bioavailability:* No data

*Excreted unchanged (urine) (urine):* 1%

*Serum half-life (normal/ESRD):* 7 hrs/no data

*Plasma protein binding:* 90%

*Volume of distribution ( $V_d$ ):* 6.1 L/kg

"Usual dose" assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

**Primary Mode of Elimination:** Hepatic  
**Dosage Adjustments\***

CrCl 50–80 mL/min	No change
CrCl 10–50 mL/min	No change
CrCl < 10 mL/min	No change
Post-HD or PD dose	No change
CVWH/CVVHD/ CVVHDF dose	No change
Mild-moderate hepatic insufficiency (Child- Pugh score 5–8)	700 mg (PO) q12h without ritonavir; 450 mg (PO) q12h with ritonavir
Severe hepatic insufficiency (Child- Pugh score 9–12)	350 mg (PO) q12h without ritonavir; 300 mg (PO) q12h with ritonavir

**Antiretroviral Dosage Adjustments**

Didanosine	Administer didanosine 1 hour apart
Delavirdine	Avoid
Efavirenz	Fosamprenavir 700 mg q12h + ritonavir 100 mg q12h + efavirenz; fosamprenavir 1400 mg q24h + ritonavir 200 mg q24h + efavirenz; no data for fosamprenavir 1400 mg q12h + efavirenz
Indinavir	No data
Lopinavir/ ritonavir	Avoid
Nelfinavir	No information
Nevirapine	Avoid
Saquinavir	No information

Rifampin	Avoid
Rifabutin	Reduce usual rifabutin dose by 50% (or 75% if given with fosamprenavir plus ritonavir; max. 150 mg q48h)

**Drug Interactions:** Antiretrovirals (see dose adjustment grid, above). Contraindicated with: ergot derivatives, cisapride, midazolam, triazolam, pimozone, (flecainide and propafenone if administered with ritonavir). Dose reduction (of other drug): atorvastatin, rifabutin, sildenafil, vardenafil, ketoconazole, itraconazole. colchicine (↑ colchicine levels) Concentration monitoring (of other drug): amiodarone, systemic lidocaine, quinidine, warfarin (INR), tricyclic antidepressants, cyclosporin, tacrolimus, sirolimus. H<sub>2</sub> blockers and proton pump inhibitors interfere with absorption. Sildenafil (do not give > 25 mg/48 hrs); tadalafil (max. 10 mg/72 hrs); vardenafil (max. 2.5 mg/72 hrs). Clarithromycin (↑ QTc). May increase methadone effect/decrease methadone dose.

**Contraindications:** Rifampine, telaprevir, do not coadminister with: rifampin, lovastatin, simvastatin, St. John's wort, delavirdine.

**Adverse Effects:** Rash, Stevens-Johnson Syndrome (rare), GI upset, headache, depression, diarrhea, hyperglycemia (including worsening diabetes, new-onset diabetes, DKA), ↑ cholesterol/triglycerides (evaluate risk for coronary disease/pancreatitis), fat redistribution, ↑ SGOT/SGPT, possible increased bleeding in hemophilia. Immune reconstitution syndrome.

**Allergic Potential:** High. Fosamprenavir is a sulfonamide; use with caution in patients with sulfonamide allergies

**Safety in Pregnancy:** C

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.



**Comments:** Usually given in conjunction with ritonavir. May be taken with or without food. Fosamprenavir is a prodrug that is rapidly hydrolyzed to amprenavir by gut epithelium during absorption. Amprenavir inhibits CYP3A4. Fosamprenavir contains a sulfonamide moiety (as do darunavir and tipranavir). Fosamprenavir without ritonavir is no longer recommended in treatment naive patients due to increased development to resistance as well as cross-resistance to darunavir.

#### REFERENCES:

- Blick G, Greiger-Zanlungo P, Gret S, et al. Long-term efficacy and safety of once- daily fosamprenavir 1400 mg boosted by ritonavir 100 mg: the BOLD100 study. *Int J STD AIDS* 23:18–22, 2012
- Hughes PJ, Cretton-Scott E, Teague A, et al. Protease inhibitors for patients with HIV-1 infection. *Pharm Ther* 36:332–344, 2011.
- Thompson MA, Aberg JA, Cahn P, et al. Antiretroviral treatment of adult HIV infection: 2012 recommendations of the International Antiviral Society — USA panel. *JAMA* 308:387–402, 2012.
- Website: [www.pdr.net](http://www.pdr.net)

## Foscarnet (Foscavir)

**Drug Class:** CMV, HHV-6 Antiviral (see comments).

**Usual Dose:** HSV: 40 mg/kg (IV) q12h × 2–3 weeks; CMV: *Induction Dose*: 90 mg/kg (IV) q12h × 2 weeks, then *Maintenance Dose*: 90–120 mg/kg (IV) q24h until cured. *Relapse/reinduction dose*: 120 mg/kg (IV) q24h × 2 weeks.

#### Pharmacokinetic Parameters:

*Peak serum level*: 150 mcg/ml  
*Bioavailability*: Not applicable  
*Excreted unchanged (urine)*: 85%  
*Serum half-life (normal/ESRD)*: 2–4/25 hrs  
*Plasma protein binding*: 17%  
*Volume of distribution ( $V_d$ )*: 0.5 L/kg

**Primary Mode of Elimination:** Renal/hepatic  
**Dosage Adjustments\***

Induction	
CrCl 50–80 mL/min	40–50 mg/kg (IV) q8h
CrCl 20–50 mL/min	20–30 mg/kg (IV) q8h
CrCl < 20 mL/min	Avoid
Maintenance	
CrCl 50–80 mL/min	60–70 mg/kg (IV) q24h
CrCl 20–50 mL/min	65–80 mg/kg (IV) q48h
CrCl < 20 mL/min	Avoid
Post–HD dose	60 mg/kg (IV)
Post–HF HD dose	60 mg/kg (IV)
Post–PD dose	None
CVH/CVHDF/ CVHDF dose	Induction: 20–30 mg/kg (IV) q8h; maintenance: 65–80 mg/kg (IV) q48h
Mod. hepatic insufficiency	No change
Severe hepatic insufficiency	No change

Infusion pump must be used. Adequate hydration is recommended to prevent renal toxicity  
 † Higher doses may be considered for early reinduction due to progression of CMV retinitis, and for patients showing excellent tolerance.

**Drug Interactions:** Ciprofloxacin (↑ risk of seizures); amphotericin B, aminoglycosides, cis-platinum, cyclosporine, ritonavir, saquinavir (↑ nephrotoxicity); pentamidine IV (severe hypocalcemia reported; do not combine); zidovudine (↑ incidence/severity of anemia).

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

**Adverse Effects:** Major side effects include nephrotoxicity and tetany (from ↓ Ca<sup>++</sup>). Others include anemia, nausea, vomiting, GI upset, headache, seizures, peripheral neuropathy, hallucinations, tremors, nephrogenic DI, ↓ Ca<sup>++</sup>, ↓ Mg<sup>++</sup>, ↓ PO<sub>4</sub><sup>=</sup>, oral/genital ulcers.

**Allergic Potential:** Low

**Safety in Pregnancy:** C

**Comments:** Renal failure prevented/minimized by adequate hydration. Dilute with 150 cc normal saline per 1 gm foscarnet. Do not mix with other types of solutions. Administer by IV slow infusion ≤ 1 mg/kg/min using an infusion pump. **Highly active against CMV and HHV-6. Some activity against EBV, HSV and VZV. No activity against RSV or adenoviruses.**

Meningeal dose = usual dose.

**Cerebrospinal Fluid Penetration:**

Non-Inflamed meninges = 90%

Inflamed meninges = 100%

#### REFERENCES:

Mattes FM, Hainsworth EG, Geretti AM, et al.

A randomized, controlled trial comparing ganciclovir plus foscarnet (each at half dose) preemptive therapy of cytomegalovirus infection in transplant recipients.

J Infect Dis 189:1355–61, 2004.

Website: www.pdr.net

## Fosfomycin (Monurol)

**Drug Class:** Urinary antibiotic.

**Usual Dose:** 3 gm (PO) q24h (Take without food)

**Spectrum:** (see **Susceptibility Profiles**

pp. 198–202).

**Resistance Potential:** Low

**Pharmacokinetic Parameters:**

Peak serum level: 26 mcg/ml

Bioavailability: 37%

Excreted unchanged (urine): 60%; 40% excreted as active metabolite

Serum half-life (normal/ESRD): 5.7/50 hrs

Plasma protein binding: 3%

Volume of distribution (V<sub>d</sub>): 2 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\***

CrCl 10–50 mL/min	No change
CrCl < 10 mL/min	Use with caution
Post–HD dose	3 gm (PO)
Post–PD dose	1 gm (PO)
CVH/CVHD/ CVHDF dose	No change
Moderate—severe hepatic insufficiency	No change

**Drug Interactions:** Antacids, metoclopramide (↓ fosfomycin effect).

**Adverse Effects:** Nausea, vomiting, diarrhea, GI upset, ↑ SGOT/SGPT, thrombocytosis.

**Allergic Potential:** Low

**Safety in Pregnancy:** B

**Comments:** **Useful for GNB cystitis/CAB** also active against VSE/VRE. Treat cystitis as single dose in females. **One of the few oral antibiotics active against most strains of MDR**

**P. aeruginosa, MDR Klebsiella pneumoniae, and MDR Acinetobacter baumannii in urine**

(cystitis/CAB). Treatment of MDR GNB cystitis/CAB (change Foley *before* beginning therapy). Use 3 g (PO) q 3 days × 2–3 doses. For prostatitis due to MDR GNB, use fosfp,ucom 3 g (PO) q2 days × 30 days. Urine levels > 128 mcg/ml × 36–48 (see **Antibiotic Pearls & Pitfalls**. p. 520).

\*Usual dose\* assumes normal renal/hepatic function. \*For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

## REFERENCES:

Auer S, Wojna A, Hell M. Oral treatment options for ambulatory patients with urinary tract infections caused by extended-spectrum-beta-lactamase-producing *Escherichia coli*. *Antimicrob Agents Chemother* 54:4006–4008, 2010.

Dinh A, Salomon J, Bru JP, et al. Fosfomycin: efficacy against infections caused by multidrug-resistant bacteria. *Scand J Infect Dis* 44:182–9, 2012.

Falagas ME, Kastoris AC, Kapaskelis AM, et al. Fosfomycin for the treatment of multidrug-resistant, including extended-spectrum  $\beta$ -lactamase producing, Enterobacteriaceae infections: a systematic review. *Lancet Infect Dis* 10:43–50, 2010.

Gardiner BJ, Mahony AA, Ellis AG. Is fosfomycin a potential treatment alternative for multidrug-resistant gram-negative prostatitis? *Clin Infect Dis* 58:e101-105, 2014.

Liu HY, Lin HC, Lin YC, et al. Antimicrobial susceptibilities of urinary extended-spectrum beta-lactamase-producing *Escherichia coli* and *Klebsiella pneumoniae* to fosfomycin and nitrofurantoin in a teaching hospital in Taiwan. *J Microbiol Immunol Infect* 44:364–368, 2011.

Livermore DM, Warner M, Mushtaq S, et al. What remains against carbapenem-resistant Enterobacteriaceae? Evaluation of chloramphenicol, ciprofloxacin, colistin, fosfomycin, minocycline, nitrofurantoin, temocillin and tigecycline. *Int J Antimicrob Agents* 37:415–419, 2011.

Pontikis K, Karaiskos I, Bastani S, et al. Outcomes of critically ill intensive care unit patients treated with fosfomycin for infections due to pandrug-resistant and extensively drug-resistant carbapenemase-producing Gram-negative bacteria. *Int J Antimicrob Agents* 43:52-59, 2014.

Website: [www.pdr.net](http://www.pdr.net)

## Ganciclovir (Cytovene)

**Drug Class:** CMV, HSV, HHV-6 Antiviral (see comments), nucleoside inhibitor/analogue.

**Usual Dose:** **Induction Dose:** 5 mg/kg (IV) q12h  $\times$  2 weeks; **Maintenance Dose:** 5 mg/kg (IV) q24h or 1 gm (PO) q8h.

**Pharmacokinetic Parameters:**

*Peak serum level:* 8.3 (IV)/1.2 (PO) mcg/ml

*Bioavailability:* 5%

*Excreted unchanged (urine):* 90%

*Serum half-life (normal/ESRD):* 3.6/28 hrs

*Plasma protein binding:* 1%

*Volume of distribution ( $V_d$ ):* 0.74 L/kg

**Primary Mode of Elimination:** Renal  
Dosage Adjustments\*

CrCl 50–70 mL/min	2.5 mg/kg (IV) q12h (induction); 2.5 mg/kg (IV) q24h (maintenance); 500 mg (PO) q8h
CrCl 25–50 mL/min	2.5 mg/kg (IV) q24h (induction); 1.25 mg/kg (IV) q24h (maintenance); 500 mg (PO) q12h
CrCl 10–25 mL/min	1.25 mg/kg (IV) q24h (induction); 0.625 mg/kg (IV) q24h (maintenance); 500 mg (PO) q24h
CrCl < 10 mL/min	1.25 mg/kg (IV) 3x/week (induction); 0.625 mg/kg (IV) 3x/week (maintenance); 500 mg (PO) 3x/week
Post–HD dose	1.25 mg/kg (IV) (induction); 0.625 mg/kg (IV) (maintenance); 500 mg (PO)
Post–PD dose	Same dose as Post–HD
CVWH/CVWHD/ CVWDF dose	Same dose as CrCl 50–70 mL/min
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	No change

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose  $\times$  1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

**Drug Interactions:** Cytotoxic drugs (may produce additive toxicity: stomatitis, bone marrow depression, alopecia); imipenem (↑ risk of seizures); probenecid (↑ ganciclovir levels); zidovudine (↓ ganciclovir levels, ↑ zidovudine levels, possible neutropenia); didanosine (↑ didanosine levels); cyclosporine, amphotericin, (↑ nephrotoxicity); mycophenolate mofetil (↑ mycophenolate mofetil and ganciclovir levels in renal insufficiency); tenofovir (↑ tenofovir, ganciclovir levels).

**Adverse Effects:** Headaches, hallucinations, seizures/tremor (dose related), drug fever/rash, diarrhea, nausea/vomiting, GI upset, leukopenia, thrombocytopenia, anemia, retinal detachment. Ventricular arrhythmias (VT/torsades de pointes).

**Allergic Potential:** High

**Safety in Pregnancy:** C

**Comments:** Induction dose usually given IV.

Maintenance dose may be given IV or PO. For PO therapy use valganciclovir (not PO ganciclovir). Reduce dose with neutropenia/thrombocytopenia. Bioavailability increased with food: 5% fasting; 6–9% with food; 28–31% with fatty food.

**Highly active against CMV, HHV-6, and HSV. Some activity against VZV, EBV, and adenoviruses. No activity against RSV**

Meningeal dose = usual dose.

**Cerebrospinal Fluid Penetration:** 70%

#### REFERENCES:

- Jacobsen T, Sipointis N. Drug interactions and toxicities associated with the antiviral management of cytomegalovirus infection. *Am J Health Syst Pharm* 67:1417–1425, 2010.
- Matthews T, Boehme R. Antiviral activity and mechanism of action of ganciclovir. *Rev Infect Dis* 10:490–4, 1988.
- Singh N. Preemptive therapy for cytomegalovirus with oral ganciclovir after liver transplantation. *Transplantation* 73:1977–78, 2002.

Tayebi-Khosroshahi H, Makhdami N, Heris HK, et al.

Universal prophylaxis with ganciclovir preparation is not necessary in all kidney allograft recipients. *Transplant Proc* 43:547–50, 2011

Tokimasa S, Hara J, Osugi Y, et al. Ganciclovir is effective for prophylaxis and treatment of human herpesvirus-6 in allogeneic stem cell transplantation.

*Bone Marrow Transplant* 29:595–8, 2002.

Website: www.pdr.net

## Gentamicin (Garamycin)

**Drug Class:** Aminoglycoside.

**Usual Dose:** 240 mg or 5 mg/kg (IV) q24h (preferred over q8h dosing) (see comments).

**Spectrum:** (see **Susceptibility Profiles** pp. 198–202).

**Resistance Potential:** High (*P. aeruginosa*, aerobic GNBs)

**Pharmacokinetic Parameters:**

*Peak serum levels:* 4–8 mcg/ml (q8h dosing); 16–24 mcg/ml (q24h dosing)

*Bioavailability:* Not applicable

*Excreted unchanged (urine):* 95%

*Serum half-life (normal/ESRD):* 2.5/48 hrs

*Plasma protein binding:* < 5%

*Volume of distribution ( $V_d$ ):* 0.3 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\***

CrCl 50–80 mL/min	120 mg (IV) q24h or 2.5 mg/kg (IV) q24h
CrCl 10–50 mL/min	120 mg (IV) q48h or 2.5 mg/kg (IV) q48h
CrCl < 10 mL/min	80 mg (IV) q48h or 1.25 mg/kg (IV) q48h
Post-HD dose	80 mg (IV) or 1 mg/kg (IV)

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

Post-PD dose	40 mg (IV) or 0.5 mg/kg (IV)
Post-HFHD dose	120 mg (IV) or 2.5 mg/kg
Post-dose	120 mg (IV) or 2.5 mg/kg (IV)
CWH/CVHD/ CWHDF dose	120 mg (IV) q48h or 2.5 mg/kg (IV)
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	No change

**Drug Interactions:** Amphotericin B, cephalothin, cyclosporine, enflurane, methoxyflurane, NSAIDs, polymyxin B, radiographic contrast, vancomycin (↑ nephrotoxicity); cis-platinum (↑ nephrotoxicity, ↑ ototoxicity); loop diuretics (↑ ototoxicity); neuromuscular blocking agents, magnesium sulfate (↑ apnea, prolonged paralysis); non-polarizing muscle relaxants (↑ apnea).

**Adverse Effects:** Neuromuscular blockade with rapid infusion/absorption. Nephrotoxicity only with prolonged/extremely high serum trough levels; may cause reversible non-oliguric renal failure (ATN). Ototoxicity associated with prolonged/extremely high peak serum levels (usually irreversible); Cochlear toxicity (1/3 of ototoxicity) manifests as decreased high frequency hearing, but deafness is unusual. Vestibular toxicity (2/3 of ototoxicity) develops before ototoxicity, and typically manifests as tinnitus.

**Safety in Pregnancy:** D

**Comments:** **Synergy dose:** 2.5 mg/kg (IV) q24h or 120 mg (IV) q24h. **Single daily dosing virtually eliminates nephrotoxic/ototoxic potential.** Incompatible with solutions containing  $\beta$ -lactams, erythromycin,

chloramphenicol, furosemide, sodium bicarbonate. IV infusion should be given slowly over 1 hour. **May be given IM. Avoid intraperitoneal infusion due to risk of neuromuscular blockade.** Avoid intratracheal/aerosolized intrapulmonary instillation, which predisposes to antibiotic resistance.  $V_d$  increases with edema/ascites, trauma, burns, cystic fibrosis; may require ↑ dose.  $V_d$  decreases with dehydration, obesity; may require ↓ dose.

**Renal cast counts are the best indicator of aminoglycoside nephrotoxicity, not serum creatinine.** Dialysis removes ~ 1/3 of gentamicin from serum. **CAPD dose:** 2–4 mg/L dialysate (IP) with each exchange.

**Therapeutic Serum Concentrations** (for therapeutic efficacy, *not toxicity*):  
Peak (q24h/q8h dosing) = 16–24/8–10 mcg/ml  
Trough (q24h/q8h dosing) = 0/1–2 mcg/ml

**Intrathecal (IT) dose:** 5 mg (IT) q24h.

**Cerebrospinal Fluid Penetration:**

Non-Inflamed meninges = 0%

Inflamed meninges = 20%

**Bile Penetration:** 30%

(also see **Antibiotic Pearls & Pitfalls** p. 514).

#### REFERENCES:

- Bruss JB. Lack of evidence associating nephrotoxicity with low-dose gentamicin for Staphylococcus aureus bacteremia and endocarditis. Clin Infect Dis 49:806, 2009.
- Cosgrove SE, Vigliani GA, Campion M, et al. Initial low-dose gentamicin for Staphylococcus aureus bacteremia and endocarditis is nephrotoxic. Clin Infect Dis 48:713–721, 2009.
- Cunha BA. Aminoglycosides: Current role in antimicrobial therapy. Pharmacotherapy 8:334–50, 1988.
- Edson RS, Terrell CL. The aminoglycosides. Mayo Clin Proc 74:519–28, 1999.
- Freeman CD, Nicolau DP, Belliveau PP, et al. Once-daily dosing of aminoglycosides: Review and recommendations for clinical practice. J Antimicrob Chemother 39:677–86, 1997.

"Usual dose" assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose  $\times$  1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

Krol V, Cunha BA, Schoch PE, Klein NC. Appropriateness of empiric gentamicin and vancomycin therapy for bacteremias in chronic dialysis outpatient units in the era of antibiotic resistance. *J Chemother* 18:490–3, 2006.

Munita JM, Arias CA, Murray BE. Enterococcus faecalis infective endocarditis: Is it time to abandon aminoglycosides? *Clin Infect Dis* 56:1269–1272, 2013.

Roushan MRH, Mohraz M, Hajiahmadi M, et al. Efficacy of gentamicin plus doxycycline versus streptomycin plus doxycycline in the treatment of brucellosis in humans. *Clin Infect Dis* 42:1075–1080, 2006.

Snydman, DR, McDermott LA, Jacobus NV. Evaluation of in vitro interaction of daptomycin with gentamicin or beta-lactam antibiotics against Staphylococcus aureus and Enterococci by FIC index and timed-kill curves. *J Chemother* 17:614–21, 2005.

Veinstein A, Venisse N, Badin J, et al. Gentamicin in hemodialyzed critical care patients: early dialysis after administration of a high dose should be considered. *Antimicrob Agents Chemo* 57:977–982, 2013.

Website: www.pdr.net

## Griseofulvin (Fulvicin, Grifulvin, Ultra, Gris-PEG, Grisactin)

**Drug Class:** Antifungal.

**Usual Dose:** 500 mg–1 gm (PO) q24h (microsize); 330–375 mg (PO) q24h (ultramicrosize).

**Pharmacokinetic Parameters:**

*Peak serum level:* 1–2 mcg/mL

*Bioavailability:* 50%

*Excreted unchanged (urine):* 1%

*Serum half-life (normal/ESRD):* 9/22 hrs

*Plasma protein binding:* 84%

*Volume of distribution ( $V_d$ ):* No data

**Primary Mode of Elimination:** Hepatic

**Dosage Adjustments\***

CrCl < 10 mL/min	No change
Post-HD or PD dose	None
CVWH/CVWHD/ CVWHDF dose	No change
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	Use with caution

**Drug Interactions:** Alcohol (↑ griseofulvin toxicity); barbiturates (↓ griseofulvin levels); oral contraceptives, warfarin (↓ interacting drug levels).

**Adverse Effects:** Photosensitivity reactions, rash, headache, nausea, vomiting, diarrhea, angular stomatitis, glossitis, leukopenia.

Allergic Potential: Moderate

**Safety in Pregnancy:** C

**Comments:** May exacerbate SLE/acute intermittent porphyria. Take microsize griseofulvin with fatty meal to ↑ absorption to ~ 70%. Ultramicrosize griseofulvin is absorbed 1.5 times better than microsize griseofulvin.

### REFERENCES:

Trepanier EF, Amsden GW. Current issues in onychomycosis. *Ann Pharmacotherapy* 32:204–14, 1998.  
Website: www.pdr.net

## Imipenem/Cilastatin (Primaxin)

**Drug Class:** Carbapenem.

**Usual Dose:** 1 gm (IV) q6h (see comments).

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

**Spectrum:** (see *Susceptibility Profiles*)

pp. 198–202).

**Resistance Potential:** High (*P. aeruginosa*, ↑ prevalence of MRSA)**Pharmacokinetic Parameters:***Peak serum level:* 42–116 mcg/ml*Bioavailability:* Not applicable*Excreted unchanged (urine):* 70%*Serum half-life (normal/ESRD):* 1/4 hrs*Plasma protein binding:* 20% / 40% (*cilastatin*)*Volume of distribution ( $V_d$ ):* 0.2 L/kg**Primary Mode of Elimination:** Renal**Dosage Adjustments\***

CrCl 40–70 mL/min	500 mg (IV) q6h
CrCl 20–40 mL/min	250 mg (IV) q8h
CrCl < 20 mL/min <sup>†</sup>	250 mg (IV) q12h
Post-HD dose	250 mg (IV)
Post-PD dose	250 mg (IV)
CVWH/CVHD/CVHDF dose	500 mg (IV) q8h
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	No change

† Avoid if CrCl ≤ 5 mL/min unless dialysis is instituted within 48 hours.

**Drug Interactions:** Cyclosporine (↑ cyclosporine levels); ganciclovir (↑ risk of seizures); probenecid (↑ imipenem levels); valproic acid (↓ seizure threshold).

**Adverse Effects:** Seizures, phlebitis.

**Allergic Potential:** Low

**Safety in Pregnancy:** C

**Comments:** Imipenem:cilastatin (1:1). Infuse 1 gm (IV) over 40–60 minutes. Imipenem is renally metabolized by dehydropeptidase I; cilastatin is an inhibitor of this enzyme, effectively preventing the metabolism of imipenem. Imipenem/cilastatin can be given IM (IM absorption: imipenem 75%; cilastatin 100%). **Fully susceptible organisms dose: 500 mg (IV) q6h; Less susceptible organisms (e.g., *P. aeruginosa*) dose: 1 gm (IV) q6–8h.** Incompatible in solutions containing vancomycin or metronidazole. **Seizures more likely in renal insufficiency/high doses (> 2 gm/d).** Inhibits endotoxin release from gram-negative bacilli. **Very low incidence of cross reactions with β-lactams.**

**Cerebrospinal Fluid Penetration:**

Non-Infamed meninges = 10%

Infamed meninges = 15%

**Bile Penetration:** 1%(also see **Antibiotic Pearls & Pitfalls** p. 511).**REFERENCES:**

- Balfour JA, Bryson HM, Brogden RN. Imipenem/cilastatin: An update of its antibacterial activity, pharmacokinetics, and therapeutic efficacy in the treatment of serious infections. *Drugs* 51:99–136, 1996.
- Barza M. Imipenem: First of a new class of beta-lactam antibiotics. *Ann Intern Med* 103:552–60, 1985.
- Fish DN, Teitelbaum, I, Abraham E. Pharmacokinetics and pharmacodynamics of imipenem during continuous renal replacement therapy in critically ill patients. *Antimicrobial Agents and Chemotherapy* 49:2421–2428, 2005.
- Garbino J, Villiger P, Caviezel A, et al. A randomized prospective study of cefepime plus metronidazole with imipenem-cilastatin in the treatment of intra-abdominal infections. *Infection* 35:161–166, 2007.

\*Usual dose\* assumes normal renal/hepatic function. \*For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

Helinger WC, Brewer NS. Carbapenems and monobactams: Imipenem, meropenem, and aztreonam. *Mayo Clin Proc* 74:420–34, 1999.

Lamoth F, Erard V, Asner S, et al. High imipenem blood concentrations associated with toxic encephalopathy in a patient with mild renal dysfunction. *Int J Antimicrob Agents* 34:386–388, 2009.

Lautenbach E, Weiner MG, Nachamkin I, et al. Imipenem resistance among *Pseudomonas aeruginosa* Isolates: risk factors for infection and impact of resistance on clinical and economic outcomes. *Infection Control and Hosp Epidemiol* 27:893–900, 2006.

Maravi-Poma E, Gener J, Alvarez-Lerma F, et al. Spanish group for the study of septic complications in severe acute pancreatitis. Early antibiotic treatment (prophylaxis) of septic complications in severe acute necrotizing pancreatitis: a prospective, randomized, multicenter study comparing two regimens with imipenem-cilastatin. *Intensive Care Med* 29:1974–80, 2003.

Miliani K, L'Heriteau F, Lacave L, et al. Imipenem and ciprofloxacin consumption as factors associated with high incidence rates of resistance *Pseudomonas aeruginosa* in hospitals in northern France. *J Hosp Infect* 77:343–347, 2011.

Saltoglu N, Dalkiran A, Tetiker T, et al. Piperacillin/tazobactam versus imipenem/cilastatin for severe diabetic foot infections: a prospective, randomized clinical trial in a university hospital. *Clin Microbiol Infect* 16:1252–1257, 2010.

Website: [www.pdr.net](http://www.pdr.net)

## Indinavir (Crixivan) IDV

**Drug Class:** HIV protease inhibitor.

**Usual Dose:** 800 mg (PO) q8h or 800 mg (PO) q12h with ritonavir 100–200 mg (PO) q12h.

**Pharmacokinetic Parameters:**

*Peak serum level:* 252 mcg/mL

*Bioavailability:* 65% (77% with food)

*Excreted unchanged (urine):* < 20%

*Serum half-life (normal/ESRD):* 2 hrs/no data

*Plasma protein binding:* 60%

*Volume of distribution ( $V_d$ ):* No data

**Primary Mode of Elimination:** Hepatic

**Dosage Adjustments\***

CrCl < 10 mL/min	No change
Post–HD dose	None
Post–PD dose	None
CVWH/CVWHD/ CVWHD dose	No change
Moderate hepatic insufficiency	600 mg (PO) q8h
Severe hepatic insufficiency	400 mg (PO) q8h

## Antiretroviral Dosage Adjustments

Atazanavir	Avoid
Didanosine	Administer didanosine 1 hour apart
Delavirdine	Indinavir 600 mg q8h
Efavirenz	Indinavir 1000 mg q8h
Lopinavir/ritonavir	Indinavir 600 mg q12h
Nelfinavir	Limited data for indinavir 1200 mg q12h + nelfinavir 1250 mg q12h
Nevirapine	Indinavir 1000 mg q8h
Ritonavir	Indinavir 800 mg q12h + ritonavir 100–200 mg q12h
Saquinavir	No information
Rifampin	Avoid combination
Rifabutin	Indinavir 1000 mg q8h; rifabutin 150 mg q24h or 300 mg 2–3x/week

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.



**Drug Interactions:** Antiretrovirals, rifabutin, rifampin (see dose adjustment grid, above); astemizole, terfenadine, benzodiazepines, cisapride, ergot alkaloids, statins, calcium channel blockers (↑ calcium channel blocker levels); carbamazepine, phenobarbital, phenytoin (↓ indinavir levels, ↑ anticonvulsant levels; monitor); tenofovir (↓ indinavir levels, ↑ tenofovir levels); clarithromycin, erythromycin, telithromycin (↑ indinavir and macrolide levels); didanosine (administer indinavir on empty stomach 1 hour apart); ethinyl estradiol, norethindrone (↑ interacting drug levels; no dosage adjustment); grapefruit juice (↓ indinavir levels); itraconazole, ketoconazole (↑ indinavir levels); sildenafil (↑ or ↓ sildenafil levels; do not exceed 25 mg in 48 hrs), tadalafil (max. 10 mg/72 hrs), vardenafil (max 2.5 mg/72 hrs); theophylline (↓ theophylline levels); fluticasone nasal spray (avoid concomitant use).

**Contraindications:** rifapentine, St. John's wort

**Adverse Effects:** Nephrolithiasis, nausea, vomiting, diarrhea, anemia, leukopenia, headache, insomnia, hyperglycemia (including worsening diabetes, new-onset diabetes, DKA), ↑ SGOT/SGPT, ↑ indirect bilirubin (2° to drug-induced Gilbert's syndrome; inconsequential), fat redistribution, lipid abnormalities (evaluate risk of coronary disease/pancreatitis), abdominal pain, night sweats, possible ↑ bleeding in hemophilia, dry skin, chelitis, paronychia.

**Allergic Potential:** Low

**Safety in Pregnancy:** C

**Comments:** Renal stone formation may be prevented/minimized by adequate hydration (1–3 liters water daily); ↑ risk of nephrolithiasis with alcohol. Take 1 hour before or 2 hours after meals (may take with skim milk or low fat meal). Separate dosing with ddl by 1 hour.

**Cerebrospinal Fluid Penetration:** 16%

#### REFERENCES:

- Antinori A, Giancola MI, Griseri S, et al. Factors influencing virological response to antiretroviral drugs in cerebrospinal fluid of advanced HIV-1-infected patients. *AIDS* 16:1867–76, 2002.
- Deeks SG, Smith M, Holodniy M, et al. HIV-1 protease inhibitors: A review for clinicians. *JAMA* 277:145–53, 1997.
- DiCenzo R, Forrest A, Fischl MA, et al. Pharmacokinetics of indinavir and nelfinavir in treatment-naive, human immunodeficiency virus-infected subjects. *Antimicrob Agents Chemother* 48:918–23, 2004.
- Hughes PJ, Cretton-Scott E, Teague A, et al. Protease inhibitors for patients with HIV-1 infection. *Pharm Ther* 36:332–344, 2011.
- Justesen US, Andersen AB, Klitgaard NA, et al. Pharmacokinetic interaction between rifampin and the combination of indinavir and low-dose ritonavir in HIV-infected patients. *Clin Infect Dis* 38:426–9, 2004.
- McDonald CK, Kuritzkes DR. Human immunodeficiency virus type 1 protease inhibitors. *Arch Intern Med* 157:951–9, 1997.
- Thompson MA, Aberg JA, Cahn P, et al. Antiretroviral treatment of adult HIV infection: 2012 recommendations of the International Antiviral Society — USA panel. *JAMA* 308:387–402, 2012.
- Website: [www.pdr.net](http://www.pdr.net)

## Isavuconazole (Cresemba)

**Drug Class:** Azole antifungal

**Usual Dose:** **Loading dose:** 200 mg (IV) q8h × 48 hrs, then **Maintenance dose:** 200 mg (IV/PO) q24h.

**Pharmacokinetic Parameters:**

*Peak serum level:* 1.85 – 2.56 mcg/ml

*Bioavailability (IV/PO):* 98%

*Excreted unchanged:* no data

*Serum half-life (normal/ESRD):* IV  $t_{1/2}$  = 76–104 hrs;

PO  $t_{1/2}$  = 56–77 hrs/no data

*Plasma protein binding:* 98%

*Volume of distribution ( $V_d$ ):* 4.4–7.7 L/kg

"Usual dose" assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

**Primary Mode of Elimination:** Hepatic

**Dosage Adjustments\***

CrCl < 30 mL/min	No change
Post-HD dose	No change
Post-PD dose	No change
CVVH dose	No change
Mild/Moderate Hepatic insufficiency	200 mg (IV) q8h × 48 hrs, then 100 mg (IV/PO) q24h
Severe hepatic insufficiency	Avoid use

**Drug Interactions:** CYP3A4 substrate; Concomitant use of CYP3A4 inhibitors and inducers is not recommended. No clinically relevant interactions with warfarin or cyclosporine.

**Adverse Effects:** Nausea, vomiting, diarrhea, pyrexia, headache, constipation, dyspnea, cough, febrile neutropenia, chills, fatigue, ↑ AST/ALT, hypokalemia, hypersensitivity reactions, infusion related reactions, and shortening of the QT interval.

**Allergic Potential:** Low (potential anaphylaxis/severe cutaneous reactions 1.9%)

**Safety in Pregnancy:** Unknown

**Comments:** Water soluble azole, which does not require cyclodextrin for IV solubility. Isavuconazonium, is a prodrug (water soluble) that is rapidly hydrolyzed in the blood, to the active moiety, isavuconazole (lipid soluble). Available as both IV and PO formulations. Only azole that does not prolong the QT interval. Following reconstitution, IV isavuconazonium

may spontaneously hydrolyze and precipitate as insoluble isavuconazole; an inline filter is recommended to remove precipitates. **Highly active against *Candida albicans*; non-albicans *Candida*, *Aspergillus*; Active against *Cryptococcus*; Some activity against *Mucor*, *Zygomycetes*.**

**Cerebrospinal Fluid Penetration:** < 10%

**Bile Penetration:** Good

**REFERENCES:**

- Faici DR, Pasqualotto AC. Profile of isavuconazole and its potential in the treatment of severe invasive fungal infections. *Infection and Drug Resistance*. 6:163-174, 2013.
- Gregson L, Goodwin J, Johnson A, et al. In vitro susceptibility of *Aspergillus fumigatus* to isavuconazole: correlation with itraconazole, voriconazole, and posaconazole. *Antimicrob Agents Chemo*. 57:5778-5780, 2013.
- Katragkou A, McCarthy M, Meletiadis J, et al. In vitro combination of isavuconazole with micafungin or amphotericin B deoxycholate against medically important molds. *Antimicrob Agents Chemo*. 58:6934-6937, 2014.
- Livormore J, Hope W. Evaluation of the pharmacokinetics and clinical utility of isavuconazole for treatment of invasive fungal infections. *Expert Opinion on Drug Metab & Tox*. 8:759-765, 2012.
- Schmitt-Hoffmann A, Roos B, Heep M, et al. Single-ascending-dose pharmacokinetics and safety of the novel broad-spectrum antifungal triazole BAL4815 after intravenous infusions (50, 100, and 200 milligrams) and oral administrations (100, 200, and 400 milligrams) of its prodrug, BAL8557, in healthy volunteers. *Antimicrob Agents Chemo*. 50:279-285, 2006.
- Schmitt-Hoffmann A, Roos B, Maeres J, et al. Multiple-dose pharmacokinetics and safety of the new antifungal triazole BAL4815 after intravenous infusion and oral administration of its prodrug, BAL8557, in healthy volunteers. *Antimicrob Agents Chemo*. 50:286-293, 2006.

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

Schmitt-Hoffmann A, Roos B, Spickermann J, et al.

Effect of mild and moderate liver disease on the pharmacokinetics of isavuconazole after intravenous and oral administration of a single dose of the prodrug BAL8557. *Antimicrob Agents Chemo*. 53:4885-4890, 2009.

Seyedmousavi S, Venweij PE, Mouton JW. Isavuconazole, a broad-spectrum triazole for the treatment of systemic fungal diseases. *Expert Review Anti-infective Therapy*. 13:9-27, 2015.

Thompson GR, 3rd, Wederhold NP. Isavuconazole: a comprehensive review of spectrum of activity of a new triazole. *Mycopathologia*. 170:291-313, 2010.

Website: www.pdr.net

## Isoniazid (INH)

**Drug Class:** Anti-TB drug.

**Usual Dose:** 300 mg (PO) q24h or 5 mg/kg (see comments).

### Pharmacokinetic Parameters:

*Peak serum level:* 7 mcg/ml

*Bioavailability:* 90%

*Excreted unchanged (urine):* 50–70%

*Serum half-life (normal/ESRD):* 1/1 hr

*Plasma protein binding:* 15%

*Volume of distribution ( $V_d$ ):* 0.75 L/kg

**Primary Mode of Elimination:** Hepatic

### Dosage Adjustments\*

CrCl < 10 mL/min	No change
Post-HD dose	300 mg
Post-PD dose	300 mg
CWVH/CVWHD/CVWHDf dose	None
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	Use with caution

**Drug Interactions:** Alcohol, rifampin (↑ risk of ↑ SGOT/SGPT); alfentanil (↑ duration of alfentanil effect); aluminum salts (↓ isoniazid absorption); carbamazepine, phenytoin (↑ interacting drug levels); itraconazole (↓ itraconazole levels); warfarin (↑ INR).

**Adverse Effects:** ↑ SGOT/SGPT, drug fever/rash, age-dependent hepatotoxicity (after age 60), drug-induced ANA/SLE, hemolytic anemia, neuropsychiatric changes in the elderly. Transient/reversible ↑ SGOT/SGPT frequently occur early (< 3 weeks) after INH use. If ↑ SGOT/SGPT, monitor levels twice weekly until levels peak, then monitor levels weekly until levels return to within normal range. D/C INH if ↑ SGOT/SGPT ≥ 10 × upper limit of normal. Fulminant hepatic failure (rare).

**Allergic Potential:** Low

**Safety in Pregnancy:** C

**Comments:** Administer with 50 mg of pyridoxine daily to prevent peripheral neuropathy. Increased blood pressure/rash with tyramine-containing products, e.g., cheese/wine. ↑ hepatotoxicity in slow acetylators.

**Slow acetylator dose:** 150 mg (PO) q24h.

**TB D.O.T. dose:** 15 mg/kg or 900 mg (PO) 3x/week.

Meningeal dose = usual dose.

### Cerebrospinal Fluid Penetration:

Non-Inflamed meninges = 90%

Inflamed meninges = 90%

### REFERENCES:

CDC. Severe Isoniazid-associated liver injuries among persons being treated for latent tuberculosis infection. *MMWR* 59:224–229, 2010.

Ena J, Valls V. Short-course therapy with rifampin plus isoniazid, compared with standard therapy with isoniazid, for latent tuberculosis infection: A meta-analysis. *Clin Infect Dis* 40:670–6, 2005.

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

- McIllerson H, Wash P, Burger A, et al. Determinants of rifampin, isoniazid, pyrazinamide and ethambutol pharmacokinetics in a cohort of tuberculosis patients. *Antimicrob Agents Chemother* 50:1170–77, 2006.
- Pina JM, Clotel L, Ferrer A, et al. Cost-effectiveness of rifampin for 4 months and isoniazid for 9 months in the treatment of tuberculosis infection. *Eur J Clin Microbiol Inf Dis* 32:647–655, 2013.
- Schaller A, Sun Z, Yang Y, et al. Salicylate reduces susceptibility of *Mycobacterium tuberculosis* to multiple antituberculosis drugs. *Antimicrob Agents Chemother* 46:2533–9, 2002.
- Schlossberg D. An approach to drug-induced liver injury during tuberculosis therapy. *Infect Dis Clin Pract* 18:107–108, 2010.
- Schlossberg D. Treatment of multi-drug resistant tuberculosis. *Antibiotics for Clinicians* 9:317–321, 2005.
- Van Scoy RE, Wilkowske CJ. Antituberculous agents. *Mayo Clin Proc* 67:179–87, 1992.
- Website: [www.pdr.net](http://www.pdr.net)

## Itraconazole (Sporanox)

**Drug Class:** Antifungal.

**Usual Dose:** 200 mg (PO, capsules or solution, solution produces better absorption) q12h or q24h (Take without food). Depending on disease, due to enhanced drug delivery, IV therapy begins with 200 mg IV q12h × 2 days (4 doses) and then continues with only 200 mg (IV) q24h; PO follow-up to IV therapy for serious infection is at 200 mg PO q12h. Each IV dose should be infused over 1 hour (see comments).

### Pharmacokinetic Parameters:

- Peak serum level:* 2.8 mcg/ml  
*Bioavailability:* 55% (capsules)/90% (solution)  
*Excreted unchanged (urine):* 1%  
*Serum half-life (normal/ESRD):* 21–64/35 hrs  
*Plasma protein binding:* 99.8%  
*Volume of distribution ( $V_d$ ):* 10 L/kg

**Primary Mode of Elimination:** Hepatic; metabolized predominantly by the cytochrome P450 3A4 isoenzyme system (CYP3A4)

### Dosage Adjustments\*

CrCl > 30 mL/min	No change
CrCl < 30 mL/min	No change for (PO); avoid (IV) due to ↑ cyclodextrin
Post-HD dose	100 mg (IV/PO)
Post-PD dose	None
CVWH/CVHHD/ CVHDF dose	No change
Moderate hepatic insufficiency	No change <sup>†</sup>
Severe hepatic insufficiency	Use with caution

† ↑  $t_{1/2}$  of itraconazole in patients with hepatic insufficiency should be considered when given with medications metabolized by P450 isoenzymes. Also see Adverse Effects for information regarding patients who develop liver dysfunction.

**Drug Interactions:** *Itraconazole may* ↑ *plasma levels of:* alfentanil, buspirone, busulfan, carbamazepine, cisapride, cyclosporine, digoxin, dihydropyridines, docetaxel, dofetilide, methylprednisolone, oral hypoglycemics (↑ risk of hypoglycemia), pimozone, quinidine, rifabutin, saquinavir, sirolimus, tacrolimus, trimetrexate, verapamil, vinca alkaloids, warfarin; alprazolam, diazepam, midazolam, triazolam (↑ sedative/hypnotic effects); atorvastatin, lovastatin, simvastatin (↑ risk of rhabdomyolysis); indinavir, ritonavir, saquinavir; coadministration of oral midazolam, triazolam, lovastatin, or simvastatin with itraconazole is contraindicated; coadministration of cisapride, pimozone, quinidine, amiodarone, or dofetilide with itraconazole is contraindicated due to the risk of ↑ QT<sub>c</sub>/life-threatening ventricular arrhythmias. *Decreased*

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

*itraconazole levels may occur with:* antacids, carbamazepine, H<sub>2</sub>-receptor antagonists, isoniazid, nevirapine, phenobarbital, phenytoin, proton pump inhibitors, rifabutin, rifampin; coadministration of rifampin with itraconazole is not recommended. *Increased itraconazole levels may occur with:* clarithromycin, erythromycin, indinavir, ritonavir.

**Adverse Effects:** ≥ 2%: nausea, diarrhea, vomiting, headache, abdominal pain, bilirubinemia, rash, ↑ SGPT/SGOT, hypokalemia, ↑ serum creatinine. Rarely, itraconazole has been associated with serious hepatotoxicity (liver failure/death). If liver disease develops, discontinue treatment, perform liver function testing, and reevaluate risk/benefit of further treatment. Use itraconazole with caution in patients with ↑ liver enzymes, active liver disease, or previous drug-induced hepatotoxicity. Life-threatening ventricular arrhythmias/sudden death have occurred in patients using cisapride, pimozide, or quinidine concomitantly with itraconazole; coadministration of these drugs with itraconazole is contraindicated. Use itraconazole with caution in patients with ventricular dysfunction. IV itraconazole may cause transient, asymptomatic ↓ in ejection fraction for ≤ 12 hours. If CHF develops, consider discontinuation of itraconazole.

**Allergic Potential:** Low

**Safety in Pregnancy:** C

**Comments:** **Oral itraconazole:** Requires gastric acidity for absorption. When antacids are required, administer ≥ 1 hour before or 2 hours after itraconazole capsules. Oral solution is better absorbed without food; capsules are better absorbed with food. Capsule bioavailability is food dependent: 40% fasting/90% postprandial. For oral therapy, bioavailability of 10 ml of solution without food = 100 mg capsule with food. Administer with a cola beverage in

patients with achlorhydria or taking H<sub>2</sub>-receptor antagonists/other gastric acid suppressors. While oral solution and capsules can be interchanged for treatment of systemic disease if adequate blood levels are achieved, oral solution produces more reliable blood levels and is definitely preferred for oral/esophageal candidiasis where the local effect of the solution on the infection seem helpful. **IV itraconazole:** Hydroxypropyl-β-cyclodextrin stabilizer in IV formulation accumulates in renal failure. IV itraconazole should not be used in patients with CrCl < 30 mL/min; if possible, use the oral preparation. Infuse 60 ml of dilute solution (3.33 mg/ml = 200 mg itraconazole, pH ~ 4.8) IV over 60 minutes, using infusion set provided. After administration, flush the infusion set with 15–20 ml of normal saline injection. The compatibility of IV itraconazole with flush solutions other than normal saline is unknown. **Highly active against C. albicans, non-albicans Candida, Cryptococcus, Histoplasmosis, Blastomycosis, Paracoccidiomycosis, Sporotrichosis. Active against dermatophytes. No activity against Pseudoallescheria/Scedosporium, Fusaria or Mucor.**

**Cerebrospinal Fluid Penetration:** < 10%

**Bile Penetration:** 3–18%

#### REFERENCES:

- Boogaerts M, Winston DJ, Bow EJ, et al. Intravenous and oral itraconazole versus intravenous amphotericin B deoxycholate as empirical antifungal therapy for persistent fever in neutropenic patients with cancer who are receiving broad-spectrum antibacterial therapy. A randomized controlled trial. *Ann Intern Med* 135:412–22, 2001.
- Bruggemann RJM, Alffenaar JWC, Blijlevens NMA, et al. Clinical relevance of the pharmacokinetic interactions of azole antifungal drugs with coadministered agents. *Clin Infect Dis* 48:1441–1458, 2009.

"Usual dose" assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

- Calvopina M, Guevara AG, Armijos RX, et al. Itraconazole in the treatment of New World mucocutaneous leishmaniasis. *Int J Dermatol* 43:659–63, 2004.
- Caputo R. Itraconazole (Sporanox) in superficial and systemic fungal infections. *Expert Rev Anti Infect Ther* 1:531–42, 2004.
- Conte JE Jr, Golden JA, Kipps J, et al. Intrapulmonary pharmacokinetics and pharmacodynamics of itraconazole and 14-hydroxyitraconazole steady state. *Antimicrob Agents Chemother* 48:3823–7, 2004.
- Dominguez-Gil Hurlle A, Sanchez Navarro A, Garcia Sanchez MJ. Therapeutic drug monitoring of itraconazole and the relevance of pharmacokinetic interactions. *Clin Microbiol and Infection* 12:597–106, 2006.
- Glasmacher A, Prentice A. Current experience with itraconazole in neutropenic patients: a concise overview of pharmacological properties in use in prophylactic and empirical antifungal therapy. *Clin Microbiol and Infection* 12:584–90, 2006.
- Go J, Cunha BA. Itraconazole. *Antibiotics for Clinicians* 3:61–70, 1999.
- Horusseau JL, Dekker AW, et al. Itraconazole oral solution for primary prophylaxis of fungal infections in patients with hematological malignancy and profound neutropenia: a randomized, double-blind, double-placebo, multicenter trial comparing itraconazole and amphotericin B. *Antimicrob Agents Chemother* 44:1887–93, 2000.
- Marks DI, Pagliuca A, Kibbler CC, et al. Voriconazole versus itraconazole for antifungal prophylaxis following allogeneic haematopoietic stem cell transplantation. *Br J Haematol* 155:318–327, 2011.
- Mattiuzzi GN, Alvarado G, Giles FJ, et al. Open-label, randomized comparison of itraconazole versus caspofungin for prophylaxis in patients with hematologic malignancies. *Antimicrob Agents Chemother* 50:143–7, 2006.
- Phillips EJ. Itraconazole was as effective as amphotericin B for fever and neutropenia in cancer and led to fewer adverse events. *ACP J Club* 136:58, 2002.
- Rubin MA, Carroll KC, Cahill BC. Caspofungin in combination with itraconazole for the treatment of invasive aspergillosis in humans. *Clin Infect Dis* 34:1160–1, 2002.
- Sabatelli F, Patel R, Mann PA, et al. In vitro activities of posaconazole, fluconazole, itraconazole, voriconazole, and amphotericin B against a large collection of clinically important molds and yeasts. *Antimicrob Agents Chemother* 50:2009–15, 2006.
- Terrell CL. Antifungal agents Part II. The azoles. *Mayo Clin Proc* 74:78–100, 1999.
- Urnsak M, Ilkit M, Evruke C, et al. Clinical and mycological efficacy of single-day oral treatment with itraconazole (400 mg) in acute vulvovaginal candidosis. *Mycoses* 47:422–7, 2004.
- Winston DJ. Itraconazole vs. fluconazole for antifungal prophylaxis in allogeneic stem-cell transplant patients 12:591–96, 2006.
- Winston DJ, Busuttill RW. Randomized controlled trial of oral itraconazole solution versus intravenous/oral fluconazole for prevention of fungal infections in liver transplant recipients. *Transplantation* 74:688–95, 2002.
- Website: [www.pdr.net](http://www.pdr.net)

## Ketoconazole (Nizoral)

**Drug Class:** Antifungal.

**Usual Dose:** 200 mg (PO) q24h.

**Pharmacokinetic Parameters:**

*Peak serum level:* 3.5 mcg/ml

*Bioavailability:* 82%

*Excreted unchanged (urine):* 2–4%

*Serum half-life (normal/ESRD):* 6/20 hrs

*Plasma protein binding:* 99%

*Volume of distribution (V<sub>d</sub>):* 2 L/kg

**Primary Mode of Elimination:** Hepatic

**Dosage Adjustments\***

CrCl < 40 mL/min	No change
Post-HD dose	None
Post-PD dose	None
CVWH/CVWHD/ CVWHDf dose	No change
Moderate hepatic insufficiency	Use with caution
Severe hepatic insufficiency	Avoid

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

**Drug Interactions:** Astemizole, cisapride, terfenadine, amiodarone (may ↑ QT interval, torsades de pointes); carbamazepine, INH (↓ ketoconazole levels); cimetidine, famotidine, nizatidine, ranitidine, omeprazole, INH (↓ ketoconazole absorption); cyclosporine, digoxin, loratadine, tacrolimus (↑ interacting drug levels with possible toxicity); didanosine (↓ ketoconazole levels); midazolam, triazolam (↑ interacting drug levels, ↑ sedative effects); oral hypoglycemics (severe hypoglycemia); phenytoin, rifabutin, rifampin (↓ ketoconazole levels, ↑ interacting drug); statins (↑ statin levels; rhabdomyolysis reported); warfarin (↑ INR).

**Adverse Effects:** Nausea, vomiting, abdominal pain, pruritus.

**Allergic Potential:** Low

**Safety in Pregnancy:** C

**Comments:** Dose-dependent reduction in gonadal (androgenic) function. Decreased cortisol production with doses ≥ 800 mg/day, but **does not result in adrenal insufficiency**. Give oral doses with citric juices.

**Highly active against dermatophytes, Malassezia, Candida albicans. Some activity against Aspergillus, Coccidiomycosis, Paracoccidiomycosis, Histoplasmosis. No activity against Pseudoallescheria/Scedosporium, Fusaria or Mucor.**

**Cerebrospinal Fluid Penetration:** < 10%

#### REFERENCES:

- Allen LV. Ketoconazole oral suspension. *US Pharm* 18:98–9, 1993.
- Kaur IP, Kakkar S. Topical delivery of antifungal agents. *Expert Opin Drug Deliv* 7:1303–1327, 2010.
- Terrell CL. Antifungal agents: Part II. The azoles. *Mayo Clin Proc* 74:78–100, 1999.
- Zhao Y, Li L, Wang JJ, et al. Cutaneous malasseziasis: four case reports of atypical dermatitis and onychomycosis caused by Malassezia. *Int J Dermatol* 49:141–145, 2010.
- Website: www.pdr.net

## Lamivudine (Epivir) 3TC

**Drug Class:** HIV NRTI (nucleoside reverse transcriptase inhibitor); HBV antiviral.

**Usual Dose:** 150 mg (PO) q12h or 300 mg (PO) q24h (HIV); 100 mg (PO) q24h (HBV).

#### Pharmacokinetic Parameters:

*Peak serum level:* 1.5 mcg/mL

*Bioavailability:* 86%

*Excreted unchanged (urine):* 71%

*Serum half-life (normal/ESRD):* 5–7/20 hrs

*Plasma protein binding:* 36%

*Volume of distribution (V<sub>d</sub>):* 1.3 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\* (HIV/HBV dose)**

CrCl 30–50 mL/min	150 mg (PO) q24h /100 mg (PO), then 50 mg (PO) q24h
CrCl 15–30 mL/min	100 mg (PO) q24h /100 mg (PO), then 25 mg (PO) q24h
CrCl 5–15 mL/min	50 mg (PO) q24h /35 mg (PO), then 15 mg (PO) q24h
CrCl < 5 mL/min	25 mg (PO) q24h /35 mg (PO), then 10 mg (PO) q24h
Post-HD dose	25 mg (PO)/10 mg (PO)
Post-PD dose	None
CVWH/CVWHD/ CVWHD dose	100 mg (PO) q24h /100 mg (PO), then 25 mg (PO) q24h
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	Use with caution

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

**Drug Interactions:** Didanosine, zalcitabine (↑ risk of pancreatitis); TMP-SMX (↑ lamivudine levels); zidovudine + lamivudine not recommended.

**Adverse Effects:** Lamivudine tablets and oral solution used to treat HIV infection contain a higher dose of lamivudine than Epiriv-HBV® tablets and oral solution used to treat chronic HBV. Severe acute exacerbations of HBV have been reported in patients who are co-infected with HBV and HIV and have discontinued lamivudine. Hepatic function should be monitored closely for at least several months in patients who discontinue lamivudine. Most common include headache, dizziness, malaise, nausea nasal congestion, cough, rash, neuropathy, leucopenia. Pancreatitis and or lactic acidosis with hepatic steatosis (rare but potentially life threatening toxicity with NRTI's). Immune reconstitution syndrome can occur.

**Allergic Potential:** Low

**Safety in Pregnancy:** C

**Comments:** Potential cross resistance with didanosine. Prevents development of AZT resistance and restores AZT susceptibility. May be taken with or without food. Although effective against HBV, lamivudine is no longer a first-line option because of the development of resistance in 70% of treated patients.

**Cerebrospinal Fluid Penetration:** 15%

#### REFERENCES:

- Diaz-Brito V, Leon A, Knobel H, et al. Post-exposure prophylaxis for HIV-infection: a clinical trial comparing lopinavir/ritonavir versus atazanavir each with zidovudine/lamivudine. *Antivir Ther* 17:337-46, 2012.
- Leung N. Lamivudine for chronic hepatitis B. *Expert Rev Anti Infect Ther* 2:173-80, 2004.
- Liaw YF, Sung JY, Chow WC, et al. Lamivudine for patients with chronic hepatitis B and advanced liver disease. *N Engl J Med* 351:1521-31, 2004.

Marrone A, Zampino R, D'Onofrio M, et al. Combined interferon plus lamivudine treatment in young patients with dual HBV (HBeAg positive) and HCV chronic infection. *J Hepatol* 41:1064-5, 2004.

Nishijima T, Tsukada K, Teruya K, et al. Efficacy and safety of once-daily ritonavir-based antiretroviral therapy in low-body weight treatment-naïve patients with HIV infections. *AIDS* 13;26:649-51 2012.

Perry CM, Faulds D. Lamivudine. A review of its antiviral activity, pharmacokinetic properties and therapeutic efficacy in the management of HIV infection. *Drugs* 53:657-80, 1997.

Sax PE, Tierney C, Collier AC, et al. Abacavir/lamivudine versus tenofovir DF/emtricitabine as part of combination regimens for initial treatment of HIV: final results. *J Infect Dis* 204:1191-201, 2011.

Schmilovitz-Weiss H, Ben-Ari Z, Sikuler E, et al. Lamivudine treatment for acute severe hepatitis B: a pilot study. *Liver Int* 24:547-51, 2004.

Sheng YJ, Liu JY, Tong SW, et al. Lamivudine plus adefovir combination therapy versus entecavir monotherapy for lamivudine-resistant chronic hepatitis B: a systematic review and meta-analysis. *Virology* 43:393, 2011.

Thompson MA, Aberg JA, Cahn P, et al. Antiretroviral treatment of adult HIV infection: 2012 recommendations of the International Antiviral Society — USA panel. *JAMA* 308:387-402, 2012.

Website: www.pdr.net

## Lamivudine + Zidovudine (Combivir) 3TC/ZDV

**Drug Class:** HIV NRTIs combination.

**Usual Dose:** Combivir tablet = 150 mg lamivudine + 300 mg zidovudine. Usual dose = 1 tablet (PO) q12h.

**Pharmacokinetic Parameters:**

*Peak serum level:* 2.6/1.2 mcg/mL

*Bioavailability:* 82/60%

*Excreted unchanged (urine):* 86/64%

*Serum half-life (normal/ESRD):* [6/1.1]/[20/2.2] hrs

*Plasma protein binding:* <36/<38%

*Volume of distribution (V<sub>d</sub>):* 1.3/1.6 L/kg

"Usual dose" assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.



**Primary Mode of Elimination:** Renal

**Dosage Adjustments\***

CrCl 50–80 mL/min	No change
CrCl 10–50 mL/min	Avoid
CrCl < 10 mL/min	Avoid
Post-HD dose	Avoid
Post-PD dose	Avoid
CWVH/CVWHD/CVWHDf dose	Avoid
Moderate hepatic insufficiency	Avoid
Severe hepatic insufficiency	Avoid

**Drug Interactions:** Atovaquone (↑ zidovudine levels); stavudine (antagonist to stavudine; avoid combination); ganciclovir, doxorubicin (neutropenia); tipranavir (↓ zidovudine levels); TMP-SMX (↑ lamivudine and zidovudine levels); vinca alkaloids (neutropenia).

**Adverse Effects:** Zidovudine, has been associated with hematologic toxicity including neutropenia and severe anemia. Prolonged use of zidovudine has been associated with symptomatic myopathy. Lactic acidosis and severe hepatomegaly with steatosis, including fatal cases. Severe acute exacerbations of hepatitis B have been reported in patients who are co-infected with hepatitis B virus (HBV) and HIV and have discontinued lamivudine. Nausea, vomiting, diarrhea, anorexia, insomnia fever/chills, headache, malaise/fatigue, peripheral neuropathy and pancreatitis. Immune reconstitution syndrome and redistribution/accumulation of body fat have been reported in patients treated with combination antiretroviral therapy.

**Allergic Potential:** Low

**Safety in Pregnancy:** C.

**Cerebrospinal Fluid Penetration:**

Lamivudine = 12%; Zidovudine = 60%

**REFERENCES:**

- [No authors listed.] Drugs for AIDS and associated infections. *Med Lett Drug Ther* 35:79–86, 1993.
- Hirsch MS, D'Aquila RT. Therapy for human immunodeficiency virus infection. *N Engl J Med* 328:1685–95, 1993.
- McLeod GX, Hammer SM. Zidovudine: Five years later. *Ann Intern Med* 117:487–510, 1992.
- Panel on Antiretroviral Guidelines for Adults and Adolescents. Department of Health and Human Services. Guidelines for the use of antiretroviral agents in HIV-infected adults and adolescents. <http://aidsinfo.nih.gov/ConsentFiles/AdultandAdolescentGL.pdf>, 2012.
- Thompson MA, Aberg JA, Cahn P, et al. Antiretroviral treatment of adult HIV infection: 2012 recommendations of the International Antiviral Society — USA panel. *JAMA* 308:387–402, 2012.
- Website: [www.pdr.net](http://www.pdr.net)

**Leflunomide (Arava)**

**Drug Class:** CMV antiviral.

**Usual Dose:** 100 mg/day (PO) × 3 days, then 20 mg (PO) q24h.

**Pharmacokinetic Parameters:**

*Peak serum level:* 8 mcg/mL

*Bioavailability:* 80%

*Excreted unchanged:* 0%

*Serum half-life (normal/ESRD):* 14 days/no data

*Plasma protein binding:* 99%

*Volume of distribution (V<sub>d</sub>):* 0.13 L/kg

**Primary Mode of Elimination:** Hepatic/renal

**Dosage Adjustments\***

CrCl 15–30 mL/min	Use with caution
CrCl < 15 mL/min	Use with caution
Post-HD dose	Use with caution
Post-PD dose	Use with caution
CVWH/CVWHD/ CVWHDf dose	Use with caution
Moderate—severe hepatic insufficiency	Avoid if ↑ SGOT/ SGPT > 2 × n or pre-existing liver disease

\*Usual dose\* assumes normal renal/hepatic function. \*For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

**Drug Interactions:** May inhibit CYP 2C9; rifampin (↑ levels of active metabolite, M1); hepatotoxic drugs.

**Adverse Effects:** Headache, nausea, diarrhea, hypertension; hepatotoxicity, bone marrow suppression, rash, Stevens-Johnson Syndrome, toxic epidermal necrolysis, interstitial lung disease.

**Allergic Potential:** Low

**Safety in Pregnancy:** X

**Comments:** Alternate therapy of CMV infection after 1<sup>st</sup> line therapies fail; do not use with pre-existing liver disease; metabolized to active metabolite (M1); not recommended in patients with severe, uncontrolled infections, bone marrow dysplasia or severe immunodeficiency, prolonged administration leads to significant accumulation, (drug elimination procedure with cholestyramine described in PI). Screen for TB before starting leflunomide; avoid live vaccinations while on leflunomide; monitor LFT's and CBCs. **Discontinue if ↑ SGOT/SGPT > 3 × n.**

#### REFERENCES:

- Avery RK, et al. Utility of leflunomide in the treatment of complex cytomegalovirus syndromes. *Transplantation* 90:419–26, 2010.
- Chacko B, John GT. Leflunomide for cytomegalovirus: bench to bedside. *Transpl Infect Dis* 14:111–20, 2012.
- Chon WJ, Josephson MA. Leflunomide in renal transplantation. *Expert Rev Clin Immunol* 7:273–281, 2011.
- Henao-Martinez AF, Weinberg A, Waldman WJ, et al. Successful treatment of acyclovir-resistant herpes simplex virus type 2 proctitis with leflunomide in an HIV-infected man. *J Clin Virol* 54:276–278, 2012.
- John GT, et al. Leflunomide therapy for cytomegalovirus disease in renal allograft recipients. *Transplantation* 77:1460–61, 2004.
- Website: www.pdr.net

**Spectrum:** (see **Susceptibility Profiles** pp. 198–202).

**Resistance Potential:** Low

**Pharmacokinetic Parameters:**

*Peak serum level:* 5–8 mcg/ml

*Bioavailability:* 99%

*Excreted unchanged (urine):* 87%

*Serum half-life (normal/ESRD):* 7 hrs/40 hrs

*Plasma protein binding:* 30%

*Volume of distribution (V<sub>d</sub>):* 1.3 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\*** (based on 750 mg/500 mg (IV/PO) q24h)

CrCl > 50 mL/min	No change
CrCl 20–50 mL/min	750 mg (IV/PO) q48h /500 mg (IV/PO), then 250 mg (IV/PO) q24h
CrCl < 20 mL/min	750 mg (IV/PO), then 500 mg (IV/PO) q48h /500 mg (IV/PO), then 250 mg (IV/PO) q48h
HD/CAPD dose	500 mg (IV/PO) q48h
Post–HD dose	None
Post–HFHD dose	250 mg (IV/PO)
Post–PD dose	250 mg (IV/PO)
CVWH/CVWHD/CVWHDf dose	500 mg (IV/PO) q48h/250 mg (IV/PO) q24h
HD q 48–72h dose	500 mg (IV/PO) q HD*
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	No change

\*Give at end of HD.

## Levofloxacin (Levaquin)

**Drug Class:** Respiratory quinolone.

**Usual Dose:** 500–750 mg (IV/PO) q24h (see comments).

**Drug Interactions:** Al<sup>+</sup>, Fe<sup>+</sup>, Mg<sup>+</sup>, Zn<sup>+</sup> antacids (↓ absorption of levofloxacin if taken together); NSAIDs (CNS stimulation); probenecid (↑ levofloxacin levels); warfarin (↑ INR).

"Usual dose" assumes normal renal/hepatic function. \*For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

**Adverse Effects:** Headache, insomnia, delirium (rare) nausea, constipation, delirium (rare) *C. difficile* diarrhea/colitis. **Does not cause seizures.** FQ use has an increased risk of tendinitis/tendon rupture. Risk is highest in the elderly, those on steroids, and those with heart, lung, or renal transplants. Peripheral neuropathy may occur early and may be permanent.

**Allergic Potential:** Low

**Safety in Pregnancy:** C

**Comments:** Take 2 hours before or after aluminum/magnesium-containing antacids. Does not increase digoxin concentrations. May ↑ QTc interval, particularly in patients with predisposing conditions or medications e.g., amiodarone. **Acute bacterial sinusitis dose:** 750 mg (PO) q24h × 5 days. **CAP dose:** 750 mg (IV/PO) q24h × 5 days. **Nosocomial pneumonia dose:** 750 mg (IV/PO) q24h × 7–14 days. **cSSII dose:** 750 mg (IV/PO) q24h × 7–14 days. **TB dose** (alternate drug in MDR TB regimen): 500 mg (PO) q24h. **Complicated UTI/acute pyelonephritis dose:** 750 mg (IV/PO) q24h × 5 days.

↑ **incidence of *C. difficile* diarrhea/colitis with PPIs (for patients on PPIs during FQ therapy, D/C PPIs or switch to H<sub>2</sub> blocker for duration of FQ therapy).**

Effective against oral anaerobes/above the waist but not anaerobes below the waist e.g. *B. fragilis*.

**Cerebrospinal Fluid Penetration:** 16%

**Bile Penetration:** 100%

(also see **Antibiotic Pearls & Pitfalls** p. 516).

#### REFERENCES:

Alvarez-Lerma F, Grau S, Alvarez-Beltran. Levofloxacin in the treatment of ventilator-associated pneumonia. *Clin Microbiol Infect* 12:81–92, 2006.

Bucaneve G, Micozzi A, Menichetti F, et al. Levofloxacin to prevent bacterial infection in patients with cancer and neutropenia. *N Engl J Med* 353:977–87, 2005.

Chou HW, Wang JL, Chang CH, et al. Risk of severe dysglycemia among diabetic patients receiving levofloxacin, ciprofloxacin, or moxifloxacin in Taiwan. *Clin Infect Dis* 57:971–980, 2013.

Cunha BA, Lenopoli S, Hage JE. Klebsiella pneumoniae septic wrist arthritis successfully treated with ertapenem and levofloxacin. *J Chemother* 23:376–7, 2011.

File TM Jr, Milkovich G, Tenenber AM, et al. Clinical implications of 750 mg, 5-day levofloxacin for the treatment of community-acquired pneumonia. *Curr Med Res Opin* 20:1473–81, 2004.

Garcia-Vazquez E, Mensa J, Sarasa M, et al. Penetration of levofloxacin into the anterior chamber (aqueous humor) of the human eye after intravenous administration. *Eur J Clin Microbiol Infect Dis* 26:137–140, 2007.

Klausner HA, Brown P, Peterson J, et al. Double-blind, randomized comparison of levofloxacin 750 mg once daily for 5 days versus ciprofloxacin for 10 days in the treatment of acute pyelonephritis. *Curr Med Res Opin* 23:2637–2645, 2007.

Kuriyama T, Williams DW, Yanagisawa M, et al. Antimicrobial susceptibility of 800 anaerobic isolates from patients with dentoalveolar infection to 13 oral antibiotics. *Oral Microbiol Immunol* 22:285–288, 2007.

Lee CK, Boyle MP, Diener-West M, et al. Levofloxacin pharmacokinetics in adult cystic fibrosis. *Chest* 131:796–802, 2007.

Marchetti F, Viale P. Current and future perspectives for levofloxacin in severe *Pseudomonas aeruginosa* infections. *J Chemotherapy* 15:315–322, 2003.

Maurin M, Raoult D. Bacteriostatic and bactericidal activity of levofloxacin against *Rickettsia rickettsii*, *Rickettsia conorii*, 'Israel spotted fever group rickettsii' and *Coxiella burnetii*. *Journal of Antimicrobial Chemotherapy* 39:725–30, 1997.

Nightingale CH, Grant EM, Quintiliani R. Pharmacodynamics and pharmacokinetics of levofloxacin. *Chemotherapy* 46:6–14, 2000.

Nista EC, Candelli M, Zocco, MA, et al. Levofloxacin-based triple therapy in first-line treatment for *Helicobacter pylori* eradication. *Am J Gastroenterol* 101:1985–90, 2006.

Ogrendik M. Levofloxacin treatment in patients with rheumatoid arthritis receiving methotrexate. *South Med J* 100:135–139, 2007.

Pankey GA, Ashcraft DS. In vitro synergistic/additive activity of levofloxacin with meropenem against *Stenotrophomonas maltophilia*. *Diagn Microbiol Infect Dis* 67:297–300, 2010.

Pankuch GA, Seifert H, Appelbaum PC. Activity of doripenem with and without levofloxacin, amikacin, and colistin against *Pseudomonas aeruginosa* and *Acinetobacter baumannii*. *Diagn Microbiol Infect Dis* 67:191–197, 2010.

- Pea F, Marioni G, Pavan F, et al. Penetration of levofloxacin into paranasal sinuses mucosa of patients with chronic rhinosinusitis after a single 500 mg oral dose. *Pharmacol Res* 55:38–41, 2007.
- Poole M, Anon J, Paglia M, et al. A trial of high dose, short-course levofloxacin for the treatment of acute bacterial sinusitis. *Otolaryngol Head Neck Surg* 134:10–17, 2006.
- Pugh RJ, Cooke RPD, Dempsey G. Short course antibiotic therapy for Gram-negative hospital-acquired pneumonia in the critically ill. *J of Hosp Infect* 74:337–343, 2010.
- Samonis G, Koutsounaki E, Karageorgopoulos DE, et al. Empirical therapy with ceftazidime combined with levofloxacin or once-daily amikacin for febrile neutropenia in patients with neoplasia: a prospective comparative study. *Eur J Clin Microbiol Infect Dis* 31:1389–1398, 2012.
- Tsuruoka S, Yokota N, Hayasaka T, et al. Pharmacokinetics of multiple-dose levofloxacin in hemodialysis patients. *Am J Kidney Dis* 58:498–499, 2011.
- Website: www.pdr.net

## Linezolid (Zyvox)

**Drug Class:** Oxazolidinone.

**Usual Dose:** 600 mg (IV/PO) q12h.

**Spectrum:** (see **Susceptibility Profiles** pp. 198–202).

**Resistance Potential:** Low

**Pharmacokinetic Parameters:**

*Peak serum level:* 15–21 mcg/ml

*Bioavailability:* 100% (IV and PO)

*Excreted unchanged (urine):* 30%

*Serum half-life (normal/ESRD):* 6.4/7.1 hrs

*Plasma protein binding:* 31%

*Volume of distribution ( $V_d$ ):* 0.64 L/kg

**Primary Mode of Elimination:** Hepatic/  
Metabolized

**Dosage Adjustments\***

CrCl < 10 mL/min	No change
Post-HD dose	600 mg (IV/PO)
Post-PD dose	None
CVH/CVHD/CVHDF dose	No change
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	Use with caution

**Drug Interactions:** Pseudoephedrine, tyramine-containing foods (↑ risk of hypertensive crisis); serotonergic agents, e.g., SSRIs, MAOIs, St. John's wort, ritonavir (↑ risk of serotonin syndrome). Serotonin syndrome: fever, delirium, hypertension, tremor/clonus, hyperreflexia.

**Adverse Effects:** Mild, readily reversible thrombocytopenia, anemia, or leukopenia may occur usually after  $\geq 2$  weeks of therapy. Lactic acidosis (rare), optic/peripheral neuropathy ( $> 28$  days of therapy, rare). If used for  $> 4$  weeks, ↑ risk of myelosuppression or blindness (irreversible) particularly in those with optic nerve disorders or glaucoma. Monitor visual function if symptomatic or therapy  $\geq 2$  weeks; monitor CBC weekly if therapy  $> 2$  weeks. Hypoglycemia may occur ( $> 7$  days) with IV/PO linezolid.

**Allergic Potential:** Low

**Safety in Pregnancy:** C

**Comments:** *Ideal for IV-to-PO switch programs.* Unlike vancomycin, **linezolid use does not increase VRE prevalence.** Unlike

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose  $\times 1$  followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

quinupristin/dalfopristin, linezolid is active against *E. faecalis* (VSE). **Effective oral therapy for MRSA, MRSE, and *E. faecium* (VRE) infections.** *Alternate therapy for Nocardia and Listeria infections.*

Meningeal dose = usual dose.

**Cerebrospinal Fluid Penetration:** 70%

**Bile Penetration:** 200%

(also see **Antibiotic Pearls & Pitfalls** p. 517).

#### REFERENCES:

- Aeziokoro CO, Cannon JP, Pachucki CT, et al. The effectiveness and safety of oral linezolid for the primary and secondary treatment of osteomyelitis. *J Chemother* 17:643–50, 2005.
- Bassetti M, Baguneid M, Bouza E, et al. European perspective and update on the management of complicated skin and soft tissue infections due to methicillin-resistant *Staphylococcus aureus* after more than 10 years of experience with linezolid. *Clin Microb Infect*. 4:3-18, 2014.
- Bassetti M, Vitale F, Melica G, et al. Linezolid in the treatment of gram-positive prosthetic joint infections. *J Antimicrob Chemother* 55:387–390, 2005.
- Cannon JP, Pachucki CT, Aneziokoro CO and Lentino JR. The effectiveness and safety of oral linezolid as primary or secondary treatment of bloodstream infections. *Infect Dis Clin Pract* 14:221–226, 2006.
- Carroll W, Choi H, Min S, et al. Rhabdomyolysis in patient treated with linezolid for extensively drug-resistant tuberculosis. *Clin Infect Dis* 54:1624–7, 2012.
- Castro P, Soriano A, Escrich C, Villalba G, et al. Linezolid treatment of ventriculoperitoneal shunt infection without implant removal. *Eur J Clin Microbiol Infect Dis* 24:603–06, 2005.
- Chang KC, Leung CC, Daley CL. Linezolid for multidrug-resistant tuberculosis. *Lancet Infect Dis* 12:502–3, 2012.
- Colli A, Compodonico R, Gherli T. Early switch from vancomycin to oral linezolid for treatment of gram-positive heart valve endocarditis. *Ann Thorac Surg* 84:87–91, 2007.
- Conte JE Jr, Golden JA, Dippes J, et al. Intrapulmonary pharmacokinetics of linezolid. *Antimicrob Agents Chemother* 46:1475–80, 2000.
- Cunha BA. Antimicrobial therapy of multidrug-resistant *Streptococcus pneumoniae*, vancomycin-resistant enterococci, and methicillin-resistant *Staphylococcus aureus*. *Med Clin N Am* 90:1165–82, 2006.
- Cunha BA. Oral antibiotic therapy of serious systemic infections. *Med Clin N Am* 90:1197–1222, 2006.
- Estes KS, Derendorf H. Comparison of the pharmacokinetic properties of vancomycin, linezolid, tigecycline, and daptomycin. *Eur J. Med Res* 15:533–543, 2010.
- Falagas ME, Manta KG, Ntziora F, et al. Linezolid for the treatment of patients with endocarditis: a systematic review of published evidence. *J Antimicrob Chemother* 58:273–280, 2006.
- Hill EE, Herijgers P, Herregods MC, Peetermans WE. Infective endocarditis treated with linezolid: case report and literature review. *Eur J Clin Microbiol Infect Dis* 25:202–4, 2006.
- Hoyo I, Martinez-Pastor J, Garcia-Ramiro S, et al. Decreased serum linezolid concentrations in two patients receiving linezolid and rifampicin due to bone infections. *Scand J Infect Dis* 44:548–50, 2012.
- Joson J, Grover C, Downer C, et al. Successful treatment of methicillin-resistant *Staphylococcus aureus* mitral valve endocarditis with sequential linezolid and telavancin monotherapy following daptomycin failure. *J Antimicrob Chemother* 66:2186–2188, 2011.
- Johnson JR. Linezolid versus vancomycin for methicillin-resistant *Staphylococcus aureus* infections. *Clin Infect Dis* 36:236–7, 2003.
- Kaka AS, Rueda AM, Shelburne SA 3<sup>rd</sup>, Hamill RJ. Bactericidal activity of orally available agents against methicillin-resistant *Staphylococcus aureus*. *J Antimicrob Chemother* 58:680–3, 2006.
- Kruse AJ, Peederman SM, Bet PM, Debets-Oseenkopp YJ. Successful treatment with linezolid of meningitis due to methicillin-resistant *Staphylococcus epidermidis* refractory to vancomycin treatment. *Eur J Clin Microb Infect* 215:135–7, 2006.
- Kutscha-Lissberg F, Hebler U, Muhr G, et al. Linezolid penetration into bone and joint tissues infected with methicillin-resistant staphylococci. *Antimicrob Agents Chemother* 47:3964–6, 2003.
- Lauridsen TK, Bruun LE, Rasmussen RV, et al. Linezolid as rescue treatment for left-sided infective endocarditis: an observational, retrospective, multicenter study. *Eur J Clin Microbiol Infect Dis* 31:2567–2574, 2012.
- Lawrence KR, Adra M, Gillman PK. Serotonin toxicity associated with the use of linezolid: a review of postmarketing data. *Clin Infect Dis* 42:1578–83, 2006.

- Moreillon P, Wilson WR, Leclercq R, et al. Single-dose oral amoxicillin or linezolid for prophylaxis of experimental endocarditis due to vancomycin-susceptible and vancomycin-resistant *Enterococcus* faecalis. *Antimicrob Agents Chemother* 51:1661–1665, 2007.
- Moylett EH, Pacheco SE, Brown-Elliott BA, et al. Clinical experience with linezolid for the treatment of nocardia infection. *Clin Infect Dis* 36:313–8, 2003.
- Myriantefs P, Markantonis SL, Vlachos K, et al. Serum and cerebrospinal fluid concentrations of linezolid in neurosurgical patients. *Antimicrob Agents Chemother* 50:3971–76, 2006.
- Nathani N, Iles P, Elliott TS. Successful treatment of MRSA native valve endocarditis with oral linezolid therapy: a case report. *J Infect* 51:213–215, 2005.
- Nguyen S, Pasquet A, Legout L, et al. Efficacy and tolerance of rifampin-linezolid compared with rifampin-cotrimoxazole combinations in prolonged oral therapy for bone and joint infections. *Clin Microbiol Infect Dis* 15:1163–1169, 2009.
- Ravindran V, John J, Kaye GC, et al. Successful use of oral linezolid as a single active agent in endocarditis unresponsive to conventional antibiotic therapy. *J Infect* 47:164–6, 2003.
- Rao N, Ziran BH, Hall RA, et al. Successful treatment of chronic bone and joint infections with oral linezolid. *Clin Orthop* 427:67–71, 2004.
- Schechter GF, Scott C, True L, et al. Linezolid in the treatment of multidrug-resistant tuberculosis. *Clin Infect Dis* 50:49–55, 2010.
- Siegel RE. Linezolid to decrease length of stay in the hospital for patients with methicillin-resistant *Staphylococcus aureus* infection. *Clin Infect Dis* 36:124–5, 2003.
- Souli M, Pontikis K, Chryssouli Z, et al. Successful treatment of right-sided prosthetic valve endocarditis due to methicillin-resistant teicoplanin-heteroresistant *Staphylococcus aureus* with linezolid. *Eur J Clin Microbiol Infect Dis* 24:760–762, 2005.
- Stein GE, Schooley SL, Havlicek DH, et al. Outpatient intravenous antibiotic therapy compared with oral linezolid in patients with skin and soft tissue infections: a pharmaco-economic analysis. *Infectious Dis in Clin Practice* 16:235–240, 2008.
- Stevens DL, Herr D, Lampiris H, et al. Linezolid versus vancomycin for the treatment of methicillin-resistant *Staphylococcus aureus* infections. *Clin Infect Dis* 34:1481–90, 2002.
- Taylor JJ, Wilson JW, Estes LL. Linezolid and serotonergic drug Interactions: a retrospective survey. *Clin Infect Dis* 43:180–7, 2006.
- Valerio M, Pedromingo M, Munoz P, et al. Potential protective role of linezolid against *Clostridium difficile* infection. *Int J Antimicrob Agents* 39:414–9, 2012.
- Villani P, Pegazzi MB, Marubbi F, et al. Cerebrospinal fluid linezolid concentrations in postneurosurgical central nervous system infections. *Antimicrob Agents Chemother* 46:936–7, 2002.
- Viswanathan P, Iarikov D, Wassel R, Davidson A, Nambiar S. Hypoglycemia in patients treated with linezolid. *Clin Infect Dis* 59:634–635, 2014.
- Xu HB, Jiang RH, Li L, et al. Linezolid in the treatment of MDR-TB: a retrospective clinical study. *Int J Tuberc Lung Dis* 16:358–63, 2012.
- Zhang L, Pang Y, Yu X, et al. Linezolid in the treatment of extensively drug-resistant tuberculosis. *Infection*. 42:705–711, 2014.
- Website: [www.pdr.net](http://www.pdr.net)

## Lopinavir + Ritonavir (Kaletra) LPV/RTV

**Drug Class:** HIV protease inhibitor combination.

**Usual Dose:** Therapy-naive: 400/100 mg q12h or 800/200 mg q24h (if <3 lopinavir resistance—associated substitutions).

**Pharmacokinetic Parameters:**

*Peak serum level:* 9.6/≤ 1 mcg/mL

*Bioavailability:* No data

*Excreted unchanged (urine):* 3%

*Serum half-life (normal/ESRD):* 5–6/5–6 hrs

*Plasma protein binding:* 99%

*Volume of distribution (V<sub>d</sub>):* No data/0.44 L/kg

**Primary Mode of Elimination:** Hepatic

“Usual dose” assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

**Dosage Adjustments\***

CrCl < 10 mL/min	No change
Post-HD dose	None
Post-PD dose	None
CWH/CVWH/ CVHDF dose	No change
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	Avoid

**Antiretroviral Dosage Adjustments**

Fosamprenavir	Avoid
Delavirdine	No data
Efavirenz	Consider lopinavir/ ritonavir 500/125 mg q12h
Indinavir	Indinavir 600 mg q12h
Nelfinavir	Same as for efavirenz
Nevirapine	Same as for efavirenz
Rifabutin	Max. dose of rifabutin 150 mg qod or 3 times per week
Saquinavir	Saquinavir 1000 mg q12h

**Drug Interactions:** Antiretrovirals, rifabutin, (see dose adjustment grid, above); Rifampin may reduce lopinavir levels leading to drug failure and should be avoided. Sildenafil, tadalafil, and vardenafil levels will be greatly increased and their dosage must be decreased. Antiarrhythmics such as amiodarone, bepredil, lidocaine and quinidine may increase levels. Once daily dosing not recommended in pregnancy or for patients receiving efavirenz, nevirapine, fosamprenavir, nelfi

navir, carbamazepine, phenytoin, or phenobarbital. Tacrolimus, cyclosporine and rapamycin levels can be altered leading to increased toxicity. Methadone levels may be decreased leading to withdrawal reactions. Fluticasone levels may increase leading to Cushing like syndrome, astemizole, terfenadine, benzodiazepines, cisapride, ergotamine, flecainide, pimozide, propafenone, rifampin, statins, tenofovir (↓ lopinavir levels, ↑ tenofovir levels). ↓ effectiveness of oral contraceptives. Insufficient data on other drug interactions listed for ritonavir alone. At end of drug interaction add lopinavir/resolution contains alcohol and can lead to a disulfiram reaction when administered with metronidazole clarithromycin (↑ QT<sub>c</sub>).

**Contraindications:** St. John's wort, boceprevir, telaprevir, rifapentine.

**Adverse Effects:** Diarrhea (very common), headache, nausea, vomiting, asthenia, ↑ SGOT/SGPT, hepatotoxicity, abdominal pain, pancreatitis, paresthesias, hyperglycemia (including worsening diabetes, new-onset diabetes, DKA), ↑ cholesterol/triglycerides (evaluate risk for coronary disease, pancreatitis), ↑ CPK, ↑ uric acid, fat redistribution, possible increased bleeding in hemophilia. Oral solution contains 42.4% alcohol. Immune reconstitution syndrome and redistribution/accumulation of body fat have been reported in patients treated with combination antiretroviral therapy.

**Allergic Potential:** Low

**Safety in Pregnancy:** C

**Comments:** Tablet formulation does not require refrigeration and may be taken with or without food. With oral solution, Lopinavir serum concentrations with moderately fatty meals are increased 54%.

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

**REFERENCES:**

- Decloedt EH, McIlleron H, Smith P, et al. Pharmacokinetics of lopinavir in HIV-infected adults receiving rifampin with adjusted doses of lopinavir-ritonavir tablets. *Antimicrob Agents Chemother* 55:3195–3200, 2011.
- Diaz-Brito V, Leon A, Knobel H, et al. Post-exposure prophylaxis for HIV-infection: a clinical trial comparing lopinavir/ritonavir versus atazanavir each with zidovudine/lamivudine. *Antivir Ther* 17:337–46, 2012.
- Panel on Antiretroviral Guidelines for Adults and Adolescents. Department of Health and Human Services. Guidelines for the use of antiretroviral agents in HIV-infected adults and adolescents. <http://aidsinfo.nih.gov/ConsentFiles/AdultandAdolescentGL.pdf>, 2012.
- Thompson MA, Aberg JA, Cahn P, et al. Antiretroviral treatment of adult HIV infection: 2012 recommendations of the International Antiviral Society — USA panel. *JAMA* 308:387–402, 2012.
- Young J, Schafer J, Fux CA, et al. Renal function in patients with HIV starting therapy with tenofovir and either efavirenz, lopinavir or atazanavir. *AIDS* 26:567–75, 2012  
Website: [www.pdr.net](http://www.pdr.net)

**Loracarbef (Lorabid)**

- Drug Class:** 2<sup>nd</sup> generation oral cephalosporin.
- Usual Dose:** 400 mg (PO) q12h (see comments).
- Spectrum:** (see **Susceptibility Profiles** pp. 191–197).
- Resistance Potential:** Low
- Pharmacokinetic Parameters:**
- Peak serum level:* 14 mcg/mL
- Bioavailability:* 90%
- Excreted unchanged (urine):* 90%
- Serum half-life (normal/ESRD):* 1.2/32 hrs
- Plasma protein binding:* 25%
- Volume of distribution (V<sub>d</sub>):* 0.35 L/kg

**Primary Mode of Elimination:** Renal  
**Dosage Adjustments\***

CrCl 10–50 mL/min	200 mg (PO) q24h
CrCl < 10 mL/min	200 mg (PO) q72h
Post–HD dose	400 mg (PO)
Post–PD dose	200 mg (PO)
CVWH/CVWHD/ CVWHD dose	200 mg (PO) q12h
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	No change

**Drug Interactions:** None.

**Adverse Effects:** Drug fever/rash, *C. difficile* diarrhea/colitis.

**Allergic Potential:** Low

**Safety in Pregnancy:** B

**Comments:** Take 1 hour before or 2 hours after meals. **CAP dose:** 400 mg (PO) q12h. **Sinusitis/tonsillitis dose:** 200 mg (PO) q12h.

**Cerebrospinal Fluid Penetration:** < 10% (also see **Antibiotic Pearls & Pitfalls** p. 510).

**REFERENCES:**

- Bandak SI, Turnak MR, Allen BS, et al. Assessment of the susceptibility of *Streptococcus pneumoniae* to cefaclor and loracarbef in 13 cases. *J Chemother* 12:299–305, 2000.
- Gooch WM 3<sup>rd</sup>, Adelglass J, Kelsey DK, et al. Loracarbef versus clarithromycin in children with acute otitis media with effusion. *Clin Infect Dis* 21:711–22, 1999.

\*Usual dose\* assumes normal renal/hepatic function. \*For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.



Paster RZ, McAdoo MA, Keyserling CH. A comparison of a five-day regimen of cefdinir with a seven-day regimen of loracarbef for the treatment of acute exacerbations of chronic bronchitis. *Int J Clin Pract* 64:293–9, 2000.

Vogel F, Ochs HR, Wettich K, et al. Effect of step-down therapy of ceftriaxone plus loracarbef versus parenteral therapy of ceftriaxone on the intestinal microflora in patients with community-acquired pneumonia. *Clin Microbiol Infect* 7:376–9, 2001.

Website: www.pdr.net

## Maraviroc (Selzentry) MVC

**Drug Class:** HIV chemokine receptor 5 (CCR5) antagonist.

**Usual Dose:** 150 mg, 300 mg, or 600 mg (PO) q12h, depending on concomitant medications (see below), in CCR5-tropic HIV-1 isolates. (available in 150 mg and 300 mg tablets).

**Pharmacokinetic Parameters:**

*Peak serum level:* 266–618 mcg/mL

*Bioavailability:* 23–33%

*Excreted unchanged:* 20% (urine); 76% (feces)

*Serum half-life (normal/ESRD):* 14–18 hrs/not studied

*Plasma protein binding:* 76%

*Volume of distribution ( $V_d$ ):* 194 L

**Primary Mode of Elimination:** Fecal/renal

**Dosage Adjustments\***

CrCl > 30 mL/min with potent CYP3A4 inhibitor	150 mg (PO) q12h
CrCl > 30 mL/min with potent CYP3A4 inducer but no inhibitor	600 mg (PO) q12h
CrCl < 30 mL/min with potent CYP3A4 inducer or inhibitor	Avoid

CrCl < 30 mL/min without potent CYP3A4 inducer or inhibitor	300 mg (PO) q12h
Post–HD dose	None
Post–PD dose	None
CVVH/CVVHD/CVVHDF dose	No data
Moderate hepatic insufficiency with potent CYP3A4 inhibitor	Use with caution
Severe hepatic insufficiency	Avoid

### Antiretroviral Dosage Adjustments

Protease inhibitors (except tipranavir/ritonavir), delavirdine, ketoconazole, itraconazole, clarithromycin, nefazodone, telithromycin	150 mg (PO) q12h
Tipranavir/ritonavir, nevirapine, all NRTIs and enfuvirtide	300 mg (PO) q12h
Efavirenz, rifampin, etravirine, carbamazepine, phenobarbital, phenytoin	600 mg (PO) q12h

**Drug Interactions:** Maraviroc is a substrate of CYP3A and P-glycoprotein and is likely to be modulated by inhibitors and inducers of these enzymes/transporters.

**Adverse Effects:** Hepatotoxicity has been reported. A systemic allergic reaction (e.g., pruritic rash, eosinophilia, or elevated IgE) prior to the development of hepatotoxicity may occur. Other Adverse Effects: cough, infection, upper respiratory tract infection, rash, pyrexia, dizziness, abdominal pain, musculoskeletal symptoms (joint/muscle pain). Orthostatic hypotension with severe renal insufficiency.

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

**Allergic Potential:** Low

**Safety in Pregnancy:** B

**Comments:** Indicated for treatment-experienced adult patients infected with only cellular chemokine receptor (CCR) 5-tropic HIV-1 virus detectable who have evidence of viral replication and HIV-1 strains resistant to multiple antiretroviral agents. Used in combination with other antiretroviral agents. Phenotype test is needed to confirm infection with CCR5-tropic HIV-1 (also known as "R5 virus"). May increase risk of infection (especially HSV) and neoplasms.

**Cerebrospinal Fluid Penetration:** No data

#### REFERENCES:

- Brown KC, Patterson KB, Malone SA, et al. Single and multiple dose pharmacokinetics of maraviroc in saliva, semen, and rectal tissue of healthy HIV-negative men. *J Infect Dis* 203:1484–1490, 2011.
- Lazzarin A, Than S, Valluri SR, et al. Safety profile of maraviroc in patients coinfecting with HIV-1 and hepatitis B or C included in the maraviroc expanded access program. *HIV Clin Trials* 13:83–9, 2012.
- Okoli C, Siccardi M, Thomas-William S, et al. Once Daily maraviroc 300 mg or 150 mg in combination with ritonavir-boosted darunavir 800/100 mg. *J Antimicrob Chemother* 67: 671–4, 2012.
- Parra J, Portilla J, Pulido F, et al. Clinical utility of maraviroc. *Clin Drug Investig* 31:527–42, 2011.
- Thompson MA, Aberg JA, Cahn P, et al. Antiretroviral treatment of adult HIV infection: 2012 recommendations of the International Antiviral Society — USA panel. *JAMA* 308:387–402, 2012.
- Wasmuth JC, Rockstroh Jk, Hardy WD. Drug safety evaluation of maraviroc for the treatment of HIV infection. *Expert Opin Drug Saf* 11: 161–74, 2012. Website: www.pdr.net

## Meropenem (Merrem/Meronem)

**Drug Class:** Carbapenem.

**Usual Dose:** 1 gm (IV) q8h (see comments).

**Spectrum:** (see *Susceptibility Profiles* pp. 198–202).

**Resistance Potential:** Low

**Pharmacokinetic Parameters:**

*Peak serum level:* 49 mcg/ml

*Bioavailability:* Not applicable

*Excreted unchanged (urine):* 70%

*Serum half-life (normal/ESRD):* 1/7 hrs

*Plasma protein binding:* 2%

*Volume of distribution ( $V_d$ ):* 0.35 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\*** (based on 1 gm dose; for 500 mg dosing use half of dose shown in grid)

CrCl 25–50 mL/min	1 gm (IV) q12h
CrCl 10–25 mL/min	500 mg (IV) q12h
CrCl < 10 mL/min	500 mg (IV) q24h
Post-HD dose	500 mg (IV)
Post-HFHD dose	500 mg (IV)
Post-PD dose	500 mg (IV)
CVWH/CVWHD/ CVWHD dose	1 gm (IV) q12h
HD q 48–72h dose	2 gm (IV) q HD*
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	No change

\*Give at end of HD.

**Drug Interactions:** Probenecid (↑ meropenem half-life by 40%), Valproic acid (↓ seizure threshold).

**Adverse Effects:** Rarely, mild infusion site inflammation. No seizures. **No cross allergenicity with penicillins/β-lactams; safe to use in patients with a history of penicillin anaphylactic/other allergic reactions.**

"Usual dose" assumes normal renal/hepatic function. \*For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

**Allergic Potential:** Low

**Safety in Pregnancy:** B

**Comments:** *MDR GNBs, cystic fibrosis/*

*meningitis dose: 2 gm (IV) q8h; uSSIs or UTIs*

**dose:** 500 mg (IV) q8h. Inhibits endotoxin release

from gram-negative bacilli. **Meropenem is the**

**only carbapenem that may be given by IV**

**bolus injection over**

**3–5 minutes.** Na<sup>+</sup>Content = 3.92 mEq/g

Meningeal dose = 2 gm (IV) q8h.

**Cerebrospinal Fluid Penetration:**

Non-Inflamed meninges = 10%

Inflamed meninges = 15%

**Bile Penetration:**

Without obstruction = 75%

With obstruction = 40%

(also see **Antibiotic Pearls & Pitfalls** p. 511).

#### REFERENCES:

Berman SJ, Fogarty CM, Fabian T, et al. Meropenem monotherapy for the treatment of hospital-acquired pneumonia: results of a multicenter trial. *J Chemother* 16:362–71, 2004.

Cheng AC, Fisher DA, Anstey NM, et al. Outcomes of patients with melioidosis treated with meropenem. *Antimicrob Agents Chemother* 48:1763–5, 2004.

Conte JE Jr, Golden JA, Kelley MG, Zurlinden E. Intrapulmonary pharmacokinetics and pharmacodynamics of meropenem. *Int J Antimicrob Agents* 26:449–56, 2005.

Cunha BA. The safety of meropenem in elderly and renally impaired patients. *Intern J Antimicrob Ther* 10:109–117, 1998.

Cunha BA. The use of meropenem in critical care. *Antibiotics for Clinicians* 4:59–66, 2000.

Cunha BA. *Pseudomonas aeruginosa*: resistance and therapy. *Semin Respir Infect* 17:231–9, 2002.

Cunha BA. Meropenem: Current uses in critical care. *Antibiotics for Clinicians* 11:433–444, 2007.

Cunha BA. Meropenem: lack of cross reactivity in penicillin allergic patients. *Journal of Chemotherapy* 20:233–237, 2008.

Doh K, Woo H, Hur J, et al. Population pharmacokinetics of meropenem in burn patients. *J Antimicrob Chemother* 65:2428–2435, 2010.

Hagihara M, Kuti JL, Nicolau DP. Predicting doripenem susceptibility based on meropenem and imipenem interpretation for *Pseudomonas aeruginosa*. *Diagn Microbiol Infect Dis* 72:58–62, 2012.

Kitzes-Cohen R, Farin D, Piva G, et al. Pharmacokinetics and pharmacodynamics of meropenem in critically ill patients. *Int J Antimicrob Agents* 19:105–10, 2002.

Louie A, Grasso C, Bahniuk N, et al. The combination of meropenem and levofloxacin is synergistic with respect to both *Pseudomonas aeruginosa* kill rate and resistance suppression. *Antimicrob Agents Chemother* 54:2646–2654, 2010.

Manfredi R, Sabbatani S, Marinacci G, et al. *Listeria monocytogenes* meningitis and multiple brain abscesses in an immunocompetent host. Favorable response to combination linezolid-meropenem treatment. *Journal of Chemotherapy* 18:331–33, 2009.

Mattoes HM, Kuti JL, Drusano GL, et al. Optimizing antimicrobial pharmacodynamics: dosage strategies for meropenem. *Clin Ther* 26:1187–98, 2004.

Pankey GA, Ashcraft DS. In vitro synergistic/additive activity of levofloxacin with meropenem against *Stenotrophomonas maltophilia*. *Diagn Microbiol Infect Dis* 67:297–300, 2010.

Perez-Simon JA, Garcia-Escobar I, Martinez J, et al. Antibiotic prophylaxis with meropenem after allogeneic stem cell transplantation. *Bone Marrow Transplant* 33:183–7, 2004.

Romano A, Viola M, Gueant-Rodriguez R, et al. Tolerability of meropenem in patients with IgE-mediated hypersensitivity of penicillins. *Ann Intern Med* 146:266–69, 2007.

Scheetz MH, Bolon MK, Esterly JS, et al. Life-years gained with meropenem over ciprofloxacin in penicillin-allergic patients with gram-negative bacilli sepsis: results of a probabilistic model. *Pharmacotherapy* 31:469–79, 2011.

Sodhi M, Axtell SS, Callahan J, et al. Is it safe to use carbapenems in patients with a history of allergy to penicillin? *J Antimicrob Chemother* 54:1155–7, 2004.

Taccone FS, Cotton F, Roisin S, et al. Optimal meropenem concentrations to treat multidrug-resistant *Pseudomonas aeruginosa* septic shock. *Antimicrob Agents Chemother* 56:2129–31, 2012.

<sup>†</sup>Usual dose<sup>†</sup> assumes normal renal/hepatic function. <sup>\*</sup>For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

Toya T, Nannya Y, Narukawa K, et al. A comparative analysis of meropenem and doripenem in febrile patients with hematologic malignancies: a single-center retrospective study. *Jpn J Infect Dis* 65:228–32, 2012.

Yamada Y, Ikawa K, Nakamura K, et al. Prostatic penetration of meropenem after intravenous administration in patients undergoing transurethral resection of the prostate. *J Chemother* 23:179–180, 2011.

Website: [www.pdr.net](http://www.pdr.net)

## Methenamine hippurate (Hiprex, Urex) Methenamine mandelate (Mandelamine)

**Drug Class:** Urinary antiseptic.

**Usual Dose:** 1 gm (PO) q6h (hippurate); 1 gm (PO) q6h (mandelate).

**Spectrum:** All gram + and gram – uropathogens.

**Resistance Potential:** None

**Pharmacokinetic Parameters:**

*Peak serum level: Not applicable*

*Bioavailability: 90%*

*Excreted unchanged (urine): 90%*

*Serum half-life (normal/ESRD): 4 hrs/no data*

*Plasma protein binding: Not applicable*

*Volume of distribution ( $V_d$ ): Not applicable*

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\***

CrCl 50–80 mL/min	Avoid
CrCl 10–50 mL/min	Avoid
CrCl < 10 mL/min	Avoid
Post-HD dose	Avoid
Post-PD dose	Avoid
CVVH/CVVHD/CVHDF dose	Avoid
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	Avoid

**Drug Interactions:** Acetazolamide, sodium bicarbonate, thiazide diuretics (↓ antibacterial effect if ↑ urinary pH ≥ 5.5).

**Adverse Effects:** GI upset.

**Allergic Potential:** Low

**Safety in Pregnancy:** C

**Comments:** Take with food to decrease GI upset. Effectiveness depends on maintaining an acid urine (pH ≤ 5.5) with acidifying agents (e.g., ascorbic acid). **Useful only for catheter-associated bacteriuria (CAB)**, not UTIs. **Forms formaldehyde in acid urine; resistance does not develop.**

### REFERENCES:

Cunha BA, Comer JB, Pharmacokinetic considerations in the treatment of urinary tract infections. *Conn Med* 43:347–53, 1979.

Klinge D, Mannisto P, Mantyla R, et al. Pharmacokinetics of methenamine in healthy volunteers. *J Antimicrob Chemother* 9:209–16, 1982.

Musher DM, Griffith DP. Generation of formaldehyde from methenamine: Effect of pH and concentration, and antibacterial effect. *Antimicrob Agents Chemother* 6:708–11, 1974.

Musher DM, Griffith DP, Templeton GB. Further observations of the potentiation of the antibacterial effect of methenamine by acetoxyhydroxamic acid. *J Infect Dis* 133:564–67, 1976.

Schiotz HA, Guttu K. Value of urinary prophylaxis with methenamine in gynecologic surgery. *Acta Obstet Gynecol Scand* 81:743–6, 2002.

Website: [www.pdr.net](http://www.pdr.net)

## Metronidazole (Flagyl)

**Drug Class:** Nitroimidazole antiparasitic/antibiotic.

**Usual Dose:** 1 gm (IV) q24h (**PK dose**): 500 mg (IV/PO) q6–8h (**PI dose**) (see comments).

**Spectrum:** (see **Susceptibility Profiles** pp. 186–190).

PI = package insert

PK = pharmacokinetic dose

**Resistance Potential:** Low (H. pylori, Gardnerella, Prevotella)

**Pharmacokinetic Parameters:**

Peak serum level: 26 (IV)/12 (PO) mcg/mL

Bioavailability: 100%

Excreted unchanged (urine): 20%

Serum half-life (normal/ESRD): 8/14 hrs

Plasma protein binding: 20%

Volume of distribution ( $V_d$ ): 0.25–0.85 L/kg

**Primary Mode of Elimination:** Hepatic

**Dosage Adjustments\***

CrCl 10–50 mL/min	No change
CrCl < 10 mL/min	500 mg (IV) q24h/ 250 mg (PO) q12h
Post-HD dose	1 gm (IV) 500 mg (PO)
Post-PD dose	1 gm (IV) 500 mg (PO)
CWH/CWHD/ CWHDF dose	No change
Mild-moderate hepatic insufficiency	No change
Severe hepatic insufficiency	500 mg (IV/PO) q24h

**Drug Interactions:** Alcohol (disulfiram-like reaction); disulfiram (acute toxic psychosis); warfarin (↑ INR); phenobarbital, phenytoin (↑ metronidazole metabolism).

**Adverse Effects:** Encephalopathy, metallic taste, aseptic meningitis, seizures, peripheral neuropathy. With oral formulation, nausea, vomiting, GI upset. May discolor urine brown.

**Allergic Potential:** Low

**Safety in Pregnancy:** B (avoid in 1<sup>st</sup> trimester)

**Comments:** For intra-abdominal/pelvic sepsis (**plus anti-aerobic GNB drug**). **q24h dosing is preferred to q6h/q8h dosing because of long half-life** ( $t_{1/2} = 8$  h).

**C. difficile diarrhea dose:** 250 mg (PO) q6h.

**C. difficile colitis dose:** 500 mg (IV/PO) q6–8h or 1 gm (IV) q24h.

**For severe C. difficile colitis, add anti-GNB coverage**, e.g., ertapenem or meropenem (for microscopic/macrosopic perforation/peritonitis). **Use increases VRE prevalence. Metronidazole misses some anaerobes eg, Propionibacteria, Actinobacillus, Eikenella, and limited activity against Actinomyces, Mobiluncus and Clostridium ramosum.**

Meningeal dose = usual dose.

**Cerebrospinal Fluid Penetration:**

Non-Inflamed meninges = 30%

Inflamed meninges = 100%

(also see **Antibiotic Pearls & Pitfalls** p. 519).

**REFERENCES:**

- Bouza E, Burillo A, Munoz P. Antimicrobial therapy of Clostridium difficile-associated diarrhea. Med Clin N Am 90:1141–63, 2006.
- Canete R, Rodriguez P, Mesa L, et al. Albendazole versus metronidazole in the treatment of adult giardiasis: a randomized, double-blind, clinical trial. Curr Med Res Opin 28:149–54, 2012.
- Clay PG, Graham MR, Lindsey CC, et al. Clinical efficacy, tolerability, and cost savings associated with the use of open-label metronidazole plus ceftriaxone once daily compared with ticarcillin/clavulanate every 6 hours as empiric treatment for diabetic lower-extremity infections in older males. Am J Geriatr Pharmacother 2:181–9, 2004.
- Cunha BA. Oral Antibiotic Therapy of Serious Systemic Infections. Med Clin N Am 90:1197–1222, 2006.
- Desai JA, Dobson J, Melanson M, et al. Metronidazole-induced encephalopathy: case report and review of MRI findings. Can J Neurol Sci 38:512–513, 2011.
- Falagas ME, Gorbach SL. Clindamycin and metronidazole. Med Clin North Am 79:845–67, 1995.
- Freeman CD, Klutman NE. Metronidazole: A therapeutic review and update. Drugs 54:679–708, 1997.

PI = package insert

PK = pharmacokinetic

Freeman CD, Nightingale CH, Nicolau DP, et al. Serum bactericidal activity of ceftriaxone plus metronidazole against common intra-abdominal pathogens. *Am J Hosp Pharm* 51:1782–7, 1994.

Garbino J, Villiger P, Caviezel A, et al. A randomized prospective study of cefepime plus metronidazole with imipenem-cilastatin in the treatment of intra-abdominal infections. *Infection* 35:161–166, 2007.

Ishikawa T, Okamura S, Oshimoto H, et al. Metronidazole plus ciprofloxacin therapy for active Crohn's disease. *Intern Med* 42:318–21, 2003.

Metzger R, Swenson BR, Bonatti H, et al. Identification of risk factors for the development of Clostridium difficile-associated diarrhea following treatment of polymicrobial surgical infections. *Ann Surg* 251:722–727, 2010.

Musher DM, Aslam S, Logan N, et al. Relatively poor outcome after treatment of Clostridium difficile colitis with metronidazole. *Clinical Infectious Disease* 40:1586–90, 2005.

Vasa CV, Glatt AE. Effectiveness and appropriateness of empiric metronidazole for Clostridium difficile diarrhea. *Am J Gastroenterol* 98:354–8, 2003.

Wang S, Cunha BA, Hamid NS, et al. Intravenous metronidazole: once daily vs. multiple dosing in intra-abdominal infections. *Journal of Chemotherapy* 19:410–416, 2007.

Website: www.pdr.net

## Mezlocillin (Mezlin)

**Drug Class:** Anti-pseudomonal penicillin.

**Usual Dose:** 3 gm (IV) q6h.

**Resistance Potential:** Low

**Pharmacokinetic Parameters:**

*Peak serum level:* 300 mcg/ml

*Bioavailability:* Not applicable

*Excreted unchanged (urine):* 65%

*Serum half-life (normal/ESRD):* 1.1/4 hrs

*Plasma protein binding:* 30%

*Volume of distribution ( $V_d$ ):* 0.18 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\***

CrCl > 30 mL/min	No change
CrCl 10–30 mL/min	3 gm (IV) q8h
CrCl < 10 mL/min	2 gm (IV) q8h
Post–HD dose	3 gm (IV)
Post–PD dose	None
CVH/CVHD/ CVHDF dose	3 gm (IV) q8h
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	3 gm (IV) q12h

**Drug Interactions:** Aminoglycosides (inactivation of mezlocillin in renal failure); warfarin (↑ INR); oral contraceptives (↓ oral contraceptive effect); cefoxitin (↓ mezlocillin effect).

**Adverse Effects:** Drug fever/rash, E. multiforme/ Stevens–Johnson Syndrome, anaphylactic reactions (hypotension, laryngospasm, bronchospasm), hives; serum sickness. **Dose-dependent inhibition of platelet aggregation is minimal/absent** (usual dose is less than carbenicillin).

**Allergic Potential:** Low

**Safety in Pregnancy:** B

**Comments:** Dose-dependent half-life ( $t_{1/2}$ ).  $\text{Na}^+$  content = 1.8 mEq/g.

**Cerebrospinal Fluid Penetration:** < 10%

**Bile Penetration:** 3000%

(also see **Antibiotic Pearls & Pitfalls** p. 508).

### REFERENCES:

Wright AJ. The penicillins. *Mayo Clin Proc* 74:290–307, 1999.

Website: www.pdr.net

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

## Micafungin (Mycamine)

**Drug Class:** Echinocandin antifungal.

**Usual Dose:**

Candidemia:

100 mg (IV) q24h.

Esophageal candidiasis:

150 mg (IV) q24h.

Prophylaxis of candidal infections in

stem cell transplants:

50 mg (IV) q24h.

**Pharmacokinetic Parameters:**

Peak serum level: 5.1 mcg/ml (50 mg); 11.2 mcg/ml (100 mg); 16.4 mcg/ml (150 mg)

Bioavailability: Not applicable

Excreted unchanged (urine): 11%

Serum half-life (normal/ESRD): 10–15 hrs/no change

Plasma protein binding: > 99%

Volume of distribution ( $V_d$ ): 0.39 L/kg

**Primary Mode of Elimination:** Liver

**Dosage Adjustments**

CrCl < 10 mL/min	No change
Post-HD dose	None
Post-PD dose	None
CVWH/CVHD/CVHDF dose	No change
Mild or Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	Use with caution

**Drug Interactions:** Sirolimus AUC ↑ by 21% with no change in C<sub>max</sub> (monitor for sirolimus toxicity), nifedipine AUC, and C<sub>max</sub> ↑ by 18% and 42%, respectively (monitor for nifedipine toxicity).

**Adverse Effects:** Nausea, vomiting, leukopenia, anemia, thrombocytopenia, ↑ SGOT/SGPT, ↑ BUN/↑ creatinine, rash.

**Allergic Potential:** Low

**Safety in Pregnancy:** C

**Comments:** Administer by IV infusion over one hour—do not give as a bolus as more rapid infusions may result in more frequent histamine mediated reactions. Do not mix or co-use with other medications; flush intravenous line with 0.9% saline prior to infusion.

**Highly active against *Candida albicans* and non-albicans *Candida* (↓ susceptibility to *C. parasilosis*), and *Hansenula*.**

**Some activity against *Aspergillus*. No activity against *Cryptococcus*, *Fusarium*, *Pseudoallescheria/Scedosporium*, *Trichosporon*, *Rhodotulula* or *Mucor*.**

Most cost effective parenteral antifungal for empiric therapy of systemic *Candida*/*Aspergillus* infections.

**Cerebrospinal Fluid Penetration:** < 1%

### REFERENCES:

- Andes D, Safdar N. Efficacy of micafungin for the treatment of candidemia. *Eur J Clin Microbiol Infect Dis* 24:662–664, 2005.
- Chandraseka PH, Sobel JD. Micafungin: A New Echinocandin. *Clin Infect Dis* 42:1171–78, 2006.
- Cornely OA, Marty FM, Stucker F, et al. Efficacy and safety of micafungin for treatment of serious *Candida* infections in patients with or without malignant disease. *Mycoses* 54:838–47, 2011
- Denning DW, Marr KA, Lau WM, et al. Micafungin (FK463), alone or in combination with other systemic antifungal agents, for the treatment of acute invasive aspergillosis. *Journal of Infection* 53:337–349, 2006.
- Higashiyama Y, Kohno S. Micafungin: a therapeutic review. *Expert Rev Anti Infect Ther* 2:345–55, 2004.
- Ikeda F, Tanaka S, Ohki H, et al. Role of micafungin in the antifungal armamentarium. *Curr Med Chem* 14:1263–1275, 2007.
- Isumikawa K, Ohtsu Y, Kawabata M, et al. Clinical efficacy of micafungin for chronic pulmonary aspergillosis. *Med Mycol* 45:273–278, 2007.
- Joseph JM, Jain R, Danziger LH. Micafungin: a new echinocandin antifungal. *Pharmacotherapy* 27:53–67, 2007.
- Kofia G, Ruhnke M. Pharmacology and metabolism of anidalfungin, caspofungin, and micafungin in the

“Usual dose” assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

- treatment of invasive candidosis: review of the literature. *Eur J Med Res* 16:159–166, 2011.
- Kohno S, Izumikawa K, Ogawa K, et al. Intravenous micafungin versus voriconazole for chronic pulmonary aspergillosis. *J Infect* 61:410–418, 2011.
- Kuse ER, Chetchotisakd P, da Cunha CA, et al. Micafungin versus liposomal amphotericin B for candidaemia and invasive candidosis: a phase III randomised double-blind trial. *Lancet* 369:1519–1527, 2007.
- Nakagawa Y, Ichii Y, Saeki Y, et al. Plasma concentration of micafungin in patients with hematologic malignancies. *J Infect Chemother* 13:39–45, 2007.
- Okugawa S, Ota Y, Tatsuno K, et al. A case of invasive central nervous system aspergillosis treated with micafungin with monitoring of micafungin concentrations in the cerebrospinal fluid. *Scand J Infect Dis* 39:344–346, 2007.
- Ostrosky-Zeichner L, Kontoyiannis D, Raffalli J, et al. International, open-label, noncomparative, clinical trial of micafungin alone and in combination for treatment of newly diagnosed and refractory candidemia. *Eur J Clin Microbiol Infect Dis* 24:654–661, 2005.
- Pappas PG, Rotstein CM, Betts RF, et al. Micafungin versus caspofungin for treatment of candidemia and other forms of invasive candidiasis. *Clin Infect Dis* 45:883–893, 2007.
- Pawlitcz D, Young M, Klepser M. Micafungin. *Formulary* 38:354–67, 2003.
- Pettengell K, Mynhardt J, et al. Successful treatment of oesophageal candidiasis by micafungin: a novel systemic antifungal agent. *Aliment Pharmacol* 20:475–481, 2004.
- Toubai T, Tanaka J, Ota S, et al. Efficacy and safety of micafungin in febrile neutropenic patients treated for hematological malignancies. *Intern Med* 46:3–9, 2007.
- Wiederhold NP, Lewis JS 2<sup>nd</sup>. The echinocandin micafungin: a review of the pharmacology, spectrum of activity, clinical efficacy and safety. *Expert Opin Pharmacother* 8:1155–1166, 2007.
- Yamaguchi M, Kurokawa T, Ishiyama K, et al. Efficacy and safety of micafungin as an empirical therapy for invasive fungal infections in patients with hematologic disorders: a multicenter, prospective study. *Ann Hematol* 90:1209–17, 2011.
- Yoshida M, Tamura K, Imamura M, et al. Efficacy and safety of micafungin as an empirical antifungal therapy for suspected fungal infection in neutropenic patients with hematological disorders. *Ann Hematol* 91:449–57, 2012.
- Wilke M. Treatment and prophylaxis of invasive candidiasis with anidulafungin, caspofungin, and micafungin and its impact on use and costs: review of the literature. *Eur J Med Res* 16:180–186, 2011.
- Website: www.pdr.net

## Minocycline (Minocin)

**Drug Class:** 2<sup>nd</sup> generation tetracycline.

**Usual Dose:** 100 mg (IV/PO) q12h or 200 mg (IV/PO) q24h (see comments).

**Spectrum:** (see **Susceptibility Profiles** pp. 186–190).

**Resistance Potential:** Low

**Pharmacokinetic Parameters:**

*Peak serum level:* 4 mcg/mL

*Bioavailability:* 95%

*Excreted unchanged (urine):* 10%

*Serum half-life (normal/ESRD):* 15/18–69 hrs

*Plasma protein binding:* 75%

*Volume of distribution (V<sub>d</sub>):* 1.5 L/kg

**Primary Mode of Elimination:** Hepatic

**Dosage Adjustments\***

CrCl < 10 mL/min	No change
Post-HD/PD dose	None
CWVH/CWVHD/CWVHDF dose	No change
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	100 mg (IV/PO) q24h

**Drug Interactions:** Antacids, Al<sup>3+</sup>, Ca<sup>2+</sup>, Fe<sup>2+</sup>, Mg<sup>2+</sup>, Zn<sup>2+</sup>, multivitamins, sucralfate (↓ minocycline absorption); isotretinoin (pseudotumor cerebri); warfarin (↑ INR).

**Adverse Effects:** Nausea, GI upset if not taken with food, hyperpigmentation of skin with prolonged use, vestibular toxicity (dizziness), photosensitivity (rare), drug induced SLE.

**Allergic Potential:** Low

**Safety in Pregnancy:** D

**Comments:** Infuse slowly over 1 hour. Also **effective against Nocardia**.

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CWVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.



**Effective IV/PO against serious systemic**

**MSSA/MRSA infections**, e.g., MSSA/MRSA ABE, osteomyelitis, or meningitis.

Tetracycline susceptibilities do not predict minocycline effectiveness against MSSA, MRSA, and *Acinetobacter* sp.

**Even if MSSA/MRSA susceptible to doxycycline, use minocycline more reliably effective.**

For neuroborreliosis minocycline may be preferable to doxycycline

Meningeal dose = usual dose

**Cerebrospinal Fluid Penetration:**

Non-inflamed meninges = 50%

Inflamed meninges = 50%

**Bile Penetration:** 1000%

(also see **Antibiotic Pearls & Pitfalls**

p. 512).

**REFERENCES:**

- Bishburg E, Shah M, Chan T. Use of intravenous minocycline for treatment of methicillin-resistant staphylococcus aureus (MRSA) and resistant Gram-negative organisms. *Inf Dis in Clin Prac* 22:26-31, 2014.
- Bouza E. New therapeutic choices for infections caused by methicillin-resistant Staphylococcus aureus. *Clin Microbiol Infect Dis* 15:44-52, 2009.
- Copeland KFT, Brooks JI. A Novel use for an old drug: the potential for minocycline as anti-HIV adjuvant therapy. *J Infect Dis* 201:1115-1117, 2010.
- Cukras CA, Petrou P, Chew EY, et al. oral minocycline for the treatment of diabetic macular edema (DME): results of a phase I/II clinical study. *Invest Ophthalmol Vis Sci* 53:3865-74, 2012.
- Cunha BA. Minocycline is a Reliable Effective Oral Option to Treat Methicillin Resistant *S. aureus* (MRSA) Skin & Soft Tissue Infections (SSTIs) Including Doxycycline Treatment Failures. *Int Antimicrob Agents*. 43:386-387, 2014.
- Cunha BA. Pharmacoeconomic advantages of oral minocycline for the therapy of methicillin-resistant Staphylococcus aureus (MRSA) Skin and Soft Tissue Infections (SSTIs). *Euro Clin Microb & Infect Dis*. 33:1869-1871, 2014.
- Cunha BA. Minocycline, often forgotten but preferred to trimethoprim-sulfamethoxazole or doxycycline for the treatment of community-acquired methicillin-resistant Staphylococcus aureus skin and soft-tissue infections. *Int J Antimicrob Agents* 42:497-499, 2013.

- Cunha BA. Minocycline versus doxycycline for methicillin-resistant Staphylococcus aureus (MRSA): in vitro susceptibility versus in vivo effectiveness. *Int J Antimicrob Agents* 35:517-8, 2012.
- Cunha BA. Minocycline versus doxycycline for CA-MRSA: in vitro versus in vivo differences. *International Journal of Antimicrobial Agents* 35:517-518, 2010.
- Cunha BA. Minocycline vs. doxycycline for the antimicrobial therapy of Lyme neuroborreliosis. *Clin Infect Dis* 30:237-238, 2000.
- Elkayam O, Yaron M, Caspi D. Minocycline-induced autoimmune syndromes: an overview. *Semin Arthritis Rheum* 28:392-397, 1999.
- Golf DA, Kaye KS. Minocycline: An Old Drug for a New Bug: Multidrug-Resistant *Acinetobacter baumannii*. *Clin Infect Dis* 59:S365-S366, 2014.
- Ritchie DJ, Garavaglia-Wilson A. A Review of Intravenous Minocycline for Treatment of Multidrug-Resistant *Acinetobacter* Infections. *Clin Infect Dis* 59:S374-S380, 2014.
- Schwartz BS, Graber CJ, Diep BA, et al. Doxycycline, not minocycline, induces its own resistance in multidrug-resistant community-associated methicillin-resistant Staphylococcus aureus clone USA300. *Clin Infect Dis* 48:1483-1484, 2009.
- Website: [www.pdr.net](http://www.pdr.net)

**Moxifloxacin (Avelox)**

**Drug Class:** Respiratory quinolone.

**Usual Dose:** 400 mg (IV/PO) q24h (see comments).

**Spectrum:** (see **Susceptibility Profiles** pp. 198-202).

**Resistance Potential:** Low

**Pharmacokinetic Parameters:**

*Peak serum level:* 4.4 (IV)/4.5 (PO) mcg/mL

*Bioavailability:* 90%

*Excreted unchanged (urine):* 20%

*Serum half-life (normal/ESRD):* 12/12 hrs

*Plasma protein binding:* 50%

*Volume of distribution (V<sub>d</sub>):* 2.2 L/kg

**Primary Mode of Elimination:** Hepatic

**Dosage Adjustments\***

CrCl < 10 mL/min	No change
Post-HD dose	None
Post-PD dose	None

CVH/CVHD/CVHDF dose	No change
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	No change

**Drug Interactions:** Al<sup>++</sup>, Fe<sup>++</sup>, Mg<sup>++</sup>, Zn<sup>++</sup> antacids, citrate/citric acid, dairy products (↓ absorption of fluoroquinolones only if taken together); amiodarone, procainamide, sotalol (may ↑ QT<sub>c</sub> interval, torsade de pointes), rifampicin (↓ moxifloxacin levels).

**Adverse Effects:** *C. difficile* diarrhea/colitis. May ↑ QT<sub>c</sub> interval; avoid taking with medications, e.g., amiodarone that prolong the QT<sub>c</sub> interval, and in patients with cardiac arrhythmias/heart block. **Does not cause seizures.** FQ use has an increased risk of tendinitis/tendon rupture. Risk is highest in the elderly, those on steroids, and those with heart, lung or renal transplants. Peripheral neuropathy may occur early and may be permanent.

**Allergic Potential:** Low

**Safety in Pregnancy:** C

**Comments:** **Only quinolone with anti-*B. fragilis* activity. FQ with the highest activity against *S. pneumoniae*, VSE, and MSSA.**

Metabolized to microbiologically-inactive glucuronide (M1)/sulfate (M2) conjugates. Take 4 hours before or 8 hours after calcium or magnesium containing antacids or didanosine. No interactions with oral hypoglycemics. C8-methoxy group increases activity and decreases resistance potential. **TB dose** (*alternate drug in MDR TB drug regimens*): 400 mg (PO) q24h.

↑ **incidence of *C. difficile* diarrhea/colitis with PPIs (for patients on PPIs during FQ therapy, D/C PPIs or switch to H<sub>2</sub> blocker for duration of FQ therapy).**

**Cerebrospinal Fluid Penetration:** < 10%

**Bile Penetration:** 93%

(also see **Antibiotic Pearls & Pitfalls** p. 516).

#### REFERENCES:

- Al-Nawas B, Walter C, Morbach T. Clinical and microbiological efficacy of moxifloxacin versus amoxicillin/clavulanic acid in severe odontogenic abscesses: a pilot study. *Eur J Clin Microbiol Infect Dis* 28:75–82, 2009.
- Arrieta JR, Galgano AS, Sakano E, et al. Moxifloxacin vs. amoxicillin/clavulanate in the treatment of acute sinusitis. *Am J Otolaryngol* 28:78–82, 2007.
- Balfour JA, Wiseman LR. Moxifloxacin. *Drugs* 57:363–73, 1999.
- Bassetti M, Rosso R, Tosi C, et al. Moxifloxacin in the treatment of hospitalized community acquired pneumonia in HIV-infected subjects. *J Chemother* 19:104–105, 2007.
- Brunkhorst FM, Oppert M, Marx G, et al. Effect of empirical treatment ith moxifloxacin and meropenem vs meropenem on sepsis-related organ dysfunction in patients with severe sepsis: a randomized trial. *JAMA* 307:2309–9, 2012.
- Cheadle W, Lee JT, Napolitano LM, et al. Clinical update on the use of moxifloxacin in the treatment of community-acquired complicated intra-abdominal infections. *Surg Infect* 11:487–494, 2010.
- Cheon JH, Kim N, Lee DH, et al. Efficacy of moxifloxacin-based triple therapy as second-line treatment for *Helicobacter pylori* infection. *Helicobacter* 11:46–51, 2006.
- Chou HW, Wang JL, Chang CH, et al. Risk of severe dysglycemia among diabetic patients receiving levofloxacin, ciprofloxacin, or moxifloxacin in Taiwan. *Clin Infect Dis* 57:971–980, 2013.
- Cox H, Ford N, Keshavjee S, et al. Rational use of moxifloxacin for tuberculosis treatment. *Lancet Infect Dis* 11: 259–260, 2011.
- Cunha BA. Oral antibiotic therapy of serious systemic infections. *Med Clin N Am* 90:1197–1222, 2006.
- Ewig S, Hecker H, Suttorp N, et al. Moxifloxacin monotherapy versus B-lactam mono or combination therapy in hospitalized patients with community-acquired pneumonia. *J Infect* 62: 218–225, 2011.
- Fouad M, Gallagher JC. Moxifloxacin as an alternative or additive therapy for treatment of pulmonary tuberculosis. *Ann Pharmacother*, 45:1439–1444, 2011.
- Gillespie SH, Crook AM, McHugh TD, et al. Four-month moxifloxacin-based regimens for drug-sensitive tuberculosis. *NEJM* 371:1577–1587, 2014.

"Usual dose" assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose x 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

- Giordano P, Song J, Pertel P, et al. Sequential intravenous/oral moxifloxacin versus intravenous piperacillin-tazobactam followed by oral amoxicillin-clavulanate for the treatment of complicated skin and skin structure infection. *Int J Antimicrob Agents* 26:357–65, 2005.
- Haggerty CL, Ness RB. Newest approaches to treatment of pelvic inflammatory disease: a review of recent randomized clinical trials. *Clin Infect Dis* 44:953–960, 2007.
- Hariprasad SM, Shah GK, Mieler WF, Feiner L, Blinder KJ, et al. Vitreous and aqueous penetration of orally administered moxifloxacin in humans. *Arch Ophthalmol* 124:178–82, 2006.
- Hoeffken G, Talan D, Larsen LS, et al. Efficacy and safety of sequential moxifloxacin for treatment of community-acquired pneumonia associated with atypical pathogens. *Eur J Clin Microbiol Infect Dis* 23:772–5, 2004.
- Jindani A, Harrison TS, Nunn AJ, et al. High-dose rifapentine with moxifloxacin for pulmonary tuberculosis. *NEJM* 371:1599–1608, 2014.
- Katz E, Larsen LS, Fogarty CM, et al. Safety and efficacy of sequential I.V. to P.O. moxifloxacin versus conventional combination therapies the treatment of community-acquired pneumonia in patients requiring initial I.V. therapy. *J Emerg Med* 27:554–9, 2004.
- Lapi F, Wilchesky M, Abbas K, et al. Fluoroquinolones and the risk of serious arrhythmia: A population-based study. *Clin Infect Dis* 55:1457–65, 2012.
- Lode H, Grossman C, Choudhri S, et al. Sequential IV/PO moxifloxacin treatment of patients with severe community-acquired pneumonia. *Respir Med* 97:1134–42, 2003.
- Lobera T, Audicana MT, Alarcon E, et al. Allergy to quinolones: low cross-reactivity to levofloxacin. *J Investig Allergol Clin Immunol* 20: 607–611, 2010.
- Malincarne L, Ghebtegzabher M, Moretti MV, et al. Penetration of moxifloxacin into bone in patients undergoing total knee arthroplasty. *J Antimicrob Chemother* 57:950–4, 2006.
- Malingoni MA, Song J, Herrington J, et al. Randomized controlled trial of moxifloxacin compared with piperacillin-tazobactam and amoxicillin-clavulanate for the treatment of complicated intra-abdominal infections. *Ann Surg* 244:204–11, 2006.
- Mendel C, Springsklee M. Moxifloxacin for tuberculosis. *Lancet Infect Dis* 176–17, 2012.
- Mu YP, Liu RL, Wang LQ, et al. Moxifloxacin monotherapy for treatment of complicated intra-abdominal infections: a meta-analysis of randomized controlled trials. *Int J Clin* 66:210–7, 2012.
- Nijland HM, Ruslami R, Suroto AJ, et al. Rifampicin reduces plasma concentrations of moxifloxacin in patients with tuberculosis. *Clin Infect Dis* 45:1001–1007, 2007.
- Pranger AD, Van Altena R, Aarnoutse RE, et al. Evaluation of moxifloxacin for the treatment of tuberculosis: 3 years of experience. *Eur Respir J* 38: 884–894, 2011.
- Rijnders BJ. Moxifloxacin for community-acquired pneumonia. *Antimicrob Agents Chemother* 47:444–445, 2003.
- Schulze MH, Keller C, Müller A, et al. Rickettsia rypsi infection with interstitial pneumonia in a traveler treated with moxifloxacin. *J Clin Microb* 49:741–743, 2011.
- Solnick, JV. Treatment with moxifloxacin versus standard therapy for community-acquired pneumonia. *10 Clinics* 41:1697–1705, 2005.
- Stass, H, Rink AD, Delesen H, et al. Pharmacokinetics and peritoneal penetration of moxifloxacin in peritonitis. *J Antimicrob Chemother* 58:693–6, 2006.
- Torres A, Muir JF, Corris P, et al. Effectiveness of oral moxifloxacin in standard first-line therapy in community-acquired pneumonia. *Eur Respir J* 21:135–43, 2003.
- Walter S, Kuchenbecker J, Banditt P, et al. Concentration of moxifloxacin in serum and human aqueous humor following a single 400 mg oral dose. *J Cataract Refract Surg* 33:553–555, 2007.
- Welte T, Petermann W, Schurmann D, et al. Treatment with sequential intravenous or oral moxifloxacin was associated with faster clinical improvement that the standard therapy for hospitalized patients with community-acquired pneumonia who received initial parenteral therapy. *Clin Infect Dis* 41:1697–705, 2005.
- Zervos M, Martinez FJ, Amsden GW, et al. Efficacy and safety of 3-day azithromycin versus 5-day moxifloxacin for the treatment of acute bacterial exacerbations of chronic bronchitis. *Int J Antimicrob Agents* 29:56–61, 2007.
- Website: [www.pdr.net](http://www.pdr.net)

## Nafcillin (Unipen)

**Drug Class:** Anti-staphylococcal (MSSA) penicillin.

**Usual Dose:** 2 gm (IV) q4h.

**Spectrum:** (see **Susceptibility Profiles** pp. 186–190).

**Resistance Potential:** Low

"Usual dose" assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

**Pharmacokinetic Parameters:**

Peak serum level: 80 mcg/ml

Bioavailability: Not applicable

Excreted unchanged (urine): 10–30%

Serum half-life (normal/ESRD): 0.5/4 hrs

Plasma protein binding: 90%

Volume of distribution ( $V_d$ ): 0.24 L/kg**Primary Mode of Elimination:** Hepatic**Dosage Adjustments\***

CrCl < 10 mL/min	No change
Post-HD/PD dose	None
CVWH/CVVDH/ CVVHDF dose	No change
Moderate—severe hepatic insufficiency	No change

**Drug Interactions:** Cyclosporine (↓ cyclosporine levels); nifedipine, warfarin (↓ interacting drug effect).

**Adverse Effects:** Not hepatotoxic. Drug fever/rash, leukopenia, interstitial nephritis.

**Allergic Potential:** High

**Safety in Pregnancy:** B

**Comments:** Avoid oral formulation (not well absorbed/erratic serum levels).

Meningeal dose = usual dose.

**Cerebrospinal Fluid Penetration:**

Non-Inflamed meninges = 1%

Inflamed meninges = 20%

**Bile Penetration:** 100%

(also see **Antibiotic Pearls & Pitfalls** p. 509).

**REFERENCES:**

Donowitz GR, Mandell GL. Beta-lactam antibiotics. *N Engl J Med* 318:419–26 490–500, 1993.

Youngster I, Shenoy ES, Hooper D, et al. Comparative

Evaluation of the Tolerability of Cefazolin and Nafcillin for Treatment of Methicillin-Susceptible Staphylococcus aureus Infections in the Outpatient Setting. *Clin Infect Dis* 59:369–375, 2014.

Wright AJ. The penicillins. *Mayo Clin Proc* 74:290–307, 1999.

Website: www.pdr.net

**Nelfinavir (Viracept) NFV**

**Drug Class:** HIV protease inhibitor.

**Usual Dose:** 1250 mg (PO) q12h (two 625-mg tablets per dose) with meals, or five 250-mg tabs or 750 mg (three 250-mg tabs) (PO) q8h.

**Pharmacokinetic Parameters:**

Peak serum level: 35 mcg/mL

Bioavailability: 20–80%

Excreted unchanged (urine): 1–2%

Serum half-life (normal/ESRD): 4 hrs/no data

Plasma protein binding: 98%

Volume of distribution ( $V_d$ ): 5 L/kg**Primary Mode of Elimination:** Hepatic**Dosage Adjustments\***

CrCl < 10 mL/min	No change
Post-HD dose	None
Post-PD dose	None
CVWH/CVVDH/ CVVHDF dose	No change
Mild hepatic insufficiency	No change
Moderate—severe hepatic insufficiency	Use with caution

**Antiretroviral Dosage Adjustments**

Delavirdine	No data (monitor for neutropenia)
Efavirenz	No change
Indinavir	Limited data for nelfinavir 1250 mg q12h + indinavir 1200 mg q12h
Lopinavir/ritonavir	Nelfinavir 1000 mg q12h or lopinavir/r 600/150 mg q12h
Nevirapine	No data

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

Ritonavir	No data
Saquinavir	Saquinavir 1200 mg q12h
Rifampin	Avoid
Rifabutin	Nelfinavir 1250 mg q12h; rifabutin 150 mg q24h or 300 mg 2–3x/ week

**Drug Interactions:** Antiretrovirals, rifabutin, rifampin (see dose adjustment grid, above); amiodarone, quinidine, astemizole, terfenadine, benzodiazepines, cisapride, ergot alkaloids, statins, St. John's wort (avoid if possible); carbamazepine, phenytoin, phenobarbital (↓ nelfinavir levels, ↑ anticonvulsant levels; monitor); caspofungin (↓ caspofungin levels, may ↓ caspofungin effect); clarithromycin, erythromycin, telithromycin (↑ nelfinavir and macrolide levels); didanosine (dosing conflict with food; give nelfinavir with food 2 hours before or 1 hour after didanosine); itraconazole, voriconazole, ketoconazole (↑ nelfinavir levels); lamivudine (↑ lamivudine levels); methadone (may require ↑ methadone dose); oral contraceptives, zidovudine (↓ zidovudine levels); sildenafil (↑ or ↓ sildenafil levels; do not exceed 25 mg in 48 hrs, tadalafil (max. 10 mg/72 hrs, vardenafil (max. 2.5 mg/72 hrs).

**Adverse Effects:** Impaired concentration, nausea, abdominal pain, secretory diarrhea, ↑ SGOT/SGPT, rash, ↑ cholesterol/triglycerides (evaluate risk for coronary disease/pancreatitis), fat redistribution, hyperglycemia (including worsening diabetes, new-onset diabetes, DKA), possible increased bleeding in hemophilia.

**Allergic Potential:** Low

**Safety in Pregnancy:** B

**Comments:** Take with food (absorption increased 300%). 625-mg tablet available.

**Cerebrospinal Fluid Penetration:**  
Undetectable

#### REFERENCES:

- Albrecht MA, Bosch RJ, Hammer SM, et al. Nelfinavir, efavirenz, or both after the failure of nucleoside treatment of HIV infection. *N Engl J Med* 345:398–407, 2001.
- Go J, Cunha BA. Nelfinavir: a review. *Antibiotics for Clinicians* 4:17–23, 2000.
- Hughes PJ, Cretton-Scott E, Teague A, et al. Protease inhibitors for patients with HIV-1 infection. *Pharm Ther* 36:332–344, 2011.
- Perry CM, Benfield P. Nelfinavir. *Drugs* 54:81–7, 1997.
- Saber P, Ranatunga DK, Quesenberry CP, et al. Clinical implications of the nelfinavir-proton pump inhibitor drug interaction in patients with human immunodeficiency virus. *Pharmacotherapy* 31:253–61, 2011.
- Simpson KN, Luo MP, Chumney E, et al. Cost-effective of lopinavir/ritonavir versus nelfinavir as the first-line highly active antiretroviral therapy regimen for HIV infection. *HIV Clin Trials* 5:294–304, 2004.
- Thompson MA, Aberg JA, Cahn P, et al. Antiretroviral treatment of adult HIV infection: 2012 recommendations of the International Antiviral Society — USA panel. *JAMA* 308:387–402, 2012.
- Walmsley S, Bernstein B, King M, et al. Lopinavir-ritonavir versus nelfinavir for the initial treatment of HIV infection. *N Engl J Med* 346:2039–46, 2002.
- Website: [www.pdr.net](http://www.pdr.net)

## Nevirapine (Viramune) NVP

**Drug Class:** HIV NNRTI (non-nucleoside reverse transcriptase inhibitor).

**Usual Dose:** 200 mg (PO) q24h × 2 weeks, then 200 mg (PO) q12h.

**Pharmacokinetic Parameters:**

*Peak serum level:* 0.9–3.6 mcg/mL

*Bioavailability:* 90%

*Excreted unchanged (urine):* 5%

"Usual dose" assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

Serum half-life (normal/ESRD): 40 hrs/no data

Plasma protein binding: 60%

Volume of distribution ( $V_d$ ): 1.4 L/kg

**Primary Mode of Elimination:** Hepatic

**Dosage Adjustments\***

CrCl 10–50 mL/min	No change
CrCl < 20 mL/min	Use with caution
Post-HD dose	200 mg (PO)
Post-PD dose	None
CVVH/CVVHD/ CVVHDF dose	No change
Moderate hepatic insufficiency	Use with caution
Severe hepatic insufficiency	Avoid

#### Antiretroviral Dosage Adjustments

Delavirdine	No data
Efavirenz	No data
Indinavir	Indinavir 1000 mg q8h
Lopinavir/ritonavir (l/r)	Consider l/r 600/150 mg q12h in PI-experienced patients
Nelfinavir	No data
Ritonavir	No change
Saquinavir	No data
Rifampin	Avoid
Rifabutin	Use with caution

**Drug Interactions:** Antiretrovirals, rifabutin, rifampin (see dose adjustment grid, above); carbamazepine, phenobarbital, phenytoin (monitor anticonvulsant levels); caspofungin (↓ caspofungin levels), itraconazole (↓ itraconazole/↑ nevirapine levels), voriconazole

(↑ nevirapine levels); ethinyl estradiol (↓ ethinyl estradiol levels; use additional/alternative method); methadone (↓ methadone levels; titrate methadone dose to effect); tacrolimus (↓ tacrolimus levels) clarithromycin (↓ clarithromycin levels), rifampentine (avoid).

**Adverse Effects:** Severe, life-threatening, and in some cases fatal hepatotoxicity, particularly in the first 18 weeks, has been reported in patients treated with nevirapine. In some cases, patients presented with non-specific prodromal signs or symptoms of hepatitis and progressed to hepatic failure. These events are often associated with rash. Female gender and higher CD<sub>4</sub> counts at initiation of therapy place patients at increased risk; women with CD<sub>4</sub> counts > 250 cells/mm<sup>3</sup>, including pregnant women receiving nevirapine in combination with other antiretrovirals for the treatment of HIV infection, are at the greatest risk. However, hepatotoxicity associated with nevirapine use can occur in both genders, all CD<sub>4</sub> counts and at any time during treatment. Severe, life-threatening skin reactions, including fatal cases, have occurred in patients treated with nevirapine. These have included cases of Stevens-Johnson Syndrome, toxic epidermal necrolysis, and hypersensitivity reactions characterized by rash, constitutional findings, and organ dysfunction. Extra vigilance is warranted during the first 6 weeks of therapy, which is the period of greatest risk of these events. Do not restart nevirapine following severe hepatic, skin or hypersensitivity reactions. Take out all the rest under adverse reactions.

**Allergic Potential:** High

**Safety in Pregnancy:** B

**Comments:** Absorption not affected by food. Not to be used for post-exposure prophylaxis because of potential for fatal hepatitis.

**Cerebrospinal Fluid Penetration:** 45%

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

## REFERENCES:

- Bogner JR. Nevirapine extended-release for the treatment of HIV-1 infection. *Expert Rev Anti Infect Ther* 10:21–30, 2012.
- Coster LO, Kumar PN. Contemporary role of nevirapine in HIV treatment. *AIDS Rev* 14:132–44, 2012.
- Herzmann C, Karcher H. Nevirapine plus zidovudine to prevent mother-to-child transmission of HIV. *N Engl J Med* 351:2013–5, 2004.
- Milinkovic A, Martinez E. Nevirapine in the treatment of HIV. *Expert Rev Anti Infect Ther* 2:367–73, 2004.
- Nachman S, McIntyre J. Nevirapine-based therapy after suppression: can we switch. *Lancet Infect Dis* 12:498–9, 2012.
- Panel on Antiretroviral Guidelines for Adults and Adolescents. Department of Health and Human Services. Guidelines for the use of antiretroviral agents in HIV-infected adults and adolescents. <http://aidsinfo.nih.gov/ConsentFiles/AdultandAdolescentGL.pdf>, 2012.
- Schafer JJ, Ravi S, Rowland EV, et al. The expanding class of non-nucleoside reverse transcriptase inhibitors for the treatment of HIV-1 infection. *Pharm Ther* 36:46–364, 2011.
- Thompson MA, Aberg JA, Cahn P, et al. Antiretroviral treatment of adult HIV infection: 2012 recommendations of the International Antiviral Society — USA panel. *JAMA* 308:387–402, 2012.
- Urial A, Lewthwaite P. Malaria therapy in HIV: drug interactions between nevirapine and quinine. *Int J STD AIDS* 22:768, 2011.
- Website: [www.pdr.net](http://www.pdr.net)

## Nitazoxanide (Alinia)

**Drug Class:** Antiprotozoal; anti-*C. difficile* antibiotic.

**Usual Dose:** 500 mg (PO) q12h.

**Pharmacokinetic Parameters:**

*Peak serum level:* 9.1–10.6 mcg/mL; tizoxanile/tizoxanile glucuronide (metabolites)

*Bioavailability:* No data

*Excreted unchanged (urine):* < 10%

*Serum half-life (normal/ESRD):* No data

*Plasma protein binding:* 99%

*Volume of distribution ( $V_d$ ):* No data

**Primary Mode of Elimination:** Hepatic (66%); renal (34%)

**Dosage Adjustments\***

CrCl 50–80 mL/min	No change
CrCl 10–50 mL/min	250 mg (PO) q12h
CrCl < 10 mL/min	Use with caution
Post–HD dose	None
Post–PD dose	None
CVVH/CVVHD/ CVVHDF dose	No change
Mild/moderate	No change
Mild hepatic insufficiency	No change
Moderate—severe hepatic insufficiency	Use with caution

**Drug Interactions:** Avoid in patients with hypersensitivity to salicylates.

**Adverse Effects:** Dizziness, headache, nausea/vomiting, abdominal pain, flu-like syndrome, yellow discoloration of sclera/urine.

**Allergic Potential:** Low

**Safety in Pregnancy:** B

**Comments:** Nitazoxanide concentrates in GI tract but is not present in serum. Absorption greatly increased with food.

(↑ plasma levels of metabolites ~ 50%). Contains 1.4 gm sucrose/5 mL of reconstituted suspension.

**Preferred therapy for *Cryptosporidia* and giardiasis. Also effective against *Cyclospora* and *Isopora*. Effective for *C. difficile* diarrhea/colitis (including metronidazole/vancomycin failures). Also useful for *H. pylori* or *B. hominis* and *Norovirus* diarrhea.**

## REFERENCES:

- Aslam S, Hamill RJ, Musher DM. Treatment of Clostridium difficile-associated disease: old therapies and new strategies. *Lancet Infect Dis* 5:549–57, 2005.

\*Usual dose\* assumes normal renal/hepatic function. \*For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

- Basu PP, Rayapudi K, Pacana T, et al. A randomized study comparing levofloxacin, omeprazole, nitazoxanide, and doxycycline versus triple therapy for the eradication of *Helicobacter pylori*. *AM J Gastroenterol* 106:1970–5, 2011.
- Bouza E, Burillo A, Munoz P. Antimicrobial therapy of *Clostridium difficile*-associated diarrhea. *Med Clin N Am* 90:1141–63, 2006.
- Freeman J, Baines SD, Todhunter SL, et al. Nitazoxanide is active against *Clostridium difficile* strains with reduced susceptibility to metronidazole. *J Antimicrob Chemother* 66:1407–1408, 2011.
- Gilles HM, et al. Treatment of intestinal parasitic infections: a review of nitazoxanide. *Trends in Parasitology* 18:95–7, 2002.
- Gurgen J, Hogan D, Grace E, et al. Nitazoxanide in the treatment of chronic cutaneous leishmaniasis resistant to traditional sodium stibogluconate. *J Am Acad Dermatol* 64: 202–203, 2011.
- Jung KS, Park JJ, Chon YE, et al. Risk factors for treatment failure and recurrence after metronidazole treatment for *Clostridium difficile*-associated diarrhea. *Gut Liver* 4:332–337, 2011.
- Musher DM, Logan N, Hamill RJ, et al. Nitazoxanide for the treatment of *Clostridium difficile* colitis. *Clin Infect Dis* 43:421–30, 2006.
- White CA Jr. Nitazoxanide: a new broad spectrum antiparasitic agent. *Expert Rev Anti Infect Ther* 2:43–9, 2004.
- Website: www.pdr.net

## Nitrofurantoin (Macrochantin, Macrobid)

**Drug Class:** Urinary antiseptic.

**Usual Dose:** 100 mg (PO) q12h.

**Spectrum:** (see **Susceptibility Profiles** pp. 198–202).

**Resistance Potential:** Low

**Pharmacokinetic Parameters:**

*Peak serum level:* 1 mcg/mL

*Bioavailability:* 80%

*Excreted unchanged (urine):* 25%

*Serum half-life (normal/ESRD):* 0.5/1 hrs

*Plasma protein binding:* 40%

*Volume of distribution ( $V_d$ ):* 0.8 L/kg

**Primary Mode of Elimination:** Renal  
**Dosage Adjustments\***

CrCl 50–80 mL/min	100 mg (PO) q12h
CrCl 30–50 mL/min	100 mg (PO) q24h
CrCl < 30 mL/min	Avoid
Post–HD dose	Avoid
Post–PD dose	Avoid
CVH/CVHD/ CVHDF dose	Avoid
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	No change

**Drug Interactions:** Antacids, magnesium (↓ nitrofurantoin absorption); probenecid (↑ nitrofurantoin levels).

**Adverse Effects:**

Acute hypersensitivity reactions (reversible):

pneumonitis, ALHA

Chronic reactions (irreversible): chronic hepatitis, peripheral neuropathy, interstitial fibrosis.

**Allergic Potential:** Moderate

**Safety in Pregnancy:** B

**Comments:** For CAB/lower UTIs only.

**Useful for asymptomatic bacteriuria of pregnancy;** no transplacental transfer. Chronic toxicities associated with prolonged use/renal insufficiency. Optimal effectiveness if CrCl > 60 mL/min. Usually effective if CrCl > 30 mL/min. Avoid if CrCl < 30 mL/min. **Active against most aerobic GNB uropathogens (including many ESBL + strains), except *P. aeruginosa*, *Serratia marcescens*, and *Proteus* sp.** **Effective for VSE/VRE catheter associated bacteriuria (CAB)/lower UTIs** (also see **Antibiotic Pearls & Pitfalls** p. 516).

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.



## REFERENCES:

- Auer S, Wojna A, Hell M. Oral treatment options for ambulatory patients with urinary tract infections caused by extended-spectrum-beta-lactamase-producing *Escherichia coli*. *Antimicrob Agents Chemother* 54:4006–4008, 2010.
- Cunha BA. Prophylaxis for recurrent urinary tract infections: nitrofurantoin, not trimethoprim-sulfamethoxazole or cranberry juice. *Arch Intern Med* 172:82–3, 2012.
- Cunha BA, Schoch P, Hage JE. Oral therapy of catheter-associated bacteriuria (CAB) in the era of antibiotic resistance: nitrofurantoin revisited. *J Chemother* 24:122–4, 2012.
- Cunha BA. Prophylaxis for Recurrent Urinary Tract Infections (UTIs): Nitrofurantoin, Not TMP-SMX or Cranberry Juice. *Archives of Internal Medicine* 172:82, 2012.
- Cunha BA, Schoch PE, Hage JR. Nitrofurantoin: preferred empiric therapy for community-acquired-lower urinary tract infections. *Mayo Clin Proc* 86:1243–4, 2011.
- Cunha BA. Nitrofurantoin: Preferred Empiric Therapy for Community Acquired and Nosocomial Lower Urinary Tract Infections (UTIs). *Mayo Clinic Proceedings* 86: 1243–1244, 2011.
- Cunha BA. New uses for older antibiotics: nitrofurantoin, amikacin, colistin, polymyxin B, doxycycline, and minocycline revisited. *Med Clin North Am* 90:1089–107, 2006.
- Cunha BA. Nitrofurantoin: An update. *OB/GYN* 44:399–406, 1989.
- Cunha BA. Nitrofurantoin: Bioavailability and therapeutic equivalence. *Adv Ther* 5:54–63, 1988.
- Livermore DM, Warner M, Mushtaq S, et al. What remains against carbapenem-resistant Enterobacteriaceae? Evaluation of chloramphenicol, ciprofloxacin, colistin, fosfomicin, minocycline, nitrofurantoin, temocillin and tigecycline. *Int J Antimicrob Agents* 37:415–9, 2011.
- McKinnel JA, Stollenwerk NS, Jung CW, et al. Nitrofurantoin compares favorably to recommended agents as empirical treatment of uncomplicated urinary tract infections in a decision and cost analysis. *Mayo Clin Proc* 86:480–8, 2011.
- Mendoza-Valdes A, Rosete A, Rios Bueno E, Frentzel A, et al. Antimicrobial and clinical efficacy of nitrofurantoin in the treatment of acute lower urinary tract infections in adults. *Med Klin* 105:698–704, 2010.
- Website: [www.pdr.net](http://www.pdr.net)

## Ofloxacin (Oflox)

**Drug Class:** Quinolone.

**Usual Dose:** 400 mg (IV/PO) q12h (see comments).

**Spectrum:** (see *Susceptibility Profiles* pp. 198–202).

**Resistance Potential:** Low

**Pharmacokinetic Parameters:**

*Peak serum level:* 5.5–7.2 mcg/ml

*Bioavailability:* 95%

*Excreted unchanged (urine):* 90%

*Serum half-life (normal/ESRD):* 6/40 hrs

*Plasma protein binding:* 32%

*Volume of distribution (V<sub>d</sub>):* 2 L/kg

**Primary Mode of Elimination:** Renal  
**Dosage Adjustments\***

CrCl 20–50 mL/min	400 mg (IV/PO) q24h
CrCl < 20 mL/min	200 mg (IV/PO) q24
Post–HD dose	200 mg (IV/PO)
Post–PD dose	200 mg (IV/PO)
CVWH/CVWHD/ CVWHDf dose	300 mg (IV/PO) q24h
Mild-moderate hepatic insufficiency	No change
Severe hepatic insufficiency	400 mg (IV/PO) q24h

**Drug Interactions:** Al<sup>3+</sup>, Ca<sup>2+</sup>, Fe<sup>2+</sup>, Mg<sup>2+</sup>, Zn<sup>2+</sup> antacids, citrate/citric acid, dairy products (↓ absorption of ofloxacin only if taken together); cimetidine (↑ ofloxacin levels); cyclosporine (↑ cyclosporine levels); NSAIDs (CNS stimulation); probenecid (↑ ofloxacin levels); warfarin (↑ INR).

**Adverse Effects:** Drug fever/rash, mild neuroexcitatory symptoms. *C. difficile* diarrhea/colitis. FQ use has an increased risk of tendinitis/tendon rupture. Risk in highest in the elderly, those on steroids, and those with heart, lung or renal transplants.

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

**Allergic Potential:** Low

**Safety in Pregnancy:** C

**Comments:** H<sub>2</sub> antagonist increases half-life by ~ 30%. Levofloxacin has improved pharmacokinetics/pharmacodynamics and greater antimicrobial activity. Take ofloxacin 2 hours before or after calcium/magnesium containing antacids. **PPNG dose:** 400 mg (PO) × 1 dose. **NGU/cervicitis dose:** 300 mg (PO) q12h × 1 week.

**Cerebrospinal Fluid Penetration:** < 10%

**Bile Penetration:** 1500%

(also see **Antibiotic Pearls & Pitfalls** p. 516).

#### REFERENCES:

Abسالon J, Domenico PD, Ortega AM, Cunha, BA. The antibacterial activity of ofloxacin versus ciprofloxacin against *Pseudomonas aeruginosa* in human urine. *Adv Ther* 13:191–4, 1996.

Kumar A, Girdhar BK. Multibacillary leprosy: follow up observations on 19 patients treated with 12 doses of rifampicin, ofloxacin and minocycline therapy in Agra. *Indian J Derm Venereol Leprol*. 80:156-159, 2014.

Monk JP, Campoli-Richards DM. Ofloxacin: A review of its antibacterial activity, pharmacokinetics properties, and therapeutic use. *Drugs* 33:346–91, 1987.

Website: www.pdr.net

## Oritavancin (Orbactiv)

**Drug Class:** Lipoglycopeptide

**Usual Dose:** 1200 mg (IV) × 1 dose

**Spectrum:** (see **Susceptibility Profiles** pp. 198-202)

**Resistance Potential:** Low

**Pharmacokinetic Parameters:**

*Peak serum level:* 138 mcg/ml

*Bioavailability:* Not applicable

*Excreted unchanged (urine/feces):* < 5%/< 1%

*Serum half-life (normal/ESRD):* 245 hr/No data

*Plasma protein binding:* 85%

*Volume of distribution (Vd):* 1.25 L/kg

**Primary Mode of Elimination:** renal

#### Dosage Adjustments\*

CrCl < 30 mL/min	No change (use with caution)
Post-HD dose	No change (use with caution)
Post-PD dose	No change (use with caution)
CVVH dose	No data
Hepatic insufficiency	No change

**Drug Interactions:** ↑ warfarin levels, monitor INR. Oritavancin is a weak inhibitor of CYP2C19 & CYP2C9; it is a weak inducer of CYP3A4.

**Adverse Effects:** Dizziness, headache, nausea, vomiting, skin abscesses, diarrhea, *C. difficile*, phlebitis, increased AST/ALT, hypersensitivity reactions (monitor closely patients with known hypersensitivity to glycopeptides)

**Allergic Potential:** Low

**Safety in Pregnancy:** C

**Comments:** Oritavancin artificially prolongs coagulation tests: activated clotting time (ACT); artificial effects last for 48 hours for PTT, 24 hours for PT and 24 hours for INR. Oritavancin is not removed during hemodialysis. Useful for VRE infections. Infuse slowly over 3 hours; if infusion reaction develops, slow the rate or interrupt infusion and then resume infusion upon resolution. *Use of unfractionated heparin is contraindicated for 48 hours after oritavancin administration.* ↑ incidence of osteomyelitis vs. vancomycin; institute alternate therapy if osteomyelitis is known or suspected osteomyelitis. Oritavancin may adhere to *C. difficile* spores, potentially preventing early inhibition of germinated cells; this may reduce the risk of recurrent *C. difficile*.

**Cerebrospinal Fluid Penetration:**

Non-inflamed meninges = 1%

Inflamed meninges = 10%

(also see **Antibiotic Pearls & Pitfalls** P. 198)

\*Usual dose\* assumes normal renal/hepatic function. \*For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

## REFERENCES:

- Arias CA, Mendes RE, Stilwell MG, et al. Unmet needs and prospects for oritavancin in the management of vancomycin-resistant enterococcal infections. *Clin Infect Dis*. 54 Suppl 3:S233-238, 2012.
- Belley A, Arhin FF, Sarmiento I, et al. Pharmacodynamics of a simulated single 1,200-milligram dose of oritavancin in an in vitro pharmacokinetic/pharmacodynamic model of methicillin-resistant staphylococcus aureus infection. *Antimicrob Agents Chemother*. 57:205-211, 2013.
- Chilton CH, Freeman J, Baines SO, et al. Evaluation of the effect of oritavancin on *Clostridium difficile* spore germination, outgrowth and recovery. *J Antimicrob Chemother*. 68:2078-2082, 2013.
- Corey GR, Good S, Jiang H, et al. Single-Dose Oritavancin Versus 7-10 Days of Vancomycin in the Treatment of Gram-Positive Acute Bacterial Skin and Skin Structure Infections: The SOLO II Noninferiority Study. *Clin Infect Dis*. 60:254-262, 2015.
- Corey GR, Kabler H, Mehra P, et al. Single-dose oritavancin in the treatment of acute bacterial skin infections. *N Engl J Med*. 370:2180-2190, 2014.
- Kumar A, Mann HJ, Keshitgarpour M, et al. In vitro characterization of oritavancin clearance from human blood by low-flux, high-flux, and continuous renal replacement therapy dialyzers. *Int J Artif Organs*. 34:1067-1074, 2011.
- Lin G, Pankuch G, Appelbaum PC, et al. Antistaphylococcal activity of oritavancin and its synergistic effect in combination with other antimicrobial agents. *Antimicrob Agents Chemother*. 58:6251-6254, 2014.
- Mendes RE, Farrell OJ, Sader HS, et al. Activity of oritavancin against Gram-positive clinical isolates responsible for documented skin and soft-tissue infections in European and US hospitals (2010-13). *J Antimicrob Chemother*. 70:498-504, 2015.
- Mendes RE, Sader HS, Flamm RK, et al. Oritavancin activity against *Staphylococcus aureus* causing invasive infections in U.S. and European hospitals: a 5-year international surveillance program. *Antimicrob Agents Chemother*. 58:2921-2924, 2014.
- Mendes RE, Sader HS, Flamm RK, et al. Activity of oritavancin tested against uncommonly isolated Gram-positive pathogens responsible for documented

infections in hospitals worldwide. *J Antimicrob Chemother*. 69:1579-1581, 2014.

Tice A. Oritavancin: a new opportunity for outpatient therapy of serious infections. *Clin Infect Dis*. 54 Suppl 3:S239-243, 2012.

Website: [www.pdr.net](http://www.pdr.net)

## Osetemavir (Tamiflu)

**Drug class:** Neuraminidase inhibitor.

**Usual dose:** Treatment: 75 mg (PO) q12h × 5 days; Prophylaxis: 75 mg (PO) q24h × 10 days.

**Pharmacokinetic Parameters:**

*Peak serum level:* 65.2 mcg/L

*Bioavailability:* 75%

*Excreted unchanged:* 1%

*Serum half-life (normal/ESRD):* 1–3 hours/no data  
*plasma protein binding:* 42%

*Volume of distribution (V<sub>d</sub>):* 23–26 L/kg

**Primary Mode of Elimination:** Renal (oseltamivir carboxylate)

**Dosage Adjustments\***

CrCl > 60 mL/min	No adjustment
CrCl 30–60 mL/min	30 mg (PO) q12h (treatment) 30 mg (PO) q24h (prophylaxis)
CrCl 10–30 mL/min	30 mg (PO) q24h
Post-HD dose	30 mg (PO)
Post-PD dose	30 mg (PO)
CVVH/CVVHD/ CVVHDF dose	30 mg (PO) q24h (treatment) 30 mg (PO) q24h (prophylaxis)
Moderate—severe hepatic insufficiency	Use with caution

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

**Drug Interactions:** Live attenuated influenza vaccine 2 weeks before or 2 days after oseltamivir.

**Adverse Effects:** nausea, vomiting, serious skin reactions, diplopia, visual disturbances, palpitations, atrial fibrillation, supraventricular/ventricular tachycardias neuropsychiatric events.

**Allergic Potential:** Low

**Safety in Pregnancy:** C

**Comments:** Begin prophylaxis within 3 days of exposure.

#### REFERENCES:

- Agrawal R, Rewatkar PV, Kokil GR, et al. Oseltamivir: a first line defense against swine flu. *Med Chem* 6:247–251, 2010.
- Chen H, Cheung CL, Tai H, et al. Oseltamivir-resistant influenza A pandemic (H1N1) 2009 virus, Hong Kong, China. *Emerg Infect Dis* 15:1970–1972, 2009.
- Cho D, Hossain M, Liew P, et al. Side effects of oseltamivir in end-stage renal failure patients. *Nephrol Dial Transplant* 26:2339–2344, 2011.
- Couch RB, Davis BR. Has oseltamivir been shown to be effective for treatment of H5N1 influenza? *J Infect Dis* 202:1149–1151, 2010.
- Jackson RJ, Cooper KL, Tappenden P, et al. Oseltamivir, zanamivir and amantadine in the prevention of influenza: A systematic review. *J Infect* 62:14–25, 2011.
- Lee VJ, Yap J, Maurer-Stroh S, et al. Investigation causes of oseltamivir chemoprophylaxis failures during influenza A (H1N1-2009) outbreaks. *J Clin Virology* 50:104–108, 2011.
- Sophia NG, Cowling BJ, Fang VJ, et al. Effects of oseltamivir treatment on duration of clinical illness and viral shedding and household transmission of influenza virus. *Clin Infect Dis* 50:707–714, 2010.
- Sotos JG. Emergence of oseltamivir-resistant pandemic H1N1 virus during prophylaxis. *N Engl J Med* 361:2296–2298, 2009.
- Website: www.pdr.net

## Oxacillin (Prostaphlin)

**Drug Class:** Anti-staphylococcal (MSSA) penicillin.

**Usual Dose:** 1–2 gm (IV) q4h.

Spectrum: (see **Susceptibility Profiles** pp. 186–190).

**Resistance Potential:** Low

**Pharmacokinetic Parameters:**

*Peak serum level:* 43 mcg/ml

*Bioavailability:* Not applicable

*Excreted unchanged (urine):* 39–66%

*Serum half-life (normal/ESRD):* 0.5/1 hrs

*Plasma protein binding:* 94%

*Volume of distribution ( $V_d$ ):* 0.2 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\*** (based on 2 gm q4h)

CrCl 10–80 mL/min	No change
CrCl < 10 mL/min	No change
Post–HD dose	None
Post–PD dose	None
CVWH/CVWHD/ CVWHDf dose	No change
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	No change

**Drug Interactions:** Cyclosporine (↓ cyclosporine levels); nifedipine, warfarin (↓ interacting drug effect).

**Adverse Effects:** Drug fever/rash, leukopenia, ↑ SGOT/SGPT, hepatitis.

**Allergic Potential:** High

**Safety in Pregnancy:** B

**Comments:** Avoid oral formulation (not well absorbed/erratic serum levels).

Meningeal dose = 2 gm (IV) q4h.

**Cerebrospinal Fluid Penetration:**

Non-inflamed meninges = 1%

Inflamed meninges = 10%

**Bile Penetration:** 25%

(also see **Antibiotic Pearls & Pitfalls** p. 509).

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

**REFERENCES:**

- Al-Homaidhi H, Abdel-Haq NM, El-Baba M, et al. Severe hepatitis associated with oxacillin therapy. *South Med J* 95:650–2, 2002.
- Donowitz GR, Mandell GL. Beta-lactam antibiotics. *N Engl J Med* 318:419–26 and 318:490–500, 1993.
- Wright AJ. The penicillins. *Mayo Clin Proc* 74:290–307, 1999.
- Website: www.pdr.net

## Pegylated Interferon alfa-2a (Pegasy)

**Drug Class:** HCV antiviral.

**Usual Dose:** Pegasy: 180 mcg (SQ) once weekly × 48 weeks; Ribavirin: 500 mg (< 75 kg) or 800 mg (> 75 mg) (PO) q12h × 48 weeks (see comments).

**Pharmacokinetic Parameters:**

*Trough level:* 16 ng/ml

*Bioavailability:* No data

*Excreted unchanged (urine):* No data

*Serum half-life (normal/ESRD):* 80/100–120 hrs

*Plasma protein binding:* No data

*Volume of distribution (V<sub>d</sub>):* 8–12 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\***

CrCl 10–50 mL/min	Pegasy: 180 mcg (SQ); avoid ribavirin
CrCl < 10 mL/min	Pegasy: 135 mcg (SQ) q week; avoid ribavirin
Post-HD dose	Pegasy: 135 mcg (SQ) q week; avoid ribavirin
Post-PD dose	Pegasy: 135 mcg (SQ); avoid ribavirin
CWH/CVHD/ CWHDF dose	Avoid ribavirin

Moderate hepatic insufficiency	135 mcg if ALT ↑; Avoid ribavirin
Severe hepatic insufficiency	Avoid
Moderate depression	↓ Pegasy to 135 mcg (90 mcg in some); evaluate once weekly
Severe depression	Discontinue Pegasy; immediate psychiatric consult

**Pegasy Adjustment for Hematologic Toxicity**

Absolute neutrophil count < 750/mm <sup>3</sup>	↓ Pegasy to 135 mcg
Absolute neutrophil count < 500/mm <sup>3</sup>	Discontinue Pegasy until neutrophils > 1000/mm <sup>3</sup> . Reinstitute at 90 mcg and monitor neutrophil count
Platelets < 50,000/mm <sup>3</sup>	↓ Pegasy to 90 mcg
Platelets < 25,000/mm <sup>3</sup>	Discontinue Pegasy

**Ribavirin Adjustment for Hematologic Toxicity**

Hgb < 10 gm/dL and no cardiac disease	↓ Ribavirin to 600 mg/day <sup>†</sup>
Hgb < 8.5 gm/dL and no cardiac disease	Discontinue ribavirin
Hgb ≥ 2 gm/dL ↓ during 4-week treatment period and stable cardiac disease	↓ Ribavirin to 600 mg/day*
Hgb < 12 gm/dL despite 4 weeks at reduced dose and stable cardiac disease	Discontinue ribavirin

\* One 200 mg tablet in a.m. and two 200 mg tablets in p.m.

<sup>†</sup>Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

**Drug Interactions:** Inhibits CYP 1A2. May ↑ theophylline levels.

**Adverse Effects:** Flu-like symptoms, serious neuropsychiatric disturbances, bone marrow toxicity, nausea/vomiting/diarrhea, alopecia, hypothyroidism, ARDS.

**Allergic Potential:** Low

**Safety in Pregnancy:** Pegasys: C (ribavirin: X; also avoid in partners of pregnant women)

**Comments:** Ribavirin dose with Pegasys for HCV (genotypes 2, 3) is 800 mg/d. The optimal ribavirin dose for HCV (genotype 1) has not been determined but 800–1400 mg/d has been used. Combination therapy is more effective than monotherapy. Avoid in autoimmune hepatitis and decompensated cirrhosis. Check CBC weekly. For genotypes 2 and 3, duration of therapy is 24 weeks and ribavirin dose is 400 mg (PO) bid (rather than q12h). Ribavirin should be taken with breakfast and dinner to ↑ bioavailability/↓ nausea. Ribavirin taken after dinner may cause insomnia. Ribavirin's long serum half-life allows q24h dosing, but ↑ nausea.

**Cerebrospinal Fluid Penetration:** No data

**Bile Penetration:** No data

#### REFERENCES:

- Davis GL. Combination treatment with interferon and ribavirin for chronic hepatitis C. *Clin Liver Dis* 1:811–26, 1999.
- Ferenci P. Peginterferon alfa-2a (40KD) (Pegasys) for the treatment of patients with chronic hepatitis C. *Int J Clin Pract* 57:610–5, 2003.
- Fried MN, et al. Peg-interferon alfa-2a plus ribavirin for chronic hepatitis C virus infection. *N Engl J Med* 347:975–82, 2002.
- Kamal SM, et al. Peg-interferon alone or with ribavirin enhances HCV-specific CD<sub>4</sub> T-helper responses in patients with chronic hepatitis C. *Gastroenterology* 123:1070–83, 2002.
- Liu C-H, Liang C-C, Liu C-J, et al. Pegylated interferon alfa-2a monotherapy for hemodialysis patients with acute hepatitis c. *Clin Infect Dis* 51:541–549, 2010.

- Plosker GL, Keating GM. Peginterferon-alpha-2a (40kD) plus ribavirin: a review of its use in hepatitis C virus and HIV co-infection. *Drugs* 64:2823–43, 2004.
- Rasenack J, Zeuzem S, Feinman SV, et al. Peginterferon alpha-2a (40kD) [Pegasys] improves HR-QOL outcomes compared with unmodified interferon alpha-2a [Roferon-A]: in patients with chronic hepatitis C. *Pharmacoeconomics* 21:341–9, 2003.
- Torriani FJ, Rodriguez-Torres M, Rockstroh JK, et al. Peginterferon alfa-2a plus ribavirin for chronic hepatitis C virus infection in HIV-infected patients. *N Engl J Med* 351:438–50, 2004.
- Website: [www.pdr.net](http://www.pdr.net)

## Pegylated Interferon alfa-2b (Peg-Intron)

**Drug Class:** HCV antiviral.

**Usual Dose:** See pp. 97–98.

**Pharmacokinetic Parameters:**

*Peak serum level:* 30 IU/ml / 3680 ng/ml

*Bioavailability:* No data/64%

*Excreted unchanged (urine):* No data/17%

*Serum half-life (normal/ESRD):* [40 hrs/no data] / [298 hrs/no data]

*Plasma protein binding:* No data

*Volume of distribution (V<sub>d</sub>):* 0.99 L/kg

**Primary Mode of Elimination:** Renal/renal

**Dosage Adjustments\***

CrCl 30–50 mL/min	give 75% of Peg-Intron dose; avoid ribavirin
CrCl < 30 mL/min	give 50% of Peg-Intron dose; avoid ribavirin
Post–HD dose	Avoid ribavirin
Post–PD dose	Avoid ribavirin
CVWH/CVWHD/ CVWHDf dose	Avoid ribavirin

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

Moderate hepatic insufficiency	None
Severe hepatic insufficiency	Avoid
Moderate depression	↓ Peg-Intron by 50%; evaluate once weekly
Severe depression	Discontinue both drugs; immediate psychiatric consult

### Dosage Adjustments for Hematologic Toxicity

Hgb < 10 gm/dL	↓ ribavirin by 200 mg/day
Hgb < 8.5 gm/dL	Permanently discontinue both drugs
WBC < $1.5 \times 10^9/L$	↓ Peg-Intron by 50%
WBC < $1.0 \times 10^9/L$	Permanently discontinue both drugs
Neutrophils < $0.75 \times 10^9/L$	↓ Peg-Intron by 50%
Neutrophils < $0.5 \times 10^9/L$	Permanently discontinue both drugs
Platelets < $80 \times 10^9/L$	↓ Peg-Intron by 50%
Platelets < $50 \times 10^9/L$	Permanently discontinue both drugs
Hgb $\geq 2$ gm/dL ↓ during 4-week treatment period and stable cardiac disease	↓ Peg-Intron by 50% and ribavirin by 200 mg/day
Hgb < 12 gm/dL despite ribavirin dose reduction	Permanently discontinue both drugs

**Drug Interactions:** Nucleoside analogs; fatal/non-fatal lactic acidosis. May ↑ levels of

theophylline. Overlapping toxicity with other bone marrow suppressants.

**Adverse Effects:** Hemolytic anemia, psychological effects, cytopenias, pulmonary symptoms, pancreatitis, hypersensitivity reactions, ↑ triglycerides, flu-like symptoms, nausea/vomiting, alopecia, rash, hypothyroidism.

**Allergic Potential:** Low

**Safety in Pregnancy:** Peg-Intron: C (ribavirin: X; also avoid in partners of pregnant women)

**Comments:** Avoid in autoimmune hepatitis. Check CBC pretreatment and at 2 and 4 weeks (must adjust doses for ↓ WBC, ↓ neutrophils, ↓ platelets, ↓ hemoglobin). Poorly tolerated in decompensated cirrhosis or recurrent hepatitis C after transplant. Be alert for severe depression/psychiatric changes.

**Cerebrospinal Fluid Penetration:** No data

**Bile Penetration:** No data

### REFERENCES:

- Alfaleh FZ, Hadad Q, Khuroo MS, et al. Peginterferon alpha-2b plus ribavirin compared with interferon alpha-2b plus ribavirin for initial treatment of chronic hepatitis C in Saudi patients commonly infected with genotype 4. *Liver Int* 24:568–74, 2004.
- Fried MN. Side effects of therapy of hepatitis C and their management. *Hepatology* 36:5237–44, 2002.
- Jaeckel E, Cornberg M, Wedemeyer H, et al. Treatment of acute hepatitis C with interferon alpha-2b. *N Engl J Med* 345:1452–7, 2001.
- Wright TL. Treatment of patients with hepatitis C and cirrhosis. *Hepatology* 36:S185–91, 2002.
- Website: [www.pdr.net](http://www.pdr.net)

## Penicillin G (various)

**Drug Class:** Natural penicillin.

**Usual Dose:** 2–4 mu (IV) q4h (see comments).

**Spectrum:** (see *Susceptibility Profiles* pp. 186–190).

"Usual dose" assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose  $\times 1$  followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

**Resistance Potential:** Low

**Pharmacokinetic Parameters:**

Peak serum level: 20–40 mcg/ml

Bioavailability: Not applicable

Excreted unchanged (urine): 80%

Serum half-life (normal/ESRD): 0.5/5.1 hrs

Plasma protein binding: 60%

Volume of distribution ( $V_d$ ): 0.3 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\***

CrCl 50–80 mL/min	2–4 mu (IV) q4h
CrCl 10–50 mL/min	1–2 mu (IV) q4h
CrCl < 10 mL/min	1 mu (IV) q8h
Post-HD dose	2 mu (IV)
Post-PD dose	0.5 mu (IV)
CVWH/CVVHD/CVVHDF dose	2 mu (IV) q6h
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	No change

**Drug Interactions:** Probenecid (↑ penicillin G levels).

**Adverse Effects:** Drug fever/rash, E. multiforme/Stevens–Johnson Syndrome; anaphylactic reactions (hypotension, laryngospasm, bronchospasm), hives, serum sickness.

**Allergic Potential:** High

**Safety in Pregnancy:** B

**Comments:** Incompatible in solutions containing erythromycin, aminoglycosides, calcium bicarbonate, or heparin.

Penicillin G (potassium): K<sup>+</sup> content = 1.7 mEq/g;

Na<sup>+</sup> content = 0.3 mEq/g.

Penicillin G (sodium): Na<sup>+</sup> content = 2 mEq/g. **1°**,

**2°**, early latent syphilis dose: PCN benzathine

2.4 mu (IM) × 1 dose. **Late latent syphilis dose:** PCN benzathine 2.4 mu (IM) q8h once weekly × 3. **Neurosyphilis dose:** PCN G 4 mu (IV) q4h × 2 weeks.

Meningeal dose = 4 mu (IV) q4h.

**Cerebrospinal Fluid Penetration:**

Non-inflamed meninges ≤ 1%

Inflamed meninges = 5%

**Bile Penetration:** 500%

(also see **Antibiotic Pearls & Pitfalls** p. 508).

**REFERENCES:**

- Falco V, Almirante B, Jordano Q, et al. Influence of penicillin resistance of outcome in adult patients with invasive pneumococcal pneumonia. Is penicillin useful against intermediately resistant strains? J Antimicrob Chemother 54:481–8, 2004.
- Giachetto G, Pirez MC, Nanni L, et al. Ampicillin and penicillin concentration in serum and pleural fluid of hospitalized children with community-acquired pneumonia. Pediatr Infect Dis J 7:625–9, 2004.
- Sandoe JA, Patel PA, Baig MW, et al. What is the effect of penicillin dosing interval on outcomes in streptococcal infective endocarditis? J antimicrob Chemo 68:2660–2663, 2013.
- Wendel Jr GD, Sheffield JS, Hollier LM, et al. Treatment of syphilis in pregnancy and prevention of congenital syphilis. Clin Infect Dis 35(Suppl2):S200–9, 2002.
- Wright AJ. The penicillins. Mayo Clin Proc 74:290–307, 1999.
- Website: www.pdr.net

**Penicillin V (various)**

**Drug Class:** Natural penicillin.

**Usual Dose:** 500 mg (PO) q6h.

(Take without food).

**Spectrum:** (see **Susceptibility Profiles** pp. 186–190).

**Resistance Potential:** Low

**Pharmacokinetic Parameters:**

Peak serum level: 5 mcg/ml

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.



*Bioavailability:* 60%

*Excreted unchanged (urine):* 80%

*Serum half-life (normal/ESRD):* 0.5/8 hrs

*Plasma protein binding:* 70%

*Volume of distribution ( $V_d$ ):* 0.5 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\***

CrCl 10–50 mL/min	No change
CrCl < 10 mL/min	250–500 mg (PO) q6h
Post-HD dose	250 mg (PO)
Post-PD dose	250 mg (PO)
CVWH/CVHD/ CWHDF dose	500 mg (PO) q6h
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	No change

**Drug Interactions:** Probenecid (↑ penicillin V levels).

**Adverse Effects:** Drug fever/rash, E. multiforme/Stevens-Johnson Syndrome, anaphylactic reactions (hypotension, laryngospasm, bronchospasm), hives, serum sickness C. difficile diarrhea/colitis.

**Allergic Potential:** High

**Safety in Pregnancy:** B

**Comments:** Jarisch–Herxheimer reactions when treating spirochetal infections, e.g., Lyme disease, syphilis, yaws.

**Cerebrospinal Fluid Penetration:** < 10%

(also see **Antibiotic Pearls & Pitfalls** p. 508).

#### REFERENCES:

Bisno AL, Gerber MA, Swaltney JM, et al. Practice guidelines for the diagnosis and management of group

A streptococcal pharyngitis. Infectious Diseases Society of America. Clin Infect Dis 35:113–25, 2002.

Chiou CC, Does penicillin remain the drug of choice for pneumococcal pneumonia in view of emerging in vitro resistance? Clin Infect Dis 42:234–7, 2006.

Donowitz GR, Mandell GL. Beta-lactam antibiotics. N Engl J Med 318:419–26 and 490–500, 1993.

Klugman KP, Yu VL. No Impact of Penicillin Resistance on Mortality. Clin Infect Dis 43:261, 2006.

Peterson LR. Penicillin for treatment of pneumococcal pneumonia: does in vitro resistance really matter? Clin Infect Dis 42:224–33, 2006.

Portier H, Filipecki J, Weber P, et al. Five day clarithromycin modified release versus 10 day penicillin V for group A streptococcal pharyngitis: a multi-centre, open-label, randomized study. J Antimicrob Chemother 49:337–44, 2002.

Wright AJ. The penicillins. Mayo Clin Proc 74:290, 1999.

Website: www.pdr.net

## Pentamidine (Pentam 300, NebuPent)

**Drug Class:** Antiparasitic.

**Usual Dose:** 4 mg/kg (IV) q24h (see comments).

**Pharmacokinetic Parameters:**

*Peak serum level:* 0.6–1.5 mcg/mL

*Bioavailability:* Not applicable

*Excreted unchanged (urine):* 50%

*Serum half-life (normal/ESRD):* 6.4/90 hrs

*Plasma protein binding:* 69%

*Volume of distribution ( $V_d$ ):* 5 L/kg

**Primary Mode of Elimination:** Metabolized

**Dosage Adjustments\***

CrCl < 10 mL/min	No change
Post-HD dose	None
Post-PD dose	None
CVWH/CVHD/CWHDF dose	No change
Moderate—severe hepatic insufficiency	No change

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

**Drug Interactions:** Alcohol, valproic acid, didanosine, ( $\uparrow$  risk of pancreatitis); foscarnet (severe hypocalcemia reported; do not combine); amphotericin B, aminoglycosides, capreomycin, cis-platinum, colistin, methoxyflurane, polymyxin B, vancomycin, other nephrotoxic drugs ( $\uparrow$  nephrotoxicity).

**Adverse Effects:** Severe hypotension, hypocalcemia, hypoglycemia, increase in creatinine, pancreatitis, severe leukopenia, anemia and thrombocytopenia. May increase the QT interval with IV administration and cause ventricular tachycardia. Severe hypoglycemic episodes may result in later diabetes mellitus. IV extravasation may cause severe skin necrosis and IM injections can be severely painful and should be avoided if possible. The following tests should be carried out at regular intervals; Daily blood urea nitrogen and serum creatinine, blood glucose, complete blood count and platelet count as well as every other day bilirubin, alkaline phosphatase, AST (SGOT), and ALT (SGPT). An electrocardiogram should be done at regular intervals to calculate the  $QT_c$  interval. If creatinine starts to rise the dose should be cut back to 3 or 2 mg/kg. The medication should be held if hypoglycemia occurs and or if the  $QT_c$  interval increases.

**Allergic Potential:** High

**Safety in Pregnancy:** C

**Comments:** Well absorbed IM, but painful.

Administer IV slowly in  $D_5W$  over 1 hour, not saline/may be used at a dose of 300 mg aerosol q month to prevent aerosol pentamidine. Caution should be used with inhaled pentamidine especially around pregnant employees as it can induce spontaneous abortion. Special attention should be made to the proper aerosolization of medication so that it is distributed evenly through the lung. Adverse effects with aerosolized

pentamidine include chest pain, arrhythmias, dizziness, wheezing, coughing, dyspnea, headache, anorexia, nausea, diarrhea, rash, pharyngitis. **If PCP patient also has pulmonary TB, aerosolized pentamidine treatments may expose medical personnel to TB via droplet inhalation.** Nebulizer dose: Inhaled pentamidine isethionate (NebuPent) 300 mg monthly via Respigard II nebulizer can be used for PCP prophylaxis. **Nebulized pentamidine less effective than IV/IM pentamidine and is not effective against extrapulmonary PCP.** Use with caution in renal disease. Subtle creatinine changes may be associated with severe toxicity.

**Cerebrospinal Fluid Penetration:** < 10%

#### REFERENCES:

- Goa KL, Campoli-Richards DM. Pentamidine isethionate: A review of its antiprotozoal activity, pharmacokinetic properties and therapeutic use in *Pneumocystis carinii* pneumonia. *Drugs* 33:242–58, 1987.
- Guerin PJ, Alar P, Sundar S, et al. Visceral leishmaniasis: current status of control, diagnosis, and treatment, and a proposed research and development agenda. *Lancet Infect Dis* 2:494–501, 2002.
- Lionakis MS, Lewis RE, Samonis G, et al. Pentamidine is active in vitro against *Fusarium* species. *Antimicrob Agents Chemother* 47:3252–9, 2003.
- Monk JP, Benfield P. Inhaled pentamidine: An overview of its pharmacological properties and a review of its therapeutic use in *Pneumocystis carinii* pneumonia. *Drugs* 39:741–56, 1990.
- Rodriguez M, Fishman JA. Prevention of infection due to *Pneumocystis* spp. in human immunodeficiency virus-negative immunocompromised patients. *Clin Microbiol Rev* 17:770–82, 2004.
- Sattler FR, Cowan R, Nielsen DM, et al. Trimethoprim-sulfamethoxazole compared with pentamidine for treatment of *Pneumocystis carinii* pneumonia in the acquired immunodeficiency syndrome. *Ann Intern Med* 109:280–7, 1988.
- Website: [www.pdr.net](http://www.pdr.net)

<sup>†</sup>Usual dose assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose  $\times$  1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

## Peramivir (Rapivab)

**Drug Class:** Neuroaminidase inhibitor.

**Usual Dose:** 600 mg (IV) over 30 minutes × 1 dose.

**Pharmacokinetic Parameters:**

*Peak serum level:* 17–44 mcg/mL/13 mcg/mL

*Bioavailability:* not applicable

*Excreted unchanged:* no data

*Serum half-life (normal/ESRD):* 7–20 hrs/no data

*Plasma protein binding:* no data

*Volume of distribution ( $V_d$ ):* no data

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\***

**Drug Interactions:** No data.

**Adverse Effects:** Diarrhea, nausea, vomiting, hallucinations, delirium, abnormal behavior, drug rash E. multiforme, anaphylaxis.

**Allergic Potential:** Low (do not use if severely allergic to other neuroaminidase inhibitors)

**Safety in Pregnancy:** C

**Comments:** For treatment of admitted adult and pediatric patients with confirmed influenza A.

**The patients not responding to either oral or inhaled antiviral therapy or when drug delivery not possible or suboptimal by another route.** For optimal activity give within 2 days of influenza onset.

**Highly active against influenza A.**

**Cerebrospinal Fluid Penetration:** No data

**REFERENCES:**

Arya V, Carter WW, Robertson SM. The role of clinical pharmacology in supporting the emergency use authorization of an unapproved anti-influenza drug, peramivir. *Clin Pharmacol Ther* 88:587–589, 2010.

Bazan JA, Bauer KA, Hollister AS, et al. Peramivir pharmacokinetics in two critically ill adults with 2009 H1N1 influenza A concurrently receiving continuous renal replacement therapy. *Pharmacotherapy* 30:1016–1020, 2010.

Castillo R, Holland LE, Bolz DA. Peramivir and its use in H1N1 influenza. *Drugs Today (Barc)* 46:399–408 2010.

Hernandez JE, Adiga R, Armstrong R, et al. Clinical experience in adults and children treated with intravenous peramivir for 2009 influenza A (H1N1) under an emergency IND program in the United States. *Clin Infect Dis* 52:695–706, 2011.

Kohn S, Kida H, Mizuguchi M, et al. Efficacy and safety of intravenous peramivir for treatment of seasonal influenza virus infection. *Antimicrob Agents Chemother* 54:4568–4574, 2010.

Kohn S, Kida H, Mizuguchi M, et al. Intravenous peramivir for treatment of influenza A and B virus infection in high-risk patients. *Antimicrob Agents Chemother* 55:2803–2812, 2011.

Mancuso CE, Gabay MP, Steinke LM, et al. Peramivir: an intravenous neuraminidase inhibitor for the treatment of 2009 H1N1 influenza. *Ann Pharmacother* 44:1240–1249, 2010.

Moss RB, Davey RT, Steigbigel RT, et al. Targeting pandemic influenza: a primer on influenza antivirals and drug resistance. *J Antimicrob Chemother* 65:1086–1093, 2010.

Nasu T, Ogawa D, Wada J, et al. Peramivir for severe influenza infection in a patient with diabetic nephropathy. *Am J Respir Crit Care Med* 182:1209–1210, 2010.

Nguyen HTM, Sheu TG, Mishin VP, et al. Assessment of pandemic and seasonal influenza A (H1N1) virus susceptibility to neuraminidase inhibitors in three enzyme activity inhibition assays. *Antimicrob Agents Chemother* 54:3671–3677, 2010.

Thomas B, Hollister AS, Muczynski KA. Peramivir clearance in continuous renal replacement therapy. *Hemodial Int* 14:339–340, 2010.

Website: [www.cdc.gov](http://www.cdc.gov)

Website: [www.pdr.net](http://www.pdr.net)

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

## Piperacillin (Pipracil)

**Drug Class:** Anti-pseudomonal penicillin.

**Usual Dose:** 3 gm (IV) q4-6h (see comments).

**Spectrum:** (see *Susceptibility Profiles* pp. 186–190).

**Resistance Potential:** Low

**Pharmacokinetic Parameters:**

*Peak serum level:* 412 mcg/ml

*Bioavailability:* Not applicable

*Excreted unchanged (urine):* 50–70%

*Serum half-life (normal/ESRD):* 1/3 hrs

*Plasma protein binding:* 16%

*Volume of distribution ( $V_d$ ):* 0.24 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\*** (based on 4 gm q8h)

CrCl 20–50 mL/min	3 gm (IV) q8h
CrCl < 20 mL/min	3 gm (IV) q12h
Post-HD dose	1 gm (IV)
Post-PD dose	2 gm (IV)
CVVH/CVVHD/CVHDF dose	3 gm (IV) q8h
Moderate or severe hepatic insufficiency	No change

**Drug Interactions:** Aminoglycosides (inactivation of piperacillin in renal failure); warfarin (↑ INR); oral contraceptives (↓ oral contraceptive effect); cefoxitin (↓ piperacillin effect).

**Adverse Effects:** Drug fever/rash, anaphylactic reactions (hypotension, laryngospasm, broncho-spasm), hives, serum sickness, leukopenia, hemolytic anemia, *C. difficile* diarrhea/colitis.

**Allergic Potential:** High

**Safety in Pregnancy:** B

**Comments:** 75% absorbed when given IM.

Do not mix/administer with aminoglycosides. **Most active antipseudomonal penicillin against *P. aeruginosa*. Nosocomial pneumonia/*P. aeruginosa* dose:** 3 gm (IV) q4h.

Meningeal dose = usual dose.

**Cerebrospinal Fluid Penetration:**

Non-Inflamed meninges = 1%

Inflamed meninges = 30%

**Bile Penetration:** 1000%

(also see *Antibiotic Pearls & Pitfalls* p. 509).

### REFERENCES:

- Donowitz GR, Mandell GL. Beta-lactam antibiotics. *N Engl J Med* 318:419–26 and 318:490–500, 1993.
- Kim MK, Capitano B, Mattoes HM, et al. Pharmacokinetic and pharmacodynamic evaluation of two dosing regimens for piperacillin-susceptible organisms. *Pharmacotherapy* 22:569–77, 2002.
- Peacock JE, Herrington DA, Wade JC, et al. Ciprofloxacin plus piperacillin compared with tobramycin plus piperacillin as empirical therapy in febrile neutropenic patients. A randomized, double-blind trial. *Ann Intern Med* 137:120, 2002.
- Tan JS, File TM, Jr. Antipseudomonal penicillins. *Med Clin North Am* 79:679–93, 1995.
- Website: www.pdr.net

## Piperacillin/Tazobactam (Zosyn, Tazocin)

**Drug Class:** Anti-pseudomonal penicillin/ $\beta$ -lactamase inhibitor.

**Usual Dose:** 3.375 gm (IV) q6h (**PI dose**) or 4.5 gm (IV) q8h (**PK dose**) (see comments).

**Spectrum:** (see *Susceptibility Profiles* pp. 186–190).

**Resistance Potential:** Low

**Pharmacokinetic Parameters:**

*Peak serum level:* 298/34 mcg/ml

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose  $\times$  1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

*Bioavailability: Not applicable*

*Excreted unchanged (urine): 60/80%*

*Serum half-life (normal/ESRD): [1.5/8] / [1/7] hrs*

*Plasma protein binding: 30/30%*

*Volume of distribution ( $V_d$ ): 0.3/0.21 L/kg*

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\*** (also see comments)

CrCl 20–40 mL/min	2.25 gm (IV) q6h 3.375 (IV) q6h for NP
CrCl < 20 mL/min	2.25 gm (IV) q8h 2.25 (IV) q6h for NP
CrCl < 10 mL/min	2.25 gm (IV) q8h
Post-HD dose	0.75 gm (IV)
Post-PD dose	None
CVH/CVHD/ CWHDF dose	3.375 gm (IV) q6h
Moderate—severe hepatic insufficiency	No change

**Drug Interactions:** Aminoglycosides (↓ aminoglycoside levels); vecuronium (↓ vecuronium effect); probenecid (↑ piperacillin/tazobactam levels); methotrexate (↑ methotrexate levels).

**Adverse Effects:** Drug fever/rash, eosinophilia, ↓/↑ platelets, leukopenia (with prolonged use > 21 days), ↑ PT/PTT, mild transient ↑ SGOT/SGPT, insomnia, nausea, headache, constipation, hypertension, hemolytic anemia, C. difficile diarrhea/colitis. May cause false + BG.

**Allergic Potential:** High

**Safety in Pregnancy:** B

**Comments:** For *P. aeruginosa/nosocomial pneumonia (normal CrCl) use 4.5 g (IV) q6h plus amikacin 1 g (IV) q24h. P. aeruginosa/nosocomial pneumonia dosing with ↓ CrCl:*

CrCl > 40 mL/min: 4.5 gm (IV) q6h.

CrCl 20–40 mL/min: 3.375 gm (IV) q6h.

CrCl < 20 mL/min: 2.25 gm (IV) q6h.

**Hemodialysis (HD) doses:** 2.25 gm (IV) q8h; give 0.75 gm (IV) Post-HD on HD days.

**Cerebrospinal Fluid Penetration:**

Non-Inflamed meninges = 1%

Inflamed meninges = 30%

**Bile Penetration:** 6000%

(also see **Antibiotic Pearls & Pitfalls** pp. 509, 512).

#### REFERENCES:

- Bauer SR, Salem C, Connor MJ Jr, et al. Pharmacokinetics and pharmacodynamics of piperacillin-tazobactam in 42 patients treated with concomitant CRRT. *Clin J Am Soc Nephrol* 7:452–457, 2012.
- Gin A, Dilay L, Karlowsky JA, et al. Piperacillin-tazobactam: a beta-lactam/beta-lactamase inhibitor combination. *Expert Rev Anti Infect Ther* 5:365–383, 2007.
- Hayashi Y, Roberts JA, Paterson DL, et al. Pharmacokinetic evaluation of piperacillin-tazobactam. *Expert Opin Drug Metab Toxicol* 6:1017–1031, 2010.
- Jones RN, Stilwell MG, Rhomberg PR, et al. Antipseudomonal activity of piperacillin/tazobactam: more than a decade of experience from the SENTRY Antimicrobial Surveillance Program (1997–2007). *Diagn Microbiol Infect Dis* 65:331–334, 2009.
- Lau WK, Mercer D, Itani KM, et al. Randomized, open-label, comparative study of piperacillin-tazobactam administered by continuous infusion versus intermittent infusion for treatment of hospitalized patients with complicated intra-abdominal infection. *Antimicrob Agents and Chemother* 50:3556–61, 2006.
- Mah GT, Mabasa VH, Chow I, et al. Evaluating outcomes associated with alternative dosing strategies for piperacillin/tazobactam: a qualitative systematic review. *Ann Pharmacother* 46:265–267, 2012.
- Mikulska M, Furfaro E, Del Bono V, et al. Piperacillin/tazobactam seems to be no longer responsible for false-positive results of the galactomannan assay. *J Antimicrob Chemother* 67:1746–1748, 2012.

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

Mattoes HM, Capitano B, Kim MK, et al. Comparative pharmacokinetic and pharmacodynamic profile of piperacillin/tazobactam 3.375 G Q4H and 4.5 G Q6H. *Chemotherapy* 458:59–63, 2002.

Minnaganti VR, Cunha BA. Piperacillin/tazobactam. *Antibiotics for Clinicians* 3:101–8, 1999.

Saltoglu N, Dalkiran A, Tetiker T, et al. Piperacillin/tazobactam versus imipenem/cilastatin for severe diabetic foot infections: a prospective, randomized clinical trial in a university hospital. *Clin Microbiol Infect* 16:1252–1257, 2010.

Yamagishi Y, Terada M, Ohki E, et al. Investigation of the clinical breakpoints of piperacillin-tazobactam against infections caused by *Pseudomonas aeruginosa*. *J Infect Chemother* 18:127–129, 2012.

Website: www.pdr.net

## Polymyxin B

**Drug Class:** Cell membrane altering antibiotic.

**Usual Dose:** 1–1.25 mg/kg (IV) q12h (1 mg = 10,000 units) (see comments).

**Spectrum:** (see *Susceptibility Profiles* pp. 198–202).

**Pharmacokinetic Parameters:**

*Peak serum level:* 8 mcg/ml

*Bioavailability:* Not applicable

*Excreted unchanged (urine):* 60%

*Serum half-life (normal/ESRD):* 6/48 hrs

*Plasma protein binding:* < 10%

*Volume of distribution ( $V_d$ ):* No data

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\***

CrCl 20–50 mL/min	1 mg/kg (IV) q12h
CrCl 5–20 mL/min	0.5 mg/kg (IV) q12h
CrCl < 5 mL/min	0.2 mg/kg (IV) q12h
Post-HD/PD dose	No information

CVVH/CVVHD/ CVHDF dose	0.5 mg/kg (IV) q12h
Moderate—severe hepatic insufficiency	No change

**Drug Interactions:** Amphotericin B, amikacin, gentamicin, tobramycin, vancomycin (↑ nephrotoxicity).

**Adverse Effects:** Renal failure. Neurotoxicity associated with very prolonged/high serum levels; neuromuscular blockade potential with renal failure/neuromuscular disorders.

**Resistance Potential:** Low

**Allergic Potential:** Low

**Safety in Pregnancy:** B

**Comments:** Inhibits endotoxin release from gram-negative bacilli. Avoid intraperitoneal infusion due to risk of neuromuscular blockade. Increased risk of reversible non-oliguric renal failure (ATN) when used with other nephrotoxic drugs. No ototoxic potential. May be given IM with procaine, but painful. **Nebulizer dose for multidrug resistant *P. aeruginosa* in cystic fibrosis/bronchiectasis:** 80 mg in saline via aerosol/nebulizer q8h (for recurrent infection use 160 mg). **Intrathecal (IT) polymyxin B dose:** 5 mg (50,000 u) q24h × 3 days, then q48h × 2 weeks. Dissolve 50 mg (500,000 u) into 10 ml for IT administration.

**Cerebrospinal Fluid Penetration:** < 10%

(also see *Antibiotic Pearls & Pitfalls* p. 520).

### REFERENCES:

- Cavaillon JM. Polymyxin B for endotoxin removal in sepsis. *Lancet Infect Dis* 11:426–7, 2011.
- Falagas ME, Kasiakou SK, Colistin: the revival of polymyxins for the management of multidrug-resistant gram-negative bacterial infections. *Clin Infect Dis* 40:1333–41, 2005.

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

Kassamali Z, Rotschafer JC, Jones RN, et al. Polymyxins: wisdom does not always come with age. *Clin Infect Dis* 57:877-884, 2013.

Kunin CM. The Pharmacokinetics of Polymyxin B are Dependent on Binding and Release from Deep-Tissue Compartments. *Clin Infect Dis* 48:842-844, 2009.

Kwa AL, Abdelraouf K, Low JG, et al. Pharmacokinetics of polymyxin B in a patient with renal insufficiency: a case report. *Clin Infect Dis* 52:1280-1, 2011.

Li J, Nation RL. Old Polymyxins are Back: Is Resistance Close? *Clin Infect Disease* 43:663-64, 2006.

Menzies D, Minnaganti VR, Cunha BA. Polymyxin B. *Antibiotics for Clinicians* 4:33-40, 2000.

Pankey GA, Ashcraft DS. Detection of synergy using the combination of polymyxin B with either meropenem or rifampin against carbapenemase-producing *Klebsiella pneumoniae*. *Diagn Microbiol Infect Dis* 70:561-4, 2011.

Parchuri S, Mohan S, Young S, Cunha BA. Chronic ambulatory peritoneal dialysis associated peritonitis ESBP producing *Klebsiella pneumoniae* successfully treated with polymyxin B. *Heart & Lung* 34:360, 2005.

Segal-Maurer S, Mariano N, Qavi A, et al. Successful treatment of ceftazidime-resistant *Klebsiella pneumoniae* ventriculitis with intravenous meropenem and intraventricular polymyxin B: Case report and review. *Clin Infect Dis* 28:1134-8, 1999.

Singh M, Vilsaint M, Quale J. Dose-related clinical efficacy of polymyxin B in a critically ill patient. *Infect Dis in Clin Prac* 21:131-132, 2013.

Website: www.pdr.net

## Posaconazole (Noxafil)

**Drug Class:** Triazole antifungal.

**Usual Dose:** Prophylaxis of invasive *Aspergillus* and *Candida* infections in high-risk patients:<sup>†</sup>

200 mg (PO) q8h; oropharyngeal candidiasis: 100 mg (PO) q12h × 1 day, then 100 mg (PO) q24h × 13 days; for fluconazole- and/or itraconazole-resistant strains, use 400 mg (PO) q12h with duration of therapy based on underlying condition and clinical response.

**Pharmacokinetic Parameters:**

*Peak serum level:* 3 mcg/mL

*Bioavailability:* Absorption is ↑ 2-6-fold by food

*Excreted unchanged (urine):* < 1%

*Excreted unchanged (feces):* 66%

*Serum half-life (normal/ESRD):* 35/35 hrs

*Plasma protein binding:* 98.2%

*Volume of distribution (V<sub>d</sub>):* 1774 L

**Primary Mode of Elimination:** Hepatic

**Dosage Adjustments\***

CrCl < 50 mL/min	No change
Post-HD dose	Use IV with caution (PO levels variable)
Post-PD dose	Use IV with caution (PO levels variable)
CVWH/CVHD/ CVHDF dose	Use IV with caution (PO levels variable)
Moderate—severe hepatic insufficiency	No change

**Drug Interactions:** Posaconazole is metabolized by hepatic glucuronidation. Inducers (e.g., rifampin, phenytoin) may alter disposition. Amiodarone, rifabutin, phenytoin, cimetidine (↓ posaconazole levels 50%). Posaconazole inhibits hepatic CYP3A4 and can increase levels of drugs metabolized by this enzyme. QT prolonging drugs terfenadine, astemizole, cisapride, pimozone, halofantrine, quinidine (↑ interacting drug levels, ↑ risk of cardiac arrhythmias); ergot (↑ ergot levels); statins (↑ statin levels and risk of rhabdomyolysis); vinca alkaloids (↑ vinca alkaloid levels and risk of neurotoxicity); cyclosporine, tacrolimus, sirolimus, midazolam and other benzodiazepines metabolized by CYP3A4, calcium channel blockers metabolized by CYP3A4 (diltiazem, verapamil, nifedipine, nisoldipine); digoxin, sulfonylureas, ritonavir, indinavir (↑ interacting drug levels); rifampin, omperazole (↓ posaconazole levels).

<sup>†</sup>Usual dose assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

† (graft-vs-host disease, prolonged neutropenia from chemotherapy) ≥ 13 years of age.

**Adverse Effects:** Nausea, vomiting, diarrhea, headache, ↑ SGOT/SGPT; ↑ QTC. Avoid oral solution or capsules with PMH of CHF or CHF.

**Allergic Potential:** Low

**Safety in Pregnancy:** C

**Comments:** Do not crush/chew tablets. Available as a 40 mg/mL oral suspension. Take with a meal or 240 mL of a nutritional supplement. Blood levels not increased with higher dosages. **Highly active against *Candida albicans*, most non-*albicans Candida*, *Aspergillus*, *Cryptococcus*, *Histoplasmosis*, *Blastomycosis*, *Sporotrichosis*, *Paracoccidiomycosis*. Some activity against *Coccidiomycosis*, *Fusarium*, *C. glabrata*, and *Mucor*. No activity against *Pseudoallescheria* / *Scedosporium*.**

**Cerebrospinal Fluid Penetration:** > 10%

**Bile Penetration:** 76%

#### REFERENCES:

- Alffenaar JWC, van Assen S, van der Werf TS. Omeprazole significantly reduces posaconazole serum through level. *Clin Infect Dis* 48:839–840, 2009.
- Bruggemann RJM, Alffenaar JWC, Blijlevens NMA, et al. Clinical relevance of the pharmacokinetic interactions of azole antifungal drugs with coadministered agents. *Clin Infect Dis* 48:1441–1458, 2009.
- Cornely OA, Helfgott D, Langston A, et al. Pharmacokinetics of different dosing strategies of oral posaconazole in patients with compromised gastrointestinal function and who are at high risk for invasive fungal infection. *Antimicrob Agents Chemother* 56:2652–8, 2012.
- Cornely OA, Maertens J, Winston DJ, et al. Posaconazole vs. fluconazole or itraconazole prophylaxis in patients with neutropenia. *N Engl J Med* 356:348–59, 2007.
- Courtney R, Sansone A, Smith W, et al. Posaconazole pharmacokinetics, safety, and tolerability in subjects with varying degrees of chronic renal disease. *J Clin Pharmacol* 45:185–192, 2005.
- Dodds Ashley ES, Alexander BD. Posaconazole. *Drugs Today* 41:393–400, 2005.
- Ezzet F, Wexler D, Courtney R, et al. Oral bioavailability of posaconazole in fasted healthy subjects-comparison between three regimens and basis for clinical dosage recommendations. *Clin Pharmacokinet* 44:211–220, 2005.
- Felton TW, Baxter C, Moore CB, et al. Efficacy and safety of posaconazole for chronic pulmonary aspergillosis. *Clin Infect Dis* 51:1383–1391, 2010.
- Girimenia C, Frustaci AM, Gentile G, et al. Posaconazole prophylaxis during front-line chemotherapy of acute myeloid leukemia: a single-center real-life experience. *Haematologica* 97:560–7, 2012.
- Green MR, Woolley JE. Optimising absorption of posaconazole. *Mycoses* 54:775–779, 2011.
- Greenberg RN, Mullane K, van Burik JA, et al. Posaconazole as salvage therapy for zygomycosis. *Antimicrob Agents Chemother* 50:126–33, 2006.
- Gubbins PO, Krishna G, Sansone-Parsons A, et al. Pharmacokinetics and safety of oral posaconazole in neutropenic stem cell transplant recipients. *Antimicrob Agents Chemother* 50:1993–9, 2006.
- Herbrecht R, Nivoix Y, Fohrer C, et al. Management of systemic fungal infections: alternatives to itraconazole. *J Antimicrob Chemother* 56:139–148, 2005.
- Hoening M, Raggam RB, Salzer HJ, et al. Posaconazole plasma concentrations and invasive mould infections in patients with haematological malignancies. *Int J Antimicrob Agents* 39:510–3, 2012.
- Hohmann C, Kang EM, Jancel T: Rifampin and posaconazole coadministration leads to decreased serum posaconazole concentrations. *Clin Infect Dis* 50:939–940, 2010.
- Howard SJ, Lestner JM, Sharp A, et al. Pharmacokinetics and pharmacodynamics of posaconazole for invasive pulmonary aspergillosis: clinical implications for antifungal therapy. *J Infect Dis* 203:1324–1332, 2011.
- Karthaum M. Prophylaxis and treatment of invasive aspergillosis and voriconazole, posaconazole, and caspofungin: review of the literature. *Eur J Med Res* 16:145–152, 2011.
- Keating GM, Posaconazole. *Drugs* 65:1553–69, 2005.
- Kim MM, Vikram HR, Kusne S, et al. Treatment of refractory coccidioidomycosis with voriconazole or posaconazole. *Clin Infect Dis*, 53:1060–6, 2011.
- Kontoyannis DP, Lewis RE. Posaconazole prophylaxis in hematologic cancer. *N Engl J Med* 24:2214–2215, 2007.
- Lipp HP. Posaconazole: clinical pharmacokinetics and drug interactions. *Mycoses* 54:32–38, 2011.

"Usual dose" assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.



Miceli MH, Diaz JA, Lee SA. Emerging opportunistic yeast infections. *Lancet Infectious Diseases* 11: 142–151, 2011.

Mehta AK, Langston AA. Use of posaconazole in the treatment of invasive fungal infections. *Expert Rec Hematol* 2:619–630, 2009.

Pound MW, Townsend ML, Dimondi V, et al. Overview of treatment options for invasive fungal infections. *Med Mycol* 49:561–580, 2011.

Raad II, Graybill JR, Bustamante AB, et al. Safety of long-term oral posaconazole use in the treatment of refractory invasive fungal infections. *Clin Infect Dis* 42:1726–34, 2006.

Raad II, Hachem RY, Herbrecht R, et al. Posaconazole as salvage treatment for invasive fusariosis in patients with underlying hematologic malignancy and other conditions. *Clin Infect Dis* 42:1398–403, 2006.

Restrepo A, Tobon A, Clark B, et al. Salvage treatment of histoplasmosis with posaconazole. *Salvage treatment of histoplasmosis with posaconazole. J Infect* 54:319–327, 2007.

Ullmann AJ, Cornely OA, Burchardt A, et al. Pharmacokinetics, safety, and efficacy of posaconazole in patients with persistent febrile neutropenia or refractory invasive fungal infection. *Antimicrob Agents Chemother* 50:658–66, 2006.

Ullmann AJ, Lipton JH, Vesole DH, et al. Posaconazole or fluconazole for prophylaxis in severe graft-versus-host disease. *N Engl J Med* 356:335–47, 2007.

Walsh TJ, Raad I, Patterson TF, et al. Treatment of invasive Aspergillosis with posaconazole in patients who are refractory to or intolerant of conventional therapy: an externally controlled trial. *Clin Infect Dis* 44:2–12, 2007.

Website: [www.pdr.net](http://www.pdr.net)

## Primaquine

**Drug Class:** Antimalarial/Anti-PCP.

**Usual Dose:** Antimalarial dose: 15 mg (base 26.3 mg) (PO) q24h. **PCP dose:** 15–30 mg (PO) q24h plus clindamycin 600 mg (IV) or 300 mg (PO) q6h.

### Pharmacokinetic Parameters:

*Peak serum level:* 30–100 mcg/ml

*Bioavailability:* 90%

*Excreted unchanged (urine):* 3.6%

*Serum half-life (normal/ESRD):* 3.7–9.6/3.7–9.6 hrs

*Plasma protein binding:* 75%

*Volume of distribution ( $V_d$ ):* No data

**Primary Mode of Elimination:** Hepatic  
**Dosage Adjustments\***

CrCl < 80 mL/min	No change
Post-HD/Post-PD dose	None
CVWH/CWHD/CWHDF dose	No change
Hepatic insufficiency	No change

**Drug Interactions:** Avoid in patients receiving bone marrow suppressive drugs (↑ risk of agranulocytosis), alcohol (↑ GI side effects), quinacrine (↑ primaquine levels/toxicity).

**Adverse Effects:** Nausea/vomiting, headache, abdominal pain, leukopenia (dose dependent), agranulocytosis, pruritus. Hemolytic anemia in G6PD deficiency; methemoglobinemia in NADH reductase deficiency.

Allergic Potential: Low

Safety in Pregnancy: D

**Comments:** Take with food to ↓ GI side effects. Use with caution in G6PD deficiency. Use with caution in NADH reductase deficiency.

**Cerebrospinal Fluid Penetration:** No data

### REFERENCES:

- Baird JK. Effectiveness of antimalarial drugs. *N Engl J Med* 352:1565–77, 2005.
- Chen LH, Keystone JS. New strategies for the prevention of malaria in travelers. *Infect Dis Clin North Am* 19:185–210, 2005.
- Korraa H, Saadeh C. Options in the management of pneumonia caused by *Pneumocystis carinii* in patients with acquired immune deficiency syndrome and

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

intolerance to trimethoprim/sulfamethoxazole. *South Med J* 89:272–7, 1996.

Safrin S, Finkelstein DM, Feinberg J, et al. Comparison of three regimens for treatment of mild to moderate *Pneumocystis carinii* pneumonia in patients with AIDS. A double-blind randomized trial of oral TMP-SMX, dapsone-TMP & clinda-primaquine. *Ann Intern Med* 124:792–802, 1996.

Warren E, George S, You J, Kazanjian P. Advances in the treatment and prophylaxis of *Pneumocystis carinii* pneumonia. *Pharmacotherapy* 17:900–16, 1997.

Website: www.pdr.net

## Pyrazinamide (PZA)

**Drug Class:** TB drug.

**Usual Dose:** 25 mg/kg (PO) q24h (max. 2 gm) (see comments).

**Pharmacokinetic Parameters:**

*Peak serum level:* 30–50 mcg/ml

*Bioavailability:* 90%

*Excreted unchanged (urine):* 10%

*Serum half-life (normal/ESRD):* 9/26 hrs

*Plasma protein binding:* 10%

*Volume of distribution ( $V_d$ ):* 0.9 L/kg

**Primary Mode of Elimination:** Hepatic

**Dosage Adjustments\***

CrCl < 30 mL/min	25 mg/kg (PO) 3x/week
Post-HD dose	25 mg/kg (PO) or 1 gm (PO)
Post-PD dose	None
CVVH/CVVHD/ CVVHDF dose	25 mg/kg (PO) 3x/week
Moderate hepatic insufficiency	Use with caution
Severe hepatic insufficiency	Avoid

**Drug Interactions:** INH, rifabutin, rifampin (may ↑ risk of hepatotoxicity).

**Adverse Effects:** Drug fever/rash, malaise, nausea, vomiting, anorexia, ↑ SGOT/SGPT, ↑ uric acid, sideroblastic anemia.

**Allergic Potential:** Low

**Safety in Pregnancy:** C

**Comments:** Avoid in patients with gout (may precipitate acute attacks). **TB D.O.T. dose:** 4 gm (PO) 2x/week or 3 gm (PO) 3x/week.

Meningeal dose = usual dose.

**Cerebrospinal Fluid Penetration:** 100%

### REFERENCES:

- Ahn C, Oh KH, Kim K, et al. Effect of peritoneal dialysis on plasma and peritoneal fluid concentrations of isoniazid, pyrazinamide, and rifampin. *Perit Dial Int* 23:362–7, 2003.
- Davidson PT, Le HQ. Drug treatment of tuberculosis 1992. *Drugs* 43:651–73, 1992.
- [No authors listed.] Drugs for tuberculosis. *Med Lett Drugs Ther* 35:99–101, 1993.
- Furin J, Nardell EA. Multidrug-resistant tuberculosis: An update on the best regimens. *J Respir Dis* 27:172–82, 2006.
- Iseman MD. Treatment of multidrug-resistant tuberculosis. *N Engl J Med* 329:784–91, 1993.
- Ijaz K, McElroy PD, Navin TR. Short course rifampin and pyrazinamide compared with isoniazid for latent tuberculosis infection: a cost-effectiveness analysis based on a multicenter clinical trial. *Clin Infect Dis* 39:289, 2004.
- McIllerson H, Wash P, Burger A, et al. Determinants of rifampin, isoniazid, pyrazinamide and ethambutol pharmacokinetics in a cohort of tuberculosis patients. *Antimicrob Agents Chemother* 50:1170–77, 2006.
- Schlossberg D. An approach to drug-induced liver injury during tuberculosis therapy. *Infect Dis Clin Pract* 18:107–108, 2010.
- Van Scoy RE, Wilkowske CJ. Antituberculous agents. *Mayo Clin Proc* 67:179–87, 1992.
- Website: www.pdr.net

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

## Pyrimethamine (Daraprim)

**Drug Class:** Antiparasitic.

**Usual Dose:** 75 mg (PO) q24h (see comments).

**Pharmacokinetic Parameters:**

*Peak serum level:* 0.4 mcg/ml

*Bioavailability:* 90%

*Excreted unchanged (urine):* 25%

*Serum half-life (normal/ESRD):* 96/96 hrs

*Plasma protein binding:* 87%

*Volume of distribution ( $V_d$ ):* 2.5 L/kg

**Primary Mode of Elimination:** Hepatic

**Dosage Adjustments\***

CrCl < 10 mL/min	Use with caution
Post-HD dose	None
Post-PD dose	25 mg (PO)
CWH/CVWHD/ CWHDF dose	No change
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	Use with caution

**Drug Interactions:** Folic acid (↓ pyrimethamine effect); lorazepam (↑ risk of hepatotoxicity); sulfamethoxazole, trimethoprim, TMP-SMX (↑ risk of thrombocytopenia, anemia, leukopenia).

**Adverse Effects:** Megaloblastic anemia, leukopenia, thrombocytopenia, ataxia, tremors, seizures.

**Allergic Potential:** Low

**Safety in Pregnancy:** C

**Comments:** Antacids decrease absorption.

**Toxoplasmosis dose:** 200 mg (PO) × 1 dose, then 50–75 mg/kg (PO) q24h (with folic acid 20 mg PO q24h plus either sulfadiazine or clindamycin).

**Cerebrospinal Fluid Penetration:** 10–25%

### REFERENCES:

- [No authors listed.] Drugs for Parasitic Infections. Med Lett Drugs Ther 40: 1–12, 2000.
- Lemnge MM, Ali AS, Malecela EK, et al. Therapeutic efficacy of sulfadoxine pyrimethamine and amodiaquine among children with uncomplicated Plasmodium falciparum malaria in Zanzibar, Tanzania. Am. J. Trop. Med Hyg 73:681–5, 2005.
- Podzamczar D, Salazar A, Jimenez J, et al. Intermittent trimethoprim-sulfamethoxazole compared with dapsone-pyrimethamine for the simultaneous primary prophylaxis of Pneumocystis pneumonia and toxoplasmosis in patients infected with HIV. Ann Intern Med 122:755–61, 1995.
- Porter SB, Sande MA. Toxoplasmosis of the central nervous system in the acquired-immunodeficiency syndrome. N Engl J Med 327:1643–8, 1992.
- Website: www.pdr.net

## Quinine sulfate

**Drug Class:** Antimalarial.

**Usual Dose:** 650 mg (PO) q8h (see comments).

**Pharmacokinetic Parameters:**

*Peak serum level:* 3.8 mcg/ml

*Bioavailability:* 80%

*Excreted unchanged (urine):* 5%

*Serum half-life (normal/ESRD):* 7/14 hrs

*Plasma protein binding:* 95%

*Volume of distribution ( $V_d$ ):* 3 L/kg

**Primary Mode of Elimination:** Renal/hepatic

**Dosage Adjustments\***

CrCl 10–50 mL/min	650 mg (PO) q12h
CrCl < 10 mL/min	650 mg (PO) q24h
Post-HD dose	None
Post-PD dose	650 mg (PO)
CWH/CVWHD/ CWHDF dose	650 mg (PO) q24h

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVW = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

Mild-moderate hepatic insufficiency	No change
Severe hepatic insufficiency	325 mg (PO) q12h; use caution

**Drug Interactions:** Amiodarone, aluminum-based antacids ( $\downarrow$  quinidine absorption); astemizole, cisapride, terfenadine ( $\uparrow$  interacting drug levels, torsade de pointes; avoid); cimetidine, ritonavir ( $\uparrow$  quinidine toxicity: headache, deafness, blindness, tachycardia); cyclosporine ( $\downarrow$  cyclosporine levels); digoxin ( $\uparrow$  digoxin levels); dofetilide, flecainide (arrhythmias); mefloquine (seizures, may  $\uparrow$  QT interval, torsade de pointes, cardiac arrest,  $\downarrow$  mefloquine efficacy); metformin ( $\uparrow$  risk of lactic acidosis); pancuronium, succinylcholine, tubocurarine (neuromuscular blockade); warfarin ( $\uparrow$  INR).

**Adverse Effects:** Drug fever/rash,  $\uparrow$  QT<sub>c</sub> interval, arrhythmias, drug-induced SLE, lightheadedness, diarrhea, abdominal discomfort, nausea, vomiting, cinchonism with chronic use. Avoid in patients with G6PD deficiency.

**Allergic Potential:** High

**Safety in Pregnancy:** C

**Comments:** **PO malaria dose:** 650 mg (PO) q8h (plus doxycycline 100 mg PO q12h)  $\times$  3–7 days). **IV malaria dose:** quinine hydrochloride 600 mg (IV) q8h  $\times$  3–7 days, or quinidine gluconate 10 mg/kg (IV)  $\times$  1 dose (infuse over 1–2 hours) then 0.02 mg/kg/min (IV)  $\times$  72 hours or until parasitemia  $<$  1%.

**Cerebrospinal Fluid Penetration:** 2–5%

#### REFERENCES:

Corpelet C, Vacher P, Coudor F, et al. Role of quinine in life-threatening *Babesia divergens* infection successfully treated with clindamycin. *Eur J Clin Microbiol Infect Dis* 24:74–75, 2005.

Croft AM, Herxheimer A. Tolerability of antimalaria drugs. *Clin Infect Dis* 34:1278; discussion 1278–9, 2002.

Drugs for Parasitic Infections. Med Letter. March, 2000.

Panisko DM, Keystone JS. Treatment of malaria. *Drugs* 39:160–89, 1990.

Wylter DJ. Malaria: Overview and update. *Clin Infect Dis* 16:449–56, 1993.

Website: www.pdr.net

## Quinupristin/dalfopristin (Synercid)

**Drug Class:** Streptogramin.

**Usual Dose:** 7.5 mg/kg (IV) q8h.

**Spectrum:** (see **Susceptibility Profiles** pp. 198–202).

**Resistance Potential:** Low

**Pharmacokinetic Parameters:**

*Peak serum level:* 3.2/8 mcg/ml

*Bioavailability:* Not applicable

*Excreted unchanged:* 20% (urine); 80% (feces) Serum

*half-life (normal/ESRD):* [3.1/1]/[3.1/1] hrs

*Plasma protein binding:* 55/15%

*Volume of distribution (V<sub>d</sub>):* 0.45/0.24 L/kg

**Primary Mode of Elimination:** Hepatic

**Dosage Adjustments\***

CrCl $<$ 10 mL/min	No change
Post-HD dose	None
Post-PD dose	None
CVWH/CVWHD/CVWHDf dose	No change
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	Use with caution

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose  $\times$  1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl  $<$  10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

**Drug Interactions:** Amiodarone, amlodipine (↑ amlodipine toxicity); astemizole, cisapride (may ↑ QT interval, torsades de pointes); carbamazepine (↑ carbamazepine toxicity; ataxia, nystagmus, diplopia, headache, seizures); cyclosporine, delavirdine, indinavir, nevirapine (↑ interacting drug levels); diazepam, midazolam (↑ interacting drug effect); diltiazem, felodipine, isradipine (↑ interacting drug toxicity: dizziness, hypotension, headache, flushing); disopyramide (↑ disopyramide toxicity: arrhythmias, hypotension, syncope); docetaxel (↑ interacting drug toxicity: neutropenia, anemia, neuropathy); lidocaine (↑ lidocaine toxicity: neurotoxicity, arrhythmias, seizures); methylprednisolone (↑ methylprednisolone toxicity: myopathy, diabetes mellitus, Cushing's syndrome); nicardipine, nifedipine, nimodipine (↑ interacting drug toxicity: dizziness, hypotension, flushing, headache); statins (↑ risk of rhabdomyolysis).

**Adverse Effects:** Pain, inflammation, and swelling at infusion site (dose related), severe/prolonged myalgias, hyperbilirubinemia. Hepatic insufficiency increases concentration (AUC) of metabolites by 180%/50%.

**Allergic Potential:** Low

**Safety in Pregnancy:** B

**Comments:** Administer in D<sub>5</sub>W or sterile water, not in saline. Requires central IV line for administration. **Not effective against *E. faecalis* (VSE). May be useful for daptomycin resistant MSSA/MRSA infections.**

**Cerebrospinal Fluid Penetration:** < 10% (also see **Antibiotic Pearls & Pitfalls** p. 518).

#### REFERENCES:

- Blondeau JM, Sanche Se. Quinupristin/dalfopristin. Expert Opin Pharmacother 3:1341–64, 2002.  
 Bouza E. New therapeutic choices for infections caused by methicillin-resistant *Staphylococcus aureus*. Clin Microbiol Infect Dis 15:44–52, 2009.

- Bryson HM, Spencer CM. Quinupristin/dalfopristin. Drugs 52:406–15, 1996.  
 Chant C, Ryback MH. Quinupristin/dalfopristin (RP 59500): A new streptogramin antibiotic. Ann Pharmacother 29:1022–7, 1995.  
 Goff DA, Sierawski SJ. Clinical experience of quinupristin-dalfopristin for the treatment of antimicrobial-resistant gram-positive infections. Pharmacotherapy 22:748–58, 2002.  
 Griswold MW, Lomaestro BM, Briceland LL. Quinupristin-dalfopristin (RP 59500): An injectable streptogramin combination. Am J Health Syst Pharm 53:2045–53, 1996.  
 Kim MK, Nicolau DP, Nightingale CH, et al. Quinupristin/dalfopristin: A treatment option for vancomycin-resistant enterococci. Conn Med 64:209–12, 2000.  
 Klastersky J. Role of quinupristin/dalfopristin in the treatment of Gram-positive nosocomial infections in haematological or oncological patients. Cancer Treat Rev 29:431–40, 2003.  
 Nadler H, Dowzicky MJ, Feger C, et al. Quinupristin/dalfopristin: A novel selective-spectrum antibiotic for the treatment of multi-resistant and other gram-positive pathogens. Clin Microbiol Newslett 21:103–12, 1999.  
 Nailor MD, Sobel JD. Antibiotics for gram-positive bacterial infection: vancomycin, teicoplanin, quinupristin/dalfopristin, oxazolidinones, daptomycin, telavancin, and ceftaroline fosamil. Med Clin North Am 95:723–742, 2011.  
 Scotton PG, Rigoli R, Vaglia A. Combination of quinupristin/dalfopristin and glycopeptide in severe methicillin-resistant staphylococcal infections failing previous glycopeptide regimen. Infection 30:161–3, 2002.  
 Website: www.pdr.net

## Raltegravir (Isentress) RAL

**Drug Class:** HIV integrase inhibitor.

**Usual Dose:** 400 mg (PO) q12h.

**Pharmacokinetic Parameters:**

*Peak serum level:* 6.5 μM

*Bioavailability:* ~ 32% (20–43%)

*Excreted unchanged:* 51% (feces); 9% (urine)

*Serum half-life (normal/ESRD):* 9–12 hrs/no data

*Plasma protein binding:* 83%

*Volume of distribution (V<sub>d</sub>):* not studied

"Usual dose" assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

**Primary Mode of Elimination:** Fecal/renal

**Dosage Adjustments\***

CrCl < 10 mL/min	No change
Post-HD dose	None
Post-PD dose	None
CVWH/CVWHD/ CWHDF dose	No change
Mild/moderate hepatic insufficiency	No change
Severe hepatic insufficiency	Use with caution

**Antiretroviral Dosage Adjustments**

Atazanavir	No change
Atazanavir/ritonavir	No change
Efavirenz	No change
Rifampin	raltegravir 800 mg q12h
Ritonavir	No change
Tenofovir	No change
Tipranavir/ritonavir	No change

**Drug Interactions:** Rifampin (↓ raltegravir levels, use with caution). In-vitro, raltegravir does not inhibit CYP1A2, CYP2B6, CYP2C8, CYP2C9, CYP2C19, CYP2D6 or CYP3A and does not induce CYP3A4. In addition, raltegravir does not inhibit P-glycoprotein-mediated transport. Raltegravir is therefore not expected to affect the pharmacokinetics of drugs that are substrates of these enzymes or P-glycoprotein (e.g., protease inhibitors, NNRTIs, methadone, opioid analgesics, statins, azole antifungals, proton pump inhibitors, oral contraceptives, anti-erectile dysfunction agents).

**Adverse Effects:** Nausea, headache, diarrhea, pyrexia, rash, stevens-johnson Syndrome, TEN.

**Allergic Potential:** Low

**Safety in Pregnancy:** C

**Comments:** May be taken with or without food. CPK elevations, myopathy and rhabdomyolysis have been reported—use with caution in patients at increased risk for myopathy or rhabdomyolysis, such as those receiving concomitant medications e.g., statins. Raltegravir is indicated for treatment-naïve/ experienced adult patients who have evidence of viral replication and HIV-1 strains resistant to multiple antiretroviral agents. Slight increase in cancer in severely immunocompromised hosts.

**Cerebrospinal Fluid Penetration:** No data

**REFERENCES:**

- Carey D, Pett SL, Bloch M, et al. A randomized study of pharmacokinetics, efficacy, and safety of 2 raltegravir plus atazanavir strategies in ART-treated adults. *J Acquir Immune Defic Syndr* 60:143–9, 2012.
- Hicks C, Gulick RM. Raltegravir: The First HIV Type 1 Integrase Inhibitor. *Clin Infect Dis* 48:931–940, 2009.
- Iwamoto M, Wenning LA, Petry AS, et al. Safety, tolerability, and pharmacokinetics of raltegravir after single and multiple doses in healthy subjects. *Clin Pharmacol Ther* 83: 293–299, 2008.
- Palmisano L. Role of integrase inhibitors in the treatment of HIV disease. *Expert Rev Anti Infect Ther* 5:67–75, 2007.
- Thompson MA, Aberg JA, Cahn P, et al. Antiretroviral treatment of adult HIV infection: 2012 recommendations of the International Antiviral Society — USA panel. *JAMA* 308:387–402, 2012.
- Website: [www.pdr.net](http://www.pdr.net)

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

## Ribavirin (Rebetol) (Copegus)

**Drug Class:** RSV, HCV Antiviral; (see comments).

**Usual Dose:** 600 mg (PO) q12h (see comments).

**Pharmacokinetic Parameters:**

Peak serum level: 0.07–0.28 mcg/ml

Bioavailability: 64%

Excreted unchanged (urine): 40%

Serum half-life (normal/ESRD): 120 hrs/no data

Plasma protein binding: 0%

Volume of distribution ( $V_d$ ): 10 L/kg

**Primary Mode of Elimination:** Hepatic

**Dosage Adjustments\***

CrCl 50–80 mL/min	No change
CrCl < 50 mL/min	Avoid
Post-HD dose	Avoid
Post-PD dose	Avoid
CWH/CVWHD/CVWHDf dose	Avoid
Mild-moderate hepatic insufficiency	No change
Severe hepatic insufficiency	Avoid

**Drug Interactions:** Ribavirin may antagonize the *in vitro* antiviral activity of stavudine and zidovudine against HIV. Pegylated interferon (Pegasys) in HIV patients ( $\downarrow$  CD<sub>4</sub> counts).

**Adverse Effects:** Hemolytic anemia. The anemia associated with ribavirin therapy may result in worsening of cardiac disease that has led to fatal and non-fatal myocardial infarctions. Patients with a history of significant cardiac disease should not be treated with ribavirin. If hemoglobin level falls < 10 g/dL reduce ribavirin dose to 600 mg daily. If hemoglobin falls < 8.5 g/dL discontinue

ribavirin. When combined with interferon adverse effects includes severe depression and suicidal ideation, bone marrow suppression, autoimmune/infectious disorders, pulmonary dysfunction, pancreatitis, and diabetes. Drug fever/rash, nausea, vomiting, GI upset, leukopenia, hyperbilirubinemia, hemolytic anemia,  $\uparrow$  uric acid.

**Allergic Potential:** Low

**Safety in Pregnancy:** X (avoid pregnancy during therapy and for 6 months after completion of therapy in both female patients and male partners of taking ribavirin)

**Comments:** **Nebulizer dose for RSV:** 20 mg/ml aerosolized over 12 hours administered once daily  $\times$  3–7 days. Also as activity against Lassa fever and HPS. **Transplant Dose:**

**Loading Dose:** 30 mg/kg (IV) (not to exceed 2g), then 16 mg/kg (IV) q6h  $\times$  4 days (not to exceed 1g); **Maintenance Dose:** 8 mg/kg (IV) (not to exceed 500 mg)  $\times$  3–6 days. **Highly active against HCV and RSV. Some activity against, HEV, adenoviruses and arboviral hemorrhagic fever viruses. No activity against EBV, CMV, HBV, HDV.** For Crimean-Congo hemorrhagic fever give ribavirin initial **Loading Dose** 30 mg/kg (PO)  $\times$  1 then **Maintenance Dose:** 15 mg/kg (PO) q6h  $\times$  4 days, then give 7.5 mg/kg (PO) q8h  $\times$  6 days.

**Cerebrospinal Fluid Penetration:** 70%

### REFERENCES:

- Khalid O, Bacon BR. Management of the treatment-experienced patient infected with Hepatitis C virus genotype 1: options and considerations. *Clin Liver Dis* 15:573–583, 2011.
- Razonable RR. Antiviral drugs for viruses other than human immunodeficiency virus. *Mayo Clin Proc* 86:1009–1026, 2011.
- Vanpouille C, Lisco A, Introni A, et al. Exploiting the anti-HIV-1 activity of acyclovir: suppression of primary and drug-resistant HIV isolates and potentiation of the activity by ribavirin. *Antimicrob Agents Chemother* 56:2604–2611, 2012.
- Website: [www.pdr.net](http://www.pdr.net)

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose  $\times$  1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVW = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

## Rifabutin (Mycobutin)

**Drug Class:** MAI drug.

**Usual Dose:** 5 mg/kg or 300 mg (PO) q24h (see comments).

**Pharmacokinetic Parameters:**

Peak serum level: 0.38 mcg/ml

Bioavailability: 20–50%

Excreted unchanged (urine): 10%

Serum half-life (normal/ESRD): 45/45 hrs

Plasma protein binding: 85%

Volume of distribution ( $V_d$ ): 9.3 L/kg

**Primary Mode of Elimination:** Hepatic

**Dosage Adjustments\***

CrCl < 30 mL/min	150 mcg (PO) q24h
CrCl < 10 mL/min	No change
Post-HD dose	None
Post-PD dose	None
CVWH/CVWHD/CVWHDf dose	None
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	No change

**Drug Interactions:** Atovaquone, amprenavir, indinavir, nelfinavir, ritonavir, clarithromycin, erythromycin, telithromycin, fluconazole, itraconazole, ketoconazole (↓ interacting drug levels, ↑ rifabutin levels); beta-blockers, clofibrate, cyclosporine, enalapril, oral contraceptives, quinidine, sulfonyleureas, tocainide, warfarin (↓ interacting drug effect); corticosteroids (↑ corticosteroid requirement); delavirdine (↓ delavirdine levels, ↑ rifabutin levels; avoid); digoxin, phenytoin, propafenone, theophylline, zidovudine (↓ interacting drug

levels); methadone (↓ methadone levels, withdrawal); mexiletine (↑ mexiletine clearance); protease inhibitors (↓ protease inhibitor levels, ↑ rifabutin levels; caution).

**Adverse Effects:** Headache, nausea, vomiting. ↑ SGOT/SGPT, leukopenia, anemia, thrombocytopenia, drug fever/rash. Brown/orange discoloration of body fluids.

**Allergic Potential:** High

**Safety in Pregnancy:** B

**Comments:** Avoid in leukopenic patients with WBC ≤ 1000 cells/mm<sup>3</sup>. Always use as part of a multi-drug regimen, never as monotherapy.

**Rifabutin doses when co-administered with antiretrovirals:** 450 mg (PO) q24h with EFV; 150 mg (PO) q24h with AVP, IDV, NFV, FPN; 150 mg (PO) q48h with RTV/LPV combination

Meningeal dose = usual dose.

**Cerebrospinal Fluid Penetration:** 50–70%

### REFERENCES:

- Benson CA, Williams PL, Cohn DL, and the ACTG 196/CPCRA 009 Study Team. Clarithromycin or rifabutin alone or in combination for primary prophylaxis of Mycobacterium avium complex disease in patients with AIDS: A randomized, double-blinded, placebo-controlled trial. *J Infect Dis* 181:1289–97, 2000.
- Centers for Disease Control and Prevention. Notice to readers: Updated guidelines for the use of rifabutin or rifampin for the treatment and prevention of tuberculosis among HIV-infected patients taking protease inhibitors or nonnucleoside reverse transcriptase inhibitors. *MMWR* 49:183–189, 2000.
- Choong K, Looke D. Cross-reaction to rifabutin after rifampicin induced flu-like syndrome and thrombocytopenia. *Scand J Infect Dis* 43:238, 2011.
- Finch CK, Chrisman CR, Baciewicz AM, et al. Rifampin and rifabutin drug Interactions: an update. *Arch Intern Med* 162:985–92, 2002.

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.



Horne DJ, Spitters C, Narita M. Experience with rifabutin replacing rifampin in the treatment of tuberculosis. *Int J Tuberc Lung Dis* 15:1485–9, 2011.

Hoy J, Mijch A, Sandland M, et al. Quadruple-drug therapy for *Mycobacterium avium*-intracellular bacteremia in AIDS patients. *J Infect Dis* 161:801–5, 1990.

Nightingale SD, Cameron DW, Gordin FM, et al. Two controlled trials of rifabutin prophylaxis against *Mycobacterium avium* complex infection in AIDS. *N Engl J Med* 329:828–33, 1993.

Nishizawa T, Suzuki H, Matsuzaki J, et al. *Helicobacter pylori* resistance to rifabutin in the last 7 years. *Antimicrob Agents Chemother* 55:5374–5, 2011.

Sax PE, Cohen CJ, Kuritzkes DR. *HIV Essentials*, 5th edition, Jones & Bartlett, Sudbury, MA, 2011. pp. 123–125.

Zhang J, Shu L, Stonier M, et al. Determination of rifabutin dosing regimen when administered in combination with ritonavir-boosted atazanavir. *J Antimicrob Chemother* 66:2075–82, 2011.

Website: [www.pdr.net](http://www.pdr.net)

## Rifampin (Rifadin, Rimactane)

**Drug Class:** Antibiotic/TB drug.

**Usual Dose:** 600 mg (PO) q24h (TB dose); 300 mg (PO) q12h (antibiotic dose)

**Spectrum:** (see **Susceptibility Profiles** pp. 186–190).

**Resistance Potential:** High (aerobic GNBs with non-TB monotherapy)

Pharmacokinetic Parameters:

Peak serum level: 7 mcg/ml

Bioavailability: 95%

Excreted unchanged (urine): 15%

Serum half-life (normal/ESRD): 3.5/11 hrs

Plasma protein binding: 80%

Volume of distribution ( $V_d$ ): 0.93 L/kg

**Primary Mode of Elimination:** Hepatic

### Dosage Adjustments\*

CrCl < 10 mL/min	No change
Post-HD dose	None
Post-PD dose	None
CVH/CVHD/CVHDF dose	No change
Moderate hepatic insufficiency	No change; use caution
Severe hepatic insufficiency	Avoid

**Drug Interactions:** Amprenavir, indinavir, nelfinavir (↑ rifampin levels); beta-blockers, clofibrate, cyclosporine, oral contraceptives, quinidine, sulfonyleureas, tocainamide, warfarin (↓ interacting drug effect); caspofungin (↓ caspofungin levels, may ↓ caspofungin effect); clarithromycin, ketoconazole (↑ rifampin levels, ↓ interacting drug levels); corticosteroids (↑ corticosteroid requirement); delavirdine (↑ rifampin levels, ↓ delavirdine levels; avoid); disopyramide, itraconazole, phenytoin, propafenone, theophylline, methadone, nelfinavir, ritonavir, tacrolimus, drugs whose metabolism is induced by rifampin, e.g., ACE inhibitors, dapsone, diazepam, digoxin, diltiazem, doxycycline, fluconazole, fluvastatin, haloperidol, nifedipine, progestins, triazolam, tricyclics, zidovudine (↓ interacting drug levels); fluconazole, rifampin (↓ posaconazole levels) TMP-SMX (↑ rifampin levels); INH (INH converted into toxic hydrazine); mexiletine (↑ mexiletine clearance); nevirapine (↓ nevirapine levels; avoid).

**Adverse Effects:** Dizziness, Flu-like symptoms, ↑ SGOT/SGPT (rarely severe hepatitis), drug fever/rash, thrombocytopenia. Red/orange discoloration of body secretions.

**Allergic Potential:** Moderate

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

**Safety in Pregnancy:** C

**Comments:** Potent CYP 3A4 inducer. Contraindicated in HIV. For anti-TB prophylaxis/therapy, monitor potential hepatotoxicity with serial SGOT/SGPTs weekly  $\times$  3, then monthly  $\times$  3. Take 1 hour before or 2 hours after meals. **TB D.O.T. dose:** 10 mg/kg or 600 mg (PO) 2–3 $\times$ /week. **Do not use as monotherapy for *S. aureus*. Of no proven clinical advantage when combined with another anti-MSSA/MRSA antibiotic. With another anti-staphylococcal drug dose:** 300 mg (PO) q12h. 300 mg (PO) q8h may be used for *S. aureus* PVE or *Brucella* SBE.

**Nasal carriage dose:** 600 mg (PO) q5h  $\times$  72 hours.

Meningeal dose = usual dose.

**Cerebrospinal Fluid Penetration:**

Non-Inflamed meninges = 50%

Inflamed meninges = 50%

**Bile Penetration:** 7000%

**REFERENCES:**

Cascio A, Scarlata F, Giordano S, et al. Treatment of human brucellosis with rifampin plus minocycline. *J Chemother* 15:248–52, 2003.

Castahneira D, Rifampin. *Antibiotics for Clinicians* 6:89–100, 2001.

Curry SR, Marsh JW, Shutt KA, et al. High frequency of rifampin resistance identified in an epidemic *clostridium difficile* clone from a large teaching hospital. *Clin Infect Dis* 48:425–430, 2009.

Czekaj J, Dinh A, Moldovan A, et al. Efficacy of a combined oral clindamycin-rifampicin regimen for therapy of staphylococcal osteoarticular infections. *Scand J Infect Dis* 43:962–967, 2011.

Davidson PT, Le HQ. Drug treatment of tuberculosis 1992. *Drugs* 43:651–73, 1992.

Durante-Mangoni E, Signoriello G, Andini R, et al. Colistin and rifampicin compared with colistin alone for the treatment of serious infections due to extensively drug-resistant *A. cinetobacter baumannii*. *Clin Infect Dis* 57:349–358, 2013.

Eisen D, Denholm JS. Recommendations for rifampicin therapy of staphylococcal infection in IDSA prosthetic joint infection guidelines are not supported by available literature. *Clin Infect Dis* 57:159–2013.

Ena J, Valls V. Short-course therapy with rifampin plus isoniazid compared with standard therapy with isoniazid, for latent tuberculosis infection: a meta-analysis. *Clin Infect Dis* 40:670–6, 2005.

Euga G, Murillo O, Fernandez-Sabe N, et al. Treatment of chronic staphylococcal osteomyelitis: intravenous or oral? how long? *Antimicrob Agents Chemother* 53:2672–2676, 2009.

Gauthier TP. Rifampicin plus colistin in the era of extensively drug-resistant *acinetobacter baumannii* infections. *Clin Infect Dis* 57:359–362, 2013.

Gomez J, Canovas E, Banos V, et al. Linezolid plus rifampin as a salvage therapy in prosthetic joint infections treated without removing the implant. *Antimicrob Agents Chemother* 55:4308–4310, 2011.

Hohmann C, Kang EM, Jancel T. Rifampin and posaconazole coadministration leads to decreased serum posaconazole concentrations. *Clin Infect Dis* 50:939–940, 2010.

Hoyo I, Martinez-Pastor J, Garcia-Ramiro S, et al. Decreased serum linezolid concentrations in two patients receiving linezolid and rifampicin due to bone infections. *Scand J Infect Dis* 44:548–50, 2012.

Ijaz K, Jereb JA, Lambert LA, et al. Severe or fatal injury in 50 Patients in the United States taking rifampin and pyrazinamide for latent tuberculosis infection. *Clin Infect Dis* 42:346–56, 2006.

Ijaz K, McElroy PD, Navin TR. Short course rifampin and pyrazinamide compared with isoniazid for latent tuberculosis infection: a cost-effectiveness analysis based on a multicenter clinical trial *Clin Infect Dis* 39:289, 2004.

Johnson JR. Rifampin and methicillin-resistant *Staphylococcus aureus* bone and joint infections. *Clin Infect Dis* 53:98–99, 2011.

Krause PJ, Corrow CL, Bakken JS. Successful treatment of human granulocytic ehrlichiosis in children using rifampin. *Pediatrics*. 112:e252–3, 2003.

"Usual dose" assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose  $\times$  1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

Lundstrom TS, Sobel JD. Vancomycin, trimethoprim-sulfamethoxazole, and rifampin. *Infect Dis Clin North Am* 91:747-67, 1995.

Nguyen S, Pasquet A, Legout L, et al. Efficacy and tolerance of rifampin-linezolid compared with rifampin-cotrimoxazole combinations in prolonged oral therapy for bone and joint infections. *Clin Microbiol Infect Dis* 15:1163-1169, 2009.

Pina JM, Clotet L, Ferrer A, et al. Cost-effectiveness of rifampin for 4 months and isoniazid for 9 months in the treatment to tuberculosis infection. *Eur J Clin Microbiol Inf Dis* 32:647-655, 2013.

Sax PE, Cohen CJ, Kuritzkes DR. *HIV Essentials*, 5<sup>th</sup> edition Jones & Bartlett, Sudbury, MA, 2012.

Schlossberg D: An approach to drug-induced liver injury during tuberculosis therapy. *Infect Dis Clin Pract* 18:107-108, 2010.

Schlossberg D, Treatment of multi-drug resistant tuberculosis. *Antibiotics for Clinicians* 9:317-321, 2005.

Van Scoy RE, Wilkowske CJ. Antituberculous agents. *Mayo Clin Proc* 67:179-87, 1992.

Vesely JJ, Pien FD, Pien BC. Rifampin, a useful drug for nonmycobacterial infections. *Pharmacotherapy* 18:345-57, 1998.

Website: [www.pdr.net](http://www.pdr.net)

## Rilpivirine (Edurant) RPV

**Drug class:** HIV NNRTI.

**Usual dose:** 25 mg (PO) q24h (with meal).

**Pharmacokinetic Parameters:**

*Peak serum level:* 1 mcg/L

*Bioavailability:* unknown

*Excreted unchanged:* 1%

*Serum half-life (normal/ESRD):* 50/50 hrs

*Plasma protein binding:* 99%

*Volume of distribution (V<sub>d</sub>):* unknown

**Primary Mode of Elimination:** Hepatic

**Dosage Adjustments\***

CrCl <10 mL/min	No change
Post-HD dose	None
Post-PD dose	None
CVWH/CVWHD/CVWHDf dose	No change
Mild to moderate hepatic insufficiency	No change
Severe hepatic insufficiency	Use with caution

**Drug Interactions:** Clarithromycin (↓ clarithromycin levels). Any drug that induces or inhibits CYP3A may ↑/↓ rilpivirine levels.

**Contraindications:** carbamazepine, oxycarbazepine, phenobarbital, phenytoin, rifabutin, rifampin, rifapentine, proton pump inhibitors, dexamethasone (*except* single dose), St. John's Wort.

**Adverse Effects:** Nausea, vomiting, abdominal pain, fatigue, headache, dizziness, depressive disorders, insomnia, abnormal dreams, rash, ↑ QTC.

**Allergic Potential:** Low

**Safety in Pregnancy:** B

**Comments:** Antacid and H2 blockers should be separated. **Do not administer with other NNRTI's.**

### REFERENCES:

- Miller CD, Crain J, Tran V, et al. Rilpivirine: A new addition to the anti-HIV-1 armamentarium. *Drugs Today* 47:5-15, 2011.
- Molina JM, Cahn P, Grinsztejn B, et al. Rilpivirine versus efavirenz with tenofovir and emtricitabine in treatment-naïve adults infected with HIV-1 (ECHO): A phase 3

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

randomised double-blind active-controlled trial. *Lancet* ID 16:238–246, 2011.

Schrijvers R, Desimie BA, Debyser Z. Rilipivirine: A step forward in tailored HIV treatment. *Lancet* ID 16:201–203, 2011.

Website: [www.pdr.net](http://www.pdr.net)

## Rimantadine (Flumadine)

**Drug Class:** Antiviral (influenza)

**Usual Dose:** 100 mg (PO) q12h (see comments).

**Pharmacokinetic Parameters:**

*Peak serum level:* 0.7 mcg/ml

*Bioavailability:* 90%

*Excreted unchanged (urine):* 25%

*Serum half-life (normal/ESRD):* 25/38 hrs

*Plasma protein binding:* 40%

*Volume of distribution ( $V_d$ ):* 4.5 L/kg

**Primary Mode of Elimination:** Hepatic

**Dosage Adjustments\***

CrCl 10–50 mL/min	No change
CrCl < 10 mL/min	100 mg (PO) q24h
Post-HD dose	None
Post-PD dose	None
CVVH/CVVHD/ CWHDF dose	No change
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	100 mg (PO) q24h

**Drug Interactions:** Alcohol (↑ CNS effects); benzotropine, trihexyphenidyl, scopolamine (↑ interacting drug effect: dry mouth, ataxia, blurred vision, slurred speech, toxic psychosis); cimetidine

(↓ rimantadine clearance); CNS stimulants (additive stimulation); digoxin (↑ digoxin levels); trimethoprim (↑ rimantadine and trimethoprim levels).

**Adverse Effects:** Dizziness, headache, insomnia, anticholinergic effects (blurred vision, dry mouth, orthostatic hypotension, urinary retention, constipation).

Allergic **Potential:** Low

Safety in Pregnancy: C

**Comments:** Less anticholinergic side effects than amantadine. Patients ≥ 60 years old or with a history of seizures should receive 100 mg (PO) q24h. **Influenza dose (prophylaxis):** 100 mg (PO) q12h for duration of exposure/outbreak. **Influenza dose (therapy):** 100 mg (PO) q12h × 7 days. **May improve peripheral airway function/oxygenation in severe influenza pneumonia. Highly active against influenza A. No activity against other viruses.**

**Cerebrospinal Fluid Penetration:** < 10%

### REFERENCES:

- Dolin R, Reichman RC, Madore HP, et al. A controlled trial of amantadine and rimantadine in the prophylaxis of Influenza A infection. *N Engl J Med* 307:580–4, 1982.
- Gravenstein S, Davidson HE. Current strategies for management of influenza in the elderly population. *Clin Infect Dis* 35:729–37, 2002.
- Jefferson T, Deeks JJ, Demicheli V, et al. Amantadine and rimantadine for preventing and treating influenza A in adults. *Cochrane Database Syst Rev* 3:CD001169, 2004.
- Keating MR. Antiviral agents. *Mayo Clin Proc* 67:160–78, 1992.
- Schmidt AC. Antiviral therapy for influenza: a clinical and economic comparative review. *Drugs* 64:2031–46, 2004.
- Wintermeyer SM, Nahata MC. Rimantadine: A clinical perspective. *Ann Pharmacotherapy* 29:299–310, 1995.
- Website: [www.pdr.net](http://www.pdr.net)

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

## Ritonavir (Norvir) RTV

**Drug Class:** HIV protease inhibitor.

**Usual Dose:** 100–400 mg (PO) q12h–q24h (see comments).

**Pharmacokinetic Parameters:**

*Peak serum level:* 11 mcg/ml

*Bioavailability:* No data

*Excreted unchanged (urine):* 3.5%

*Serum half-life (normal/ESRD):* 4 hrs/no data

*Plasma protein binding:* 99%

*Volume of distribution ( $V_d$ ):* 0.41 L/kg

**Primary Mode of Elimination:** Hepatic

**Dosage Adjustments\***

CrCl < 10 mL/min	No change
Post-HD dose	None
Post-PD dose	None
CWVH/CVWHD/ CWVHDF dose	None
Mild-moderate hepatic insufficiency	No change
Severe hepatic insufficiency	Avoid

**Antiretroviral Dosage Adjustments**

Atazanavir	Ritonavir 100 mg q24h + atazanavir 300 mg q24h (with food)
Delavirdine	Delavirdine: no change; ritonavir: No data
Efavirenz	Ritonavir 600 mg q12h (500 mg q12h for intolerance)
Fosamprenavir	Fosamprenavir 1400 mg + ritonavir 200 mg q24h

Indinavir	Ritonavir 100–200 mg q12h + indinavir 800 mg q12h, or 400 mg q12h of each drug
Nelfinavir	Ritonavir 400 mg q12h + nelfinavir 500–750 mg q12h
Nevirapine	No change
Saquinavir	Ritonavir 400 mg q12h + saquinavir 400 mg q12h
Ketoconazole	Caution; do not exceed ketoconazole 200 mg q24h
Rifampin	Avoid
Rifabutin	Rifabutin 150 mg q48h or 3x/week

**Drug Interactions:** Antiretrovirals, rifabutin, rifampin (see dose adjustment grid, above); saquinavir (↑ QTc/torsades de pointes, ↑ PR intervals) alprazolam, diazepam, estazolam, flurazepam, midazolam, triazolam, zolpidem, meperidine, propoxyphene, piroxicam, quinidine, amiodarone, encainide, flecainide, propafenone, astemizole, bepridil, bupropion, cisapride, clorazepate, clozapine, pimozide, St. John's wort, terfenadine (avoid); alfentanil, fentanyl, hydrocodone, tramadol, disopyramide, lidocaine, mexiletine, erythromycin, clarithromycin, warfarin, dronabinol, ondansetron, metoprolol, pindolol, propranolol, timolol, amlodipine, diltiazem, felodipine, isradipine, nicardipine, nifedipine, nimodipine, nisoldipine, nitrendipine, verapamil, etoposide, paclitaxel, tamoxifen, vinblastine, vincristine, loratadine, tricyclic antidepressants, paroxetine, nefazodone, sertraline, trazodone, fluoxetine, venlafaxine, fluvoxamine, cyclosporine, tacrolimus, chlorpromazine, haloperidol, perphenazine, risperidone, thioridazine, clozapine,

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

pimozide, methamphetamine (↑ interacting drug levels); voriconazole (↓ voriconazole levels); telithromycin (↑ ritonavir levels); codeine, hydromorphone, methadone, morphine, ketoprofen, ketorolac, naproxen, diphenoxylate, oral contraceptives, theophylline (↓ interacting drug levels); carbamazepine, phenytoin, phenobarbital, clonazepam, dexamethasone, prednisone (↓ ritonavir levels, ↑ interacting drug levels; monitor anticonvulsant levels); metronidazole (disulfiram-like reaction); tenofovir, tobacco (↓ ritonavir levels); sildenafil (do not exceed 25 mg in 48 hrs); tadalafil (max. 10 mg/72 hrs); vardenafil (max. 2.5 mg/72 hrs).

**Adverse Effects:** Anorexia, anemia, leukopenia, hyperglycemia (including worsening diabetes, new-onset diabetes, DKA), ↑ cholesterol/triglycerides (evaluate risk for coronary disease/pancreatitis), fat redistribution, ↑ CPK, nausea, vomiting, diarrhea, abdominal pain, circumoral/extremity paresthesias, ↑ SGOT/SGPT, pancreatitis, taste perversion, possible increased bleeding in hemophilia, ↑ incidence of renal stones with ritonavir boosted atazanavir.

**Allergic Potential:** Low

**Safety in Pregnancy:** B

**Comments:** Usually used at low dose (100–400 mg/day) as pharmacokinetic “booster” of other PIs. GI intolerance decreases over time. Take with food if possible (serum levels increase 15%, fewer GI side effects). Separate dosing from ddl by 2 hours.

**Cerebrospinal Fluid Penetration:** < 1%

#### REFERENCES:

- Cameron DW, Japour AJ, Xu Y, et al. Ritonavir and saquinavir combination therapy for the treatment of HIV infection. *AIDS* 13:213–224, 1999.
- DeJeus E, Rockstroh JK, Henry K, et al. Co-Formulated elvitegravir, cobicistat, emtricitabine and tenofovir disoproxil fumarate versus ritonavir-boosted atazanavir plus co-formulated emtricitabine and tenofovir

disoproxil fumarate for initial treatment of HIV-infection: a randomized, double-blind, phase 3, non-inferiority trial. *Lancet* 379–2429–38, 2012.

- Diaz-Brito V, Leon A, Knobel H, et al. Post-exposure prophylaxis for HIV-infection: a clinical trial comparing lopinavir/ritonavir versus atazanavir each with zidovudine/lamivudine. *Antivir Ther* 17:337–46, 2012.
- Kaul DR, Cinti SK, Carver PL, et al. HIV protease inhibitors: Advances in therapy and adverse reactions, including metabolic complications. *Pharmacotherapy* 19:281–98, 1999.
- Lea AP, Faulds D. Ritonavir. *Drugs* 52:541–6, 1996.
- McDonald CK, Kuritzkes DR. Human immunodeficiency virus type 1 protease inhibitors. *Arch Intern Med* 157:951–9, 1997.
- McDonald C, Uy J, Hu W, et al. Clinical significance of hyperbilirubinemia among HIV-1-infected patients treated with atazanavir/ritonavir through 96 weeks in the CASTLE study. *AIDS patient care STDS* 26–259–64, 2012.
- Nishijima T, Tsukada K, Teruya K, et al. Efficacy and safety of once-daily ritonavir-based antiretroviral therapy in low-body weight treatment-naïve patients with HIV infections. *AIDS* 13:26:649–51 2012.
- Pillero PJ. Interaction between ritonavir and stains. *Am J Med* 112:510–1, 2002.
- Rockwood N, Mandalia S, Bower M, et al. Ritonavir-boosted atazanavir exposure is associated with an increased rate of renal stones compared with efavirenz, ritonavir-boosted lopinavir and ritonavir-boosted darunavir. *AIDS* 25:1671–3, 2011.
- Thompson MA, Aberg JA, Cahn P, et al. Antiretroviral treatment of adult HIV infection: 2012 recommendations of the International Antiviral Society — USA panel. *JAMA* 308:387–402, 2012.
- Website: [www.pdr.net](http://www.pdr.net)

## Saquinavir (Invirase) SQV

**Drug Class:** HIV protease inhibitor.

**Usual Dose:** 1000 mg (PO) q12h (see comments) with ritonavir 100 mg (PO) q12h.

**Pharmacokinetic Parameters:**

*Peak serum level: 0.07 mcg/mL*

“Usual dose” assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

*Bioavailability: hard-gel (4%)*

*Excreted unchanged (urine): 13%*

*Serum half-life (normal/ESRD): 13 hrs/no data*

*Plasma protein binding: 98%*

*Volume of distribution ( $V_d$ ): 10 L/kg*

**Primary Mode of Elimination:** Hepatic

**Dosage Adjustments\***

CrCl < 10 mL/min	No change
Post-HD dose	None
Post-PD dose	None
CWH/CVHD/ CWHDF dose	No change
Moderate hepatic insufficiency	Use with caution
Severe hepatic insufficiency	Avoid

#### Antiretroviral Dosage Adjustments

Darunavir	Avoid
Delavirdine	No data
Efavirenz	Avoid use as sole PI
Indinavir	No data
Lopinavir/ritonavir 2 capsules q12h	Saquinavir 1 gm q12h
Nelfinavir	Saquinavir 1 gm q12h or 1200 mg q12h
Nevirapine	No data
Ritonavir	Ritonavir 100 mg q12h + saquinavir 1 gm q12h
Rifampin	Avoid
Rifabutin	Avoid

**Drug Interactions:** Antiretrovirals, rifabutin, rifampin (see dose adjustment grid, above); astemizole, terfenadine, benzodiazepines, cisapride,

ergotamine, statins, St. John's wort (avoid if possible); carbamazepine, phenytoin, phenobarbital, dexamethasone, prednisone (↓ saquinavir levels, ↑ interacting drug levels; monitor anticonvulsant levels); clarithromycin, erythromycin, telithromycin (↑ saquinavir and macrolide levels); grapefruit juice, itraconazole, voriconazole, ketoconazole (↑ saquinavir levels); sildenafil (do not give > 25 mg/48 hrs); tadalafil (max. 10 mg/72 hrs), vardenafil (max. 2.5 mg/72 hrs), ritonavir (↑ QTc, ↑ PR intervals), PPIs (↑ saquinavir levels). Atovaquone/proguanil (↑ saquinavir levels).

**Contraindications:** Rifapentine, trazadone.

**Adverse Effects:** Anorexia, headache, anemia, leukopenia, hyperglycemia (including worsening diabetes, new-onset diabetes, DKA), ↑ cholesterol/triglycerides (evaluate risk for coronary disease/pancreatitis), ↑ SGOT/SGPT, hyperuricemia, fat redistribution, night sweats, possible increased bleeding in hemophilia.

**Allergic Potential:** Low

**Safety in Pregnancy:** B

**Comments:** Take with food. Avoid garlic supplements, which ↓ saquinavir levels ~ 50%. Boosted dose: 1 gm saquinavir/100 mg ritonavir (PO) q12h. Preferred formulation is 500 mg hard-gel capsule (Invirase 500). Soft-gel capsules (Fortovase) no longer available.

**Cerebrospinal Fluid Penetration:** < 1%

#### REFERENCES:

- Cameron DW, Japour AJ, Xu Y, et al. Ritonavir and saquinavir combination therapy for the treatment of HIV infection. *AIDS* 13:213–224, 1999.
- Hughes PJ, Cretton-Scott E, Teague A, et al. Protease inhibitors for patients with HIV-1 infection. *Pharm Ther* 36:332–344, 2011.

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

Panel on Antiretroviral Guidelines for Adults and Adolescents. Department of Health and Human Services. Guidelines for the use of antiretroviral agents in HIV-infected adults and adolescents. [http://aidsinfo.nih.gov/ConsentFiles/Adult and Adolescent\\_GL.pdf](http://aidsinfo.nih.gov/ConsentFiles/Adult_and_Adolescent_GL.pdf), 2012.

Perry CM, Noble S. Saquinavir soft-gel capsule formation: A review of its use in patients with HIV infection. *Drugs* 55:461–86, 1998.

Thompson MA, Aberg JA, Cahn P, et al. Antiretroviral treatment of adult HIV infection: 2012 recommendations of the International Antiviral Society — USA panel. *JAMA* 308:387–402, 2012.

Tomassi C, Bellagamba R, Tempestilli M, et al. Marked increase in etravirine and saquinavir plasma concentrations during atovaquone/proguanil prophylaxis. *Malar J* 10:141, 2011.

Website: [www.pdr.net](http://www.pdr.net)

## Spectinomycin (Spectam, Trobicin)

**Drug Class:** Aminocyclitol.

**Usual Dose:** 2 gm (IM) × 1 dose.

**Pharmacokinetic Parameters:**

*Peak serum level:* 100 mcg/ml

*Bioavailability:* Not applicable

*Excreted unchanged (urine):* 80%

*Serum half-life (normal/ESRD):* 1.6/16 hrs

*Plasma protein binding:* 20%

*Volume of distribution ( $V_d$ ):* 0.25 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\***

CrCl < 10 mL/min	No change
Post-HD dose	None
Post-PD dose	None
CVVH/CVVHD/CVHDF dose	None

Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	No change

**Drug Interactions:** None.

**Adverse Effects:** Local pain at injection site.

**Allergic Potential:** Low

**Safety in Pregnancy:** B

**Comments:** Ineffective in pharyngeal GC (poor penetration into secretions).

**Cerebrospinal Fluid Penetration:** < 10%

### REFERENCES:

Fiumara NJ. The treatment of gonococcal proctitis:

An evaluation of 173 patients treated with 4 gm of spectinomycin. *JAMA* 239:735–7, 1978.

Holloway WJ. Spectinomycin. *Med Clin North Am* 66:169–173, 1995.

McCormack WM, Finland M. Spectinomycin. *Ann Intern Med* 84:712–16, 1976.

Tapsall J. Current concepts in the management of gonorrhoea. *Expert Opin Pharmacother* 3:147–57, 2002.

Website: [www.pdr.net](http://www.pdr.net)

## Stavudine (Zerit) d4T

**Drug Class:** HIV NRTI (nucleoside reverse transcriptase inhibitor).

**Usual Dose:** ≥ 60 kg: 40 mg (PO) q12h;  
< 60 kg: 30 mg (PO) q12h.

**Pharmacokinetic Parameters:**

*Peak serum level:* 4.2 mcg/mL

*Bioavailability:* 86%

*Excreted unchanged (urine):* 40%

*Serum half-life (normal/ESRD):* 1.0/5.1 hrs

*Plasma protein binding:* 0%

*Volume of distribution ( $V_d$ ):* 0.5 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\*** ≥ 60 kg/[≤ 60 kg]

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.



CrCl 50–80 mL/min	40 mg (PO) q12h [30 mg (PO) q12h]
CrCl 25–50 mL/min	20 mg (PO) q12h [15 mg (PO) q12h]
CrCl ~ 10–25 mL/min	20 mg (PO) q24h [15 mg (PO) q24h]
Post-HD dose	20 mg (PO) [15 mg (PO)]
Post-PD dose	No data
CWVH/CVWHD/ CWVHDF dose	20 mg (PO) q24h [15 mg (PO) q24h]
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	No change

**Drug Interactions:** Ribavirin (↓ stavudine efficacy, ↑ risk of lactic acidosis); dapson, INH, other neurotoxic agents (↑ risk of neuropathy), didanosine (↑ risk of neuropathy, lactic acidosis). Avoid combining zidovudine with stavudine.

**Adverse Effects:** Lactic acidosis and severe hepatomegaly with steatosis, including fatal cases, have been reported with the use of nucleoside analogues alone or in combination, including stavudine and other antiretrovirals. Fatal lactic acidosis has been reported in pregnant women who received the combination of stavudine and didanosine with other antiretroviral agents. The combination of stavudine and didanosine should be used with great caution during pregnancy and only when necessary. Drug fever/rash, nausea, vomiting, GI upset, diarrhea, headache, insomnia, dose dependent peripheral neuropathy, myalgias, pancreatitis, ↑ SGOT/SGPT, ↑ cholesterol, facial fat pad wasting, lipodystrophy,

thrombocytopenia, leukopenia, lactic acidosis with hepatic steatosis. There is an increased risk of hepatotoxicity may occur in patients treated with stavudine in combination with didanosine and hydroxyurea. Immune reconstitution syndrome can occur.

**Allergic Potential:** Low

**Safety in Pregnancy:** C

**Comments:** Pancreatitis may be severe/fatal. Avoid coadministration with AZT or ddC. Decrease dose in patients with peripheral neuropathy to 20 mg (PO) q12h. Pregnant women may be at increased risk for lactic acidosis/liver damage when stavudine is used with didanosine (ddl). WHO recommends 30 mg q12h regardless of body weight.

**Cerebrospinal Fluid Penetration:** 30%

#### REFERENCES:

- Barasconi E, Boubaker K, Junghans C, et al. Abnormalities of body fat distribution in HIV-infected persons treated with antiretroviral drugs: The Swiss HIV Cohort Study. *J Acquir Immune Defic Syndr* 31:50–5, 2002.
- Dudley MN, Graham KK, Kaul S, et al. Pharmacokinetics of stavudine in patients with AIDS and AIDS-related complex. *J Infect Dis* 166:480–5, 1992.
- Miller KD, Cameron M, Wood LV, et al. Lactic acidosis and hepatic steatosis associated with use of stavudine: Report of four cases. *Ann Intern Med* 133:192–196, 2000.
- Thompson MA, Aberg JA, Cahn P, et al. Antiretroviral treatment of adult HIV infection: 2012 recommendations of the International Antiviral Society — USA panel. *JAMA* 308:387–402, 2012.

Website: [www.pdr.net](http://www.pdr.net)

## Streptomycin

**Drug Class:** Aminoglycoside.

**Usual Dose:** 15 mg/kg (IM) q24h or 1 gm (IM) q24h (see comments).

"Usual dose" assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

**Pharmacokinetic Parameters:***Peak serum level:* 25–50 mcg/ml*Bioavailability:* Not applicable*Excreted unchanged (urine):* 90%*Serum half-life (normal/ESRD):* 2.5/100 hrs*Plasma protein binding:* 35%*Volume of distribution ( $V_d$ ):* 0.26 L/kg**Primary Mode of Elimination:** Renal**Dosage Adjustments\***

CrCl 10–50 mL/min	15 mg/kg (IM) q72h or 1 gm (IM) q72h
CrCl < 10 mL/min	15 mg/kg (IM) q72h or 1 gm (IM) q96h
Post-HD dose	15 mg/kg (IM) or 1 gm (IM) 2–3 x/week
Post-PD dose	15 mg/kg (IM) or 1 gm (IM) or 20–40 mg/ml in dialysate q24h
CVVH/CVVHD/ CVVHDF dose	15 mg/kg (IM) or 1 gm (IM) q96h
Mild-moderate hepatic insufficiency	No change
Severe hepatic insufficiency	No change

**Drug Interactions:** Amphotericin B, cephalothin, cyclosporine, enflurane, methoxyflurane, NSAIDs, polymyxin B, radiographic contrast, vancomycin (↑ nephrotoxicity); cis-platinum (↑ nephrotoxicity, ↑ ototoxicity); loop diuretics (↑ ototoxicity); neuromuscular blocking agents (↑ apnea, prolonged paralysis); non-polarizing muscle relaxants (↑ apnea).

**Adverse Effects:** Most ototoxic aminoglycoside (usually vestibular ototoxicity); least nephrotoxic aminoglycoside.

**Allergic Potential:** Low**Safety in Pregnancy:** D**Comments:** May be given IV slowly over 1 hour.**TB D.O.T. dose:** 20–30 mg/kg (IM) 2–3x/week.**Tularemia dose:** 1 gm (IV/IM) q12h. **Plague dose:** 2 gm (IV/IM) q12h.**Cerebrospinal Fluid Penetration:** 20%**REFERENCES:**

- Akaho E, Maekawa T, Uchinashi M. A study of streptomycin blood level information of patients undergoing hemodialysis. *Biopharm Drug Dispos* 23:47–52, 2002.
- Davidson PT, Le HQ. Drug treatment of tuberculosis 1992. *Drugs* 43:651–73, 1992.
- Kim-Sing A, Kays MB, Vivien EJ, et al. Intravenous streptomycin use in a patient infection with high-level gentamicin-resistant *Streptococcus faecalis*. *Ann Pharmacother* 27:712–4, 1993.
- Morris JT, Cooper RH. Intravenous streptomycin: A useful route of administration. *Clin Infect Dis* 19:1150–1, 1994.
- Roushan MRH, Mohraz M, Hajiahmadi M, et al. Efficacy of gentamicin plus doxycycline versus streptomycin plus doxycycline in the treatment of brucellosis in humans. *Clin Infect Dis* 42:1075–80, 2006.
- Van Scoy RE, Wilkowske CJ. Antituberculous agents. *Mayo Clin Proc* 67:179–87, 1992.
- Ormerod P. The clinical management of the drug-resistant patient. *Ann NY Acad Sci* 953:185–91, 2001.
- Website: [www.pdr.net](http://www.pdr.net)

**Tedizolid (Sivextro)****Drug Class:** Oxazolidinone**Usual Dose:** 200 mg (IV) q24h × 6 days; 200 mg (PO) q24h × 6 days**Spectrum:** (see **Susceptibility Profiles** pp. 198–202)

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

**Resistance Potential:** Low

**Pharmacokinetic Parameters:**

Peak serum level: 2.2–3 mcg/ml

Bioavailability: 91%

Excreted unchanged: < 3%

Serum half-life (normal/ESRD): 11 h/11 h

Plasma protein binding: 70–90%

Volume of distribution ( $V_d$ ): 1–1.14 L/kg

**Primary Mode of Elimination:** Hepatic

**Dosage Adjustments\***

CrCl < 30 mL/min	No change
Post-HD dose	No change
Post-PD dose	No change
CVWH dose	No change
Hepatic insufficiency	No change

**Drug Interactions:** None

**Adverse Effects:** Headache, dizziness, nausea, vomiting, diarrhea, C. difficile, anemia, thrombocytopenia, pancytopenia

**Allergic Potential:** Low

**Safety in Pregnancy:** C

**Comments:** Useful for treatment of linezolid resistant staphylococci. ↓ incidence of thrombocytopenia compared to linezolid (2.3% vs. 4.9%). Only compatible with 0.9% NaCl; infuse over 1 hour. PO dose may be taken with or without food. Other anti-staphylococcal antibiotics are preferred, in the setting of febrile neutropenia.

**Cerebrospinal Fluid Penetration:** No data

## REFERENCES:

Das D, Tulkens PM, Mehra P, et al. Tedizolid phosphate for the management of acute bacterial skin and skin structure infections: safety summary. *Clin Infect Dis.* 58;Suppl1:S51–S7, 2014

Flanagan S, Fang E, Munoz KA, et al. Single- and multiple-dose pharmacokinetics and absolute bioavailability of tedizolid. *Pharmacotherapy.* 34:891–900, 2014.

Flanagan S, Minassian SL, Morris D, et al. Pharmacokinetics of tedizolid in subjects with renal or hepatic impairment. *Antimicrob Agents Chemother.* 58:6471–6476, 2014.

Housman ST, Pope JS, Russomanno J, et al. Pulmonary disposition of tedizolid following administration of once-daily oral 200-milligram tedizolid phosphate in healthy adult volunteers. *Antimicrob Agents Chemother.* 56:2627–2634, 2012.

Kisgen JJ, Mansour H, Unger NR, et al. Tedizolid: a new oxazolidinone antimicrobial. *Am J Health Syst Pharm.* 71:621–633, 2014.

Lodise TP, Fang E, Minassian SL, et al. Platelet profile in patients with acute bacterial skin and skin structure infections receiving tedizolid or linezolid: findings from the Phase 3 ESTABLISH clinical trials. *Antimicrob Agents Chemother.* 58:7198–7204, 2014

Molina-Torres CA, Barba-Marines A, Valles-Guerra O, et al. Intracellular activity of tedizolid phosphate and ACH-702 versus *Mycobacterium tuberculosis* infected macrophages. *Ann Clin Microbiol Antimicrob.* 13:13, 2014.

Moran GJ, Fang E, Corey GR, et al. Tedizolid for 6 days versus linezolid for 10 days for acute bacterial skin and skin-structure infections (ESTABLISH-2): a randomised, double-blind, phase 3, non-inferiority trial. *Lancet Infect Dis.* 14:696–705, 2014.

O’Riordan W, Green S, Mehra P, et al. Tedizolid phosphate for the management of acute bacterial skin and skin structure infections: efficacy summary. *Clin Infect Dis.* 58;Suppl1:S43–50, 2014.

Prokocimer P, De Anda C, Fang E, et al. Tedizolid phosphate vs linezolid for treatment of acute bacterial skin and skin structure infections: the ESTABLISH-1 randomized trial. *JAMA.* 309:559–569, 2013.

Rodriguez-Avil I, Culebras E, Betriu C, et al. In vitro activity of tedizolid (TR-700) against linezolid-resistant staphylococci. *J Antimicrob Chemother.* 67:167–169, 2012.

Shorr AF, Lodise TP, Corey GR, et al. Analysis of the phase 3 ESTABLISH trials: Tedizolid versus linezolid in acute bacterial skin and skin structure infection. *Antimicrob Agents Chemother.* 59:864–871, 2015.

Urbina O, Ferrandez O, Espona M, Salas E, Ferrandez I, Grau S. Potential role of tedizolid phosphate in the treatment of acute bacterial skin infections. *Drug Des Devel Ther.* 7:243–265, 2013.

Website: [www.pdr.net](http://www.pdr.net)

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

## Telavancin (Vibativ)

**Drug class:** Lipoglycopeptide.

**Usual dose:** 10 mg/kg (IV) q24h.

**Spectrum:** (see **Susceptibility Profiles** pp. 198–202).

**Resistance Potential:** Low

**Pharmacokinetic Parameters:**

*Peak serum level:* 93–108 mcg/mL

*Bioavailability:* not applicable

*Excreted unchanged:* 76%

*Serum half-life (normal/ESRD):* 6–9 hrs/no data

*Plasma protein binding:* 93%

*Volume of distribution ( $V_d$ ):* 0.133–0.145 mL/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\***

CrCl > 50 mL/min	No change
CrCl 30–50 mL/min	7.5 mg/kg (IV) q24h
CrCl 10–29 mL/min	10 mg/kg (IV) q48h
CrCl < 10 mL/min	Avoid
Post-HD dose	None
Post-PD dose	None
CVVH/CVVHD/ CVHDF dose	7.5 mg/kg (IV) q24h
Mild-moderate hepatic insufficiency	No change
Severe hepatic insufficiency	Use with caution

**Drug Interactions:** Drugs that prolong  $QT_c$  (additive effect); PT, INR, PTT ( $\uparrow$  levels); other interactions unlikely because is not metabolized through CYP450 system.

**Adverse Effects:** Headache, taste disturbance, nausea, vomiting, diarrhea,  $\uparrow$  nephrotoxicity, with CrCl < 50 mL/min “red man syndrome.”

**Allergic Potential:** Low

**Safety in Pregnancy:** C

**Comments:** Infuse over at least 60 minutes to minimize infusion-related reactions. Effective for MSSA/MRSA cSSSIs. Efficacy may be decreased with CrCl < 50 mL/min.

**Cerebrospinal Fluid Penetration:**

Inflamed meninges: 2%

Non-inflamed meninges: 1%

(also see **Antibiotic Pearls & Pitfalls** p. 519).

### REFERENCES:

- Damodaran SE, Madhan S. Telavancin: A novel lipoglycopeptide antibiotic. *J Pharmacother* 2: 135–137, 2011.
- Joson J, Grover C, Downer C, et al. Successful treatment of methicillin-resistant *Staphylococcus aureus* mitral valve endocarditis with sequential linezolid and telavancin monotherapy following daptomycin failure. *J Antimicrob Chemother* 66:2186–2188, 2011.
- Leonard SN, Szeto YG, Zolotarev M, et al. Comparative in vitro activity of telavancin, vancomycin, and linezolid against heterogeneously vancomycin-intermediate *Staphylococcus aureus* (hVISA). *Int J Antimicrob* 37:558–561, 2011.
- Marcos LA, Camins BC. Successful treatment of vancomycin-intermediate *Staphylococcus aureus* pacemaker lead infective endocarditis with telavancin. *Antimicrob Agents Chemother* 54:5376–5378, 2010.
- Nailor MD, Sobel JD. Antibiotics for gram-positive bacterial infection: vancomycin, teicoplanin, quinupristin/dalfopristin, oxazolidinones, daptomycin, telavancin, and ceftaroline fosamil. *Med Clin North Am* 95:723–742, 2011.
- Plotkin P, Patel K, Uminski A, et al. Telavancin (Vibativ), a new option for the treatment of gram-positive infections. *Pharm Ther* 36:127–138, 2011.
- Plotkin P, Patel K, Uminski A, et al. Telavancin for the treatment of methicillin-resistant *Staphylococcus aureus* J. *Antimicrob Chemother* 66:2675–2677, 2011.
- Saravolatz LD, Stein GE, Johnson LB. Telavancin: a novel lipoglycopeptide. *Clin Infect Dis* 49:1908–1914, 2009.

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose  $\times$  1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

Shams W, Walker ES, Levy F, et al. Comparative activity of telavancin and other antimicrobial agents against methicillin-resistant *Staphylococcus aureus* isolates collected from 1991 to 2006. *Chemotherapy* 56:411–416, 2010.

Stryjewski ME, Graham DR, Wilson SE, et al. Telavancin versus vancomycin for the treatment of complicated skin and skin-structure infections caused by gram-positive organisms. *Clin Infect Dis* 46:1683–1693, 2008.

Twillia JD, Gelfand MS, Cleveland KO, et al. Telavancin for the treatment of methicillin-resistant *Staphylococcus aureus* osteomyelitis. *J Antimicrob Chemother* 66:2675–2677, 2011.

Wilson SE, O'Riordan W, Hopkins A, et al. Telavancin versus vancomycin for the treatment of complicated skin and skin-structure infections associated with surgical procedures. *Am J Surg* 197:791–796, 2009. Website: www.pdr.net

## Telithromycin (Ketek)

**Drug Class:** Ketolide.

**Usual Dose:** Acute sinusitis/AECB: 800 mg (PO) q24h × 5 days. Community-acquired pneumonia: 800 mg (PO) q24h × 7–10 days. 800 mg (PO) dose taken as two 400-mg tablets (PO) at once.

**Spectrum:** (see **Susceptibility Profiles** pp.186–190).

**Resistance Potential:** Low.

**Pharmacokinetic Parameters:**

*Peak serum level:* 2.27 mcg/ml

*Bioavailability:* 57%

*Excreted unchanged (urine):* 13%

*Serum half-life (normal/ESRD):* 9.8/11 hrs

*Plasma protein binding:* 65%

*Volume of distribution ( $V_d$ ):* 2.9 L/kg

**Primary Mode of Elimination:** Hepatic

**Dosage Adjustments\***

CrCl < 30 mL/min	600 mg (PO) q24h
CrCl < 30 mL/min + hepatic impairment	400 mg (PO) q24h
Post-HD/Post-PD dose	400 mg (PO)
CVVH/CVVHD/CVVHDF dose	400 mg (PO) q24h

**Drug Interactions:** Digoxin (↑ interacting drug levels); ergot derivatives (acute ergot toxicity); itraconazole, ketoconazole (↑ telithromycin level); midazolam, triazolam (↑ interacting drug levels, sedation); oral anticoagulants (may ↑ anticoagulant effects; monitor PT/INR); simvastatin (↑ risk of rhabdomyolysis; giving simvastatin 12h after telithromycin decreases the ↑ in simvastatin levels ~ 50%); theophylline (additive nausea). CYP 3A4 inhibitor/substrate. Telithromycin is contraindicated with cisapride and pimozide.

**Adverse Effects:** Nausea, diarrhea, dizziness, syncope, eye accommodation difficulties.

**Contraindicated in patients with history of hepatitis/jaundice** (rarely fatal acute/fulminant hepatitis or acute liver failure in those with pre-existing liver disease) **or myasthenia gravis** (↑ risk of respiratory failure).

**Allergic Potential:** Low

**Safety in Pregnancy:** C

**Comments:** May take with or without food.

### REFERENCES:

- Aubier M, Aldons PM, Leak A, et al. Telithromycin is as effective as amoxicillin/clavulanate in acute exacerbations of chronic bronchitis. *Respir Med* 96:862–71, 2002.
- Brown SD, Farrell DJ. Antibacterial susceptibility among *Streptococcus pneumoniae* isolated from paediatric and adult patients as part of the PROTECT US study in

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

- 2001–2002. *Journal of Antimicrobial Chemotherapy* 54 (Suppl. S1):23–29, 2004.
- Brown SD, Rybak MJ. Antimicrobial susceptibility of *Streptococcus pneumoniae*, *Streptococcus pyogenes* and *Haemophilus influenzae* collected from patients across the USA, in 2001–2002, as part of the PROTECT US study. *Journal of Antimicrobial Chemotherapy* 54 (Suppl. S1):7–15, 2004.
- Carbon C, Moola S, Velancsics I, et al. Telithromycin 800 mg once daily for seven to ten days is an effective and well-tolerated treatment for community-acquired pneumonia. *Clin Microbiol Infect* 9:691–703, 2003.
- Dunbar LM, Carbon C, van Rensburg D, et al. Efficacy of telithromycin in community-acquired pneumonia caused by atypical and intracellular pathogens. *Infect Dis Clin Pract* 13:10–16, 2005.
- Ferguson BJ, Guzzetta RV, Spector SL, et al. Efficacy and safety of oral telithromycin once daily for 5 days versus moxifloxacin one daily for 10 days in the treatment of acute bacterial rhinosinusitis. *Otolaryngol Head Neck Surg* 131:207–14, 2004.
- Hagberg L, Torres A, van Rensburg D, et al. Efficacy and tolerability of once-daily telithromycin compared with high-dose amoxicillin for treatment of community-acquired pneumonia. *Infection* 30:378–386, 2002.
- Jacobs MR. In vivo veritas: in vitro macrolide resistance in systemic *S. pneumoniae* infections does result in clinical failure. *Clin Infect Dis* 35:565–9, 2002.
- Kim, DM, Yu KD, Lee JH, et al. Controlled trial of 5-day course of telithromycin versus doxycycline for treatment of mild to moderate scrub typhus. *Antimicrob Agents Chemother* 51:2011–2015, 2007.
- Klugman KP, Lonks JR. Hidden epidemic of macrolide-resistant pneumococci. *Emerg Infect Dis* 11:802–807, 2005.
- Low DE, Brown S, Felmingham D. Clinical and bacteriology efficacy of the ketolide telithromycin against isolates of key respiratory pathogens: a pooled analysis of phase III studies. *Clin Microbiol Infect* 10:27–36, 2004.
- Ortega M, Marco F, Almela M, et al. Activity of telithromycin against erythromycin-susceptible resistant *Streptococcus pneumoniae* is from adults with invasive infections. *Int J Antimicrob Agents* 24:616–8, 2004.
- Tellier G, Niederman MS, Nusrat R, et al. Clinical and bacteriological efficacy and safety of 5 and 7 day regimens of telithromycin once daily compared with a 10 day regimen of clarithromycin twice daily in patients with mild to moderate community-acquired pneumonia. *Journal of Antimicrobial Chemotherapy* 54:515–523, 2004.
- van Rensburg D, Fogarty C, De Salvo MC, et al. Efficacy of oral telithromycin in community-acquired pneumonia caused by resistant *Streptococcus pneumoniae*. *Journal of Infection* 51:201–5, 2005.
- Zervos MJ, Heyder AM, Leroy B. Oral telithromycin 800 mg once daily for 5 days versus cefuroxime axetil 500 mg twice daily for 10 days in adults with acute exacerbations of chronic bronchitis. *J Int Med Res* 31:157–69, 2003.

Website: [www.pdr.net](http://www.pdr.net)

## Tenofovir disoproxil fumarate (Viread) TDF

**Drug Class:** HIV nucleotide analogue.

**Usual Dose:** 300 mg (PO) q24h.

**Pharmacokinetic Parameters:**

*Peak serum level:* 0.29 mcg/mL

*Bioavailability:* 25%/39% (fasting/high fat meal)

*Excreted unchanged (urine):* 32%

*Serum half-life (normal/ESRD):* 17 hrs/no data

*Plasma protein binding:* 0.7–7.2%

*Volume of distribution ( $V_d$ ):* 1.3 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\***

CrCl 30–49 mL/min	300 mg (PO) q48h
CrCl 10–29 mL/min	300 mg (PO) 2x/week
CrCl < 10 mL/min	Avoid
Post-HD dose	300 mg q7 days

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

Post-PD dose	No data
CVWH/CVWHD/ CVWHDf dose	300 mg (PO) 2x/week
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	No change

**Drug Interactions:** Not a substrate/inhibitor of cytochrome P-450 enzymes. Didanosine (if possible, avoid concomitant didanosine due to impaired CD<sub>4</sub> response and increased risk of virologic failure); valganciclovir (↑ tenofovir levels); atazanavir, lopinavir/ritonavir (↑ tenofovir levels)(↓ atazanavir levels; use atazanavir 300 mg/ritonavir 100 mg with tenofovir); telaprevir (↑ tenofovir levels).

**Adverse Effects:** Lactic acidosis and severe hepatomegaly with steatosis, including fatal cases, with nucleoside analogs. Severe acute exacerbations of hepatitis in HBV who have discontinued HBV therapy, including tenofovir. Mild nausea, vomiting, GI upset, asthenia, headache, diarrhea renal tubular acidosis, decrease bone density occur. It is recommended that creatinine clearance be calculated in all patients prior to initiating therapy and as clinically appropriate during therapy with tenofovir. Fanconi-like Syndrome routine monitoring of calculated creatinine clearance and serum phosphorus should be performed in patients at risk for renal impairment. Immune constitution syndrome can occur.

**Allergic Potential:** Low

**Safety in Pregnancy:** B

**Comments: Active against HBV.** Eliminated by glomerular filtration/tubular secretion. May be taken with or without food. If possible, avoid concomitant didanosine (see drug interactions).

**Cerebrospinal Fluid Penetration:** No data

#### REFERENCES:

- Choi AL, Vittinghoff E, Deeks SG, et al. Cardiovascular risks associated with abacavir and tenofovir exposure in HIV-infected persons. *AIDS* 19;25:1289–98, 2011.
- Cooper RD, Wiebe N, Smith N, et al. Systematic review and Meta-analysis: Renal safety of tenofovir disoproxil fumarate in HIV-infected patients. *Clin Infect Dis* 51:496–505, 2010.
- Daudé M, Rostaing L, Sauné K, et al. Tenofovir therapy in hepatitis B virus-positive solid-organ transplant recipients. *Transplantation* 91:916–920, 2011.
- Dupouey J, Gerolami R, Solas C, et al. Adefovir and tenofovir therapy. *Clin Res Hepatol Gastroenterol* 36:e26–28, 2012.
- Gallant JE, Deresinski S, Tenofovir disoproxil fumarate. *Clin Infect Dis* 37:944–50, 2003.
- Hall AM, Hendry BM, Nitsch D, et al. Tenofovir-associated kidney toxicity in HIV-infected patients: A review of new evidence. *Am J Kidney Dis* 57:773–80, 2011.
- Maggi P, Montinaro V, Bellacosa C, et al. Tenofovir versus abacavir. *AIDS Patient Care STDS* 26:5–11, 2012.
- Panel on Antiretroviral Guidelines for Adults and Adolescents. Department of Health and Human Services. Guidelines for the use of antiretroviral agents in HIV-infected adults and adolescents. <http://aidsinfo.nih.gov/ConsentFiles/AdultandAdolescentGL.pdf>, 2012.
- Ratcliffe L, Beadsworth MD, Pennell A, et al. Managing hepatitis B/HIV co-infected: adding entecavir to truvada (tenofovir disoproxil/emtricitabine) experienced patients. *AIDS* 25:1051–1056, 2011.
- Sax PE, Tierney C, Collier AC, et al. Abacavir/lamivudine versus tenofovir DF/emtricitabine as part of combination regimens for initial treatment of HIV: final results. *AIDS clinical trials group study A5202 team* 204:1191–1201, 2011.

"Usual dose" assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

Sax PE, Tierney C, Collier AC, et al. Abacavir/lamivudine versus tenofovir DF/emtricitabine as part of combination regimens for initial treatment of HIV: final results. *J Infect Dis* 204:1191–201, 2011.

Scotto G, D'Addiego G, Giammarino A, et al. Tenofovir plus emtricitabine as rescue therapy for multidrug-resistant chronic hepatitis B. *Liver Int* 32:171–2, 2012.

Si-Ahmen SN, Pradat P, Soutendijk R, et al. Efficacy and tolerance of a combination of tenofovir disoproxil fumarate plus emtricitabine in patients with chronic hepatitis B: A European multicenter study 92: 90–95, 2011.

Thompson MA, Aberg JA, Cahn P, et al. Antiretroviral treatment of adult HIV infection: 2012 recommendations of the International Antiviral Society — USA panel. *JAMA* 308:387–402, 2012.

Website: www.pdr.net

## Terbinafine (Lamisil, Daskil)

**Drug Class:** Antifungal.

**Usual Dose:** 250 mg (PO) q24h.

**Pharmacokinetic Parameters:**

*Peak serum level:* 1 mcg/ml

*Bioavailability:* 70%

*Excreted unchanged (urine):* < 1%

*Serum half-life (normal/ESRD):* 24 hrs/no data

*Plasma protein binding:* 99%

*Volume of distribution ( $V_d$ ):* 13.5 L/kg

**Primary Mode of Elimination:** Renal/Hepatic

**Dosage Adjustments\***

CrCl ~ 50–60 mL/min	No change
CrCl < 50 mL/min	Avoid
Post-HD dose	Avoid
Post-PD dose	Avoid
CVVH/CVVHD/CVVHDF dose	Avoid
Moderate—severe hepatic insufficiency	Avoid

**Drug Interactions:** Cimetidine (↓ terbinafine clearance, ↑ terbinafine levels); phenobarbital, rifampin (↑ terbinafine clearance, ↓ terbinafine levels).

**Adverse Effects:** Drug fever/rash, lymphopenia, leukopenia, ↑ SGOT/SGPT, visual disturbances, nausea, vomiting, GI upset.

**Allergic Potential:** Low

**Safety in Pregnancy:** B

**Comments:** May cause green vision and changes in the lens/retina.

**Cerebrospinal Fluid Penetration:** < 10%

### REFERENCES:

- Abdel-Rahman SM, Nahata MC. Oral terbinafine: A new antifungal agent. *Ann Pharmacother* 31:445–56, 1997.
- Amichai B, Grunwald MH. Adverse drug reactions of the new oral antifungal agents – terbinafine, fluconazole, and itraconazole. *Int J Dermatol* 37:410–5, 1998.
- Darkes MJ, Scott LJ, Goa KL. Terbinafine: a review of its use in onychomycosis in adults. *Am J Clin Dermatol* 4:39–65, 2003.
- Jain S, Sehgal VN. Itraconazole versus terbinafine in the management of onychomycosis: an overview. *J Dermatolog Treat* 14:30–42, 2003.
- Mosquera J, Shartt A, Moore CB, et al. In vitro interaction of terbinafine with itraconazole, fluconazole, amphotericin B and 5-flucytosine against *Aspergillus* spp. *J Antimicrob Chemother* 50:189–94, 2002.
- Salo H, Pekurinen M. Cost effectiveness of oral terbinafine (Lamisil) compared with oral fluconazole (Diflucan) in the treatment of patients with toenail onychomycosis. *Pharmacoeconomics* 20:319–24, 2002.
- Trepanier EF, Amsden GW. Current issues in onychomycosis. *Ann Pharmacother* 32:204–14, 1998.
- Website: www.pdr.net

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.



## Tetracycline (various)

**Drug Class:** Tetracycline.

**Usual Dose:** 500 mg (PO) q6h  
(Take without food).

**Spectrum:** (see *Susceptibility Profiles* pp. 186–190).

**Resistance Potential:** High (*S. pneumoniae*, MSSA)

**Pharmacokinetic Parameters:**

Peak serum level: 1.5 mcg/ml

Bioavailability: 60%

Excreted unchanged (urine): 60%

Serum half-life (normal/ESRD): 8/108 hrs

Plasma protein binding: 5%

Volume of distribution ( $V_d$ ): 0.7 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\***

CrCl 50–80 mL/min	500 mg (PO) q12h
CrCl 10–50 mL/min	500 mg (PO) q24h
Post-HD/Post-PD dose	None
CWH/CWHD/ CWHDF dose	500 mg (PO) q12h
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	≤ 1 gm (PO) q24h

**Drug Interactions:** Antacids, Al<sup>++</sup>, Ca<sup>++</sup>, Fe<sup>++</sup>, Mg<sup>++</sup>, Zn<sup>++</sup>, multivitamins, sucralfate (↓ absorption of tetracycline); barbiturates, carbamazepine, phenytoin (↓ half-life of tetracycline); bicarbonate (↓ absorption and ↑ clearance of tetracycline); digoxin (↑ digoxin levels); insulin (↑ insulin effect); methoxyflurane (↑ nephrotoxicity).

**Adverse Effects:** Nausea, vomiting, GI upset, diarrhea, hepatotoxicity, vaginal candidiasis, photosensitizing reactions, benign intracranial hypertension (pseudotumor cerebri).

**Allergic Potential:** Low

**Safety in Pregnancy:** D

**Comments:** Take without food. *Hepatotoxicity dose dependent (≥ 2 gm/day), especially in pregnancy/renal failure. Avoid prolonged sun exposure. Doxycycline or minocycline preferred for all tetracycline indications.*

**Cerebrospinal Fluid Penetration:**

Non-Inflamed meninges = 5%

Inflamed meninges = 5%

**Bile Penetration:** 1000%

(also see *Antibiotic Pearls & Pitfalls* p. 512).

### REFERENCES:

- Agwuh KN, MacGowan A. Pharmacokinetics and pharmacodynamics of the tetracyclines including glycylicyclines. *J of Antimicrob Chemotherapy* 58:256–265, 2006.
- Cunha BA, Comer J, Jonas M. The tetracyclines. *Med Clin North Am* 66:293–302, 1982.
- Dahl EL, Shock JL, Shenai BR, et al. Tetracyclines specifically target the apicoplast of the malaria parasite *Plasmodium falciparum*. *Antimicrob Agents Chemother* 50:3124–31, 2006.
- Malfertheiner P, Bazzoli F, Delchier JC, et al. *Helicobacter pylori* eradication with a capsule containing bismuth subcitrate potassium, metronidazole, and tetracycline given with omeprazole versus clarithromycin-based triple therapy: a randomized, open-label, non-inferiority, phase 3 trial. *Lancet* 33:905–913, 2011.
- Pugliese A, Cunha BA. Tetracyclines. *Int J Urogyn* 5:221–7, 1994.
- Smilack JD, Wilson WE, Cocerill Fr 3<sup>rd</sup>. Tetracycline, chloramphenicol, erythromycin, clindamycin, and metronidazole. *Mayo Clin Proc* 66:1270–80, 1991.
- Website: [www.pdr.net](http://www.pdr.net)

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

## Ticarcillin (Ticar)

**Drug Class:** Anti-pseudomonal penicillin.

**Usual Dose:** 3 gm (IV) q6h.

**Spectrum:** (see *Susceptibility Profiles* pp. 186–190).

**Resistance Potential:** Low

**Pharmacokinetic Parameters:**

*Peak serum level:* 118–300 mcg/ml

*Bioavailability:* Not applicable

*Excreted unchanged (urine):* 95%

*Serum half-life (normal/ESRD):* 1/5 hrs

*Plasma protein binding:* 45%

*Volume of distribution ( $V_d$ ):* 0.2 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\***

CrCl 50–80 mL/min	No change
CrCl 10–50 mL/min	2 gm (IV) q8h
CrCl < 10 mL/min	2 gm (IV) q12h
Post-HD dose	2 gm (IV)
Post-PD dose	3 gm (IV)
CVWH/CVWHD/ CWHDF dose	2 gm (IV) q8h
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	No change

**Drug Interactions:** Aminoglycosides (inactivation of ticarcillin in renal failure); warfarin (↑ INR); oral contraceptives (↓ oral contraceptive effect); cefoxitin (↓ ticarcillin effect).

**Adverse Effects:** Drug fever/rash; E. multiforme/Stevens-Johnson Syndrome, anaphylactic reactions (hypotension, laryngospasm, bronchospasm), hives, serum sickness.

Dose-dependent inhibition of platelet aggregation is minimal/absent (usual dose is less than carbenicillin). C. difficile diarrhea/colitis.

**Allergic Potential:** High

**Safety in Pregnancy:** B

**Comments:** Administer 1 hour before or after aminoglycoside. Na<sup>+</sup> content = 5.2 mEq/g

Meningeal dose = usual dose.

**Cerebrospinal Fluid Penetration:**

Non-Inflamed meninges = 1%

Inflamed meninges = 30%

(also see *Antibiotic Pearls & Pitfalls* p. 509).

### REFERENCES:

Donowitz GR, Mandell GL. Beta-lactam antibiotics.

N Engl J Med 318:419–26 and 318:490–500, 1993.

Tan JS, File TM, Jr. Antipseudomonal penicillins. Med Clin North Am 79:679–93, 1995.

Wright AJ. The penicillins. Mayo Clin Proc 74:290–307, 1999.

Website: www.pdr.net

## Ticarcillin/Clavulanate (Timentin)

**Drug Class:** Anti-pseudomonal penicillin.

**Usual Dose:** 3.1 gm (IV) q6h.

**Spectrum:** (see *Susceptibility Profiles* pp. 186–190).

**Resistance Potential:** Low

**Pharmacokinetic Parameters:**

*Peak serum level:* 330 mcg/ml

*Bioavailability:* Not applicable

*Excreted unchanged (urine):* 95/45%

*Serum half-life (normal/ESRD):* [1/13]/[1/2] hrs

*Plasma protein binding:* 45/25%

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

Volume of distribution ( $V_d$ ): 0.2/0.3 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\***

CrCl 30–60 mL/min	3.1 gm (IV) q8h
CrCl 10–30 mL/min	3.1 gm (IV) q12h
CrCl < 10 mL/min	2 gm (IV) q12h
Post-HD dose	3.1 gm (IV)
Post-PD dose	3.1 gm (IV)
CWH/CVVDH/ CWHDF dose	3.1 gm (IV) q8h
Moderate hepatic insufficiency	If CrCl < 10 mL/min: 2 gm (IV) q24h
Severe hepatic insufficiency	If CrCl < 10 mL/min: 2 gm (IV) q24h

**Drug Interactions:** Aminoglycosides (↓ aminoglycoside levels); methotrexate (↑ methotrexate levels); vecuronium (↑ vecuronium effect).

**Adverse Effects:** Drug fever/rash, E. multiforme/Stevens-Johnson Syndrome, anaphylactic reactions (hypotension, laryngospasm, bronchospasm), hives, serum sickness. C. difficile diarrhea/colitis.

**Allergic Potential:** High

**Safety in Pregnancy:** B

**Comments:** 20% of clavulanate removed by dialysis. Na<sup>+</sup> content = 4.75 mEq/g. K<sup>+</sup> content = 0.15 mEq/g.

**Cerebrospinal Fluid Penetration:** < 10%

**Bile Penetration:**

Without obstruction = 100%

With obstruction = 10%

(also see **Antibiotic Pearls & Pitfalls** p. 509).

#### REFERENCES:

- Donowitz GR, Mandell GL. Beta-lactam antibiotics. N Engl J Med 318:419–26 and 318:490–500, 1993.
- Graham DR, Talan DA, Nichols RL, et al. Once-daily, high-dose levofloxacin versus ticarcillin-clavulanate alone or followed by amoxicillin-clavulanate for complicated skin and skin-structure infections: a randomized, open-label trial. Clin Infect Dis 35:381–9, 2002.
- Itokazu GS, Danziger LH. Ampicillin-sulbactam and ticarcillin-clavulanic acid: A comparison of their in vitro activity and review of their clinical efficacy. Pharmacotherapy 11:382–414, 1991.
- Winston LG, Charlebois ED, Pang S, et al. Impact of a formulary switch from ticarcillin-clavulanate to piperacillin-tazobactam on colonization with vancomycin-resistant enterococci. Am J Infect Control 32:462–9, 2004.
- Website: www.pdr.net

## Tigecycline (Tygacil)

**Drug Class:** Glycylcycline.

**Usual Dose:** **Loading Dose:** 100 mg (IV) × 1 dose, (PI dose) **or 200 mg (IV) × 1 dose, (PK dose) then Maintenance**

**Dose:** 50 mg (IV) q12h (PI dosing) **or 100 mg (IV) q24h (PK dosing) (see comments<sup>†</sup>).**

**Spectrum:** (see **Susceptibility Profiles** pp. 198–202).

**Resistance Potential:** Low (Acinetobacter baumannii—dose related)

**Pharmacokinetic Parameters:**

Peak serum level: 1.45 mcg/ml (100 mg dose); 0.87 mcg/ml (50 mg dose)

Bioavailability: Not applicable

Excreted unchanged (urine): 22%

Serum half-life (normal/ESRD): 42/42 hrs

Plasma protein binding: 89%

Volume of distribution ( $V_d$ ): 8 L/kg

**Primary Mode of Elimination:** Biliary  
**Dosage Adjustments\***

PI = Package insert

PK = Pharmacokinetic

CrCl < 10 mL/min	No change
Post-HD dose	None
Post-PD dose	None
CVWH/CVVHD/ CVVHDF dose	No change
Mild-moderate hepatic insufficiency	No change
Severe hepatic insufficiency (Child Pugh C)	100 mg (IV) × 1 dose, then 25 mg (IV) q12h

**Drug Interactions:** Warfarin (↑ INR). Does not inhibit and is not metabolized by CYP450.

**Adverse Effects:** N/V, dyspepsia, diarrhea, dizziness, asthenia, thrombocytosis, ↑ alkaline phosphatase, pancreatitis, (↑ amylase/lipase), ↑ LDH, ↑ BUN, ↓ total protein.

**Allergic Potential:** Low

**Safety in Pregnancy:** D

**Comments:** Tetracycline or doxycycline susceptibilities not predictive of tigecycline susceptibilities. If any question re: tetracycline/doxycycline susceptibility, request tigecycline Ettest susceptibilities.

**Higher loading (LD)/Maintenance doses (MD) suggested for serious systemic infections or infection due to MDR *Klebsiella pneumoniae*, CRE, or MDR *Acinetobacter baumannii*.**\*<sup>§†</sup> **Following the initial Loading Dose** (based on PK parameters), **tigecycline may also be given as a single daily (q24h) Maintenance Dose** (since  $t_{1/2} = 42$  hrs). **Misses only *P. aeruginosa* and most *Proteus* sp., but effective against all other GNBs, MSSA/MRSA, VSE/VRE, and *B. fragilis*.** Tigecycline rarely causes N/V if sufficiently diluted and slowly infused. If N/V occurs, ↑ infusion volume and infusion time as shown

Tigecycline Dose <sup>†</sup>	Volume	Infusion Time
100 mg	100 ml	30 min
200 mg*	250 ml	60 min
400 mg <sup>§</sup>	500 ml	120 min

\* For systemic infections.

§ For serious systemic infections or UTIs due to susceptible MDR GNBs.

† Maintenance dose → half of the loading dose.

**Cerebrospinal Fluid Penetration:** 8%

**Bile Penetration:** 3800%

(also see **Antibiotic Pearls & Pitfalls** p. 518).

#### REFERENCES:

- Babinchak T, Ellis-Grosse E, Dartois N, et al. The efficacy and safety of tigecycline for the treatment of complicated intra-abdominal infections: analysis of pooled clinical trial data. *Clin Infect Dis* 41(Suppl 5:S3):54–67, 2005.
- Baines SD, Saxton K, Freeman K, et al. Tigecycline does not induce proliferation or cytotoxin production by epidemic *Clostridium difficile* strains in a human gut model. *J Antimicrob Chemother* 58:1062–1065, 2006.
- Balandin MB, Fernandez SI, Pintado GV, et al. Tigecycline therapy for infections due to carbapenemase-producing *Klebsiella pneumoniae* in critically ill patients. *Scand J Infect Dis*. 46:175–180, 2014.
- Bendu C, Culebras E, Gomez M, et al. In vitro activity of tigecycline against *Bacteroides* species. *J Antimicrob Chemother* 56:349–52, 2005.
- Bergallo C, Josovich A, Teglia O, et al. Safety and efficacy of intravenous tigecycline in Treatment of community-acquired pneumonia: results from a double-blind randomized Phase 3 comparison study with levofloxacin. *Diagn Microbiol Infect Dis* 63:52–61, 2009.
- Betts JW, Phee LM, Woodford N, et al. Activity of colistin in combination with tigecycline or rifampicin against multidrug-resistant *Stenotrophomonas maltophilia*. *Eur J Clin Microbiol Infect Dis* 33:1565–1572, 2014.
- Bhattacharya I, Gotfried MH, Ji AJ, et al. Reassessment of tigecycline bone concentrations in volunteers

Particularly effective monotherapy for serious systemic infections with minimal no resistance potential (not due to *P. aeruginosa* or *Proteus* sp.) Prevents *C. difficile* and effective against *C. difficile* diarrhea/colitis.

- undergoing elective orthopedic procedures. *J Clin Pharm.* 54:70-74, 2014.
- Bopp LH, Baltch AL, Ritz WJ, et al. Activities of tigecycline and comparators against *Legionella pneumophila* and *Legionella micdadei* extracellularly and in human monocytederived macrophages. *Diagn Microbiol Infect Dis* 69:86-93, 2011.
- Bouza E. New therapeutic choices for infections caused by methicillin-resistant *Staphylococcus aureus*. *Clin Microbiol Infect Dis* 15:44-52, 2009.
- Bulik CC, Wiskirchen DE, Shepard A, et al. Tissue penetration and pharmacokinetics of tigecycline in diabetic patients with chronic wound infections described by using in vivo microdialysis. *Antimicrob Agents Chemother* 54:5209-4213, 2010.
- Cai Y, Wang R, Liang B, et al. Systematic review and meta-analysis of the effectiveness and safety of tigecycline for treatment of infectious disease. *Antimicrob Agents Chemother* 55:1162-1172, 2011.
- Cheong EY, Gottlieb T. Intravenous tigecycline in the treatment of severe recurrent *Clostridium difficile* colitis. *Med J. Aust* 4:374-375, 2011.
- Conde-Estévez D, Grau S, Horcajada JP, et al. Off-label prescription of tigecycline: clinical and microbiological characteristics and outcomes. *Int J Antimicrob Agents* 36:471-472, 2010.
- Cunha BA, McDermott B, Nausheen S. Single once daily high dose therapy of tigecycline in a urinary tract infection due to multi-drug resistant (MDR) *Klebsiella pneumoniae* and MDR *Enterobacter agglomerans*. *Journal of Chemotherapy* 19:753-754, 2008.
- Cunha BA. Once daily tigecycline therapy of multidrug-resistant and non-multidrug resistant gram-negative bacteremias. *J of Chemother* 19:232-33, 2007.
- Cunha BA. Antimicrobial therapy of multidrug-resistant *Streptococcus pneumoniae*, vancomycin-resistant enterococci, and methicillin-resistant *Staphylococcus aureus*. *Med Clin N Am* 90:1165-82, 2006.
- Daly MW, Riddle DJ, Ledebor NA, et al. Tigecycline for treatment of pneumonia and empyema caused by carbapenems-producing *Klebsiella pneumoniae*. *Pharmacotherapy* 27:1052-1057, 2007.
- Dandache P, Nicolau DP, Sakoulas G. Tigecycline for the treatment of multidrug-resistant *Klebsiella pneumoniae* meningitis. *Infect Dis in Clin Practice* 17:66-68, 2009.
- Dartois N, Castiang N, Gandjini H, et al. Tigecycline versus levofloxacin for the treatment of community-acquired pneumonia: European experience. *J Chemother* 20:28-35, 2008.
- Dizbay M, Kilic S, Hizel K, et al. Tigecycline: its potential for treatment of brucellosis. *Scand J Infect Dis* 39:432-434, 2007.
- Dominguez EA. Single agent therapy with tigecycline in the treatment of complicated skin and skin structure and complicated intraabdominal infections. *Infectious Diseases in Clinical Practice* 17:144-149, 2009.
- Ellis-Grosse EJ, Babinchak T, Dartois N, et al. The efficacy and safety of tigecycline in the treatment of skin and skin-structure infections: results of 2 double-blind phase 3 comparison studies with vancomycin-aztreonam. *Clin Infect Dis* 41(Suppl 5): S341-53, 2005.
- Fraise AP. Tigecycline: The answer to beta-lactam and fluoroquinolone resistance? *Journal of Infection* 53:293-300, 2006.
- Gardiner D, Dukart G, Cooper A, et al. safety and efficacy of intravenous tigecycline in subjects with secondary bacteremia: pooled results from 8 phase III clinical trials. *Clin Infect Dis* 50:229-238, 2010.
- Geerlings SE, van Donselaar-van der Pant KA, Keur I. Successful treatment with tigecycline of two patients with complicated urinary tract infections caused by extended-spectrum beta-lactamase-producing *Escherichia coli*. *J Antimicrob Chemother* 65:2048-2049, 2010.
- Hawkey P, Finch R. Tigecycline: in-vitro performance as a predictor of clinical efficacy. *Clin Microbiol Infect* 13:354-362, 2007.
- Hawser SP. Activity of tigecycline against recent European clinical isolates of *Clostridium difficile*. *Int J Antimicrob Agents* 35:97-98, 2010.
- Herperts BL, Vlamincx B, Burkhardt O, et al. Intravenous tigecycline as adjunctive or alternative therapy for severe refractory *Clostridium difficile* infection. *Clin Infect Dis* 48:1732-1735, 2009.
- Hoban DJ, Bouchillon SK, Dowdzicky MJ. Antimicrobial susceptibility of extended-spectrum beta-lactamase producers and multidrug-resistant *Acinetobacter baumannii* throughout the United States and comparative in vitro activity of tigecycline, a new glycolcylcine antimicrobial. *Diagn Microbiol Infect Dis* 57:423-428, 2007.
- Hope R, Mushtaq S, James D, et al. Tigecycline activity: low resistance rates but problematic disc breakpoints revealed by a multicentre sentinel survey in the UK. *J Antimicrob Chemother* 65:2602-2609, 2010.
- Huang V, Cheung CM, Kaatz GW, et al. Evaluation of dalbavancin, tigecycline, minocycline, tetracycline, teicoplanin and vancomycin against community-associated and multidrug-resistant hospital-associated

- meticillin-resistant *Staphylococcus aureus*. *Int J Antimicrob Agents* 35:25–29, 2010.
- Hunte CA, Abbo LM. Skin and soft tissue infections due to methicillin-resistant *Staphylococcus aureus*: Role of tigecycline. *Clin Infect Dis* 52:1468–1467, 2011.
- Jenkins I. Linezolid- and vancomycin-resistant *Enterococcus faecium* endocarditis: successful treatment with tigecycline and daptomycin. *J Hosp Med* 2:343–344, 2007.
- Jump RL, Li Y, Pultz, JF, et al. Tigecycline exhibits inhibitory activity against *Clostridium difficile* in the colon of mice and does not promote growth or toxin production. *Antimicrob. Agents Chemother* 55:546–549, 2011.
- Kuo SC, Wang FD, Fung CP, et al. Clinical experience with tigecycline as treatment for serious infections in elderly and critically ill patients. *J Microbiol Immunol Infect* 44:45–51, 2011.
- Kwon SH, Ahn HL, Han OY, et al. Efficacy and safety profile comparison of colistin and tigecycline on the extensively drug resistant *Acinetobacter baumannii*. *Biol Pharm Bull.* 37:340–360, 2014.
- Larson KC, Belliveau PP, Spooner LM. Tigecycline for the treatment of severe *Clostridium difficile* infection. *Ann Pharm* 45:1005–1010, 2011.
- Lauf L, Ozsvar Z, Mitha I, et al. Phase 3 study comparing tigecycline and ertapenem in patients with diabetic foot infections with and without osteomyelitis. *Diag Microbiol Infect Dis.* 78:469–480, 2014.
- Lu CL, Liu CY, Liao CH, et al. Severe and refractory *Clostridium difficile* infection successfully treated with tigecycline and metronidazole. *Int J Antimicrob Agents* 35:311–312, 2010.
- Marchaim D, Pogue JM, Tzuman O, et al. Major variation in MICs of tigecycline in gram-negative bacilli as a function of testing method. *J Clin Microbiol.* 52:1617–1621, 2014.
- McKeage K, Keating GM. Tigecycline: in community-acquired pneumonia. *Drugs* 68:2633–2644, 2008.
- Meagher AK, Ambrose PG, Grasela TH, et al. The pharmacokinetic and pharmacodynamic profile of tigecycline. *Clin Infect Dis* 41(Suppl 5):S333–40, 2005.
- Metan G, Alp E, Yildiz O, et al. Clinical experience with tigecycline in the treatment of carbapenem-resistant *Acinetobacter* infections. *J Chemother* 22:110–114, 2010.
- Nilsson LE, Frimodt-Moller N, Vaara M, et al. Comparative activity of tigecycline and tetracycline on Gram-negative and Gram-positive bacteria revealed by a multicentre study in four North European countries. *Scand J Infect Dis* 43:707–713, 2011.
- Nix DE, Matthias KR. Should tigecycline be considered for urinary tract infections? A pharmacokinetic re-evaluation. *J Antimicrob Chemother* 65:1311–1312, 2010.
- Nord CE, Sillerstrom E, Wahlund E. Effect of Tigecycline on Normal Oropharyngeal and Intestinal Microflora. *Antimicrob Agents Chemother* 50:3375–80, 2006.
- Noviello S, Ianniello F, Leone S, et al. In vitro activity of tigecycline: MICs, MBCs, time-kill curves and post-antibiotic effect. *J Chemotherapy* 20:577–580, 2008.
- Ozbek B, Sentruk A. Postantibiotic effects of tigecycline, colistin sulfate, and levofloxacin alone or tigecycline-colistin sulfate and tigecycline-levofloxacin combinations against *Acinetobacter baumannii*. *Chemotherapy* 56:466–471, 2010.
- Pai MP. Serum and urine pharmacokinetics of tigecycline in obese class III and normal weight adults. *J Antimicrob Chemother.* 69:190–199, 2014.
- Pallotto C, Fiorio M, D'Avolio A, et al. Cerebrospinal fluid penetration of tigecycline. *Scand J Infect Dis.* 46:69–72, 2014.
- Pankey GA. Tigecycline. *J Antimicrob Chemother* 53:470–80, 2005.
- Pavan M, Chaudhari AP, Ranganth R. Altered bioavailability of tacrolimus following intravenous administration of tigecycline. *Am J Kidney Dis*57:345, 2011.
- Polidori M, Nuccorini A, Tascini C, et al. Vancomycin-resistant *Enterococcus faecium* (VRE) bacteremia in infective endocarditis successfully treated with combination daptomycin and tigecycline. *J Chemother* 23:240–241, 2011.
- Qvist N, Warren B, Leister-Tebbe H, et al. Efficacy of tigecycline versus ceftriaxone plus metronidazole for the treatment of complicated intra-abdominal infections: results from a randomized, controlled trial. *Surg Infect* 13:102–3, 2012.
- Raad I, Hanna H, Jiang Y, et al. Comparative activities of daptomycin, linezolid, and tigecycline against catheter-related methicillin-resistant *Staphylococcus bacteremic* isolates embedded in biofilm. *Antimicrob Agents Chemother* 51:1656–1660, 2007.
- Rodvold KA, Gotfried MH, Cwik M, et al. Serum, tissue and body fluid concentrations of tigecycline after a single 100 mg dose. *J Antimicrob Chemother* 58:1221–1229, 2006.

“Usual dose” assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

- Sands M, McCarter Y, Sanchez W. Synergy testing of multidrug resistant *Acinetobacter baumannii* against tigecycline and polymyxin using an E-test methodology. *Eur J Clin Microbiol Infect Dis* 26:521–522, 2007.
- Schwab KS, Hahn-Ast C, Heinz WJ, et al. Tigecycline in febrile neutropenic patients with haematological malignancies: a retrospective case documentation in four university hospitals. *Infection* 42:97–104, 2014.
- Slover CM, Rodvold KA, Danziger LH. Tigecycline: a novel broad-spectrum antimicrobial. *Ann Pharmacother* 41:965–972, 2007.
- Stein GE, Craig WA. Tigecycline: A Critical Analysis. *Clin Infect Dis* 43:518–24, 2006.
- Stein GE, Smith C, Nicolau DP, et al. Treatment of *Clostridium difficile* infection with tigecycline. *Infectious Diseases in Clinical Practice*. 22:216–218, 2014.
- Swoboda S, Ober M, Hainer C, et al. Tigecycline for the treatment of patients with severe sepsis of septic shock: a drug use evaluation in a surgical intensive care unit. *J Antimicrob Chemother* 61:729–733, 2008.
- Taccone FS, Rodriguez-Villalobos H, DeBacker D, et al. Successful treatment of septic shock due to pan-resistant *Acinetobacter baumannii* using combined antimicrobial therapy including tigecycline. *Eur J Clin Microbiol Infect Dis* 25:257–60, 2006.
- Valve K, Vaalasti A, Anttila VJ, et al. Disseminated *Legionella pneumophila* infection in an immunocompromised patient treated with tigecycline. *Scand J Infect Dis* 42:152–155, 2010.
- Vardakas KZ, Rafailidis PL, Falagas ME. Effectiveness and safety of tigecycline: focus on use for approved indications. *Clin Infect Dis* 54:4672–4, 2012.
- Woodford N, Hill RL, Livermore DM. In vitro activity of tigecycline against carbapenem-susceptible and resistant isolates of *Klebsiella* spp. and *Enterobacter* spp. *J Antimicrob Chemother* 59:582–583, 2007.
- Xie J, Wang T, Sun J, et al. Optimal tigecycline dosage regimen is urgently needed: results from a pharmacokinetic/pharmacodynamic analysis of tigecycline by Monte Carlo simulation. *Int J Infect Dis* 18:62–67, 2014.
- Yaha D, Lador A, Paul M, et al. Tigecycline and overall mortality. *Clin Infect Dis* 55:1739, 2012.

- Zinner SH. Overview of antibiotic use and resistance: setting the stage for tigecycline. *Clin Infect Dis* 41(Suppl 5):S289–92, 2005.  
Website: www.pdr.net

## Tipranavir (Aptivus) TPV

**Drug Class:** HIV protease inhibitor.

**Usual Dose:** 500 mg (PO) q12h with ritonavir 200 mg (PO) q12h.

**Pharmacokinetic Parameters:**

*Peak serum level:* 77–94 mcg/mL

*Bioavailability:* No data

*Excreted unchanged (urine):* 44%

*Serum half-life (normal/ESRD):* 5.5–6/5.5–6 hrs

*Plasma protein binding:* 99.9%

*Volume of distribution (V<sub>d</sub>):* 7–10 L/kg

**Primary Mode of Elimination:** Hepatic

**Dosage Adjustments\***

CrCl < 10 mL/min	No change
Post-HD dose	No change
Post-PD dose	No change
CVWH/CVWH/D/ CVHDF dose	No change
Mild hepatic insufficiency	No change
Moderate—severe hepatic insufficiency	Avoid

**Drug Interactions:** Metabolized via CYP3A4 rifabutin (↑ levels), clarithromycin (↑ levels), loperamide (↓ levels), statins (↑ risk of myopathy); abacavir, saquinavir, tenofovir, zidovudine, amprenavir/RTV, lopinavir/RTV (↓ levels). Aluminum/magnesium antacids (↓ absorption 25–30%). Ritonavir (↑ risk of hepatitis). St. John's Wort (↓ tipranavir levels). Fluconazole (↑ tipranavir levels).

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

**Contraindications:** simvastatin, atorvastatin, lovastatin, rifapentine

**Adverse Effects:** Contraindicated in moderate/severe hepatic insufficiency. ↑ risk of hepatotoxicity in HIV patients co-infected with HBV/HCV.

Intracerebral hemorrhage—use with caution in patients with coagulopathies.

**Allergic Potential:** High. Tipranavir is a sulfonamide; use with caution in patients with sulfonamide allergies

**Safety in Pregnancy:** C

**Comments:** Should be taken with food. Keep refrigerated 2–8°C.

**simvastatin, atorvastatin, lovastatin, rifapentine** Increased bioavailability when taken with meals. Must be co-administered with 200 mg ritonavir. Tipranavir contains a sulfa moiety (as do darunavir and fosamprenavir).

**Cerebrospinal Fluid Penetration:** No data

#### REFERENCES:

- Croom KF, Keam SJ. Tipranavir: a ritonavir-boosted protease inhibitor. *Drugs* 65:1669–79, 2005.
- Kandula VR, Khanlou H, Farthing C. Tipranavir: a novel second-generation nonpeptidic protease inhibitor. *Expert Rev Anti Infect Ther* 3:9–21, 2005.
- Kashuba AD. Drug-drug interactions and the pharmacotherapy of HIV infection. *Top HIV Med* 13:64–9, 2005.
- Panel on Antiretroviral Guidelines for Adults and Adolescents. Department of Health and Human Services. Guidelines for the use of antiretroviral agents in HIV-infected adults and adolescents. <http://aidsinfo.nih.gov/ConsentFiles/AdultandAdolescentGL.pdf>, 2012.
- Posker GL, Figgitt DP. Tipranavir. *Drugs* 63:1611–8, 2003.
- Thompson MA, Aberg JA, Cahn P, et al. Antiretroviral treatment of adult HIV infection: 2012 recommendations of the International Antiviral Society — USA panel. *JAMA* 308:387–402, 2012.
- Website: [www.pdr.net](http://www.pdr.net)

## Tobramycin (Nebcin)

**Drug Class:** Aminoglycoside.

**Usual Dose:** 240 mg or 5 mg/kg (IV) q24h (preferred over q8h dosing) (see comments).

**Spectrum:** (see **Susceptibility Profiles** pp. 198–202).

**Resistance Potential:** High (P. aeruginosa, aerobic GNBs)

Pharmacokinetic Parameters:

*Peak serum levels:* 4–8 mcg/ml (q8h dosing);

16–24 mcg/ml (q24h dosing)

*Bioavailability:* Not applicable

*Excreted unchanged (urine):* 95%

*Serum half-life (normal/ESRD):* 2.5/56 hrs

*Plasma protein binding:* 10%

*Volume of distribution (V<sub>d</sub>):* 0.24 L/kg

**Primary Mode of Elimination:** Renal  
**Dosage Adjustments\***

CrCl 50–80 mL/min	120 mg (IV) q24h or 2.5 mg/kg (IV) q24h
CrCl 10–50 mL/min	120 mg (IV) q48h or 2.5 mg/kg (IV) q48h
CrCl < 10 mL/min	60 mg (IV) q48h or 1.25 mg/kg (IV) q48h
Post-HD dose	80 mg (IV) or 1 mg/kg (IV)
Post-HFHD dose	120 mg (IV) or 2.5 mg/kg (IV)
Post-PD dose	40 mg (IV) or 0.5 mg/ kg (IV) or 2–4 mg/L in dialysate q24h
CVWH/CVWHD/ CVWHDF dose	120 mg (IV) q48h or 2.5 mg/kg (IV)
Mild-moderate hepatic insufficiency	No change
Severe hepatic insufficiency	No change

\*Usual dose\* assumes normal renal/hepatic function. \*For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.



**Drug Interactions:** Amphotericin B, cyclosporine, enflurane, methoxyflurane, NSAIDs, polymyxin B, radiographic contrast, vancomycin (↑ nephrotoxicity); cis-platinum (↑ nephrotoxicity, ↑ ototoxicity); loop diuretics (↑ ototoxicity); neuromuscular blocking agents (↑ apnea, prolonged paralysis); non-polarizing muscle relaxants (↑ apnea).

**Adverse Effects:** Neuromuscular blockade with rapid infusion/absorption. Nephrotoxicity only with prolonged/extremely high serum trough levels; may cause reversible non-oliguric renal failure (ATN). Ototoxicity associated with prolonged/extremely high peak serum levels (usually irreversible): Cochlear toxicity (1/3 of ototoxicity) manifests as decreased high frequency hearing, but deafness is unusual. Vestibular toxicity (2/3 of ototoxicity) develops before ototoxicity, and typically manifests as tinnitus.

**Allergic Potential:** Low

**Safety in Pregnancy:** D

**Comments:** Single daily dosing greatly reduces nephrotoxic/ototoxic potential.

Incompatible with solutions containing  $\beta$ -lactams, erythromycin, chloramphenicol, furosemide, sodium bicarbonate. IV infusion should be given slowly over 1 hour. May be given IM. **Avoid intraperitoneal infusion due to risk of neuromuscular blockade. Avoid intratracheal/aerosolized intrapulmonary instillation, which predisposes to antibiotic resistance.**  $V_d$  increases with edema/ascites, trauma, burns, cystic fibrosis; may require ↑ dose.  $V_d$  decreases with dehydration, obesity; may require ↓ dose. **Renal cast counts are the best indicator of aminoglycoside nephrotoxicity, not serum creatinines.** Dialysis removes ~ 1/3 of tobramycin from serum. Tobramycin nebulizer dose: 300 mg via nebulizer q12h (**not recommended due to ↑ risk of resistance**).

**CAPD dose:** 2–4 mg/L in dialysate (I.P.) with each exchange.

**Therapeutic Serum Concentrations**

(for therapeutic efficacy, *not toxicity*):

Peak (q24h/q8h dosing) = 16–24/8–10 mcg/ ml  
Trough (q24h/q8h dosing) = 0/1–2 mcg/ml

**Synergy dose:** 120 mg (IV) q24h or 2.5 mg/kg (IV) q24h

**Intrathecal (IT) dose:** 5 mg (IT) q24h

**Cerebrospinal Fluid Penetration:**

Non-inflamed meninges = 0%

Inflamed meninges = 20%

**Bile Penetration:** 30%

(also see *Antibiotic Pearls & Pitfalls* p. 514).

#### REFERENCES:

- Begg EJ, Barclay ML. Aminoglycosides - 50 years on. *Br J Clin Pharmacol* 39:597–603, 1995.
- Buijk SE, Mouton JW, Gyssens IC, et al. Experience with a once-daily dosing program of aminoglycosides in critically ill patients. *Intensive Care Med* 28:936–42, 2002.
- Cheer SM, Waugh J, Noble S. Inhaled tobramycin (TOBI): a review of its use in the management of *Pseudomonas aeruginosa* infections in patients with cystic fibrosis. *Drugs* 63:2501–20, 2003.
- Cunha BA. Aminoglycosides: Current role in antimicrobial therapy. *Pharmacotherapy* 8:334–50, 1988.
- Dupuis LL, Sung L, Taylor T, et al. Tobramycin pharmacokinetics in children with febrile neutropenia undergoing stem cell transplantation once-daily versus thrice-daily administration. *Pharmacotherapy* 24:564–73, 2004.
- Edson RS, Terrel CL. The aminoglycosides. *Mayo Clin Proc* 74:519–28, 1999.
- Geller DE, Pistick WH, Nardella PA, et al. Pharmacokinetics and bioavailability of aerosolized tobramycin in cystic fibrosis. *Chest* 122:219–26, 2002.
- Gilbert DN. Once-daily aminoglycoside therapy. *Antimicrob Agents Chemother* 35:399–405, 1991.
- Hustinx WN, Hoepelman IM. Aminoglycoside dosage regimens: Is once a day enough? *Clin Pharmacokin* 25:427–32, 1993.
- Kahler DA, Schowengerdt KO, Fricker FJ, et al. Toxic serum trough concentrations after administration of nebulized tobramycin. *Pharmacotherapy* 23:543–5, 2003.
- Lortholary O, Tod M, Cohen Y, et al. Aminoglycosides. *Med Clin North Am* 79:761–87, 1995.
- McCormack JP, Jewesson PJ. A critical reevaluation of the "therapeutic range" of aminoglycosides. *Clin Infect Dis* 14:320–39, 1992.
- Moss RB. Long-term benefits of inhaled tobramycin in adolescent patients with cystic fibrosis. *Chest* 121:55–63, 2002.

Whitehead A, Conway SP, Etherington C, et al. Once-daily tobramycin in the treatment of adult patients with cystic fibrosis. *Eur Respir J* 19:303–9, 2002.

Wood GC, Chapman JL, Boucher BA, et al. Tobramycin bladder irrigation for treating a urinary tract infection in a critically ill patient. *Ann Pharmacother* 38:1318–9, 2004.

Website: [www.pdr.net](http://www.pdr.net)

## Trimethoprim (Proloprim, Trimpex) TMP

**Drug Class:** Folate antagonist.

**Usual Dose:** 100 mg (PO) q12h (see comments).

**Spectrum:** (see *Susceptibility Profiles* pp. 186–190).

**Resistance Potential:** High (*S. pneumoniae*, *E. coli*)

**Pharmacokinetic Parameters:**

*Peak serum level:* 2–8 mcg/ml

*Bioavailability:* 98%

*Excreted unchanged (urine):* 67%

*Serum half-life (normal/ESRD):* 8/24 hrs

*Plasma protein binding:* 44%

*Volume of distribution ( $V_d$ ):* 1.8 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\***

CrCl 10–30 mL/min	50 mg (PO) q12h
CrCl < 10 mL/min	Avoid (except for PCP, see TMP–SMX)
Post–HD dose	Avoid (except for PCP, see TMP–SMX)
Post–PD dose	100 mg (PO)
CVVH/CVVHD/ CVHDF dose	50 mg (PO) q12h
Moderate—severe hepatic insufficiency	No change

**Drug Interactions:** Azathioprine (leukopenia); amantadine, dapsone, digoxin, methotrexate, phenytoin, rifampin, zidovudine (↑ interacting drug levels, nystagmus with phenytoin); diuretics (↑ serum K<sup>+</sup> with K<sup>+</sup>-sparing diuretics, ↓ serum Na<sup>+</sup> with thiazide diuretics); warfarin (↑ INR, bleeding).

**Adverse Effects:** Folate deficiency, hyperkalemia.

**Allergic Potential:** Low

**Safety in Pregnancy:** C

**Comments:** Useful in sulfa-allergic patients unable to take TMP–SMX. **PCP**

**dose:** 5 mg/kg (PO) q8h plus dapsone 100 mg (PO) q24h.

**Meningeal Dose:** 300 mg or 5 mg/kg (PO) q6h.

**Cerebrospinal Fluid Penetration:** 40%

**Bile Penetration:** 100%

(also see *Antibiotic Pearls & Pitfalls* p. 515).

### REFERENCES:

Brogden RN, Carmine AA, Heel RC, et al. Trimethoprim:

A review of its antibacterial activity, pharmacokinetics and therapeutic use in urinary tract infections. *Drugs* 23:405–30, 1982.

Friesen WT, Hekster YA, Vree TB. Trimethoprim: clinical use and pharmacokinetics. *Drug Intelligence & Clinical Pharmacy* 15:325–30, 1981.

Neu HC. Trimethoprim alone for treatment of urinary tract infection. *Rev Infect Dis* 4:366–71, 1982.

Website: [www.pdr.net](http://www.pdr.net)

## Trimethoprim–Sulfamethoxazole (Bactrim, Septra) TMP–SMX

**Drug Class:** Folate antagonist/sulfonamide.

**Usual Dose:** 2.5–5 mg/kg (IV/PO) q6h.

**Spectrum:** (see *Susceptibility Profiles* pp. 186–190).

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

**Resistance Potential:** High (*S. pneumoniae*, *E. coli*)

**Pharmacokinetic Parameters:**

Peak serum level: 2–8/40–80 mcg/ml

Bioavailability: 98%

Excreted unchanged (urine): 67/85%

Serum half-life (normal/ESRD): (10/8)/40–80 hrs

Plasma protein binding: 44–70%

Volume of distribution ( $V_d$ ): 1.8/0.3 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\***

CrCl 15–30 mL/min	1.25–2.5 mg/kg (IV/PO) q6h
CrCl < 15 mL/min	Avoid (except for PCP use 1.25–2.5 mg/kg [IV/PO] q8h)
Post-HD dose	2.5 mg/kg (IV/PO)
Post-PD dose	0.16 mg/kg (IV/PO)
CVWH/CVWHD/ CVWHDf dose	2.5 mg/kg (IV/PO) q6h
Moderate—severe hepatic insufficiency	No change

**Drug Interactions:** *TMP component:*

Azathioprine (leukopenia); amantadine, dapsone, digoxin, methotrexate, phenytoin, rifampin, zidovudine (↑ interacting drug levels, nystagmus with phenytoin); diuretics (↑ serum  $K^+$  with  $K^+$ -sparing diuretics, ↓ serum  $Na^+$  with thiazide diuretics); warfarin (↑ INR, bleeding). *SMX component:* Cyclosporine (↓ cyclosporine levels); phenytoin (↑ phenytoin levels, nystagmus, ataxia); methotrexate (↑ antifolate activity); sulfonyleureas, thiopental (↑ interacting drug effect); warfarin (↑ INR, bleeding).

**Adverse Effects:** Common cause of drug induced aseptic meningitis.

**TMP:** Folate deficiency, hyperkalemia.

**SMX:** Leukopenia, thrombocytopenia, hemolytic anemia ± G6PD deficiency, aplastic anemia, ↑ SGOT/SGPT, severe hypersensitivity reactions (*E. multiforme*/Stevens–Johnson Syndrome), ↑ risk of hypoglycemia in chronic renal failure.

**Allergic Potential:** Very high (SMX); none (TMP)

**Safety in Pregnancy:** C

**Comments:** Drug fever/rash

Excellent bioavailability (IV = PO).

**TMP–SMX IV and PO Equivalence:**

**TMP–SMX (TMP component) 10 mg/kg (IV)**

**q24h** (70 kg patient)

= 2 SS<sup>†</sup> tablets (PO) q6h

or

= 1 DS<sup>††</sup> tablet (PO) q6h

**TMP–SMX (TMP component) 20 mg/kg (IV)**

**q24h** (70 kg patient)

= 4 SS tablets (PO) q6h

or

= 2 DS tablets (PO) q6h

† 1 SS tablet = TMP 80 mg + SMX 400 mg.

†† 1 DS tablet = TMP 160 mg + SMX 800 mg.

**In sulfa allergic patients, use TMP (SMX is the allergic component) which is equally effective as TMP–SMX.**

Meningeal dose = 5 mg/kg (IV/PO) q6h.

**Cerebrospinal Fluid Penetration:**

Non-Inflamed meninges = 40%

Inflamed meninges = 40%

**Bile Penetration:** 100%

(also see **Antibiotic Pearls & Pitfalls** p. 515).

**REFERENCES:**

Chetchoisak P, Chierakul W, et al. Trimethoprim-sulfamethoxazole versus trimethoprim-sulfamethoxazole plus doxycycline as oral eradication treatment for melioidosis: a multicentre, double-blind,

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

- non-inferiority, randomized controlled trial. *Lancet*. 1:807-814, 2014.
- Cockerill FR, Edson RS. Trimethoprim-sulfamethoxazole. *Mayo Clin Proc* 66:1260-9, 1991.
- Cunha BA. Minocycline often forgotten but preferred to trimethoprim-Sulfamethoxazole (TMP-SMX) or Doxycycline for the treatment of Community Associated Methicillin-Resistant *S. aureus* Skin & Soft Tissue Infections. *Int Antimicrob Agents*. 42:497-799, 2013.
- Cunha BA. Oral antibiotic therapy of serious systemic infections. *Med Clin N Am* 90:1197-1222, 2006.
- Cunha BA. Prophylaxis for recurrent urinary tract infections: nitrofurantoin, not trimethoprim-sulfamethoxazole or cranberry juice. *Arch Intern Med* 172:82-3, 2012.
- Euga G, Murillo O, Fernandez-Sabe N, et al. Treatment of chronic staphylococcal osteomyelitis: intravenous or oral? how long? *Antimicrob Agents Chemother* 53:2672-2676, 2009.
- Gelfand MS, Cleveland KO, Ketterer DC. Susceptibility of *Streptococcus pyogenes* to trimethoprim-sulfamethoxazole. *J Clin Micro* 51:1350-1351, 2013.
- Masters PA, O'Bryan TA, Zurlo JM, et al. Trimethoprim-sulfamethoxazole revisited. *Arch Intern Med* 163:402-10, 2003.
- Muhammed SA, Rolain JM, Le Poullain MN. Orally administered trimethoprim-sulfamethoxazole for deep staphylococcal infections. *J Antimicrob Chemother*. 69:757-760, 2014.
- Nayak SU, Simon GL. Myocarditis after trimethoprim-sulfamethoxazole treatment for ehrlichiosis. *Emerg Inf Dis* 19:1975-1977, 2013.
- Pallin DJ, Binder WD, Allen MB, et al. Clinical trial: Comparative effectiveness of cephalexin plus trimethoprim-sulfamethoxazole versus cephalexin: treatment of uncomplicated cellulitis: randomized controlled trial. *Clin Infect Dis* 56:1754-1762, 2013.
- Sanchez GV, Master RN, Bordon J. Trimethoprim-sulfamethoxazole may no longer be acceptable for the treatment of acute uncomplicated cystitis in the United States. *Clin Infect Dis* 53:316-7, 2011.
- Slack III, D, Emmons W. A case of fulminant ehrlichiosis in the setting of recent trimethoprim-sulfamethoxazole use. *Inf Dis Clin Prac* 21:141-144, 2013.
- Soeter HM, Von Gottberg A, Cohen C, et al. Trimethoprim-sulfamethoxazole prophylaxis and antibiotic non-susceptibility invasive pneumococcal disease. *Antimicrob Agents Chemother* 56:1602-5, 2012.
- Website: [www.pdr.net](http://www.pdr.net)

## Valacyclovir (Valtrex)

**Drug Class:** HSV, VZV antiviral (see comments).

**Usual Dose:**

**HSV-1/2:** Herpes labialis: 2 gm (PO) q12h x 1

day. Genital herpes: *Initial therapy*: 1 gm (PO)

q12h x 3 days. *Recurrent/intermittent therapy*

(< 6 episodes/year): normal host: 500 mg (PO)

q24h x 5 days; HIV-positive: 1 gm (PO) q12h x

7-10 days. *Chronic suppressive therapy*

(> 6 episodes/year): normal host: 1 gm (PO)

q24h x 1 year;

HIV-positive: 500 mg (PO) q12h x 1 year.

HSV-1 late onset VAP: 1 gm (PO) q8h x

10 days. Meningitis/encephalitis: **1 gm (PO) q8h**

x 10 days.\*

**VZV:** Chickenpox: 1 gm (PO) q8h x 5 days.

VZV pneumonia: 1-2 gm (PO) q8h x 10 days.

Herpes zoster (shingles) (dermatomal/

disseminated): 1 gm (PO) q8h x 7-10 days.

VZV meningitis/encephalitis: **2 gm (PO) q6h x**

10 days.\*

**Pharmacokinetic Parameters:**

*Peak serum level*: 3.7-5 mcg/ml

*Bioavailability*: 55%

*Excreted unchanged (urine)*: 1%

*Serum half-life (normal/ESRD)*: 3/14 hrs

*Plasma protein binding*: 15%

*Volume of distribution ( $V_d$ )*: 0.7 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\*** (based on 1 gm q8h)

\* Severe HSV encephalitis may require 14-21 days of therapy

CrCl 30–50 mL/min	1 gm (PO) q12h
CrCl 10–30 mL/min	1 gm (PO) q24h
CrCl < 10 mL/min	500 mg (PO) q24h
Post-HD dose	1 gm (PO)
Post-PD dose	500 mg
CWVH/CVVHD/ CWHDF dose	500 mg (PO) q24h
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	No change

**Drug Interactions:** Cimetidine, probenecid (↑ acyclovir levels).

**Adverse Effects:** Headache, nausea, diarrhea, abdominal pain, weakness. Rarely, HUS/TTP (only in HIV).

**Allergic Potential:** Low

**Safety in Pregnancy:** B

**Comments:** Converted to acyclovir in liver.

**Highly active against HSV > VZV. Some activity against CMV. No activity against EBV, RSV or adenoviruses.**

VZV IC<sub>90</sub> (mean) ~ 2 × HSV.

Meningitis/Encephalitis dose = HSV: 1 gm (PO) q8h  
VZV: 2 gm (PO) q6h

**Cerebrospinal Fluid Penetration:** 54%

#### REFERENCES:

- Acost EP, Fletcher CV. Valacyclovir. *Ann Pharmacotherapy* 31:185–91, 1997.
- Chaudhuri D, Ginn D. Valacyclovir-associated neurotoxicity in peritoneal dialysis patients. *Am J Ther* 21:2215–217, 2014.

Kotton CN, Kumar D, Caliendo AM, et al. International consensus guidelines on the management of cytomegalovirus in solid organ transplantation. *Transplantation* 89:779–795, 2010.

Mugwanya K, Baeten JM, Mugo NR, et al. High-dose valacyclovir HSV-2 suppression results in greater reduction in plasma HIV-1 levels compared with standard dose acyclovir among HIV-1/HSV-2 coinfecting persons: a randomized, crossover trial. *J Infect Dis* 204:1912–1017, 2011.

Pouplin T, Pouplin JN, Van Toi P, et al. valacyclovir for Herpes simplex encephalitis. *Antimicrob Agents Chemother* 55:3624–3626, 2011

Vigil KJ, Chemaly RF. Valacyclovir: approved and off-label uses for the treatment of herpes virus infections in immunocompetent and immunocompromised adults. *Expert Opin Pharmacother* 11:1901–1913, 2010.

Website: [www.pdr.net](http://www.pdr.net)

## Valganciclovir (Valcyte)

**Drug Class:** CMV, HSV, HHV-6 antiviral (see comments).

**Usual Dose: Induction Dose:** 900 mg (PO) q12h × 21 days, then **Maintenance Dose:** 900 mg (PO) q24h (Normal hosts: until cured; Compromised hosts: chronic suppressive therapy). 900 mg dose taken as two 450-mg tablets once daily.

**Pharmacokinetic Parameters:**

*Peak serum level:* 5.6 mcg/ml

*Bioavailability:* 59.4%

*Excreted unchanged (urine):* 90%

*Serum half-life (normal/ESRD):* 4.1/67.5 hrs

*Plasma protein binding:* 1%

*Volume of distribution (V<sub>d</sub>):* 15.3 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\***

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

CrCl 40–60 mL/min	450 mg (PO) q12h (induction), then 450 mg (PO) q24h (maintenance)
CrCl 25–40 mL/min	450 mg (PO) q24h (induction), then 450 mg (PO) q48h (maintenance)
CrCl 10–25 mL/min	450 mg (PO) q48h (induction), then 450 mg (PO) 2x/week (maintenance)
CrCl < 10 mL/min	Avoid
Post-HD dose	Avoid
Post-PD dose	Use same dose as CrCl 25–40 mL/min
CVWH/CVVDH/ CVHDF dose	Use same dose as CrCl 25–50 mL/min
Mild-moderate hepatic insufficiency	No change
Severe hepatic insufficiency	Use with caution

**Drug Interactions:** Cytotoxic drugs (may produce additive toxicity: stomatitis, bone marrow depression, alopecia); imipenem (↑ risk of seizures); probenecid (↑ valganciclovir levels); zidovudine (↓ valganciclovir levels, ↑ zidovudine levels, possible neutropenia); didanosine (↑ didanosine); cyclosporine amphotericin, (↑ nephrotoxicity); mycophenolate mofetil (↑ mycophenolate mofetil and gancyclovir levels in renal insufficiency); tenofovir (↑ tenofovir, gancyclovir levels).

**Adverse Effects:** Drug fever/rash, diarrhea, nausea, vomiting, leukopenia, anemia, thrombocytopenia, paresthesias/peripheral

neuropathy, retinal detachment, potential BM suppression.

**Allergic Potential:** Low

**Safety in Pregnancy:** C

**Comments:** Valganciclovir exposures (AUC) larger than for IV ganciclovir. Tablets should be taken with food. Valganciclovir is rapidly hydrolyzed to ganciclovir. Not interchangeable on a tablet-to-tablet basis with oral ganciclovir. Much higher bioavailability than ganciclovir capsules; serum concentration equivalent to IV ganciclovir. Indicated for induction/maintenance therapy of CMV infection. **Highly active against CMV, HHV-6, and HSV. Some activity against VZV, EBV and adenoviruses. No activity against RSV.**

Meningeal dose = usual dose.

**Cerebrospinal Fluid Penetration:** 70%

#### REFERENCES:

- Avery RK. Low-dose valganciclovir for cytomegalovirus phylaxis in organ transplantation: is less really more? *Clin Infect Dis* 52:322–324, 2011.
- Asberg A, Rollag H, Hartmann A. Valganciclovir for the prevention and treatment of CMV in solid organ transplant recipients. *Expert Opin Pharmacother* 11:1159–1166, 2010.
- Boillat Blanco N, Pascual M, Venetz JP, et al. Impact of preemptive strategy after 3 months of valganciclovir cytomegalovirus prophylaxis in kidney transplant recipients. *Transplantation* 91:251–255, 2011.
- Czock D, Scholle C, Rasche FM, et al. Pharmacokinetics of valganciclovir and ganciclovir in renal impairment. *Clin Pharmacol Ther* 72:142–50, 2002.
- Humar A, Limaye AP, Blumberg EA, et al. Extended valganciclovir prophylaxis in D+/R- kidney transplant recipients is associated with long-term reduction in cytomegalovirus disease: two-year results of the IMPACT study. *Transplantation* 90:1427–1431, 2010.
- Jacobsen T, Sipointis N. Drug interactions and toxicities associated with the antiviral management of cytomegalovirus infection. *Am J Health Syst Pharm* 67:1417–1425, 2010.

"Usual dose" assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

Kallil AC, Midru C, Florescu DF. Effectiveness of valganciclovir 900 mg versus 450 mg for cytomegalovirus prophylaxis in transplantation: direct and indirect treatment comparison meta-analysis. *Clin Infect Dis* 52:313–321, 2011.

Kotton CN, Kumar D, Caliendo AM, et al. International consensus guidelines on the management of cytomegalovirus in solid organ transplantation. *Transplantation* 89:779–795, 2010.

Lopau K, Greser A, Wanner C. Efficacy and safety of preemptive anti-CMV therapy with valganciclovir after kidney transplantation. *Clin Transplant* 21:80–85, 2007.

Luan, FL, Kommareddi M, Ojo AO. Impact of cytomegalovirus disease in D+/R- kidney transplant patients receiving 6 months low-dose valganciclovir prophylaxis. *Am J Transplant* 11:1936–1942, 2011.

Razonable RR, Paya CV. Valganciclovir for the prevention and treatment of cytomegalovirus disease in immunocompromised hosts. *Expert Rev Anti Infect Ther* 2:27–41, 2004.

Slifkin M, Doron S, Snyderman DR. Viral prophylaxis in organ transplant patients. *Drugs* 64:2763–92, 2004. Website: [www.pdr.net](http://www.pdr.net)

## Vancomycin (Vancocin)

**Drug Class:** Glycopeptide.

**Usual Dose:** 1 gm (IV) q12h (see comments).

**Spectrum:** (see **Susceptibility Profiles** pp. 198–202).

**Resistance Potential:** Low (MSSA, MRSA VSE; ↑ prevalence of VRE)

**Pharmacokinetic Parameters:**

*Peak serum level:* 63 mcg/ml

*Bioavailability:* IV (not applicable)/PO (0%)

*Excreted unchanged (urine):* 90%

*Serum half-life (normal/ESRD):* 6/180 hrs

*Plasma protein binding:* 55%

*Volume of distribution ( $V_d$ ):* 0.7 L/kg

**Primary Mode of Elimination:** Renal

**Dosage Adjustments\***

CrCl 50–80 mL/min	1 gm (IV) q12h
CrCl 10–50 mL/min	1 gm (IV) q24h
CrCl < 10 mL/min	1 gm (IV) q week
Post-HD dose	None
Post-HFHD dose	500 mg (IV)
Post-PD dose	None
CVWH/CVWHD/CVWHDf dose	1 gm (IV) q24h
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	No change

**Drug Interactions:** Aminoglycosides, amphotericin B, polymyxin B (↑ nephrotoxicity).

**Adverse Effects:** “Red man/neck syndrome” with rapid IV infusion (histamine mediated), leukopenia, thrombocytopenia, cardiac arrest, hypotension.

**Allergic Potential:** Low

**Safety in Pregnancy:** C

**Comments:** Not nephrotoxic. “Red man/neck syndrome” can be prevented/minimized by infusing IV vancomycin slowly over 1–2 hours. Intrapertoneal absorption = 40%. **IV vancomycin increases VRE prevalence. C. difficile diarrhea dose:** oral vancomycin 250 mg (PO) q6h. *If no response in 72 hours ↑ dose to 500 mg (PO) q6h to complete therapy. C. difficile diarrhea relapse dose:* 500 mg (PO) q6h × 1 month and reevaluate. **Do not taper dose.** Should C. difficile recur, treat × 2 or 3 months with 500 mg (PO) q6h. **C. difficile colitis: Oral vancomycin ineffective for C. difficile colitis.** Vancomycin doesn’t penetrate well into the CSF with non-inflamed

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

meninges. For therapeutic CSF concentrations, use IV vancomycin *plus* IT vancomycin.

### **Therapeutic Serum Concentrations**

(for therapeutic efficacy, *not* toxicity):

Peak = 25–40 mcg/ml

Trough = 5–12 mcg/ml

There are **no convincing data that vancomycin is ototoxic or nephrotoxic. CrCl, not serum levels, should be used to adjust vancomycin dosing.** Prolonged/high dose vancomycin (60 mg/kg/day or 2 gm [IV] q12h) has been useful in treating *S. aureus* osteomyelitis, *S. aureus* infections with high MICs (VISA), and infections in difficult-to-penetrate tissues without toxicity. In bone or CSF, vancomycin tissue concentrations are ~ 15% of serum levels.

**Intrathecal (IT) dose:** 20 mg (IT) in preservative free NaCl.

### **Cerebrospinal Fluid Penetration:**

Non-Inflamed meninges = <1%

Inflamed meninges = 15%

**Bile Penetration:** 50%

(also see **Antibiotic Pearls & Pitfalls** p. 517).

### **REFERENCES:**

Ajao AO, Harris AD, Roghmann MC, et al. Systematic review of measurement and adjustment for colonization pressure in studies of methicillin-resistant *Staphylococcus aureus*, vancomycin-resistant enterococci, and clostridium difficile acquisition. *Infect Control Hosp Epidemiol* 32:481–489, 2011.

Black E, Lau TT, Ensom MH. Vancomycin-induced neutropenia: is it dose- or duration-related? *Ann Pharmacother* 45:629–638, 2011.

Bhavani SM, Prakhya A, Hammel JP, et al. Cost-Effectiveness of daptomycin versus vancomycin and gentamicin for patients with methicillin-resistant *Staphylococcus aureus* bacteremia and/or endocarditis. *Clin Infect Dis* 49:691–698, 2009.

Bouza E. New therapeutic choices for infections caused by methicillin-resistant *Staphylococcus aureus*. *Clin Microbiol Infect Dis* 15:44–52, 2009.

Cantu TG, Yamanaka-Yuen NA, Lietman PS. Serum vancomycin concentrations: Reappraisal of their clinical value. *Clin Infect Dis* 18:533–43, 1994.

Cosgrove SE, Carroll KC, Perl TM. *Staphylococcus aureus* with reduced susceptibility to vancomycin. *Clin Infect Dis* 39:539–45, 2004.

Cruciani M, Gatti G, Lazzarini L, et al. Penetration of vancomycin into human lung tissue. *Journal of Antimicrobial Chemotherapy* 38:865–69, 1996.

Cunha BA. Vancomycin revisited: a reappraisal of clinical use. *Critical Care Clinics of North America* 24:394–420, 2008.

Cunha BA, Mohan SS, Hamid N, McDermott GP, Daniels P. Cost ineffectiveness of serum vancomycin levels. *Eur J Clin Microbiol Infect Dis* 13:509–511, 2007.

Cunha BA. Vancomycin serum levels: unnecessary, unhelpful, and costly. *Antibiotics for Clinicians* 8:273–77, 2004.

Cunha BA. Vancomycin. *Med Clin North Am* 79:817–31, 1995.

Cunha BA, Deglin J, Chow M, et al. Pharmacokinetics of vancomycin in patients undergoing chronic hemodialysis. *Rev Infect Dis* 3:269–72, 1981.

DelDot ME, Lipman J, Tett SE. Vancomycin pharmacokinetics in critically ill patients receiving continuous venovenous haemodiafiltration. *Br J Clin Pharmacol* 58:259–68, 2004.

Deresinski S. Vancomycin: does it still have a role as an antistaphylococcal agent? *Expert Rev Anti Infect Ther* 5:393–401, 2007.

El Amari EB, Vuagnat A, Stern R, et al. High versus standard dose vancomycin for osteomyelitis. *Scand J Infect Dis* 36:712–7, 2004.

Hidayat LK, Hsu DI, Quist R, et al. High-dose vancomycin therapy for methicillin-resistant *Staphylococcus aureus* infections. *Arch Intern Med* 166:2138–44, 2006.

Huttner A, Harbarth S. Guidelines for vancomycin use. *Clin Infect Dis* 50:616, 2010.

Leonard SN, Szeto YG, Zolotarev M, et al. Comparative in vitro activity of telavancin, vancomycin, and linezolid against heterogeneously vancomycin-intermediate *Staphylococcus aureus* (hVISA). *Int J Antimicrob* 37:558–561, 2011.

Menzies D, Goel K, Cunha BA. Vancomycin. *Antibiotics for Clinicians* 2:97–9, 1998.

Moellering RC Jr. Monitoring serum vancomycin levels: climbing the mountain because it is there? *Clin Infect Dis* 18:544–6, 1994.

<sup>†</sup>Usual dose<sup>†</sup> assumes normal renal/hepatic function. <sup>\*</sup>For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.



- Okamoto H. Vancomycin-induced immune thrombocytopenia. *N Engl J Med* 356:2537–2538, 2007.
- Patel N, Lubanski P, Ferro S, et al. Correlation between vancomycin MIC values and those of other agents against gram-positive bacteria among patients with bloodstream infections caused by methicillin-resistant *Staphylococcus aureus*. *Antimicrob Agents Chemother* 53:5141–5144, 2009.
- Pope SD, Roecker A. Vancomycin for treatment of invasive, multi-drug resistant *Staphylococcus aureus* infections. *Exp Opin Pharmacother* 8:1245–1261, 2007.
- Rangaraj G, Cleveland KO, Gelfand MS. Comparative analysis of daptomycin and vancomycin in the treatment of vertebral osteomyelitis. *Infect Dis Clin Pract* 22:2019–222, 2014.
- Ricard JD, Wolff M, Lacerade JC, et al. Levels of vancomycin in cerebrospinal fluid of adult patients receiving adjunctive corticosteroids to treat pneumococcal meningitis: a prospective multicenter observational study. *Clin Infect Dis* 44:250–255, 2007.
- Rybak MJ. The pharmacokinetic and pharmacodynamic properties of vancomycin. *Clin Infect Dis* 42:S35–9, 2006.
- Saribas S, Bagdatli Y. Vancomycin tolerance in enterococci. *Chemotherapy* 50:250–4, 2004.
- Stevens DL. The role of vancomycin in the treatment paradigm. *Clin Infect Dis* 42:S51–7, 2006.
- Strykowski ME, Szczech LA, Benjamin DK Jr, et al. Use of vancomycin or first-generation cephalosporins for the treatment of hemodialysis-dependent patients with methicillin-susceptible *Staphylococcus aureus* bacteremia. *Clin Infect Dis* 44:190–196, 2007.
- Tenover FC, Moellering RC. The rationale for revising the clinical and laboratory standards institute vancomycin minimal inhibitory concentration interpretive criteria for *Staphylococcus aureus*. *Clin Infect Dis* 44:1208–1215, 2007.
- Vandecasteele SJ, De Vriese AS. Vancomycin dosing in patients on intermittent hemodialysis. *Semin Dial* 24:50–55, 2011.
- Von Drygalski A, Curtis BR, Bougie DW, et al. Vancomycin-induced immune thrombocytopenia. *N Engl J Med* 356:904–10, 2007.
- Wang G, Hindler JF, Ward KW, et al. Increased vancomycin MICs for *Staphylococcus aureus* clinical isolates from a university hospital during a 5-year period. *Journal of Clinical Microbiology* 44:3883–86, 2006.
- Wombwell E, Young N. Intrathecal and Intraventricular Vancomycin. *Infect Dis Clin Pract* 22:141–144, 2014. Website: [www.pdr.net](http://www.pdr.net)

## Voriconazole (Vfend)

**Drug Class:** Triazole antifungal.

**Usual Dose:** IV dosing: **Loading dose:**

6 mg/kg (IV) q12h × 1 day, then

**Maintenance dose:** 4 mg/kg (IV) q12h. Can switch to weight-based PO maintenance dosing anytime while on maintenance IV dose (see comments).

PO dosing: *Weight ≥ 40 kg:* Loading dose of 400 mg (PO) q12h × 1 day, then maintenance dose of 200 mg (PO) q12h. If response is inadequate, the dose may be increased to 300 mg (PO) q12h. *Weight < 40 kg:* Loading dose of 200 mg (PO) × 1 day, then maintenance dose of 100 mg (PO) q12h. If response is inadequate, the dose may be increased to 150 mg (PO) q12h. For chronic/non-life-threatening infections, loading dose may be given PO (see comments).

**Pharmacokinetic Parameters:**

*Peak serum level:* 2.3–4.7 mcg/ml

*Bioavailability:* 96%

*Excreted unchanged (urine):* 2%

*Serum half-life (normal/ESRD):* 6/6 hrs

*Plasma protein binding:* 58%

*Volume of distribution (V<sub>d</sub>):* 4.6 L/kg

**Primary Mode of Elimination:** Hepatic  
**Dosage Adjustments\***

\*Usual dose\* assumes normal renal/hepatic function. \*For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

CrCl 50–80 mL/min	No change
CrCl 10–50 mL/min	No change PO; avoid IV
CrCl < 10 mL/min	No change PO; avoid IV
Post-HD dose	Usual dose PO; avoid IV
Post-PD dose	No data
CVWH/CVVD/ CVVHD dose	4 mg/kg (PO) q12h; avoid (IV)
Moderate hepatic insufficiency	6 mg/kg (IV) q12h × 1 day or 200 mg (PO) q12h × 1 day, then 2 mg/kg (IV) q12h or 100 mg (PO) q12h (> 40 kg)
Severe hepatic insufficiency	Use with caution

**Drug Interactions:** Benzodiazepines, vinca alkaloids (↑ interacting drug levels); carbamazepine, ergot alkaloids, rifampin, rifabutin, sirolimus, long-acting barbiturates (contraindicated with voriconazole); cyclosporine, ↓ (cyclosporine by 1/2 and monitor); efavirenz (↑ maintenance dose of voriconazole to 400 mg (PO) q12h and decrease efavirenz dose to 300mg (PO) q12h while on voriconazole); omeprazole (↑ interacting drug levels, ↓ interacting drug dose by 50%); tacrolimus (↑ tacrolimus levels, ↓ tacrolimus dose by 66%); phenytoin (↓ voriconazole levels); ↑ voriconazole dose from 4 mg/kg [IV] to 5 mg/kg [IV] and from 200 mg [PO] to 400 mg [PO]; warfarin (↑ INR); statins (↑ risk of rhabdomyolysis); dihydropyridine calcium channel blockers (hypotension); tacrolimus

(reduce tacrolimus dose by 1/3 and monitor while on voriconazole); sulfonyleureas (hypoglycemia). Voriconazole has not been studied with protease inhibitors or NNRTIs, but ↑ voriconazole levels are predicted (↑ hepatotoxicity/adverse effects).

**Adverse Effects:** ↑ SGOT/SGPT; visual events (blurring vision, ↑ brightness, pain; occurs soon after ingestion and usually resolves quickly (~ ½ hour), hallucinations, hypoglycemia, dose-dependent arrhythmias, ↑ QT<sub>c</sub> interval; rash, including Stevens-Johnson Syndrome; photosensitivity reactions (avoid direct sunlight); melanoma (↑ risk of with chronic therapy).

**Allergic Potential:** High

Safety in Pregnancy: D

**Comments:** If intolerance to therapy develops, the IV maintenance dose may be reduced to 3 mg/kg and the PO maintenance dose may be reduced in steps of 50 mg/d to a minimum of 200 mg/d q12h (weight ≥ 40 kg) or 100 mg q12h (weight < 40 kg). Non-linear kinetics (doubling of oral dose = 2.8-fold increase in serum levels). 10–15% of patients have serum levels > 6 mcg/ml. Food decreases bioavailability; take 1 hour before or after meals. Do not use IV voriconazole if CrCl < 50 mL/min to prevent accumulation of voriconazole IV vehicle, sulphobutyl ether cyclodextrin (SBECD); instead use oral formulation, which has no SBECD. Loading dose may be given PO for chronic/non-life-threatening infections. Because of visual effects, do not drive or operate machinery. Monitor LFTs before and during therapy. Take without food; suspension contains sucrose.

**Highly active against *C. albicans*, non-albicans *Candida*, *Aspergillus*, *Cryptococcus*, *Hansenula* and *Pseudoallescheria/Scedosporium*. Some activity against *Fusaria*, *Fluconazole/itraconazole resistant C. albicans*,**

"Usual dose" assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

**Coccidiomycosis, and *Penicillium marneffei*. No activity against *Rhizopus* or *Mucor*.**

Meningeal dose = usual dose.

**Cerebrospinal Fluid Penetration:** 90%

**REFERENCES:**

- Aypar E, Kendirli T, Tutar E, et al. Voriconazole-induced QT interval prolongation and torsades de pointes. *Pediatr Int* 53:761–763, 2011.
- Bruggemann RJM, Alfenaar JWC, Blijlevens NMA, et al. Clinical relevance of the pharmacokinetic interactions of azole antifungal drugs with coadministered agents. *Clin Infect Dis* 48:1441–1458, 2009.
- Camuset J, Nunes H, Dombret MC, et al. Treatment of chronic pulmonary aspergillosis by voriconazole in nonimmunocompromised patients. *Chest* 131:1435–1441, 2007.
- Capitano B, Potoski BA, Husain S, et al. Intrapulmonary penetration of voriconazole in patients receiving an oral prophylactic regimen. *Antimicrob Agents Chemother* 50:1878–80, 2006.
- Cowen EW, Nguyen JC, Miller DD, et al. Chronic phototoxicity and aggressive squamous cell carcinoma of the skin in children and adults during treatment with voriconazole. *J Am Acad Dermatol* 62:31–37, 2010.
- den Hollander JG, van Arkel C, Rijnders BJ, et al. Incidence of voriconazole hepatotoxicity during intravenous and oral treatment for invasive fungal infections. *J Antimicrob Chemother* 57:1248–50, 2006.
- Denes E, Boumediene A, Durox H, et al. Voriconazole concentrations in synovial fluid and bone tissues. *J Antimicrob Chemother* 59:818–819, 2007.
- Denning DW, Ribaud P, Milpied N, et al. Efficacy and safety of voriconazole in the treatment of acute invasive aspergillosis. *Clin Infect Dis* 34:563–71, 2002.
- Dickmeyer NJ, Kiel PJ. Dosing voriconazole in an obese patient. *Clin Infect Dis* 53:745, 2011.
- Elder T, Sieniawski M, Gossmann A, et al. Voriconazole brain tissue levels in rhinocerebral aspergillosis in a successfully treated young woman. *Int J Antimicrob Agents* 28:262–5, 2006.
- Hicheri Y, Cook G, Cordonnier C. Antifungal prophylaxis in haematology patients: the role voriconazole. *Clin Microbiol Infect* 18:1–15, 2012.
- Ho DY, Lee JD, Rosso F, et al. Treating disseminated fusariosis: amphotericin B, voriconazole or both? *Mycoses* 50:227–231, 2007.
- Hoffman HL, Rathbun RC. Review of the safety and efficacy of voriconazole. *Expert Opin Investig Drugs* 11:409–29, 2002.
- Jurkunas UV, Langston DP, Colby K. Use of voriconazole in the treatment of fungal keratitis. *Int Ophthalmol Clin* 47:47–59, 2007.
- Keirns J, Sawamoto T, Holum M, et al. Steady-state pharmacokinetics of micafungin and voriconazole after separate and concomitant dosing in healthy adults. *Antimicrob Agents Chemother* 51:787–790, 2007.
- Kim MM, Vikram HR, Kusne S, et al. Treatment of refractory coccidioidomycosis with voriconazole or posaconazole. *Clin Infect Dis* 53:1060–6, 2011.
- Leveque D, Nivoix Y, Jehl F, Herbrecht R. Clinical pharmacokinetics of voriconazole. *Int J Antimicrob Agents* 27:274–84, 2006.
- Masomoto Y, Nannya Y, Kurokawa M. Voriconazole is effective as secondary antifungal prophylaxis in leukemia patients with prior pulmonary fungal disease: case series and review of literature. *J Chemother* 23:17–23, 2011.
- Miller DD, Cowen EW, Nguyen JC, et al. Melanoma associated with long-term voriconazole therapy: a new manifestation of chronic photosensitivity. *Arch Dermatol* 146:300–304, 2010.
- Oude Lashof AM, Sobel JD, Ruhnke M, et al. Safety and tolerability of voriconazole in patients with baseline renal insufficiency and candidemia. *Antimicrob Agents Chemother* 56:3133–7, 2012.
- Pagano L, Valentini CG, Fianchi L, et al. Treatment strategies for invasive aspergillosis in neutropenic patients: voriconazole or liposomal amphotericin-B. *J Chemother* 23:5–8, 2011.
- Poza G, Montoya J, Redondo C, et al. Meningitis caused by *Pseudallescheria boydii* treated with voriconazole. *Clin Infect Dis* 30:981–2, 2000.
- Radaj J, Krouzecky A, Stehlik P, et al. Pharmacokinetic evaluation of voriconazole treatment in critically ill patients undergoing continuous venovenous hemofiltration. *Ther Drug Monit* 33:393–397, 2011.

"Usual dose" assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

Rondeau S, Couderu L, Dominique S, et al. High Frequency of voriconazole-related phototoxicity in cystic fibrosis patients. *Eur Respir J* 39:782–4, 2012.

Sambatakou H, Dupont B, Lode H, Denning DW. Voriconazole treatment for subacute invasive and chronic pulmonary aspergillosis. *Am J Med* 119:527, 2006.

Schwartz S, Ruhne M, Ribaud P, et al. Improved outcome in central nervous system aspergillosis, using voriconazole treatment. *Amer Soc of Hematology* 106:2641–5, 2006.

Shehab N, DePestel DD, Mackler ER, et al. Institutional experience with voriconazole compared with liposomal amphotericin B as empiric therapy for febrile neutropenia. *Pharmacotherapy* 27:970–979, 2007.

Smith J, Safdar N, Knasinski V, et al. Voriconazole Therapeutic Drug Monitoring. *Antimicrob Agents Chemother* 50:1570–72, 2006.

Solis-Munoz P, Lopex JC, Bernal W, et al. Voriconazole hepatotoxicity in severe liver dysfunction. *J of Inf* 66:80–86, 2013.

Supparatpinyo K, Schlamm HT. Voriconazole as therapy for systemic *Penicillium marneffeii* infections in AIDS patients. *Am J Trop Med Hyg* 77:350–353, 2007.

Tan K, Brayshaw N, Tomaszewski K, et al. Investigation of the potential relationships between plasma voriconazole concentrations and visual adverse events or liver function test abnormalities. *J Clin Pharmacol* 46:235–43, 2006.

Walsh TJ, Anaissie EJ, Denning DW, et al. Treatment of Aspergillosis: Clinical Practice Diseases Society of America. *Clin Infect Dis* 46:327–360, 2008. Website: [www.pdr.net](http://www.pdr.net)

## Zalcitabine (HIVID) ddC

**Drug Class:** HIV NRTI (nucleoside reverse transcriptase inhibitor).

**Usual Dose:** 0.75 mg (PO) q8h.

**Pharmacokinetic Parameters:**

*Peak serum level:* 0.08 mcg/ml

*Bioavailability:* 80%

*Excreted unchanged (urine):* 75%  
*Serum half-life (normal/ESRD):* 2/8.5 hrs  
*Plasma protein binding:* 0%  
*Volume of distribution ( $V_d$ ):* 0.54 L/kg

**Primary Mode of Elimination:** Renal  
**Dosage Adjustments\***

CrCl 10–50 mL/min	0.75 mg (PO) q12h
CrCl < 10 mL/min	0.75 mg (PO) q24h
Post-HD dose	No data
Post-PD dose	No data
CVWH/CVWH/D/ CVWHDF dose	0.75 mg (PO) q12h
Moderate hepatic insufficiency	No change
Severe hepatic insufficiency	No change

**Drug Interactions:** Cimetidine, probenecid, TMP-SMX (↑ zalcitabine levels); dapsone, didanosine, stavudine, INH, phenytoin, metronidazole, other neurotoxic agents or history of neuropathy (↑ risk of peripheral neuropathy); magnesium/aluminum containing antacids, metoclopramide (↓ bioavailability of zalcitabine); pentamidine IV, valproic acid, alcohol, other agents known to cause pancreatitis (↑ risk of pancreatitis).

**Adverse Effects:** Drug fever/rash, leukopenia, anemia, thrombocytopenia, hepatomegaly, hepatotoxicity/hepatic necrosis, peripheral neuropathy, pancreatitis, stomatitis, oral/genital ulcers, dysphagia, arthritis, hyperglycemia, lipotrodystrophy, wasting, night sweats, lactic acidosis with hepatic steatosis (rare, but potentially life-threatening toxicity with use of NRTIs).

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

**Allergic Potential:** High

**Safety in Pregnancy:** C

**Comments:** Foscarnet may increase toxicity. Do not use with stavudine, didanosine, or lamivudine to avoid additive toxicities. Food decreases absorption by 39%. Effective antiretroviral therapy consists of at least 3 antiretrovirals (same/different classes).

**Cerebrospinal Fluid Penetration:** 25%

#### REFERENCES:

Panel on Antiretroviral Guidelines for Adults and Adolescents. Department of Health and Human Services. Guidelines for the use of antiretroviral agents in HIV-infected adults and adolescents. <http://aidsinfo.nih.gov/ConsentFiles/AdultandAdolescentGL.pdf>, 2012.

Shelton MJ, O'Donnell AM, Morse GD. Zalcitabine. *Ann Pharmacotherapy* 27:480–9, 1993.

Thompson MA, Aberg JA, Cahn P, et al. Antiretroviral treatment of adult HIV infection: 2012 recommendations of the International Antiviral Society — USA panel. *JAMA* 308:387–402, 2012.

Website: [www.pdr.net](http://www.pdr.net)

## Zidovudine (Retrovir) ZDV Azidothymidine AZT

**Drug Class:** HIV NRTI (nucleoside reverse transcriptase inhibitor).

**Usual Dose:** 300 mg (PO) q12h or 200 mg (PO) q8h (see comments). IV solution 10 mg/mL (dose 1 mg/kg 5–6 x/day).

**Pharmacokinetic Parameters:**

*Peak serum level:* 1.2 mcg/mL

*Bioavailability:* 64%

*Excreted unchanged (urine):* 16%

*Serum half-life (normal/ESRD):* 1.1/1.4 hrs

*Plasma protein binding:* < 38%

*Volume of distribution ( $V_d$ ):* 1.6 L/kg

**Primary Mode of Elimination:** Hepatic

**Dosage Adjustments\***

CrCl < 15 mL/min	100 mg (PO) q8h or 300 mg (PO) q24h
Post-HD/PD dose	100 mg or 300 mg
CVWH/CVWHD/CVWHDf dose	300 mg (PO) q24h
Moderate—severe hepatic insufficiency	Use with caution

**Drug Interactions:** Acetaminophen, atovaquone, fluconazole, methadone, probenecid, valproic acid (↑ zidovudine levels); clarithromycin, nelfinavir, rifampin, rifabutin (↓ zidovudine levels); dapson, flucytosine, ganciclovir, interferon alpha, bone marrow suppressive/cytotoxic agents (↑ risk of hematologic toxicity); indomethacin (↑ levels of zidovudine toxic metabolite); phenytoin (↑ zidovudine levels, ↑ or ↓ phenytoin levels); ribavirin (↓ zidovudine effect; avoid).

**Adverse Effects:** Zidovudine, has been associated with hematologic toxicity including neutropenia and severe anemia. Prolonged use of zidovudine has been associated with symptomatic myopathy as well as lactic acidosis and severe hepatomegaly with steatosis, including fatal cases. May cause nausea, vomiting, GI upset, diarrhea, malaise, anorexia, macrocytosis, headachese, insomnia, blue/black nail discoloration and asthenia. Nausea, vomiting, GI upset, diarrhea, malaise, anorexia, leukopenia, severe anemia, macrocytosis, thrombocytopenia, headaches, ↑ SGOT/SGPT, hepatotoxicity, myalgias, myositis, symptomatic myopathy, insomnia, blue/black nail discoloration, asthenia, lactic acidosis with hepatic steatosis (rare, but potentially life-threatening toxicity with use of NRTI's).

**Allergic Potential:** Low

**Safety in Pregnancy:** C

**Comments:** Antagonized by ganciclovir or ribavirin. Also a component of Combivir and Trizivir. Patients on IV therapy should be switched

\*Usual dose\* assumes normal renal/hepatic function. \* For renal insufficiency, give usual dose × 1 followed by maintenance dose per CrCl. For dialysis patients, dose the same as for CrCl < 10 mL/min and give supplemental (post-HD/PD dose) immediately after dialysis. CrCl = creatinine clearance; CVWH = continuous veno-venous hemofiltration; HD/PD = hemodialysis/peritoneal dialysis.

to PO as soon as able to take oral medication. For IV administration, dilute in D5W to a concentration no greater than 4 mg/mL and infuse over 1 hour.

### Cerebrospinal Fluid Penetration: 60%

#### REFERENCES:

- Barry M, Mulcahy F, Merry C, et al. Pharmacokinetics and potential interactions amongst antiretroviral agents used to treat patients with HIV infection. *Clin Pharmacol* 36:289–304, 1999.
- Been-Tiktak AM, Boucher CA, Brun-Vezinet F, et al. Efficacy and safety of combination therapy with delavirdine and zidovudine: A European/Australian phase II trial. *Intern J Antimicrob Agents* 11:13–21, 1999.
- Diaz-Brito V, Leon A, Knobel H, et al. Post-exposure prophylaxis for HIV-infection: a clinical trial comparing

- lopinavir/ritonavir versus atazanavir each with zidovudine/lamivudine. *Antivir Ther* 17:337–46, 2012.
- McDowell JA, Lou Y, Symonds WS, et al. Multiple-dose pharmacokinetics and pharmacodynamics of abacavir alone and in combination with zidovudine in human immunodeficiency virus-infected adults. *Antimicrob Agents Chemother* 44:2061–7, 2000.
- Panel on Antiretroviral Guidelines for Adults and Adolescents. Department of Health and Human Services. Guidelines for the use of antiretroviral agents in HIV-infected adults and adolescents. <http://aidsinfo.nih.gov/ConsentFiles/AdultandAdolescentGL.pdf>, 2012.
- Thompson MA, Aberg JA, Cahn P, et al. Antiretroviral treatment of adult HIV infection: 2012 recommendations of the International Antiviral Society — USA panel. *JAMA* 308:387–402, 2012.
- Website: [www.pdr.net](http://www.pdr.net)

---

## REFERENCES AND SUGGESTED READINGS

---

### TEXTBOOKS

- Ambrose P, Nightingale AT (eds). *Principles of Pharmacodynamics*. Marcel Dekker, Inc, New York, 2001.
- Baddour L, Gorbach SL (eds). *Therapy of Infectious Diseases*. Saunders, Philadelphia, 2003.
- Bennet JE, Dolin R, Blaser MJ (Eds). *Mandell, Douglas, and Bennett's Principles and Practice of Infectious Diseases (8<sup>th</sup> Ed)*. Philadelphia Elsevier Churchill Livingstone, 2015.
- Bennet WM, Aronoff GR, Golper TA, Morrison G, Brater DC, Singer I (eds). *Drug Prescribing in Renal Failure (2<sup>nd</sup> Ed)*, American College of Physicians, Philadelphia, 2000.
- Bowden RA, Ljungman P, Snyderman DR (eds). *Transplant Infections (3<sup>rd</sup> Ed)*, Lippincott Williams & Wilkins, Philadelphia, PA, 2010.
- Bryskier A (ed). *Antimicrobial Agents*. ASM Press, Washington, D.C., 2005.
- Finch RG, Greenwood D, Norrby SR, Whitley RJ (eds). *Antibiotic and Chemotherapy*. Churchill Livingstone, United Kingdom, 2003.
- Gorbach SL, Bartlett JG, Blacklow NR (eds). *Infectious Diseases (4<sup>th</sup> Ed)*, Lippincott Williams Wilkins, Philadelphia, 2004.
- Grayson ML (ed). *Kucers The Use of Antibiotics (6<sup>th</sup> Ed)*, ASM Press, Washington, DC, 2010.
- Koff RS, *Hepatitis Essentials*. Jones & Bartlett, Sudbury, MA, 2011.
- O'Grady F, Lambert HP, Finch RG, Greenwood D (eds). *Antibiotic and chemotherapy (2<sup>nd</sup> Ed)*. Churchill Livingstone, New York, 1997.
- Pai MP, Momary KM, Rodvold KA. *Antibiotic Drug Interactions*. *Med Clin N Am* 90:1223–55, 2006.
- Piscitelli SC, Rodvold KE (eds). *Drug Interactions in Infectious Diseases*. Humana Press, Totowa, 2001.
- Ristuccia AM, Cunha BA (eds). *Antimicrobial Therapy*. Raven Press, New York, 1984.
- Sax PE, Cohen CJ, Kuritzkes DR. *HIV Essentials (6<sup>th</sup> Ed)*, Jones & Bartlett, Sudbury, MA, 2014.
- Schlossberg D (ed). *Clinical Infectious Disease (2<sup>nd</sup> Ed)*, Cambridge University Press, Cambridge, 2015.
- Schlossberg D (ed). *Current Therapy of Infectious Disease (3<sup>rd</sup> Ed)*, Mosby-Yearbook, St. Louis, 2008.
- Schlossberg D (ed). *Tuberculosis and Non-Tuberculous Mycobacterial Infections (7<sup>th</sup> Ed)*, McGraw-Hill, New York, 2011.
- Scholar EM, Pratt WB (eds). *The Antimicrobial Drugs (2<sup>nd</sup> Ed)*, Oxford University Press, New York, 2000.
- Wormser GP (ed). *AIDS (4<sup>th</sup> Ed)*, Elsevier, Philadelphia, 2004.
- Yoshikawa TT, Rajagopalan S (eds). *Antibiotic Therapy for Geriatric Patients*. Taylor & Francis, New York, 2006.
- Yu VL, Merigan, Jr. TC, Barrier SL (eds). *Antimicrobial Therapy and Vaccines 2nd Ed*, Williams & Wilkins, Baltimore, 2005.
- Zinner SH, Young LS, Acar JF, Ortiz-Neu C (eds). *New Considerations for Macrolides, Azalides, Streptogramins, and Ketolides*. Marcel Dekker, New York, 2000.

**APPENDIX**

## Malaria in Adults (United States)

Clinical Diagnosis/ <i>Plasmodium</i> species	Region Acquired	Recommended Adult Drug Doses
<b>Severe malaria</b> <sup>1,2,3,4</sup>	<b>All regions</b>	<p><b>Quinidine gluconate<sup>2</sup> plus either: Doxycycline or Clindamycin</b></p> <p><b>Quinidine gluconate:</b> 6.25 mg base/kg (=10 mg salt/kg) (IV over 1–2 hrs) then 0.0125 mg base/kg/min (=0.02 mg salt/kg/min) continuous infusion for at least 24 hours. An alternative regimen is 15 mg base/kg (=24 mg salt/kg) loading dose IV infused over 4 hours, followed by 7.5 mg base/kg (=12 mg salt/kg) infused over 4 hours q8h, starting 8 hours after the loading dose. Once parasite density &lt;1% and patient can take oral medication, complete treatment with oral quinine, dose as above. Quinidine/quinine course = 7 days in Southeast Asia; = 3 days in Africa or South America.</p> <p><b>Doxycycline:</b> Treatment as above, give 100 mg (IV/PO) q12h × 7 days.</p> <p><b>Clindamycin:</b> Treatment as above. If patient not able to take oral medication, give 10 mg base/kg (IV) then 5 mg base/kg (IV/PO) q8h × 7 days.</p>
<p><b>Uncomplicated malaria/ <i>P. falciparum</i> or</b> Species not identified (If “species not identified” is subsequently diagnosed as <i>P. vivax</i> or <i>P. ovale</i>: see <i>P. vivax</i> and <i>P. ovale</i> treat with primaquine)</p>	<p><b>Chloroquine-resistant or unknown resistance<sup>5</sup></b> (Middle Eastern countries with chloroquine-resistant <i>P. falciparum</i> include Iran, Oman, Saudi Arabia, and Yemen. Malaria acquired in the newly independent states of the former Soviet Union and Korea to date have been uniformly caused by <i>P. vivax</i> and should therefore be treated as chloroquine-sensitive infections.)</p>	<p><b>A. Atovaquone-proguanil (Malarone)<sup>6</sup></b> <i>Adult tab = 250 mg atovaquone/ 100 mg proguanil</i> 4 adult tabs (PO) q24h × 3 days <b>or</b></p>



## Malaria in Adults (United States) (cont'd)

Clinical Diagnosis/ <i>Plasmodium</i> species	Region Acquired	Recommended Adult Drug Doses
<p><b>Uncomplicated malaria/ <i>P. falciparum</i></b> (cont'd)</p>	<p><b>Chloroquine-resistant or unknown resistance</b> (cont'd)</p>	<p><b>B. Artemether-lumefantrine (Coartem)<sup>6</sup></b>  <i>1 tablet = 20 mg artemether/120 mg lumefantrine</i>            A 3-day treatment schedule with a total of 6 oral doses is recommended based on weight. The patient should receive the initial dose, followed by the second dose 8 hours later, then 1 dose (PO) q12h x 2 days.            ≥35 kg: 4 tablets per dose  <b>or</b>  <b>C. Quinine sulfate plus either: Doxycycline<sup>9</sup> or Clindamycin</b>  <b>Quinine sulfate<sup>8</sup>:</b> 542 mg base (=650 mg salt)<sup>7</sup> (PO) q8h x 3 or 7 days  <b>Doxycycline:</b> 100 mg (PO) q12h x 7 days  <b>Clindamycin:</b> 20 mg base/kg/day (PO) divided q8h x 7 days  <b>or</b>  <b>D. Mefloquine (Lariam)<sup>10</sup></b>            684 mg base (=750 mg salt) (PO) as initial dose, followed by 456 mg base (=500 mg salt) (PO) given 6–12 hours after initial dose            Total dose = 1,250 mg salt</p>
<p><b>Uncomplicated malaria/ <i>P. falciparum</i></b> or <i>Species not identified</i></p>	<p><b>Chloroquine-sensitive</b> (Central America west of Panama Canal; Haiti; the Dominican Republic; and most of the Middle East)</p>	<p><b>Chloroquine phosphate (Aralen)</b> 600 mg base (=1,000 mg salt) (PO) immediately, followed by 300 mg base (=500 mg salt) (PO) at 6, 24, and 48 hours Total dose: 1,500 mg base (=2,500 mg salt)  <b>or</b>  <b>Hydroxychloroquine (Plaquenil)</b>            620 mg base (=800 mg salt) (PO) immediately, followed by 310 mg base (=400 mg salt) (PO) at 6, 24, and 48 hours            Total dose: 1,550 mg base (=2,000 mg salt)</p>

## Malaria in Adults (United States) (cont'd)

Clinical Diagnosis/ <i>Plasmodium</i> species	Region Acquired	Recommended Adult Drug Doses
Uncomplicated malaria/ <i>P. malariae</i> or <i>P. knowlesi</i>	All regions	<b>Chloroquine phosphate:</b> Treatment as above <b>or</b> <b>Hydroxychloroquine:</b> Treatment as above
Uncomplicated malaria/ <i>P. vivax</i> or <i>P. ovale</i>	All regions (for suspected chloroquine-resistant <i>P. vivax</i> , see below)	<b>Chloroquine phosphate plus Primaquine phosphate<sup>11</sup></b> <b>Chloroquine phosphate:</b> Treatment as above <b>Primaquine phosphate:</b> 30 mg base (PO) q24h × 14 days <b>or</b> <b>Hydroxychloroquine plus Primaquine phosphate<sup>11</sup></b> <b>Hydroxychloroquine:</b> Treatment as above <b>Primaquine phosphate:</b> 30 mg base (PO) q24h × 14 days
Uncomplicated malaria/ <i>P. vivax</i>	Chloroquine-resistant <sup>12</sup> (Papua New Guinea and Indonesia)	<b>A. Quinine sulfate plus either: Doxycycline plus Primaquine phosphate<sup>11</sup></b> <b>Quinine sulfate:</b> Treatment as above <b>Doxycycline:</b> Treatment as above <b>Primaquine phosphate:</b> Treatment as above <b>or</b> <b>B. Atovaquone-proguanil plus Primaquine phosphate</b> <b>Atovaquone-proguanil:</b> Treatment as above <b>Primaquine phosphate:</b> Treatment as above <b>or</b> <b>C. Mefloquine<sup>10</sup> plus Primaquine phosphate<sup>11</sup></b> <b>Mefloquine:</b> Treatment as above <b>Primaquine phosphate:</b> Treatment as above

## Malaria in Adults (United States) (cont'd)

Clinical Diagnosis/ <i>Plasmodium</i> species	Region Acquired	Recommended Adult Drug Doses
<b>Uncomplicated malaria: alternatives for pregnant women</b> <sup>13,14,15,16</sup>	<b>Chloroquine-sensitive</b> (see uncomplicated malaria sections above for chloroquine-sensitive species by region)	<b>Chloroquine phosphate:</b> Treatment as above <b>or</b> <b>Hydroxychloroquine:</b> Treatment as above
	<b>Chloroquine resistant <i>P. falciparum</i></b> <sup>1</sup> (see sections above for regions with chloroquine resistant <i>P. falciparum</i> )	<b>Quinine sulfate plus Clindamycin</b> <b>Quinine sulfate:</b> Treatment as above <b>Clindamycin:</b> Treatment as above
<b>Uncomplicated malaria: alternatives for pregnant women (cont'd)</b>	<b>Chloroquine-resistant <i>P. vivax</i></b> (see uncomplicated malaria sections above for regions with chloroquine-resistant <i>P. vivax</i> )	<b>Quinine sulfate</b> <b>Quinine sulfate</b> <sup>2</sup> : 650 mg salt (PO) q8h x 7 days

- Persons with a positive blood smear or history of recent possible exposure and no other recognized pathology who have one or more of the following clinical criteria (impaired consciousness/coma, severe normocytic anemia, renal failure, pulmonary edema, acute respiratory distress syndrome, circulatory shock, disseminated intravascular coagulation, spontaneous bleeding, acidosis, hemoglobinuria, jaundice, repeated generalized convulsions, and/or parasitemia of >5%) are considered to have manifestations of more severe disease. Severe malaria is most often caused by *P. falciparum*.
- Patients diagnosed with severe malaria should be treated aggressively with parenteral antimalarial therapy. Treatment with IV quinidine should be initiated as soon as possible after the diagnosis has been made. Patients with severe malaria should be given an intravenous loading dose of quinidine unless they have received more than 40 mg/kg of quinine in the preceding 48 hours or if they have received mefloquine within the preceding 12 hours. Consultation with a cardiologist and a physician with experience treating malaria is advised when treating malaria patients with quinidine. During administration of quinidine, blood pressure monitoring (for hypotension) and cardiac monitoring (for widening of the QRS complex and/or lengthening of the QTc interval) should be monitored continuously and blood glucose (for hypoglycemia) should be monitored periodically. Cardiac complications, if severe, may warrant temporary discontinuation of the drug or slowing of the intravenous infusion.
- Consider exchange transfusion if the parasite density (i.e. parasitemia) is >10% OR if the patient has altered mental status, non-volume overload pulmonary edema, or renal complications. The parasite density can be estimated by examining a monolayer of red blood cells (RBCs) on the thin smear under oil immersion magnification. The slide should be examined where the RBCs are more or less touching (approximately 400 RBCs per field). The parasite density can then be estimated from the percentage of infected RBCs and should be monitored every 12 hours. Exchange transfusion should be continued until the parasite density is <1% (usually requires 8–10 units). IV quinidine administration should not be delayed for an exchange transfusion and can be given concurrently throughout the exchange transfusion.
- Pregnant women diagnosed with severe malaria should be treated aggressively with parenteral antimalarial therapy.
- There are 4 options (A, B, C, or D) available for treatment of uncomplicated malaria caused by chloroquine-resistant *P. falciparum*. Options A, B, and C are equally recommended. Because of a higher rate of severe neuropsychiatric reactions seen at treatment doses, option D (mefloquine) not recommend unless the other options cannot

be used. For option C, because there is more data on the efficacy of quinine in combination with doxycycline, these treatment combinations preferred to quinine with clindamycin.

6. Take with with food or whole milk. If patient vomits within 30 minutes of taking a dose, then they should repeat the dose.
7. US manufactured quinine sulfate capsule is in a 324 mg dosage; therefore 2 capsules should be sufficient for adult dosing. Pediatric dosing may be difficult due to unavailability of non-capsule forms of quinine.
8. For infections acquired in Southeast Asia, quinine treatment should continue for 7 days. For infections acquired elsewhere, quinine treatment should continue for 3 days.
9. Doxycycline are not indicated for use in children less than 8 years old. For children less than 8 years old with chloroquine-resistant *P. falciparum*, atovaquone-proguanil and artemether-lumefantrine are recommended treatment options; mefloquine can be considered if no other options are available. For children less than 8 years old with chloroquine-resistant *P. vivax*, mefloquine is the recommended treatment. If it is not available or is not being tolerated and if the treatment benefits outweigh the risks, atovaquone-proguanil or artemether-lumefantrine should be used instead.
10. Treatment with mefloquine is not recommended in persons who have acquired infections from Southeast Asia due to drug resistance.
11. Primaquine is used to eradicate any hypnozoites in the liver, to prevent relapses, in *P. vivax* and *P. ovale* infections. Because primaquine can cause hemolytic anemia in G6PD-deficient persons, G6PD screening must occur prior to starting treatment with primaquine. For persons with borderline G6PD deficiency or as an alternate to the above regimen, primaquine may be given 45 mg orally one time per week for 8 weeks; consultation with an expert in infectious disease and/or tropical medicine is advised if this alternative regimen is considered in G6PD-deficient persons. Primaquine must not be used during pregnancy.
12. There are three options (A, B, or C) available for treatment of uncomplicated malaria caused by chloroquine-resistant *P. vivax*. High treatment failure rates due to chloroquine-resistant *P. vivax* have been well documented in Papua New Guinea and Indonesia. Rare case reports of chloroquine-resistant *P. vivax* have also been documented in Burma (Myanmar), India, and Central and South America. Persons acquiring *P. vivax* infections outside of Papua New Guinea or Indonesia should be started on chloroquine. If the patient does not respond, the treatment should be changed to a chloroquine-resistant *P. vivax* regimen and CDC should be notified (Malaria Hotline number listed below). For treatment of chloroquine-resistant *P. vivax* infections, options A, B, and C are equally recommended.
13. For pregnant women diagnosed with uncomplicated malaria caused by chloroquine-resistant *P. falciparum* or chloroquine-resistant *P. vivax* infection, treatment with doxycycline is generally not indicated. However, doxycycline may be used in combination with quinine (as recommended for non-pregnant adults) if other treatment options are not available or are not being tolerated, and the benefit is judged to outweigh the risks.
14. Atovaquone-proguanil and artemether-lumefantrine are generally not recommended for use in pregnant women, particularly in the first trimester due to lack of sufficient safety data. For pregnant women diagnosed with uncomplicated malaria caused by chloroquine-resistant *P. falciparum* infection, atovaquone-proguanil or artemether-lumefantrine may be used if other treatment options are not available or are not being tolerated, and if the potential benefit is judged to outweigh the potential risks.
15. Because of a possible association with mefloquine treatment during pregnancy and an increase in stillbirths, mefloquine is generally not recommended for treatment in pregnant women. However, mefloquine may be used if it is the only treatment option available and if the potential benefit is judged to outweigh the potential risks.
16. For *P. vivax* and *P. ovale* infections, primaquine phosphate for radical treatment of hypnozoites should not be given during pregnancy. Pregnant patients with *P. vivax* and *P. ovale* infections should be maintained on chloroquine prophylaxis for the duration of their pregnancy. The chemoprophylactic dose of chloroquine phosphate is 300 mg base (=500 mg salt) orally once per week. After delivery, pregnant patients who do not have G6PD deficiency should be treated with primaquine.

### Based on CDC Recommendations for Malaria in the US

**CDC Malaria Hotline: weekdays 8 am – 4:30 pm EST: (770) 488-7788**

**after hours, weekends and holidays: (770) 488-7100**

## Malaria in Children (United States)

Clinical Diagnosis/ <i>Plasmodium</i> species	Region Acquired	Recommended Drugs and Pediatric Dose ( <i>pediatric dose should not exceed adult dose</i> )
Severe malaria <sup>1,2,3</sup>	All regions	<p><b>Quinidine gluconate<sup>2</sup> plus either: Doxycycline<sup>8</sup> or Clindamycin</b></p> <p><b>Quinidine gluconate:</b> Same mg/kg dosing and recommendations as for adults.</p> <p><b>Doxycycline:</b> Treatment as above. For children &lt;45 kg, give 2.2 mg/kg (IV) q12h and then switch to (PO) doxycycline (dose as above) as soon as patient can take oral medication. For children &gt;45 kg, use same dosing as for adults. Treatment × 7 days.</p> <p><b>Clindamycin:</b> Treatment as above. 10 mg base/kg loading dose (IV) followed by 5 mg base/kg (IV/PO) q8h. Treatment × 7 days.</p>
Uncomplicated malaria/ <i>P. falciparum</i> or Species not identified <sup>4</sup>	Chloroquine-resistant or unknown resistance <sup>11</sup> (Middle Eastern countries with chloroquine-resistant <i>P. falciparum</i> include Iran, Oman, Saudi Arabia, and Yemen. Of note, infections acquired in the	<p><b>A. Atovaquone-proguanil (Malarone)<sup>5</sup></b></p> <p><i>Adult tab = 250 mg atovaquone/100 mg proguanil</i></p> <p><i>Peds tab = 62.5 mg atovaquone/25 mg proguanil</i></p> <p>5–8 kg: 2 peds tabs (PO) q24h × 3d  9–10 kg: 3 peds tabs (PO) q24h × 3d  11–20 kg: 1 adult tab (PO) q24h × 3d  21–30 kg: 2 adult tabs (PO) q24h × 3d  31–40 kg: 3 adult tabs (PO) q24h × 3d  &gt;40 kg: 4 adult tabs (PO) q24h × 3d</p> <p style="text-align: center;"><b>or</b></p>

## Malaria in Children (United States) (cont'd)

Clinical Diagnosis/ <i>Plasmodium</i> species	Region Acquired	Recommended Drugs and Pediatric Dose ( <i>pediatric dose should not exceed adult dose</i> )
<p>If "species not identified" is subsequently diagnosed as <i>P. vivax</i> or <i>P. ovale</i>: see <i>P. vivax</i> and <i>P. ovale</i> (below) re. treatment with primaquine</p>	<p>Newly Independent States of the former Soviet Union and Korea to date have been uniformly caused by <i>P. vivax</i> and should therefore be treated as chloroquine-sensitive infections).</p>	<p><b>B. Artemether-lumefantrine (Coartem)<sup>5</sup></b>  <i>1 tablet = 20 mg artemether/120 mg lumefantrine</i>            A 3-day treatment schedule with a total of 6 oral doses is recommended for both adult and pediatric patients based on weight. The patient should receive the initial dose, followed by the second dose 8 hours later, then 1 dose (PO) q12h x 2 days.            5–&lt;15 kg: 1 tablet per dose            15–&lt;25 kg: 2 tablets per dose            25–&lt;35 kg: 3 tablets per dose            ≥35 kg: 4 tablets per dose  <b>or</b>  <b>C. Quinine sulfate<sup>3</sup> plus either: Doxycycline<sup>6</sup> or Clindamycin</b>  <b>Quinine sulfate<sup>6,7</sup>:</b> 8.3 mg base/kg (=10 mg salt/kg) (PO) tid x 3 or 7 days  <b>Doxycycline:</b> 2.2 mg/kg (PO) q12h x 7 days  <b>Clindamycin:</b> 20 mg base/kg/day (PO) divided q8h x 7 days  <b>or</b>  <b>D. Mefloquine (Lariam)<sup>9</sup></b>            13.7 mg base/kg (=15 mg salt/kg) (PO) as initial dose, followed by 9.1 mg base/kg (=10 mg salt/kg) (PO) given 6–12 hours after initial dose. Total dose = 25 mg salt/kg</p>

## Malaria in Children (United States) (cont'd)

Clinical Diagnosis/ <i>Plasmodium</i> species	Region Acquired	Recommended Drugs and Pediatric Dose ( <i>pediatric dose should not exceed adult dose</i> )
<b>Uncomplicated malaria/<i>P. falciparum</i></b> or <i>Species not identified</i>	<b>Chloroquine-sensitive</b> (Central America west of Panama Canal; Haiti; the Dominican Republic; and most of the Middle East)	<b>Chloroquine phosphate (Aralen)</b> 10 mg base/kg (PO) immediately, followed by 5 mg base/kg (PO) at 6, 24, and 48 hours Total dose: 25 mg base/kg <b>or</b> <b>Hydroxychloroquine (Plaquenil and generics)</b> 10 mg base/kg (PO) immediately, followed by 5 mg base/kg (PO) at 6, 24, and 48 hours Total dose: 25 mg base/kg
<b>Uncomplicated malaria/<i>P. malariae</i></b> or <i>P. knowlesi</i>	<b>All regions</b>	<b>Chloroquine phosphate:</b> Treatment as above <b>or</b> <b>Hydroxychloroquine:</b> Treatment as above
<b>Uncomplicated malaria/<i>P. vivax</i></b> or <i>P. ovale</i>	<b>All regions</b> For suspected chloroquine-resistant <i>P. vivax</i> , see below	<b>Chloroquine phosphate plus Primaquine phosphate<sup>10</sup></b> <b>Chloroquine phosphate:</b> Treatment as above <b>Primaquine:</b> 0.5 mg base/kg (PO) qd x 14 days <b>or</b> <b>Hydroxychloroquine plus Primaquine phosphate<sup>10</sup></b> <b>Hydroxychloroquine:</b> Treatment as above <b>Primaquine phosphate:</b> 0.5 mg base/kg (PO) qd x 14 days

## Malaria in Children (United States) (cont'd)

Clinical Diagnosis/ <i>Plasmodium</i> species	Region Acquired	Recommended Drugs and Pediatric Dose ( <i>pediatric dose should not exceed adult dose</i> )
<b>Uncomplicated malaria/ <i>P. vivax</i></b>	<b>Chloroquine-resistant<sup>11</sup></b> (Papua New Guinea and Indonesia)	<p><b>A. Quinine sulfate <i>plus</i> either Doxycycline<sup>8</sup> <i>plus</i> Primaquine phosphate<sup>10</sup></b>  <b>Quinine sulfate:</b> Treatment as above  <b>Doxycycline</b> Treatment as above  <b>Primaquine phosphate:</b> Treatment as above</p> <p style="text-align: center;"><b>or</b></p> <p><b>B. Atovaquone-proguanil <i>plus</i> Primaquine phosphate</b>  <b>Atovaquone-proguanil:</b> Treatment as above  <b>Primaquine phosphate:</b> Treatment as above</p> <p style="text-align: center;"><b>or</b></p> <p><b>C. Mefloquine <i>plus</i> Primaquine phosphate<sup>10</sup></b>  <b>Mefloquine:</b> Treatment as above  <b>Primaquine phosphate:</b> Treatment as above</p>

- Persons with a positive blood smear or history of recent possible exposure and no other recognized pathology who have one or more of the following clinical criteria (impaired consciousness/coma, severe normocytic anemia, renal failure, pulmonary edema, acute respiratory distress syndrome, circulatory shock, disseminated intravascular coagulation, spontaneous bleeding, acidosis, hemoglobinuria, jaundice, repeated generalized convulsions, and/or parasitemia of >5%) are considered to have manifestations of more severe disease. Severe malaria is most often caused by *P. falciparum*.
- Patients diagnosed with severe malaria should be treated aggressively with parenteral antimalarial therapy. Treatment with IV quinidine should be initiated as soon as possible after the diagnosis has been made. Patients with severe malaria should be given an intravenous loading dose of quinidine unless they have received more than 40 mg/kg of quinine in the preceding 48 hours or if they have received mefloquine within the preceding 12 hours. Consultation with a cardiologist and a physician with experience treating malaria is advised when treating malaria patients with quinidine. During administration of quinidine, blood pressure monitoring (for hypotension) and cardiac monitoring (for widening of the QRS complex and/or lengthening of the QTc interval) should be monitored continuously and blood glucose (for hypoglycemia) should be monitored periodically. Cardiac complications, if severe, may warrant temporary discontinuation of the drug or slowing of the intravenous infusion.
- Consider exchange transfusion if the parasite density (i.e. parasitemia) is >10% OR if the patient has altered mental status, non-volume overload pulmonary edema, or renal complications. The parasite density can be estimated by examining a monolayer of red blood cells (RBCs) on the thin smear under oil immersion magnification. The slide should be examined where the RBCs are more or less touching (approximately 400 RBCs per field). The parasite density can then be estimated from the percentage of infected RBCs and



should be monitored every 12 hours. Exchange transfusion should be continued until the parasite density is <1% (usually requires 8–10 units). IV quinidine administration should not be delayed for an exchange transfusion and can be given concurrently throughout the exchange transfusion.

4. There are 4 options (A, B, C, or D) available for treatment of uncomplicated malaria caused by chloroquine-resistant *P. falciparum*. Options A, B, and C are equally recommended. Because of a higher rate of severe neuropsychiatric reactions seen at treatment doses, we do not recommend option D (mefloquine) unless the other options cannot be used. For option C, because there is more data on the efficacy of quinine in combination with doxycycline or tetracycline, these treatment combinations are generally preferred to quinine in combination with clindamycin.
5. Take with with food or whole milk. If patient vomits within 30 minutes of taking a dose, then they should repeat the dose.
6. US manufactured quinine sulfate capsule is in a 324 mg dosage; therefore 2 capsules should be sufficient for adult dosing. Pediatric dosing may be difficult due to unavailability of non-capsule forms of quinine.
7. For infections acquired in Southeast Asia, quinine treatment should continue for 7 days. For infections acquired elsewhere, quinine treatment should continue for 3 days.
8. Doxycycline is not indicated for use in children less than 8 years old. For children less than 8 years old with chloroquine-resistant *P. falciparum*, atovaquone-proguanil and artemether-lumefantrine are recommended treatment options; mefloquine can be considered if no other options are available. For children less than 8 years old with chloroquine-resistant *P. vivax*, mefloquine is the recommended treatment. If it is not available or is not being tolerated and if the treatment benefits outweigh the risks, atovaquone-proguanil or artemether-lumefantrine should be used instead.
9. Treatment with mefloquine is not recommended in persons who have acquired infections from Southeast Asia due to drug resistance.
10. Primaquine is used to eradicate any hypnozoites that may remain dormant in the liver, and thus prevent relapses, in *P. vivax* and *P. ovale* infections. Because primaquine can cause hemolytic anemia in G6PD-deficient persons, G6PD screening must occur prior to starting treatment with primaquine. For persons with borderline G6PD deficiency or as an alternate to the above regimen, primaquine may be given 45 mg orally one time per week for 8 weeks; consultation with an expert in infectious disease and/or tropical medicine is advised if this alternative regimen is considered in G6PD-deficient persons. Primaquine must not be used during pregnancy.
11. There are three options (A, B, or C) available for treatment of uncomplicated malaria caused by chloroquine-resistant *P. vivax*. High treatment failure rates due to chloroquine-resistant *P. vivax* have been well documented in Papua New Guinea and Indonesia. Rare case reports of chloroquine-resistant *P. vivax* have also been documented in Burma (Myanmar), India, and Central and South America. Persons acquiring *P. vivax* infections outside of Papua New Guinea or Indonesia should be started on chloroquine. If the patient does not respond, the treatment should be changed to a chloroquine-resistant *P. vivax* regimen and CDC should be notified (Malaria Hotline number listed above). For treatment of chloroquine-resistant *P. vivax* infections, options A, B, and C are equally recommended.

### Based on CDC Recommendations for Malaria in the US

**CDC Malaria Hotline: weekdays 8 am – 4:30 pm EST: (770) 488-7788**

**after hours, weekends and holidays: (770) 488-7100**

## INDEX

Note: Tables are noted with a *t*.

- A**-flavus, endophthalmitis ..... 35  
*A. fumigatus*, endophthalmitis ..... 35  
*A. hemolyticum*, pharyngitis ..... 40, 41  
 Abacavir (ABC) ..... 526  
 Abacavir + lamivudine (Epzicom) ..... 527  
 Abacavir + lamivudine +  
   zidovudine (trizivir) ..... 528  
 ABC (abacavir) ..... 526  
 ABCD ..... 541  
 ABC/3TC (Epzicom) ..... 527  
 ABC/3TC/ZDV (Trizivir) ..... 528  
 Abdominal abnormalities (DDx)  
   abdominal wall nodules ..... 486*t*  
   abdominal wall sinus tract ..... 485*t*  
   abdominal wall tenderness ..... 485*t*  
   abdominal tenderness  
     (2° to pancreatitis) ..... 485*t*  
   ascites ..... 487*t*  
   hepatomegaly ..... 487*t*  
   hepatosplenomegaly ..... 487*t*  
   left upper abdominal quadrant  
     tenderness (splenic tenderness) ..... 486*t*  
   perumbilical purpura  
     (thumbprint sign) ..... 486*t*  
   right lower quadrant tenderness ..... 486*t*  
   right upper quadrant mass ..... 486*t*  
   right upper quadrant  
     tenderness ..... 486*t*  
   right upper quadrant tympany ..... 486*t*  
   rose spots ..... 485*t*  
   splenomegaly ..... 487*t*  
 Abdominal tenderness,  
   abdominal abnormalities (DDx), ..... 485*t*  
 Abdominal wall sinus tract,  
   abdominal abnormalities (DDx) ..... 485*t*  
 Abdominal wall tenderness,  
   abdominal abnormalities (DDx) ..... 485*t*  
 Abelcet ..... 539  
 ABLC ..... 539  
 Abortion, septic ..... 113  
 Abscess  
   brain ..... 27  
   cold ..... 492*t*  
   gallbladder wall ..... 93  
   intraabdominal ..... 102  
   intrarenal ..... 109  
   ischiorectal ..... 151  
   liver ..... 95  
   lung ..... 67  
   necrotizing ..... 136  
   pancreatic ..... 95  
   pelvic ..... 102  
   perirectal ..... 151  
   prostatic ..... 110  
   renal ..... 109  
   skin ..... 140, 141  
   tuboovarian ..... 113  
 Absidia  
   cutaneous zygomycosis ..... 290  
   pulmonary zygomycosis ..... 269, 272  
 Acanthamoeba  
   granulomatous amebic  
     encephalitis ..... 261, 263  
   keratitis ..... 32  
   meningitis ..... 22  
 A-cillin ..... 533  
 Acinetobacter baumannii, acute  
   bacterial meningitis ..... 19  
 Acinetobacter baumannii, Iloffii,  
   calcoaceticus, haemolyticus ..... 218*t*  
 Acinetobacter sp. .... 200*t*  
 Acquired resistance ..... 2  
 Actinobacillus, endocarditis ..... 72  
 Actinomyces, brain abscess ..... 27, 28  
 Actinomyces israelii ..... 236*t*  
   cutaneous actinomycosis ..... 284, 286  
   pulmonary actinomycosis ..... 268, 270  
 Actinomyces odontolyticus ..... 236*t*  
 Actinomyces sp. .... 201*t*  
 Actinomycosis  
   cavitary infiltrates  
     (thick-walled) ..... 463

- unilateral ill-defined infiltrates
  - without effusion.....439
- Acute bacterial endocarditis (ABE).....74
- Acute bacterial meningitis
  - (ABM), empiric therapy ..... 18, 387
- Acute bacterial sinusitis .....30
- Acute exacerbation of chronic
  - bronchitis (AECB).....48
  - chronic medical prophylaxis .....365t
- Acute infections. (see specific clinical syndromes)
- Acute non-bacterial meningitis.....21
- Acute otitis media.....38, 393
- Acute rheumatic fever (ARF),
  - chronic medical prophylaxis .....365t
- Acyclovir .....529
  - pediatric dose .....414
- Adefovir dipivoxil.....530
- Adenovirus
  - conjunctivitis.....33
  - transplant infections.....163
  - unilateral focal segmental/lobar
    - infiltrate with effusion .....433
  - unilateral ill-defined infiltrates
    - without effusion.....437
- Adult immunizations.....378
- Bacille Calmette Guérin (BCG).....378t
- hemophilus influenzae .....378t
- hepatitis A.....378t
- hepatitis B.....378t
- herpes zoster (HZV).....379t
- influenza .....379t
- measles .....380t
- meningococcus (invasive
  - disease).....380t
- mumps.....380t
- papilloma (human) virus.....380t
- pertussis.....380t
- pneumococcus (*S. pneumoniae*).....381t
- rubella.....381t
- tetanus-diphtheria.....381t
- varicella (VZV) chickenpox.....382t
- Adult respiratory distress syndrome
  - (ARDS), bilateral infiltrates without
    - effusion .....453
- Adverse side effects, explanation .....523
- Aerobic GNBS
  - acute pancreatitis.....94
  - endocarditis.....75
  - IV line infections .....79
  - liver abscess .....95
  - pacemaker infections.....79
  - pancreatic abscess.....95
  - surgical prophylaxis.....356t
- Aerobic isolates, clinical
  - significance, pending
    - susceptibility testing.....208t
- Aeromonas, travel prophylaxis.....372t
- Aeromonas hydrophila.....218
- African tick bite fever.....294
- Agenerase .....543
- Agent-specific antibiotic resistance.....2
- Aggregatibacter (actinobacillus)
  - actinomycetem-comitans .....218t
- AIDS. *see* HIV
- Alcaligenes xylosoxidans .....219t
- Alcoholics, pneumonia.....66
- Alinia .....656
- Alkaline phosphatase (AP),
  - elevations, liver test
    - abnormalities (DDx) .....500t
- Allergic potential, explanation.....523
- Allogeneic BMT/lung transplants,
  - transplant prophylaxis
    - (BMT/SOT).....368t
- Alopecia, localized/generalized, head
  - abnormalities (DDx) .....478t
- Alternaria alternata,
  - cutaneous alternaria.....291
- Alveolar cell carcinoma
  - bilateral infiltrates without effusion....455
  - unilateral focal segmental/lobar
    - infiltrate with effusion .....435
  - unilateral focal segmental/lobar
    - infiltrate without effusion.....429
  - unilateral ill-defined infiltrates
    - with effusion .....446
  - unilateral ill-defined infiltrates
    - without effusion.....441
- Alveolar echinococcosis.....265
- cavitary infiltrates (thick-walled).....467
- Amantadine.....531

- AmBisome .....540
- Amebiasis .....276  
travel prophylaxis.....373t  
(see also *Entamoeba histolytica*)
- Amebic cysts, cavitary infiltrates  
(thick-walled) .....467
- American trypanosomiasis .....255, 256  
(see also Chagas' disease)
- Amikacin .....532  
pearls & pitfalls .....513  
susceptibility profiles .....198t  
pediatric dose .....414
- Amikin .....532
- Aminoglycosides  
dosing .....5  
pearls & pitfalls .....513  
susceptibility profiles .....198t
- Amoxicillin .....533  
pearls & pitfalls .....508  
susceptibility profiles .....186t  
pediatric dose .....415
- Amoxicillin/clavulanic acid .....534  
pearls & pitfalls .....508  
susceptibility profiles .....186t  
pediatric dose .....415
- Amoxicillin/clavulanic acid  
ES-600 .....535
- Amoxicillin/clavulanic  
acid XR .....536
- Amoxil .....533
- Amphotec .....541
- Amphotericin B .....537  
cholesteryl sulfate complex .....541  
colloidal dispersion .....541  
lipid-associated  
formulations .....525, 525t  
lipid complex .....539  
liposomal .....540  
pediatric dose .....415
- Ampicillin .....541  
pearls & pitfalls .....508  
susceptibility profiles .....186t  
pediatric dose .....415
- Ampicillin/sulbactam .....542  
pearls & pitfalls .....512
- susceptibility profiles .....186t  
pediatric dose .....415
- Amprenavir (APV) .....543
- Amylase, elevation pleural fluid  
abnormalities (DDx) .....504t
- Anaerobes, brain  
abscess .....27
- Anaerobic isolates, clinical  
significance, pending  
susceptibility testing .....236
- Anaerobic streptococci  
complicated skin/skin structure  
tissue infections .....135  
surgical prophylaxis .....356t
- Anaplasma phagocytophilum .....260
- Anaplasmosis (HGA) .....260
- Ancef .....557
- Ancobon .....615
- Ancylostoma braziliense,  
cutaneous larva migrans .....288
- Ancylostoma duodenale, hookworm .....278
- Anemia, acute, RBC abnormalities  
(DDx) .....497t
- Angiostrongyliasis .....279
- Angiostrongylus cantonensis .....279
- Angiotensin-converting enzyme  
levels (ACE), rheumatic test  
abnormalities (DDx) .....501t
- Angular chelitis, mouth  
abnormalities (DDx) .....482t
- Anidulafungin .....545
- Animal bites .....145  
pediatric .....411  
septic arthritis secondary to .....123  
travel precautions .....372t
- Anisocoria, neurological abnormalities  
(DDx) .....494t
- Anterior cervical adenopathy, lymph node  
abnormalities (DDx) .....488t
- Anthrax  
bioterrorist .....173  
post-exposure  
prophylaxis .....362t, 363t
- Antibiotic failure .....14
- Antibiotic failure (DDx)  
apparent, causes of .....14t

- Antibiotic prescribing, pitfalls in ..... 15
- Antibiotic resistance ..... 3t
- Antibiotic resistance, combination therapy and ..... 9t
- Antibiotic selection, factors in ..... 2
- antibiotic resistance ..... 2
- cost ..... 4
- with known isolate ..... 207t
- safety profile ..... 4
- spectrum ..... 2
- tissue penetration ..... 2
- Antibiotic selection based on resistance potential ..... 207t
- Antibiotic spectrum, defined ..... 2
- Antibiotic susceptibility profiles
- aminoglycosides, fluoroquinolones
- carbapenems, others ..... 198t
- cephalosporins ..... 191t
- penicillins, macrolides, tetracyclines, others ..... 186t
- Antibiotic therapy
- bactericidal vs. bacteriostatic therapy ..... 9
- bioavailability ..... 11t
- cytochrome P450 isoenzymes ..... 524t
- dosing ..... 4
- duration of therapy ..... 11
- empiric therapy ..... 12
- (see also Empiric therapy)
- intravenous vs. oral switch therapy ..... 10
- mono vs. combination therapy ..... 11
- pearls & pitfalls ..... 507
- PK/PD considerations ..... 10t
- resistance ..... 2, 3t, 9t
- selection (isolate known) ..... 207t
- Antifungal serum tests (DDx) ..... 500t
- Antimicrobial therapy.
- see Antibiotic therapy
- Anti-nuclear antibody titers (ANA), elevation, rheumatic test abnormalities (DDx) ..... 501t
- Antiretroviral therapy ..... 312t
- Anti-smooth muscle antibodies (ASM), elevation, rheumatic test abnormalities (DDx) ..... 501t
- Aortic graft infection ..... 83
- Aphthous ulcers ..... 44, 46
- Appendicitis ..... 102
- Aptivus (tipranavir) ..... 704
- APV (amprenavir) ..... 543
- Arachnia propionica ..... 237
- Arava ..... 634
- Arboviral encephalitis ..... 26, 390
- Arcanobacterium, pharyngitis ..... 40
- Arcanobacterium (Corynebacterium) haemolyticum ... 215t
- Argyll-Robertson pupils, eye abnormalities (DDx) ..... 479t
- Arm lesions
- nodular, extremity abnormalities (DDx) ..... 490t
- verrucous ..... 490t
- Arthritis, Lyme ..... 127, 128
- Arthritis, septic ..... 121
- extremity abnormality (DDx) ..... 490t
- pediatric ..... 407
- Arthroscopy, surgical prophylaxis ..... 357t
- ART therapy, HIV infection, ..... 313t
- Ascariasis ..... 278, 280
- Ascaris lumbricoides ..... 278, 280
- Ascites, abdominal abnormalities (DDx) ..... 487t
- Aspergilloma, pulmonary ..... 266, 267
- Aspergillosis, cutaneous ..... 290, 292
- Aspergillus ..... 242t
- acute pneumonia ..... 67
- bronchopulmonary aspergillosis (BPA) ..... 270
- chronic medical prophylaxis ..... 365t
- community-acquired pneumonia ..... 55
- cutaneous aspergillosis ..... 290, 292
- febrile neutropenia ..... 158
- HIV ..... 323
- organ transplants ..... 167, 168
- pneumonia
- acute invasive ..... 268, 271
- chronic ..... 268, 271
- pulmonary aspergilloma ..... 267
- sepsis ..... 155
- transplant infections ..... 164, 165
- Aspergillus galactomannan (AG) ..... 500t

- Aspiration pneumonia ..... 65
- Asplenia ..... 157
- Asplenia/impaired splenic function,  
chronic medical prophylaxis ..... 364t
- Asymptomatic bacteriuria in pregnancy,  
chronic medical prophylaxis ..... 364t
- Atazanavir (ATV) ..... 546
- Atelectasis, unilateral focal segmental/  
lobar infiltrate without effusion ..... 426
- Athlete's foot ..... 147
- Atovaquone ..... 548
- Atovaquone + proguanil ..... 548
- Atripla [efavirenz (EFV)+(emtricitabine  
(FTC)+tenofovir (TDF)] ..... 600
- ATV (atazanavir) ..... 546
- Atypical lymphocytes, WBC  
abnormalities (DDx) ..... 496t
- Augmentin ..... 534
- ES-600 ..... 535
- XR formulation ..... 537
- Avelox ..... 650
- AV graft infection ..... 83
- Avian influenza ..... 62
- community-acquired  
pneumonia ..... 51
- post-exposure prophylaxis ..... 359t
- Avycaz ..... 559
- Axillary adenopathy, lymph node  
abnormalities (DDx) ..... 488t
- Azactam ..... 552
- Azidothymidine ..... 718
- Azithromycin ..... 549
- pearls & pitfalls ..... 518
- susceptibility profiles ..... 186t
- pediatric dose ..... 415
- AZT (zidovudine) ..... 718
- Aztreonam ..... 552
- pearls & pitfalls ..... 511
- pediatric dose ..... 417
- B**-cepacia, community-acquired  
pneumonia ..... 54
- Babesia microti ..... 258, 260
- Babesiosis ..... 258, 260
- Bacillary angiomatosis ..... 289, 291
- Bacille Calmette Guérin (BCG)  
adult immunization ..... 379t
- granulomatous hepatitis ..... 96
- sepsis ..... 155
- Bacillus anthracis ..... 215
- anthrax ..... 173, 174
- post-exposure prophylaxis ..... 360t
- Bacillus cereus  
keratitis ..... 31
- subtilis, megaterium ..... 215t
- Back abnormalities (DDx)  
D'espine's sign ..... 484t
- spinal tenderness ..... 484t
- unilateral CVA tenderness ..... 484t
- Bacteremias transplant infections ..... 166
- Bacteremia vs positive BC's ..... 13
- Bacterial conjunctivitis ..... 33
- Bacterial endophthalmitis ..... 36
- Bacterial pharyngitis ..... 41
- Bacterial resistance, to antimicrobial  
therapy ..... 2
- Bacterial sinusitis ..... 30
- Bactericidal vs. bacteriostatic antibiotics ..... 9
- Bactericidal vs. bacteriostatic therapy ..... 9
- Bacteroides fragilis ..... 240t
- acute pancreatitis ..... 94
- decubitus ulcers ..... 137
- diabetic foot ulcers ..... 138
- endometritis ..... 113
- group ..... 240t
- intraabdominal abscess ..... 101
- ischioanal/perirectal abscess ..... 151
- liver abscess ..... 95
- osteomyelitis ..... 129
- pancreatic abscess ..... 95
- PID ..... 113
- salpingitis ..... 113
- sepsis ..... 151
- septic abortion ..... 113
- surgical prophylaxis ..... 354, 355t
- tuboovarian abscess ..... 113
- Bactrim ..... 707
- Balanitis ..... 116
- Balantidiasis ..... 278
- Balantidium coli ..... 278

- Baraclude .....607
- Bartonella henselae  
 cat scratch disease.....146  
 quintana, bacilliformis .....219t
- Bartonella sp. bacillary  
 angiomatosis.....291
- Basophilia WBC  
 abnormalities (DDx) .....496t
- Bedaquiline .....53
- Beef tapeworm .....281
- Bell's palsy, neurological abnormalities  
 (DDx).....494t
- BG ( $\beta$  1, 3 D-glucan).....500t  
 susceptibility profiles .....186t
- $\beta$ -hemolytic streptococci, usual/unusual  
 susceptibility patterns .....7t
- $\beta$ -lactams. (see individual penicillins  
 cephalosporins)
- $\beta$ -lactamase inhibitor combinations,  
 pearls & pitfalls .....512
- Biaxin .....580
- Biaxin XL .....582
- Bifidobacterium sp.....237
- Biliary tract penetration explanation .....523
- Biliary tract surgery surgical  
 prophylaxis.....355t
- Bilirubin, total, elevation, liver test  
 abnormalities (DDx) .....500t
- Bioavailability  
 explanation.....521  
 oral antimicrobials.....11t
- Bioterrorist agents in adults  
 treatment .....173  
 anthrax.....173  
 botulism .....174  
 Ebola.....174  
 pneumonic plague.....174  
 smallpox .....174  
 tularemic pneumonia .....173
- BK virus, transplant infections.....162
- Blastocystis.....277
- Blastocystis hominis.....277
- Blastomyces dermatitidis  
 cutaneous blastomycosis.....284, 285  
 epididymitis .....106  
 pulmonary blastomycosis .....267
- Blood, parasites, fungi, unusual  
 organisms in .....254
- Bloody pleural fluid abnormalities  
 (DDx).....503t
- Body fluid secretions, travel  
 precautions .....372t
- Body lice.....150
- Bone infections, empiric therapy of.....121
- Bone marrow transplant (BMT),  
 transplant infections.....160
- Bordetella pertussis .....219  
 community-acquired pneumonia.....49  
 post-exposure prophylaxis.....359t
- Borrelia burgdorferi .....233  
 Lyme disease.....127  
 meningitis.....21  
 pericarditis/myocarditis.....78  
 post-exposure prophylaxis.....361t
- Borrelia recurrentis .....234t  
 relapsing fever.....257
- Botulism .....170  
 bioterrorist .....174
- Boutonneuse fever.....296
- Brachiola, microsporidia enteritis .....335
- Brain  
 cysts/mass lesions in.....261  
 parasites, fungi, unusual  
 organisms in .....261
- Brain abscess.....27  
 cardiac source.....27  
 dental source.....27  
 mastoid/otitic source.....27  
 neurosurgical procedure.....27  
 open trauma .....27  
 pulmonary source.....28  
 sinus source.....27  
 transplant infections.....164
- Bronchiectasis.....67
- Bronchiolitis obliterans with  
 organizing pneumonia (BOOP),  
 bilateral infiltrates without  
 effusion .....457
- Bronchitis, acute bacterial exacerbation  
 of chronic (AECB).....48

- Bronchogenic carcinoma, unilateral  
 ill-defined infiltrates with effusion. . . .446  
 without effusion. . . . .440
- Bronchogenic cyst, cavitory  
 infiltrates (thin-walled). . . . .472
- Bronchopulmonary aspergillosis  
 (BPA) . . . . .270
- Brownish pleural fluid  
 abnormalities (DDx) . . . . .503t
- Brucella* sp. . . . .132, 219  
 endocarditis . . . . .73  
 meningitis. . . . .22  
 septic arthritis . . . . .121
- Brugia malayi*. . . . .254
- Bullae skin . . . . .288
- Bull neck, neck abnormalities  
 (DDx). . . . .483t
- Burkholderia cepacia* . . . . .220t
- Burkholderia pseudomallei*. . . . .220t
- Bursitis septic . . . . .121, 124
- Buruli ulcer. . . . .285, 287
- C**alf tenderness, extremity  
 abnormality (DDx) . . . . .491t
- California encephalitis (CE) . . . . .25
- Camelback fever pattern, fever curves  
 (DDx). . . . .475t
- Campylobacter* sp.  
 fetus. . . . .220  
 infectious diarrhea. . . . .86  
 jejuni. . . . .221  
 HIV. . . . .333  
 travel prophylaxis. . . . .372t
- CA-MRSA. *see* Community acquired  
 MRSA (CA-MRSA)
- c-ANCA, elevation, rheumatic test  
 abnormalities (DDx) . . . . .501t
- Candida albicans*. . . . .242t  
 candidemia. . . . .161  
 catheter-associated  
 candiduria. . . . .105  
 chorioretinitis. . . . .34  
 chronic medical prophylaxis . . . . .364t  
 endophthalmitis. . . . .35  
 esophagitis. . . . .84  
 febrile neutropenia . . . . .159  
 fungal cystitis. . . . .103  
 fungal peritonitis/abscess. . . . .100  
 hepatosplenic candidiasis . . . . .96  
 IV line infection. . . . .79  
 mucocutaneous candidiasis . . . . .147  
 onychomycosis. . . . .148  
 sepsis . . . . .153  
 thrush . . . . .43  
 vaginitis, in HIV . . . . .338
- Candida chorioretinitis* . . . . .34
- Candidiasis, oropharyngeal/  
 esophageal, HIV . . . . .336
- Candida non-albicans* . . . . .243  
 candidemia. . . . .161  
 chronic medical prophylaxis . . . . .364t  
 febrile neutropenia . . . . .159  
 fungal peritonitis/abscess. . . . .100  
 IV line infection . . . . .79  
 pacemaker infection. . . . .79  
 sepsis . . . . .153  
 catheter-associated candiduria . . . . .105  
 nodular/pustular candidiasis. . . . .290  
 prosthetic valve endocarditis . . . . .76
- Candida* vaginitis, balanitis . . . . .115
- Candidemia, transplant infections . . . . .161
- Capastat . . . . .553
- CAPD-associated peritonitis . . . . .102
- Capillaria philippinensis* . . . . .279
- Capillariasis . . . . .279
- Capnocytophaga  
 canimorsus/cynodegni . . . . .235t
- Capnocytophaga canimorsus (DF2),  
 animal bite wounds . . . . .144
- Capnocytophaga ochraceus . . . . .235t
- Capnophilic isolates,  
 clinical significance, pending  
 susceptibility testing. . . . .235t
- Capreomycin. . . . .553
- Carbapenems  
 pearls & pitfalls . . . . .511  
 susceptibility profiles . . . . .198t



Cardiac conditions, indications for infective endocarditis (IE) prophylaxis .....	369t	susceptibility profiles .....	191t
Cardiac valve replacement, surgical prophylaxis .....	355t	pediatric dose .....	416
Cardiobacterium hominis .....	221	Cefixime .....	562
endocarditis .....	72	pearls & pitfalls .....	511
Cardiovascular infections, empiric therapy .....	70	susceptibility profiles .....	191t
Carpal tunnel syndrome, extremity abnormalities (DDx) .....	490t	Cefizox .....	571
Caspofungin .....	553	Cefobid .....	562
pediatric dose .....	416	Cefoperazone .....	562
Catheter associated bacteriuria (CAB) .....	104	pearls & pitfalls .....	510
Catheter associated candiduria (CAC) .....	105	susceptibility profiles .....	191t
Cat scratch disease (CSD) .....	144	Cefotan .....	564
pediatric .....	411	Cefotaxime .....	563
Cavernous vein thrombosis .....	27	pearls & pitfalls .....	510
CD <sub>4</sub> cell counts, in HIV .....	302	susceptibility profiles .....	191t
Ceclor .....	555	pediatric dose .....	416
Cedax .....	570	Cefotetan .....	564
Cefaclor .....	555	pearls & pitfalls .....	510
pearls & pitfalls .....	510	susceptibility profiles .....	191t
susceptibility profiles .....	191t	Cefoxitin .....	565
Cefadroxil .....	556	pearls & pitfalls .....	510
pearls & pitfalls .....	509	susceptibility profiles .....	191t
susceptibility profiles .....	191t	Cefpodoxime .....	565
Cefamandole .....	557	pearls & pitfalls .....	511
pearls & pitfalls .....	510	susceptibility profiles .....	191t
susceptibility profiles .....	191t	Cefprozil .....	566
Cefazolin .....	557	pearls & pitfalls .....	510
pearls & pitfalls .....	509	susceptibility profiles .....	191t
susceptibility profiles .....	191t	Ceftolozane/tazobactam .....	572
pediatric dose .....	416	Ceftaroline fosamil .....	567
Cefdinir .....	558	pearls & pitfalls .....	510
pearls & pitfalls .....	511	susceptibility profiles .....	192t
susceptibility profiles .....	191t	Ceftazidime .....	569
Cefditoren .....	559	pearls & pitfalls .....	510
pearls & pitfalls .....	511	susceptibility profiles .....	192t
susceptibility profiles .....	191t	pediatric dose .....	416
Cefepime .....	560	Ceftazidime/azibactam .....	569
pearls & pitfalls .....	511	pearls & pitfalls .....	512
susceptibility profiles .....	191t	susceptibility profiles .....	191t
Cefepime .....	560	Ceftazidime/avibactam .....	569
pearls & pitfalls .....	511	Ceftibuten .....	570
susceptibility profiles .....	191t	pearls & pitfalls .....	511
susceptibility profiles .....	191t	susceptibility profiles .....	191t

- Ceftin .....574
- Ceftizoxime .....571
- pearls & pitfalls .....510
- susceptibility profiles .....191t
- Ceftolozane/tazobactam .....572
- pearls & pitfalls .....512
- susceptibility profiles .....191t
- Ceftolozane/tazobactam .....572
- Ceftriaxone .....572
- pearls & pitfalls .....510
- susceptibility profiles .....191t
- pediatric dose .....417
- Cefuroxime .....574
- pearls & pitfalls .....510
- susceptibility profiles .....191t
- pediatric dose .....417
- Cefzil .....566
- Cellulitis
- above-the-waist .....132
- below-the-waist .....133
- facial .....29
- pediatric .....411
- periorbital .....29
- severe .....136
- Central IV line
- (CVC) infections .....81
- Cephalexin .....575
- Cephalosporins, susceptibility profiles .....191t
- Cerebral amebiasis .....262
- Cerebral coenurosis .....263
- Cerebral echinococcosis (hydatid cyst disease) .....263
- Cerebral gnathostomiasis .....263
- Cerebral nocardiosis .....262
- Cerebral paragonimiasis (lung fluke) .....262, 264
- Cerebral toxoplasmosis .....262
- Cerebrospinal fluid (CSF)
- abnormalities (DDx) .....505
- Cerebrospinal fluid penetration, explanation .....526
- Cervicitis
- gonococcal .....114
- non-gonococcal .....114, 115
- Cestodes (tapeworms), intestinal .....281
- Chagas' disease (American trypanosomiasis) .....255, 256
- blood .....256
- cerebral .....262, 264
- heart .....273
- Chancroid .....117
- post-exposure prophylaxis .....360t
- Chest abnormalities (DDx)
- chest dullness, consolidation/  
  pleural effusion .....484t
- chest wall sinuses .....484t
- costochondral tenderness .....484t
- diffuse wheezing .....484t
- shoulder tenderness .....484t
- spontaneous pneumothorax .....484t
- sternal tenderness .....484t
- trapezius tenderness .....484t
- Chest dullness, consolidation/pleural effusion, chest
- abnormalities (DDx) .....484t
- Chest wall sinuses, chest
- abnormalities (DDx) .....484t
- Chest x-ray atlas .....425
- bilateral infiltrates with effusion .....458
- infectious causes .....458
- noninfectious causes .....460
- bilateral infiltrates
- without effusion .....448
- infectious causes .....448
- non-infectious causes .....453
- cavitary infiltrates (thick-walled) .....463
- infectious causes .....464
- infectious causes based  
    on speed of cavitation .....463
- non-infectious causes .....468
- cavitary infiltrates (thin-walled) .....470
- infectious causes .....470
- non-infectious causes .....472
- unilateral focal segmental/lobar
- infiltrate with effusion .....431
- infectious causes .....431
- non-infectious causes .....434
- unilateral focal segmental/lobar
- infiltrate without effusion .....426
- infectious causes .....426
- non-infectious causes .....428
- unilateral ill-defined infiltrates with  
  effusion .....443

infectious causes	443	Cholangitis	93
non-infectious causes	444	Cholecystitis	92
unilateral ill-defined infiltrates		empysematous	92
without effusion	436	Chorioretinitis	34
infectious causes	436	eye abnormalities (DDx)	480t
non-infectious causes	440	Chromobacterium violaceum	221
Chickenpox		Chromomycosis	289
pediatric	412	Chronic bacterial sinusitis	31
post-exposure		Chronic fatigue syndrome (CFS)	42
prophylaxis	360	Chronic infections. (see	
therapy	142	specific clinical syndromes)	
Chlamydia trachomatis		Chronic mastoiditis	39
conjunctivitis	33	Chronic medical prophylaxis	364t
dysuria-pyuria		acute exacerbation of chronic	
syndrome	103	bronchitis (AECB)	365t
endometritis	113	acute rheumatic fever (ARF)	365t
epididymitis	106	asplenia/impaired splenic	
lymphogranuloma		function	364t
venereum	118	asymptomatic bacteriuria in	
PID	113	pregnancy	364t
post-exposure		neonatal group B streptococcal	
prophylaxis	360t	(GBS) infection	
salpingitis	113	(primary prevention)	365t
septic abortion	113	prophylaxis of fungal infections in	
tuboovarian abscess	113	neutropenic patients	364t
urethritis/cervicitis	114	recurrent genital herpes	
Chlamydophilia		(less than 6 episodes/year)	364t
community-acquired pneumonia	49	recurrent genital herpes	
pharyngitis	42	(more than 6 episodes/year)	365t
unilateral ill-defined		UTIs (recurrent)	364t
infiltrates without effusion	437	Chronic meningitis	22
Chlamydophilia (Chlamydia)		Chryseobacterium (Flavobacterium)	
pneumoniae	58	meningosepticum	221
community-acquired		Chyluria, urinary	
pneumonia	49	abnormalities (DDx)	503t
pharyngitis	42	Cidofovir	577
Chlamydophilia (Chlamydia) psittaci		Cipro	578
(psittacosis)		Ciprofloxacin	578
community-acquired		pearls & pitfalls	515
pneumonia	51	susceptibility profiles	198t
endocarditis	72	extended-release	579
zoonotic atypical pneumonia	60	Cipro XR	579
Chloramphenicol	576	Citrobacter sp.	222
pearls & pitfalls	512	Cladosporium/Fonsecaea,	
Chloromycetin	576	myositis	298

- Claforan .....563
- Clarithromycin .....580  
 pearls & pitfalls .....518  
 susceptibility profiles .....186t
- Clarithromycin XL .....582
- Cleocin .....583
- Clindamycin .....583  
 pearls & pitfalls .....513  
 susceptibility profiles .....186t  
 pediatric dose .....417
- Clinical and Laboratory  
 Standards Institute (CLSI) .....8
- Clonorchiasis .....273
- Clonorchis sinensis .....273
- Clostridial myonecrosis .....134, 135
- Clostridium botulinum* .....237  
 botulism .....171
- Clostridium difficile* .....238  
 diarrhea .....86  
 colitis .....86
- Clostridium difficile* diarrhea,  
 HIV .....333
- Clostridium perfringens*  
 cholecystitis .....92  
*C. septicum*, *C. novyi* .....238
- Clostridium sordelli*  
 septic abortion .....113  
 toxic shock syndrome .....169
- Clostridium* sp. complicated  
 skin/skin structure tissue infections .....134
- Clostridium tetani* .....238  
 tetanus .....170
- Clubbing, extremity  
 abnormalities (DDx) .....490t
- CNS infections, empiric therapy .....18
- CNS shunt, surgical prophylaxis .....355t
- CNS shunt infections, acute  
 bacterial meningitis .....21
- Coccidioides immitis*  
 chronic septic arthritis .....126  
 cutaneous coccidioidomycosis .....286  
 fungal arthritis .....123  
 HIV .....339  
 pulmonary coccidioidomycosis .....271
- Coccidiomycosis  
 cavitory infiltrates (thin-walled) .....470  
 cutaneous .....286  
 meningitis .....22  
 unilateral focal segmental/lobar  
 infiltrate with effusion .....432
- Coin lesions, pulmonary .....266
- Cold abscesses, cutaneous,  
 extremity abnormalities (DDx) .....492t
- Cold agglutinins, elevated,  
 serum test abnormalities (DDx) .....499t
- Colistin methanesulfonate .....584
- Colistin/Polymyxin E .....584
- Colon surgery, surgical  
 prophylaxis .....356t
- Coly-Mycin M .....584
- Combination therapy vs.  
 monotherapy .....9
- Combivir [Lamivudine (3TC) +  
 Zidovudine (ZDV)] .....633
- complicated skin/skin structure  
 tissue infections .....135
- pyomyositis/necrotizing  
 abscesses .....135
- Community-acquired  
 pneumonia (CAP) .....49
- atypical pathogens .....56  
 avian influenza ( $H_5N_1$ ,  $H_7N_9$ ) .....62  
*C. pneumoniae* .....51  
*C. psittaci* (psittacosis) .....51  
*Coxiella burnetii* (Q fever) .....51  
*Francisella tularensis* (tularemia) .....51  
 influenza A .....51  
 Legionella .....51  
*M. pneumoniae* .....51  
 swine influenza ( $H_1N_1$ ) .....51
- pertussis .....56
- typical pathogens .....56  
*B. pertussis* .....50  
*H. influenzae* .....50  
*M. catarrhalis* .....50  
*S. pneumoniae* .....50  
*S. aureus* (MSSA/MRSA) .....65

Complera [Emtricitabine (FTC) + Rilpivirine (RPV) + Tenofovir (TDF)] . . . . .	.605	Crabs . . . . .	150
Complicated skin/skin structure tissue infections (cSSSIs) . . . . .	134	Cranial nerve abnormalities, unilateral/ bilateral, neurological abnormalities (DDx) . . . . .	494t
CO-MRSA. (see Community onset MRSA)		Cresemba . . . . .	626
Concentration-dependent kinetics, antibiotics with . . . . .	10t	Crimson crescents, mouth abnormalities (DDx) . . . . .	483t
Condylomata acuminata . . . . .	118	Crixivan (Indinavir) . . . . .	625
Congestive heart failure bilateral infiltrates with effusion . . . . .	458	Cryptosporidia enteritis, HIV/AIDS . . . . .	334
unilateral ill-defined infiltrates without effusion . . . . .	443	Cryptococcomas/meningitis . . . . .	261
Conjunctival suffusion, eye abnormalities (DDx) . . . . .	479t	Cryptococcosis, cutaneous . . . . .	290
Conjunctivitis . . . . .	32	Cryptococcus meningitis . . . . .	24
bacterial . . . . .	32	Cryptococcus neoformans . . . . .	243
trachoma . . . . .	33	cryptococcomas/meningitis . . . . .	261
unilateral/bilateral, eye abnormalities (DDx) . . . . .	479t	cutaneous cryptococcosis . . . . .	290
viral . . . . .	33	meningitis . . . . .	22
Copegus . . . . .	680	HIV . . . . .	329
Corneal haziness, eye abnormalities (DDx) . . . . .	480t	pulmonary cryptococcosis . . . . .	269
Corneal ulcers, eye abnormalities (DDx) . . . . .	480t	transplant infections . . . . .	164
Coronary artery bypass graft (CABG), surgical prophylaxis . . . . .	355t	unilateral ill-defined infiltrates without effusion . . . . .	439
Corynebacterium diphtheriae . . . . .	216	Cryptosporidia . . . . .	277
diphtheria . . . . .	172	chronic watery diarrhea, HIV/AIDS . . . . .	90
pericarditis/myocarditis . . . . .	78	travel prophylaxis . . . . .	371t
pharyngitis . . . . .	40	Cryptosporidiosis . . . . .	277
post-exposure prophylaxis . . . . .	358t	CSF	
Corynebacterium hemolyticum, pharyngitis . . . . .	40	cysts/mass lesions in . . . . .	261
Corynebacterium jeikeium (JK) . . . . .	216	parasites, fungi, unusual organisms in . . . . .	261
Corynebacterium minutissimum, erythrasma . . . . .	284	CSF abnormalities (DDx)	
Costochondral tenderness, chest abnormalities (DDx) . . . . .	484t	CSF eosinophils . . . . .	506t
Coxiella burnetii (Q fever) community-acquired pneumonia . . . . .	51	CSF lactic acid levels . . . . .	506t
endocarditis . . . . .	72	CSF plasma cells . . . . .	506t
zoonotic atypical pneumonia . . . . .	60	CSF with negative Gram stain and predominantly lymphocytes/ decreased glucose . . . . .	506t
Coxsackie A virus herpangina . . . . .	44	CSF with negative Gram stain and predominantly lymphocytes/ normal glucose . . . . .	505t
pericarditis/myocarditis . . . . .	78	CSF with negative Gram stain and predominantly PMNs/decreased glucose . . . . .	505t
		highly elevated CSF protein . . . . .	506t
		purulent CSF with negative Gram stain . . . . .	505t
		RBCs in CSF . . . . .	505t

- CSF Gram stain, clinical use ..... 246t
- CSF lactic acid levels, elevation, CSF abnormalities (DDx) ..... 506t
- CSF protein, highly elevated, CSF abnormalities (DDx) ..... 506t
- Cubicin ..... 589
- Cutaneous actinomycosis ..... 284
- Cutaneous alternaria ..... 289, 291
- Cutaneous amebiasis ..... 284, 286
- Cutaneous aspergillosis ..... 290, 292
- Cutaneous blastomycosis ..... 284
- Cutaneous coccidioidomycosis ..... 284
- Cutaneous cold abscesses, extremity abnormalities (DDx) ..... 492t
- Cutaneous cryptococcosis ..... 290, 293
- Cutaneous fusarium ..... 289, 291
- Cutaneous gnathostomiasis ..... 288, 289
- Cutaneous histoplasmosis ..... 284, 285, 290, 293
- Cutaneous larva migrans (creeping eruption) ..... 288
- Cutaneous leishmaniasis (Old World/New World) ..... 283
- Cutaneous MAI ..... 285
- Cutaneous mycobacteria ..... 285
- Cutaneous nocardiosis ..... 284
- Cutaneous onchocerciasis ..... 290, 294
- Cutaneous penicillium ..... 289
- Cutaneous prototheca ..... 289, 292
- Cutaneous sporotrichosis ..... 290, 293
- Cutaneous zygomycosis (mucormycosis) ..... 290
- CVA tenderness, unilateral, back abnormalities (DDx) ..... 484t
- Cycloserine ..... 586
- Cyclospora ..... 277
- diarrhea ..... 88
- travel prophylaxis ..... 371t
- Cystic bronchiectasis, cavitory infiltrates (thin-walled) ..... 470
- Cystic fibrosis ..... 67
- Cystic lesions/masses
- CSF/brain ..... 261
- liver flukes ..... 273
- pulmonary ..... 265
- Cystitis ..... 103
- bacterial ..... 103
- candidal ..... 103
- Cystoscopy, surgical prophylaxis ..... 357t
- Cysts/masses, infected ..... 140
- Cytochrome P450 isoenzymes ..... 524t
- Cytoid bodies (cotton wool spots), eye abnormalities (DDx) ..... 480t
- Cytomegalovirus (CMV)
- bilateral infiltrates without effusion ..... 450
- colitis ..... 334
- community-acquired pneumonia ..... 55
- cytomegalovirus colitis ..... 87
- encephalitis ..... 25
- HIV infection ..... 329
- esophagitis ..... 84
- HIV infection ..... 334
- hepatitis ..... 99
- meningitis ..... 22
- pharyngitis ..... 40
- retinitis, HIV infection ..... 327, 328
- transplant infections ..... 162
- Cytovene ..... 620
- D**-actylitis, extremity abnormalities (DDx) ..... 490t
- Dalbavancin ..... 587
- pearls & pitfalls ..... 519
- susceptibility profiles ..... 198t
- Dalbavancin ..... 587
- Dalvance ..... 587
- Dapsone ..... 588
- Daptomycin ..... 589
- pearls & pitfalls ..... 517
- susceptibility profiles ..... 198t
- Daraprim ..... 674
- Darunavir ..... 591
- Daskil ..... 697
- ddC (zalcitabine) ..... 717
- ddl (didanosine) ..... 594
- Deafness
- acute, ear abnormalities (DDx) ..... 481t
- transient, neurological abnormalities (DDx) ..... 493t
- Decubitis ulcers ..... 137, 139
- Delavirdine (DLV) ..... 593

- Dental infections, severe ..... 46, 47
- Dermatophyte nail infections..... 149
- D'espines sign, back  
abnormalities (DDx) ..... 484t
- Desquamation of hands/feet,  
extremity abnormalities (DDx) ..... 493t
- d4T (stavudine) ..... 689
- Diabetes mellitus, chronic  
osteomyelitis ..... 130
- Diabetic foot ulcers ..... 138, 139
- Diarrhea, infectious..... 86  
acute watery ..... 86, 89  
C. difficile ..... 86  
chronic, watery ..... 88, 90  
typhoid/enteric fever ..... 88, 89
- Didanosine EC (ddI) ..... 594
- Dientamoeba fragilis ..... 276
- Dientamoebiasis ..... 276, 277
- Differential diagnosis ..... 474  
abdominal abnormalities ..... 485t  
back abnormalities ..... 484t  
chest abnormalities ..... 484t  
CSF abnormalities ..... 505t  
ear abnormalities ..... 481t  
extremity abnormalities ..... 490t  
eye abnormalities ..... 478t  
fever curves ..... 475t  
general appearance  
abnormalities ..... 476t  
genitourinary abnormalities ..... 489t  
head abnormalities ..... 477t  
heart abnormalities ..... 485t  
liver test abnormalities ..... 500t  
lymph node abnormalities ..... 488t  
mouth abnormalities ..... 482t  
nasal abnormalities ..... 481t  
neck abnormalities ..... 483t  
neurological abnormalities ..... 493t  
platelet abnormalities ..... 498t  
pleural fluid abnormalities ..... 503t  
RBC abnormalities ..... 497t  
rheumatic test abnormalities ..... 501t  
serum test abnormalities ..... 498t  
urinary abnormalities ..... 501t  
WBC abnormalities ..... 495t
- Diflucan ..... 614
- Digital gangrene, extremity  
abnormalities (DDx) ..... 490t
- Diphtheria ..... 170, 172  
adult immunization ..... 381t  
post-exposure prophylaxis ..... 359t  
travel prophylaxis ..... 375t
- Diphyllobothrium latum, fish  
tapeworm ..... 281
- Dirofilaria immitis, dog  
heartworm ..... 266
- Diskitis, pediatric ..... 409, 410
- Distal small bowel surgery,  
surgical prophylaxis ..... 356t
- Diverticulitis ..... 102
- DLV (delavirdine) ..... 593
- Dog heartworm ..... 266
- Dolutegravir/Abacavir/Lamivudine ..... 314
- Dolutegravir (Tivicay) DTG ..... 595
- Donovan bodies, granuloma  
inguinale ..... 118
- Doribax ..... 596
- Doripenem ..... 596  
pearls & pitfalls ..... 511  
susceptibility profiles ..... 198t
- Dosing  
adjustments, explanation ..... 522  
antibiotics, factors in ..... 4  
recommendations, explanation ..... 523
- Double quotidian fevers, fever  
curves (DDx) ..... 475t
- Double stranded DNA (DS-DNA),  
elevation, rheumatic test  
abnormalities (DDx) ..... 501t
- Doxycycline ..... 597  
pearls & pitfalls ..... 512  
susceptibility profiles ..... 186t
- Dracunculus medinensis,  
Guinea worm ..... 288
- Drug fever ..... 13
- Drug fever, clinical features ..... 13t
- Drug-induced bilateral infiltrates  
with effusion ..... 458  
without effusion ..... 448

- Drug interactions, explanations. . . . .523
- DRV (darunavir) . . . . .591
- Dry eyes, eye abnormalities (DDx), . . . . .480t
- DTG (dolutegravir) . . . . .595
- Duration of antibiotic therapy . . . . .11
- Duration of therapy . . . . .11
- Duricef. . . . .556
- Dwarf tapeworm. . . . .281
- Dysentery, acute
- E. histolytica* . . . . .91
- Shigella. . . . .91
- Dysuria-pyuria syndrome (acute urethral syndrome) . . . . .103
- E**-basidiobolus, entomorphthoromycosis . . . . .291
- E. conidiobolus*, entomorphthoromycosis . . . . .291
- Ear abnormalities (DDx)
- acute deafness . . . . .481t
- external ear lesions . . . . .481t
- Ear lesions, external, ear abnormalities (DDx) . . . . .481t
- East African trypanosomiasis . . . . .256
- Eastern equine encephalitis (EEE). . . . .25
- Ebola virus . . . . .174
- Ebstein Barr virus (EBV)
- hepatitis. . . . .99
- meningitis. . . . .21, 23
- pharyngitis . . . . .40, 42
- transplant infections. . . . .163t
- Ecalta . . . . .545
- Echinococcus granulosus
- cerebral . . . . .261, 263
- hepatic echinococcosis. . . . .274
- Echinococcus multilocularis
- alveolar echinococcosis. . . . .265
- cerebral . . . . .261, 263
- Edema, generalized, general appearance abnormalities (DDx). . . . .476t
- Edematous pancreatitis . . . . .94
- Edurant (rilpivirine) . . . . .684
- Edwardsiella tarda. . . . .222t
- Efavirenz (EFV) . . . . .599
- Efavirenz + emtricitabine + tenofovir disoproxil fumarate (Atripla) . . . . .600
- EFV (efavirenz) . . . . .599
- EFV/FTC/TDF (Atripla) . . . . .600
- Ehrlichia chaffeensis (HME) . . . . .260
- Ehrlichia ewubguum . . . . .258
- Ehrlichiosis . . . . .258
- Ehrlichiosis (HME) . . . . .260
- Eikenella corrodens . . . . .235t
- endocarditis . . . . .72
- human bite wounds. . . . .144
- septic arthritis . . . . .123
- Elephant skin, general appearance abnormalities (DDx) . . . . .477t
- Elvitegravir + Combicistat + Emtricitabine + Tenofovir (Stribild) . . . . .602
- EMB. . . . .611
- Empyema, cavitory infiltrates (thin-walled). . . . .472
- Empysematous cholecystitis . . . . .92
- Empiric antibiotic therapy, antibiotic spectrum as basis for . . . . .2
- Empiric therapy
- bone and joint infections . . . . .121
- pediatric . . . . .407
- cardiovascular infections. . . . .70
- CNS infections. . . . .18
- pediatric . . . . .387
- gastrointestinal tract infections . . . . .84
- pediatric . . . . .403
- genitourinary tract infections . . . . .103
- HEENT infections . . . . .29
- pediatric . . . . .391
- lower respiratory tract infections. . . . .48
- pediatric . . . . .396
- sexually transmitted diseases . . . . .114
- skin and soft tissue infections. . . . .132
- pediatric . . . . .411
- syndrome based. . . . .17
- vascular infections, pediatric . . . . .402
- Empyema . . . . .67
- subdural. . . . .27
- Emtricitabine (FTC) . . . . .603
- Emtricitabine + Rilpivirine + Tenofovir (Complera) . . . . .605



- Emtricitabine (FTC) + tenofovir (TDF)  
 [Truvada] .....604
- Emtriva (emtricitabine) .....603
- Encephalitis .....25
- amebic, granulomatous .....263
  - arboviral .....26
  - CMV, HIV .....328
  - HHV-6 .....26
  - HSV-1 .....26
  - Listeria .....26
  - Mycoplasma .....26
  - organ transplants .....164
  - pediatric .....389
  - primary amebic meningo-  
 encephalitis .....261
  - toxoplasma, HIV .....331
- Encephalopathy, acute, neurological  
 abnormalities (DDx) .....493t
- Endocarditis
- acute bacterial (ABE) .....75, 76
    - IV drug abusers .....75  - prosthetic valve (PVE) .....76
  - subacute bacterial (SBE) .....70
    - culture negative .....72, 74
    - GI/GU source .....71, 73
    - no obvious source .....70, 73
- Endocarditis prophylaxis for  
 above-the-waist (dental, oral,  
 esophageal, respiratory tract)  
 procedures
- IV prophylaxis .....370t
  - oral prophylaxis .....370t
- Endocarditis prophylaxis for below-the-  
 waist (genitourinary, gastrointestinal)  
 procedures
- IV prophylaxis .....371t
  - oral prophylaxis .....370t
- Endometritis .....113
- Endomyometritis .....113
- Endophthalmitis .....35
- bacterial .....35, 36
  - eye abnormalities (DDx) .....480t
  - fungal .....35, 36
  - infected lens implant .....35, 36
  - tuberculosis .....35, 36
- ENF (enfuvirtide) .....606
- Enfuvirtide (T20) .....606
- Enlarged greater auricular nerve,  
 neck abnormalities (DDx) .....483t
- Entamoeba histolytica .....274
- cerebral amebiasis .....261, 262
  - cutaneous amebiasis .....284, 286
  - dysentery .....88, 91
  - hepatic amebiasis .....274
  - liver abscess .....95
  - pulmonary amebiasis .....265
- Entecavir .....607
- Enterobacteriaceae. *see also*  
*Citrobacter, Edwardsiella;*  
*Enterobacter, Escherichia coli, Klebsiella,*  
*Proteus, Providencia, Salmonella, Serratia,*  
*Shigella*
- acute bacterial
- pyelonephritis .....107, 108
- bacterial cystitis .....103, 104
- brain abscess .....27
- CAPD-associated peritonitis .....101, 102
- chronic pyelonephritis .....108
- complicated skin/skin structure  
 tissue infections .....135
- endophthalmitis .....36
- febrile neutropenia .....158
- intraabdominal abscess .....101
- ischio-rectal/perirectal abscess .....151
- osteomyelitis .....129
- prostatitis/prostatic abscess .....110
- prosthetic valve endocarditis .....76
- renal abscess .....109
- sepsis .....151
- septic arthritis .....122, 124
- solid organ transplant .....160
- suppurative parotitis .....40
- urosepsis .....111
- Enterobacter sp. ....188
- acute bacterial meningitis .....19
  - brain abscess .....27
  - burn wounds .....145
  - sepsis .....152

- Enterobius vermicularis,  
   pinworm ..... 279, 280
- Enterococci  
   cholangitis ..... 93  
   cholecystitis ..... 92  
   chronic medical prophylaxis ..... 364t  
   gallbladder abscess ..... 93  
   liver abscess ..... 95  
   vascular graft infections ..... 83
- Enterococcus faecalis (VSE) ..... 210t  
   bone marrow transplant  
     infection ..... 160  
   catheter-associated bacteriuria ..... 104  
   cholangitis ..... 93  
   cholecystitis ..... 92  
   cystitis ..... 103  
   endocarditis ..... 71  
   gallbladder wall abscess ..... 93  
   liver abscess ..... 95  
   pyelonephritis ..... 108  
   sepsis ..... 151  
   surgical prophylaxis ..... 355t  
   urosepsis ..... 111
- Enterococcus faecium (VRE) ..... 211t  
   catheter-associated bacteriuria ..... 104  
   endocarditis ..... 71  
   pyelonephritis ..... 107  
   sepsis ..... 154  
   surgical prophylaxis ..... 357t  
   transplant infections ..... 160
- Enterocytozoon bienuesi,  
   microsporidia enteritis ..... 335
- Enteroviruses  
   meningitis ..... 21  
   travel prophylaxis ..... 373t
- Entomophthoromycosis ..... 289
- Eosinopenia (DDx) ..... 496t
- Eosinophilia, WBC abnormalities  
   (DDx) ..... 496t
- Eosinophilic folliculitis ..... 297
- Eosinophils  
   CSF abnormalities (DDx) ..... 506t  
   pleural effusion profiles in  
     various diseases (DDx) ..... 504t
- Eosinophiluria, urinary  
   abnormalities (DDx) ..... 504t
- Epidemic (louse-borne)  
   typhus ..... 294, 295
- Epidermophyton floccosum  
   onychomycosis ..... 148  
   tinea corporis ..... 147
- Epididymitis, acute and  
   chronic ..... 106
- Epididymoorchitis, genitourinary  
   abnormalities (DDx) ..... 489t
- Epiglottitis ..... 47
- Episcleritis, eye  
   abnormalities (DDx) ..... 480t
- Epistaxis, nasal  
   abnormalities (DDx) ..... 481t
- Epirochlear adenopathy,  
   lymph node abnormalities (DDx) ... 488t
- Epivir (lamivudine) ..... 632
- Epzicom [Abacavir (ABC) +  
   Lamivudine (3TC)] ..... 527
- Eraxis ..... 545
- Ertapenem ..... 608  
   pearls & pitfalls ..... 511  
   susceptibility profiles ..... 198t
- Erysipelas ..... 132, 133
- Erysipelothrix rhusiopathiae ..... 216t
- Erythema multiforme, extremity  
   abnormalities (DDx) ..... 493t
- Erythema nodosum, extremity  
   abnormalities (DDx) ..... 493t
- Erythrasma ..... 283, 284
- Erythrocyte sedimentation rate (ESR),  
   serum test abnormalities  
   (DDx) ..... 498t
- Erythroderma, general appearance  
   abnormalities (DDx) ..... 476t
- Erythromycin, pediatric dose ..... 418  
   pearls & pitfalls ..... 518  
   susceptibility profiles ..... 190t  
   pediatric dose ..... 418
- Erythromycin lactobionate, base  
   (various) ..... 609
- Eschar, extremity  
   abnormalities (DDx) ..... 492t
- Escherichia coli ..... 223  
   acute bacterial meningitis ..... 19

- catheter-associated bacteriuria . . . . .104  
 cellulitis . . . . .133  
 cholangitis . . . . .93  
 cholecystitis . . . . .92  
 decubitus ulcers . . . . .137  
 diabetic foot ulcers . . . . .138  
 dysuria-pyuria syndrome . . . . .103  
 gallbladder wall abscess . . . . .93  
 infectious diarrhea . . . . .86  
 ischemic foot ulcers . . . . .139  
 osteomyelitis . . . . .130  
 pneumonia . . . . .68  
 sepsis . . . . .152  
 skin abscesses/infected cysts . . . . .141  
 surgical prophylaxis . . . . .355t  
 travel prophylaxis . . . . .372t  
 Esophageal candidiasis, HIV . . . . .336  
 Esophagitis . . . . .84  
   fungal . . . . .84  
   viral . . . . .84, 85  
 Ethambutol . . . . .611  
 Ethionamide . . . . .611  
 Ethmoid sinus, acute bacterial  
   sinusitis . . . . .31  
 ETR (etravirine) . . . . .612  
 Etravirine (ETR) . . . . .612  
 Eubacterium sp. . . . .239t  
 EVG/COBI/FTC/TDF (Stribild) . . . . .602  
 Excreted unchanged, explanation . . . . .522  
 External otitis . . . . .37  
   benign . . . . .37  
   malignant . . . . .37  
 Extracellular debris, pleural fluid  
   abnormalities (DDx) . . . . .504t  
 Extremity abnormalities (DDx)  
   arthritis (septic/reactive) . . . . .491t  
   calf tenderness . . . . .491t  
   carpal tunnel syndrome . . . . .490t  
   clubbing . . . . .490t  
   cutaneous cold abscesses . . . . .492t  
   dactylitis . . . . .490t  
   desquamation of hands/feet . . . . .493t  
   digital gangrene . . . . .490t  
   edema of dorsum of hand/foot . . . . .490t  
   erythema multiforme . . . . .493t  
   erythema nodosum . . . . .493t  
   eschar . . . . .492t  
   foot/leg ulcers . . . . .491t  
   hyperpigmented shins . . . . .492t  
   leg edema, unilateral/bilateral . . . . .491t  
   lymphangitis . . . . .490t  
   migratory rashes . . . . .492t  
   nodular arm/leg lesions . . . . .492t  
   nodular lymphangitis . . . . .490t  
   painful leg nodules . . . . .492t  
   painless purple palm/sole lesions . . . . .490t  
   palpable purpura . . . . .492t  
   papular axillary lesions . . . . .491t  
   plantar hyperkeratoses . . . . .493t  
   purple nodules . . . . .492t  
   rash of palms/soles . . . . .492t  
   scleroderma . . . . .492t  
   splinter hemorrhages . . . . .490t  
   tender fingertips . . . . .490t  
   tender muscles . . . . .491t  
   tenosynovitis . . . . .491t  
   thigh tenderness (bilateral) . . . . .491t  
   thrombophlebitis . . . . .491t  
   verrucous foot/leg lesions . . . . .491t  
   verrucous hand/arm lesions . . . . .490t  
   wrist swelling . . . . .491t  
 Eye abnormalities (DDx)  
   Argyll-Robertson pupils . . . . .479t  
   bilateral lid edema . . . . .478t  
   bilateral upper lid edema . . . . .478t  
   chorioretinitis . . . . .480t  
   conjunctivitis, unilateral/bilateral . . . . .479t  
   conjunctival suffusion . . . . .479t  
   corneal haziness . . . . .480t  
   corneal ulcers . . . . .480t  
   cytoid bodies (cotton wool spots) . . . . .480t  
   dry eyes . . . . .480t  
   endophthalmitis . . . . .480t  
   episcleritis . . . . .480t  
   heliotrope eyelid discoloration . . . . .479t

- hemorrhagic conjunctivitis,  
unilateral/bilateral ..... 479t
- iritis ..... 479t
- keratitis ..... 480t
- periorbital edema,  
unilateral/bilateral ..... 479t
- periphlebitis (candle wax  
drippings) ..... 481t
- Roth spots ..... 481t
- scleral nodules ..... 480t
- subconjunctival hemorrhage,  
hemorrhagic conjunctivitis,  
unilateral/bilateral ..... 479t
- uveitis ..... 480t
- watery eyes ..... 480t
- Eyebrow loss, total/partial, head  
abnormalities (DDx) ..... 478t
- Eyelashes, long, head abnormalities  
(DDx) ..... 478t
- Eyelid edema, bilateral, eye  
abnormalities (DDx) ..... 478t
- Eyelid edema, bilateral upper, eye  
abnormalities (DDx) ..... 478t
- F**-acial cellulitis ..... 29
- Facial erythema, head abnormalities  
(DDx) ..... 478t
- Facial red spots, head abnormalities  
(DDx) ..... 477t
- Facial swelling (edema), head  
abnormalities (DDx) ..... 477t
- Famciclovir ..... 613
- Famvir ..... 613
- Fasciola gigantica ..... 273
- Fasciola hepatica ..... 273
- Fascioliasis ..... 273
- Fasciolopsiasis ..... 282
- Fasciolopsis buski ..... 282
- Fascitis ..... 136
- Fat emboli, bilateral infiltrates  
without effusion ..... 454
- Febrile neutropenia ..... 158
- Fecal gram stain, clinical use ..... 249t
- Feet, desquamation of, extremity  
abnormalities (DDx) ..... 493t
- Ferritin levels, elevated  
(acute/chronic), serum test  
abnormalities (DDx) ..... 499t
- Fever and chills (DDx) ..... 475t
- Fever and night sweats (DDx) ..... 475t
- Fever curves (DDx) ..... 475t
- camelback fever pattern ..... 475t
- double quotidian fevers ..... 475t
- morning (am) temperature  
spikes ..... 475t
- relative bradycardia ..... 475t
- relative tachycardia ..... 475t
- single fever spikes ..... 475t
- Filariasis  
brugia malayi ..... 254
- Wuchereria bancrofti ..... 254
- Fingertip tenderness, extremity  
abnormalities (DDx) ..... 490t
- Fish tapeworm ..... 281
- 5-FC (Flucytosine) ..... 615
- Flaccid paralysis (acute),  
neurological abnormalities  
(DDx) ..... 494t
- Flagyl ..... 645
- Flatworms, intestinal ..... 282
- Fluconazole ..... 614
- pediatric dose ..... 418
- Fluconazole-resistant Candida  
catheter-associated candiduria ..... 105
- fungal cystitis ..... 103
- thrush ..... 43
- Flucytosine ..... 615
- Flukes  
intestinal ..... 282
- liver ..... 273
- lung ..... 262, 264, 266
- Flumadine ..... 685
- Fluoroquinolones  
susceptibility profiles ..... 198t
- Flying insects, travel precautions ..... 372t
- Flying squirrel typhus ..... 294
- Fonsecaea pedrosoi/compactum,  
chromomycosis ..... 289

- Food-borne infectious disease  
(fecally-contaminated), travel  
precautions ..... 372t
- Foot edema, dorsum, extremity  
abnormality (DDx) ..... 490t
- Foot lesions, verrucous, extremity  
abnormality (DDx) ..... 491t
- Foot ulcers, extremity abnormality  
(DDx) ..... 491t
- Fortaz ..... 568
- Fosamprenavir (FPV) ..... 616
- Foscarnet ..... 618
- Foscavir ..... 618
- Fosfomicin ..... 619
- pearls & pitfalls ..... 519
- susceptibility profiles ..... 198t
- Fournier's gangrene ..... 134
- Fourth cephalosporin  
pearls & pitfalls ..... 511
- FPV (fosamprenavir) ..... 616
- Francisella tularensis (tularemia) ..... 223
- community-acquired pneumonia ..... 51
- post-exposure prophylaxis ..... 363t
- tularemia pneumonia ..... 173
- zoonotic atypical pneumonia ..... 60
- Frenal ulcer, mouth abnormalities  
(DDx) ..... 482t
- Freshwater wounds ..... 146
- Frontal sinus, acute bacterial  
sinusitis ..... 30
- FTC (emtricitabine) ..... 603
- FTC/RPV/TDF (Complera) ..... 605
- FTC/TDF (Truvada) ..... 604
- Fulvicin ..... 623
- Fungal (candida) esophagitis ..... 84
- Fungal (cryptococcal) meningitis ..... 24
- Fungal endophthalmitis ..... 36
- Fungal infections  
in neutropenic patients, chronic  
medical prophylaxis ..... 365t
- of skin/nails ..... 147
- Fungizone ..... 537
- Furunculosis ..... 140
- Fusarium, cutaneous ..... 289
- Fusarium solani, cutaneous  
fusarium ..... 289, 291
- Fusobacterium necrophorum,  
Lemierre's syndrome ..... 46
- Fusobacterium nucleatum ..... 240t
- complicated skin/skin structure  
tissue infections ..... 135
- Vincent's angina ..... 45
- Fuzeon (enfuvirtide) ..... 606
- G**-allbladder wall abscess/  
perforation ..... 93
- Ganciclovir ..... 620
- Gangrene, digital ..... 490t
- Garamycin ..... 621
- Gardnerella vaginalis ..... 115
- Gas gangrene ..... 134, 135
- Gastric perforation ..... 85
- Gastroenteritis, pediatric ..... 403
- Gastrointestinal tract infections  
empiric therapy ..... 84
- opportunistic ..... 332
- General appearance  
abnormalities (DDx)  
bullae ..... 476t
- erythroderma ..... 476t
- generalized edema/  
nephrotic syndrome ..... 476t
- hyperpigmentation ..... 476t
- hypotension/shock ..... 476t
- jaundice ..... 477t
- lizard/elephant skin ..... 477t
- rash (below waist) ..... 477t
- soft tissue crepitation ..... 476t
- urticaria ..... 477t
- vitiligo ..... 477t
- Genital herpes ..... 116
- recurrent  
less than 6 episodes/year,  
chronic medical  
prophylaxis ..... 364t
- more than 6 episodes/year,  
chronic medical prophylaxis ..... 365t
- Genital ulcers ..... 117, 489t
- Genital vesicles ..... 116
- Genital warts ..... 119
- Genitourinary abnormalities (DDx)  
epididymoorchitis ..... 489t

- groin mass ..... 489t
- perirectal fistula ..... 489t
- perirectal ulcer ..... 489t
- prostate enlargement/  
tenderness ..... 489t
- scrotal enlargement ..... 489t
- Genitourinary tract infections,  
empiric therapy ..... 103
- Gentamicin ..... 621
- pearls & pitfalls ..... 513
- susceptibility profiles ..... 198t
- pediatric dose ..... 419
- GGT (GGTP), elevation, liver test  
abnormalities (DDx) ..... 500t
- Giardia lamblia ..... 277
- diarrhea ..... 88
- travel prophylaxis ..... 371t
- Giardiasis ..... 276
- Giemsa stain, technique for ..... 246t
- Gingivitis, mouth abnormalities  
(DDx) ..... 482t
- Glucose
- decreased, pleural fluid  
abnormalities (DDx) ..... 504t
- pleural effusion profiles in  
various diseases (DDx) ..... 504
- Gnathostoma spinigerum
- cerebral ..... 261
- cutaneous gnathostomiasis ..... 288
- Gnathostomiasis, cutaneous ..... 288, 289
- Gonorrhea (GC), post-exposure  
prophylaxis ..... 360t
- Goodpasture's syndrome
- bilateral infiltrates with effusion ..... 461
- bilateral infiltrates without  
effusion ..... 453
- Gram-negative bacilli (GNB)
- aerobic ..... 218t
- anaerobic ..... 240t
- capnophilic ..... 235t
- chronic medical prophylaxis ..... 365t
- Gram-negative bacteria, septic shock ..... 155
- Gram-negative cocci ..... 213t
- aerobic, pairs ..... 214t
- Gram-positive bacilli
- aerobic ..... 215t
- anaerobic ..... 236t
- Gram-positive bacteria, septic shock ..... 155
- Gram-positive cocci (GPCs)
- aerobic ..... 210
- chains ..... 210t
- clusters ..... 208t
- pairs ..... 213t
- Gram stains
- CSF ..... 246t
- sputum ..... 248t
- technique for ..... 246t
- urine ..... 248t
- Granuloma inguinale  
(Donovanosis) ..... 118
- Granulomatous amebic  
encephalitis ..... 261, 263
- Granulomatous amebic  
meningoencephalitis ..... 23
- Granulomatous hepatitis (BCG) ..... 97
- Gray hair (premature), head  
abnormalities (DDx) ..... 477t
- Grifulvin ..... 623
- Grisactin ..... 623
- Griseofulvin ..... 623
- Gris-PEG ..... 623
- Groin mass, genitourinary  
abnormalities (DDx) ..... 489t
- Group A streptococci ..... 211
- animal bite wounds ..... 144
- burn wounds ..... 145
- cellulitis ..... 29
- complicated skin/skin structure  
tissue infections ..... 134
- decubitus ulcers ..... 137
- diabetic foot ulcers ..... 138
- human bite wounds ..... 144
- ischemic foot ulcers ..... 139
- Ludwig's angina ..... 44
- osteomyelitis ..... 130
- pharyngitis ..... 40
- septic arthritis ..... 121
- unilateral focal segmental/lobar  
infiltrate with effusion ..... 433

- Group B streptococci .....211  
 bacterial cystitis .....103  
 cellulitis .....133  
 diabetic foot ulcers .....138  
 endocarditis .....70  
 osteomyelitis .....130  
 septic arthritis .....121  
 ureprosepsis .....111
- Group C streptococci .....213  
 endocarditis .....70  
 septic arthritis .....121
- Group F streptococci .....212
- Group G streptococci .....212  
 septic arthritis .....121
- Guillain-Barré syndrome,  
 neurological abnormalities  
 (DDx) .....494t
- Guinea worm .....288
- H**-aemophilus. (see Hemophilus)
- Hafnia alvei .....224
- Hairy leukoplakia .....43
- "Halo sign," Aspergillus  
 pneumonia .....67
- HA-MRSA. *see* Hospital-acquired MRSA
- Hand edema, dorsum, extremity  
 abnormality (DDx) .....490t
- Hand lesions, verrucous .....490t
- Hands, desquamation of,  
 extremity abnormalities (DDx) .....493t
- Hantavirus (HPS), bilateral infiltrates  
 with effusion .....459
- Harvoni .....98t, 346t
- Head abnormalities (DDx)  
 facial erythema .....478t  
 facial red spots .....477t  
 facial swelling (edema) .....477t  
 generalized alopecia .....478t  
 lacrimal gland enlargement  
 (unilateral/bilateral) .....478t  
 localized alopecia .....478t  
 long eyelashes .....478t  
 parotid enlargement .....478t  
 prematurely gray hair .....477t  
 scalp nodules .....478t  
 severe seborrheic  
 dermatitis .....477t  
 temporal muscle wasting .....477t  
 total/partial eyebrow loss .....478t
- Head lice .....150
- Heart  
 Chagas' disease .....273  
 parasites, fungi, unusual  
 organisms in .....273
- Heart abnormalities (DDx)  
 heart block, acute/chronic .....485t  
 heart murmur .....485t  
 pericardial effusion .....485t  
 tachycardia .....485t
- Heart murmur, heart  
 abnormalities (DDx) .....485t
- Heart transplants, transplant  
 prophylaxis (BMT/SOT) .....368t
- HEENT infections,  
 empiric therapy .....29
- Helicobacter pylori .....224
- peptic ulcer disease .....85
- Heliotrope eyelid discoloration, eye  
 abnormalities (DDx) .....479t
- Hematuria, gross/microscopic,  
 urinary abnormalities (DDx) .....502t
- Hemiplegia/hemiparesis, neurological  
 abnormalities (DDx) .....495t
- Hemophagocytosis  
 (erythrophagocytosis), RBC  
 abnormalities (DDx) .....497t
- Hemophilus aphrophilus .....225t
- Hemophilus ducreyi  
 chancroid .....117  
 post-exposure prophylaxis .....360t
- Hemophilus influenzae .....225t  
 acute bacterial exacerbation of  
 chronic bronchitis .....48  
 acute bacterial meningitis .....18  
 acute otitis media .....38  
 adult immunization .....378t  
 brain abscess .....27  
 cellulitis .....29  
 chronic medical prophylaxis .....364t  
 community-acquired  
 pneumonia .....50

- conjunctivitis ..... 32
- endophthalmitis ..... 35
- epiglottitis ..... 47
- HIV ..... 324
- mastoiditis ..... 39
- post-exposure prophylaxis ..... 358t
- preferred prophylaxis ..... 323t
- sepsis ..... 152
- unilateral focal segmental/lobar  
infiltrate with effusion ..... 431
- unilateral focal segmental/lobar  
infiltrate without effusion ..... 426
- Hemophilus parainfluenzae* ..... 225t
- Hemophilus paraphrophilus* ..... 225t
- Hemophilus sp.* ..... 72
- Hemorrhagic conjunctivitis,  
unilateral/bilateral, eye  
abnormalities (DDx) ..... 479t
- Hemorrhagic/necrotizing  
pancreatitis ..... 94
- Hepatic amebiasis ..... 274
- Hepatic echinococcosis  
(hydatid cyst disease) ..... 274
- Hepatic insufficiency, dosing  
strategies in ..... 6t
- Hepatic surgery, surgical  
prophylaxis ..... 356t
- Hepatitis  
acute viral ..... 97t
- chronic viral ..... 97t, 98t
- CMV ..... 99
- organ transplants ..... 166
- EBV ..... 99
- granulomatous (BCG) ..... 96
- hepatitis A (HAV)  
adult immunization ..... 378t
- prophylaxis in HIV ..... 319
- post-exposure prophylaxis ..... 361t
- prophylaxis of opportunistic  
infections ..... 366t
- travel prophylaxis ..... 373t
- hepatitis B (HBV)  
adult immunization ..... 378t
- HIV ..... 344
- HIV, prophylaxis ..... 319
- HIV coinfection ..... 99, 344
- post-exposure prophylaxis ..... 361t
- prophylaxis of opportunistic  
infections ..... 366t
- hepatitis C (HCV) ..... 98
- HIV ..... 346
- post-exposure prophylaxis ..... 361t
- transplant infections ..... 166
- viral ..... 97t, 98t
- Hepatobiliary route of elimination ..... 5t
- Hepatomegaly  
abdominal abnormalities (DDx) ..... 486t
- Hepatosplenic candidiasis ..... 96
- Hepatosplenomegaly, abdominal  
abnormalities (DDx) ..... 487t
- Hepsera ..... 530
- Herpangina ..... 44, 45
- Herpes encephalitis ..... 26
- Herpes labialis ..... 44, 45
- mouth abnormalities (DDx) ..... 482t
- Herpes simplex (HSV) ..... 142
- genital, HIV ..... 340
- skin vesicles/bullae ..... 288
- Herpes simplex virus-1 (HSV-1)  
bilateral infiltrates without  
effusion ..... 449
- encephalitis ..... 25
- esophagitis ..... 84
- herpes gingivostomatitis ..... 44, 45
- herpes labialis ..... 44, 45
- herpes whitlow ..... 142, 143
- keratitis ..... 31, 32
- meningitis ..... 21
- pneumonia ..... 69
- Herpes simplex virus-2 (HSV-2)  
chronic medical prophylaxis ..... 364t
- encephalitis ..... 25
- meningitis ..... 21
- Herpes whitlow ..... 142, 143
- Herpes zoster (VZV) ..... 142, 143
- adult immunization ..... 379t
- encephalitis ..... 25
- pediatric ..... 412
- skin vesicles/bullae ..... 288
- Heterophyes heterophyes ..... 282



Heterophyiasis .....	282	overview .....	302
HHV-6		post-exposure prophylaxis (PEP) .....	366t
encephalitis .....	25	Pre-exposure prophylaxis (PrEP) .....	365
meningitis .....	21	prophylaxis of opportunistic	
pharyngitis .....	40	infections .....	366t
transplant infections .....	162	respiratory tract opportunistic	
Hickman/Broviac infection .....	79, 82	infections .....	323
High level (absolute) resistance .....	2	stages .....	303
Hiprex .....	642	thrush .....	43
Hip (total) replacement surgery,		HIV coinfections (HBV/HCV) .....	344
surgical prophylaxis .....	357t	Hoarseness, mouth abnormalities	
Histiocytosis X, cavitory infiltrates		(DDx) .....	483t
(thin-walled) .....	473	Home IV therapy, less need for .....	15
Histoplasma capsulatum .....	244t	Hookworm .....	278, 279
cutaneous histoplasmosis .....	284, 285	Hospital-acquired MRSA	
pulmonary histoplasmosis .....	267, 269	(HA-MRSA) .....	14, 14t
Histoplasmosis		Hospital-acquired pneumonia (HAP) .....	69
cavitory infiltrates (thick-walled) .....	465	Howell-Jolly bodies, RBC	
meningitis .....	22	abnormalities (DDx) .....	497t
Histoplasmosis, disseminated,		Human bite .....	144
HIV .....	341	pediatric .....	411
HIVID (zalcitabine) .....	717	Human granulocytic	
HIV .....	302	anaplasmosis (HGA) .....	258
acute infection .....	304	Human monocytic ehrlichiosis	
antiretroviral .....	312	(HME) .....	258
failure .....	316, 317t	Human papilloma virus (HPV)	
indications .....	312t	adult immunization .....	380t
initial therapy .....	315t	genital/perianal warts .....	118
therapy .....	313t	Hydradenitis suppurativa .....	141, 142
approach to testing .....	305	Hymenolepis nana, dwarf	
ELISA .....	305	tapeworm .....	281
HIV RNA viral load assays .....	307	Hyperinfection syndrome .....	165
Western blot .....	305	Hyperpigmentation, general	
ART therapy, 313t		appearance abnormalities (DDx) .....	476t
CNS opportunistic infection .....	327	Hyperpigmented shins, extremity	
coinfections .....	344	abnormalities (DDx) .....	492t
diagnosis, evaluation, treatment .....	302	Hyposplenism .....	157
HBV coinfection .....	97	Hypotension, general appearance	
HCV coinfection .....	98	abnormalities (DDx) .....	476t
initial assessment .....	308	I	
baseline laboratory testing .....	309t	-diopathic pulmonary hemosiderosis	
CD <sub>4</sub> count .....	309, 312t	(IPH), bilateral infiltrates without	
clinical evaluation .....	308	effusion .....	456
opportunistic infections		IDV (indinavir) .....	625
other .....	338		
prophylaxis .....	319t		
treatment .....	323		

- Imipenem, pediatric dose . . . . .419  
 Imipenem/cilastatin . . . . .623  
   pearls & pitfalls . . . . .511  
   susceptibility profiles . . . . .198t  
 Immunizations. (see also Adult immunizations)  
   purpose of . . . . .378  
   travel vaccines in HIV . . . . .323  
 Immunoblasts, WBC  
   abnormalities (DDx) . . . . .496t  
 Impetigo . . . . .132  
   pediatric . . . . .411  
 Indian (Bihar state) visceral leishmaniasis (Kala-azar) . . . . .275  
 Indinavir (IDV) . . . . .625  
 Infective endocarditis (IE)  
   prophylaxis, indications for . . . . .369t  
 Influenza  
   adult immunization . . . . .379t  
   avian . . . . .62  
   bilateral infiltrates without effusion . . . . .448  
   HIV, prophylaxis . . . . .319, 322  
   pneumonia . . . . .66  
     seasonal (types A/B) . . . . .62  
   post-exposure prophylaxis . . . . .358t, 359t  
   prophylaxis of opportunistic infections . . . . .366t  
 Influenza A  
   clinical presentations in adults . . . . .63  
   community-acquired pneumonia . . . . .51, 52  
   with *S. aureus* (MSSA/MRSA) CAP . . . . .65  
   sepsis . . . . .152  
 Influenza B, community-acquired pneumonia . . . . .52  
 Infraclavicular adenopathy, lymph node abnormalities (DDx) . . . . .488t  
 Inguinal adenopathy, unilateral/bilateral, lymph node abnormalities (DDx) . . . . .489t  
 INH . . . . .628  
 Intelece (etravirine) . . . . .612  
 Interferon, HIV/HCV co-infection, adverse effects . . . . .349t  
 Intermediate level (relative) resistance . . . . .2  
 Intestinal  
   cestodes . . . . .281  
   nematodes . . . . .278  
   protozoa . . . . .276  
   trematodes . . . . .282  
 Intestines, parasites, fungi, unusual organisms in . . . . .276  
 Intraabdominal abscess . . . . .100  
 Intracellular inclusion bodies in blood . . . . .258  
 Intrarenal abscess . . . . .109  
 Invanz . . . . .608  
 Invirase (saquinavir) . . . . .687  
*In vitro* data, microbiology  
   susceptibility testing limitations . . . . .7  
*In vitro* susceptibility testing  
   antibiotic-organism combinations, not predictive of *in vivo* effectiveness . . . . .7t  
   factors dependent on . . . . .7  
*In vitro* vs. *In vivo* susceptibility . . . . .7  
   lodoquinol, pediatric dose . . . . .419  
   iritis, eye abnormalities (DDx) . . . . .479t  
   isavuconazole . . . . .626  
   ischemic foot ulcers . . . . .139  
   ischiorectal abscess . . . . .151  
   Isentress (raltegravir) . . . . .678  
   Isoniazid . . . . .628  
     pediatric dose . . . . .419  
   Ispora, chronic watery diarrhea . . . . .90  
   Ispora belli . . . . .276  
     HIV . . . . .335  
   Isosporiasis . . . . .277  
   Israeli spotted fever . . . . .294  
   Itraconazole . . . . .629  
 IV drug abusers, acute bacterial endocarditis . . . . .75  
 IV line infections . . . . .81  
   pediatric . . . . .402  
 IV to PO switch therapy . . . . .10  
**J**-apanese encephalitis (JE) . . . . .25  
   travel prophylaxis . . . . .374t  
 Jaundice, general appearance  
   abnormalities (DDx) . . . . .477t

- Jock itch .....149
- Joint infections, empiric therapy.....121
- Joint prosthesis, infected.....128
- Jugular vein tenderness, neck  
abnormalities (DDx) .....483t
- K**-ala-azar.....275
- Kaletra [Lopinavir (LPV) +  
Ritonavir (RPV)].....639
- Kaposi's sarcoma, HIV infection.....302
- Keflex.....575
- Kefurox.....574
- Kefzol.....557
- Keratitis.....31
- amebic.....32
- bacterial.....31
- eye abnormalities (DDx).....480t
- HSV.....340
- viral, HSV-1.....31
- Ketek.....694
- Ketoconazole.....631
- Kingella kingae.....225t
- endocarditis.....72
- Klebsiella.....225t
- cholangitis.....93
- cholecystitis.....92
- gallbladder wall abscess.....93
- sepsis.....152
- surgical prophylaxis.....355t
- urosepsis.....112
- Klebsiella (Calymmatobacterium)  
granulomatosis, granuloma  
    inguinale.....118
- Klebsiella oxytoca.....225
- Klebsiella ozaenae,  
rhinoscleromatis.....226
- Klebsiella pneumoniae.....225
- acute bacterial meningitis.....19
- alcoholics.....66
- cavitary infiltrates (thick-walled).....463
- cellulitis.....133
- community-acquired pneumonia.....50
- decubitus ulcers.....137
- diabetic foot ulcers.....138
- lung abscess/empyema.....67
- osteomyelitis.....130
- pneumonia.....67
- sepsis.....152
- skin abscesses/infected cysts.....141
- unilateral focal segmental/lobar  
        infiltrate with effusion.....431
- unilateral focal segmental/lobar  
        infiltrate without effusion.....427
- Knee (total) replacement surgery,  
surgical prophylaxis.....357t
- L**-acrimial gland enlargement, head  
abnormalities (DDx).....478t
- Lactate dehydrogenase (LDH),  
elevated, serum test  
abnormalities (DDx).....499t
- Lactate.....500t
- serum.....500t
- CSF.....506t
- Lactobacillus sp.....239
- Lamisil.....697
- Lamivudine.....632
- Lamivudine + zidovudine  
(Combivir).....633
- Lassa fever.....178
- post-exposure prophylaxis.....363t
- LCM, meningitis.....21
- Ledipasvir/Sofosbuvir.....98t, 346t
- Leflunomide.....634
- Left upper abdominal quadrant  
tenderness (splenic tenderness),  
abdominal abnormalities (DDx).....486t
- Leg edema, unilateral/bilateral,  
extremity abnormalities  
(DDx).....491t
- Legionnaire's disease.....56, 59t
- Legionella sp.....226t
- bilateral infiltrates with effusion.....458
- bilateral infiltrates without  
        effusion.....451
- community-acquired pneumonia.....51
- endocarditis.....72
- organ transplants.....167
- transplant infections.....165

- unilateral focal segmental/lobar infiltrate with effusion .....433
- unilateral focal segmental/lobar infiltrate without effusion.....427
- unilateral ill-defined infiltrates with effusion .....444
- unilateral ill-defined infiltrates without effusion.....437
- Leg lesions, nodular, extremity abnormalities (DDx) .....492t
- Leg lesions, verrucous, extremity abnormality (DDx) .....491t
- Leg nodules, painful, extremity abnormalities (DDx) .....492t
- Leg ulcers, extremity abnormality (DDx).....491t
- Leishmania braziliensis .....283
- Leishmania donovani, visceral leishmaniasis .....275
- Leishmania major .....283
- Leishmania mexicana .....283
- Leishmaniasis
  - cutaneous (Old World/ New World) .....283
  - visceral (Kala-azar) .....275
- Leishmania tropica .....283
- Lemierre's syndrome .....46, 47
- Lens implant, infected, endophthalmitis .....35
- Leprosy .....283
- Leptospira interrogans .....226t
- Leptospirosis .....97
- Leptospirosis, meningitis .....22
- Leuconostoc .....213t
- Leukocyte esterase, urinary abnormalities (DDx) .....502t
- Leukocytosis, WBC abnormalities (DDx) .....495t
- Leukopenia, WBC abnormalities (DDx) .....495t
- Leukoplakia, mouth abnormalities (DDx).....482t
- Leukostasis, bilateral infiltrates without effusion.....456
- Levaquin .....635
- Levofloxacin .....635
  - pearls & pitfalls .....515
  - susceptibility profiles .....198t
- Lexiva (fosamprenavir) .....616
- Lice infestations, body, head, pubic .....150
- Linezolid .....637
  - pearls & pitfalls .....517
  - susceptibility profiles .....198t
  - pediatric dose .....419
- Liposomal/lipid complex, pediatric dose .....419
- Listeria, transplant infections .....164
- Listeria encephalitis .....26
- Listeria monocytogenes.....217t
  - acute bacterial meningitis .....18
  - encephalitis .....25
- Liver
  - abscess.....95
  - cystic masses .....274
  - hepatomegaly.....275
  - parasites, fungi, unusual organisms in .....273
- Liver flukes .....273
- Liver test abnormalities (DDx)
  - alkaline phosphatase (AP), elevation .....500t
  - GGT (GGTP), elevation .....500t
  - serum transaminases (SGOT/SGPT), elevation.....500t
  - total bilirubin, elevation.....500t
- Liver transplants, transplant prophylaxis (BMT/SOT) .....368t
- Lizard skin, general appearance abnormalities (DDx) .....477t
- Loa Loa (Loiasis) .....256
- Loeffler's syndrome, bilateral infiltrates without effusion.....454
- Lopinavir + ritonavir (Kaletra) .....639
- Lorabid .....641
- Loracarbef .....641
  - pearls & pitfalls .....511
  - susceptibility profiles .....192t
- Louse-borne relapsing fever (LBRF) .....257
- LPV/RTV (Kaletra).....639

- Ludwig's angina ..... 45
- Lung contusion  
 bilateral infiltrates without effusion .... 454  
 unilateral ill-defined infiltrates  
 without effusion ..... 440
- Lung fluke. *see* Paragonimus  
*westermani*
- Lungs, parasites, fungi, unusual  
 organisms in ..... 265
- Lyme arthritis ..... 127
- Lyme disease, post-exposure  
 prophylaxis ..... 361t
- Lymphadenopathy, generalized,  
 lymph node abnormalities  
 (DDx) ..... 489t
- Lymphangitic metastases  
 bilateral infiltrates with effusion ..... 462  
 unilateral ill-defined infiltrates  
 with effusion ..... 444  
 unilateral ill-defined infiltrates  
 without effusion ..... 440
- Lymphangitis ..... 132  
 extremity abnormalities (DDx) ..... 490t
- Lymph node abnormalities (DDx)  
 anterior cervical adenopathy ..... 488t  
 axillary adenopathy ..... 488t  
 epitrochlear adenopathy ..... 488t  
 generalized lymphadenopathy ..... 489t  
 infraclavicular adenopathy ..... 488t  
 inguinal adenopathy,  
 unilateral/bilateral ..... 489t  
 occipital adenopathy ..... 488t  
 periumbilical nodule ..... 488t  
 posterior cervical adenopathy,  
 unilateral/bilateral ..... 488t  
 preauricular adenopathy ..... 488t  
 supraclavicular adenopathy ..... 488t  
 ulcer-node syndromes ..... 488t
- Lymphocytes, pleural fluid  
 abnormalities (DDx) ..... 504t
- Lymphocytosis, WBC  
 abnormalities (DDx) ..... 495t
- Lymphogranuloma venereum (LGV) ..... 118
- Lymphoma  
 bilateral infiltrates with effusion ..... 462  
 bilateral infiltrates  
 without effusion ..... 455  
 cavitary infiltrates (thick-walled) ..... 468  
 unilateral focal segmental/lobar  
 infiltrate with effusion ..... 435  
 unilateral focal segmental/lobar  
 infiltrate without effusion ..... 429  
 unilateral ill-defined infiltrates  
 with effusion ..... 445  
 unilateral ill-defined infiltrates  
 without effusion ..... 441
- M**-acrobid ..... 657
- Macrodantin ..... 657
- Macrofides  
 pearls & pitfalls ..... 518  
 susceptibility profiles ..... 186t
- Malaria (worldwide) ..... 259  
 prophylaxis ..... 375t  
 severe ..... 258  
 uncomplicated ..... 259
- Malarone ..... 548
- Malassezia furfur ..... 244t  
 eosinophilic folliculitis ..... 297  
 tinea versicolor/pityriasis ..... 148
- Mandelamine ..... 645
- Mandol ..... 557
- Maraviroc (MVC) ..... 642
- Marburg virus disease ..... 178
- Mass lesions, organ  
 transplants ..... 167
- Mastitis ..... 133
- Mastoiditis ..... 39  
 pediatric ..... 394
- Maxillary sinus, acute bacterial  
 sinusitis ..... 30
- Maxipime ..... 560
- MDR  
 acute bacterial meningitis ..... 19  
 community-acquired pneumonia ..... 54  
 pneumonia ..... 69
- MDR acinetobacter, catheter-  
 associated bacteriuria ..... 104
- MDR GNBS, bacterial cystitis ..... 103

- MDR *Klebsiella*, catheter-associated  
bacteriuria.....104
- MDROs. (see Multi-drug resistant  
organisms)
- MDR *P. aeruginosa*, catheter-  
associated bacteriuria .....104
- MDRSP, acute otitis media .....38
- MDR-TB .....53
- Measles  
adult immunization.....380t  
bilateral infiltrates with effusion.....459
- Measles, mumps, rubella,  
prophylaxis.....322
- Mediastinitis .....49
- Mediterranean spotted fever .....294
- Mefoxin .....565
- Melioidosis, cavitory infiltrates  
(thick-walled).....466
- Membranous pharyngitis .....41
- Meningitis  
acute bacterial  
CNS shunt infections .....21  
elderly .....20  
empiric therapy .....18  
malignancy .....20  
normal host .....20  
pediatric .....387  
acute non-bacterial.....21  
amebic  
granulomatous.....23  
primary .....23  
aseptic/viral .....23  
chronic.....21  
cryptococcal .....24  
HIV.....329  
EVD associated ABM.....506t  
organ transplants.....167  
post-exposure prophylaxis.....358t  
transplant infections.....164  
tuberculosis .....24  
viral .....23
- Meningococcal meningitis, travel  
prophylaxis.....373t
- Meningococcus (invasive disease),  
adult immunization .....380t
- Mental confusion, neurological  
abnormalities (DDx) .....493t
- Mepron .....548
- Meronem .....643
- Meropenem .....645  
pearls & pitfalls .....511  
susceptibility profiles .....198t  
pediatric dose .....419
- MERS (Middle East Respiratory  
Syndrome) .....62
- Metagonimus yokogawai,  
heterophyiasis.....282
- Metastatic carcinoma  
bilateral infiltrates without  
effusion .....455  
cavitory infiltrates (thick-walled).....469  
unilateral ill-defined infiltrates  
with effusion .....447
- Methenamine hippurate .....645
- Methenamine mandelate .....645
- Metronidazole .....645  
pearls & pitfalls .....518  
susceptibility profiles .....186t  
pediatric dose .....419
- Mezlin .....647
- Mezlocillin .....647
- Micafungin.....648
- Microbiology susceptibility  
testing .....5  
limitations of .....6  
overview .....5
- Microfilaria in blood.....254
- Microsporidia enteritis, HIV.....335
- Microsporium canis*, tinea corporis.....147
- Migratory rashes, extremity  
abnormalities (DDx) .....492t
- Miliary (disseminated) BCG .....158
- Miliary TB .....158
- Minocin .....649
- Minocycline.....649  
pearls & pitfalls .....512  
susceptibility profiles .....186t

- Mixed aerobic/anaerobic deep soft tissue infection ..... 135
- Mobiluncus ..... 115
- Mode of elimination, explanation ..... 522
- Monobactams, pearls & pitfalls ..... 511
- Monoclonal gammopathy, serum test abnormalities (DDx) ..... 499t
- Monocytosis, WBC abnormalities (DDx) ..... 496t
- Mononeuritis multiplex, neurological abnormalities (DDx) ..... 494t
- Monospot test (false+) (DDx) ..... 499t
- Monotherapy vs. combination therapy ..... 9
- Monurol ..... 619
- Moraxella catarrhalis ..... 227
- acute bacterial exacerbation of chronic bronchitis ..... 48
- acute otitis media ..... 38
- community-acquired pneumonia ..... 49
- conjunctivitis ..... 32
- keratitis ..... 31
- unilateral focal segmental/lobar infiltrate without effusion ..... 427
- Morganella morganii ..... 227
- Morning temperature spikes, fever curves (DDx) ..... 475t
- Mouth abnormalities (DDx)
- angular cheilitis ..... 482t
- crimson crescents ..... 483t
- frenal ulcer ..... 482t
- gingivitis ..... 482t
- herpes labialis ..... 482t
- hoarseness ..... 483t
- leukoplakia ..... 482t
- oral ulcers, solitary/multiple ..... 482t
- palatal perforation ..... 483t
- palatal petechiae ..... 482t
- palatal vesicles ..... 483t
- tongue tenderness ..... 482t
- tongue ulcers ..... 482t
- tonsillar membranes ..... 483t
- tonsillar ulcers ..... 483t
- trismus ..... 482t
- uvular edema ..... 483t
- Mouth ulcers/vesicles ..... 44
- Moxifloxacin ..... 650
- pearls & pitfalls ..... 515
- susceptibility profiles ..... 198t
- MRSA. *see* Methicillin-resistant staphylococcus aureus (MRSA)
- MSSA, brain abscess ..... 27
- Mucocutaneous (local/non-disseminated) candidiasis ..... 147
- Mucor
- cutaneous zygomycosis ..... 290
- pulmonary zygomycosis ..... 269
- Mucormycosis ..... 269
- Mumps
- adult immunization ..... 380t
- meningitis ..... 21
- Murine (flea-borne) typhus ..... 294
- Muscle rigidity, general, neurological abnormalities (DDx) ..... 493t
- Muscles
- parasites, fungi, unusual organisms in ..... 283
- tenderness in, extremity abnormality (DDx) ..... 491t
- MVC (maraviroc) ..... 642
- Myambutol ..... 611
- Mycamine ..... 648
- Mycobacterium avium-intracellulare (MAI) community-acquired pneumonia ..... 53
- cutaneous ..... 285
- HIV/AIDS ..... 342
- prophylaxis ..... 319
- pneumonia ..... 66
- prophylaxis of opportunistic infections ..... 366t
- Mycobacterium chelonae, cutaneous ..... 287
- Mycobacterium fortuitum, cutaneous ..... 287
- Mycobacterium genitalium, post-exposure prophylaxis ..... 360t
- Mycobacterium haemophilum ..... 287
- cutaneous ..... 287
- Mycobacterium hominis ..... 115

- Mycobacterium kansasii pneumonia . . . . . 53  
 Mycobacterium leprae, leprosy . . . . . 283  
 Mycobacterium marinum, swimming  
   pool granuloma . . . . . 287  
 Mycobacterium scrofulaceum,  
   scrofula . . . . . 286  
 Mycobacterium tuberculosis (TB)  
   atypical, cavitary infiltrates  
     (thin-walled) . . . . . 470  
   bilateral infiltrates without  
     effusion . . . . . 450  
   cavitary infiltrates (thick-walled) . . . . . 465  
   chronic pyelonephritis . . . . . 108  
   chronic tuberculosis peritonitis . . . . . 102  
   community-acquired pneumonia . . . . . 53  
   cutaneous mycobacteria . . . . . 285  
   endophthalmitis . . . . . 35  
   epididymitis . . . . . 106  
   HIV  
     extrapulmonary . . . . . 339  
     prophylaxis . . . . . 319  
   MDR-TB . . . . . 53  
   meningitis . . . . . 22  
   miliary tuberculosis . . . . . 155  
   organ transplants . . . . . 167  
   pediatric . . . . . 400  
   pericarditis . . . . . 78  
   pneumonia . . . . . 66  
     HIV . . . . . 321  
   post-exposure prophylaxis . . . . . 359t  
   prophylaxis of opportunistic  
     infections . . . . . 366t  
   renal TB . . . . . 108  
   transplant infections . . . . . 165  
   tuberculosis arthritis . . . . . 123  
   tuberculosis osteomyelitis . . . . . 131  
   unilateral focal segmental/lobar  
     infiltrate with effusion . . . . . 432  
   unilateral ill-defined infiltrates  
     with effusion . . . . . 443  
 Mycobacterium ulcerans, Buruli ulcer . . . . . 287  
 Mycobutin . . . . . 681  
 Mycoplasma encephalitis . . . . . 26  
 Mycoplasma genitalium, urethritis/  
   cervicitis . . . . . 115  
 Mycoplasma pharyngitis . . . . . 42  
 Mycoplasma pneumoniae . . . . . 57  
   community-acquired pneumonia . . . . . 49  
   encephalitis . . . . . 25  
   pharyngitis . . . . . 41  
   pneumonia . . . . . 57  
   unilateral ill-defined infiltrates  
     without effusion . . . . . 436  
 Myocarditis . . . . . 78  
 Myoglobinuria (2° to rhabdomyolysis),  
   urinary abnormalities (DDx) . . . . . 503t  
 Myositis . . . . . 298  
**N**-meningitis, conjunctivitis . . . . . 32  
 Naegleria fowleri, meningitis . . . . . 21  
 Nafcillin . . . . . 652  
   susceptibility profiles . . . . . 186t  
   pediatric dose . . . . . 420  
 Nails, fungal infections . . . . . 147  
 Nasal abnormalities (DDx)  
   epistaxis . . . . . 481t  
   nasal septal perforation . . . . . 481t  
   nose tip gangrene . . . . . 481t  
   purple nose tip . . . . . 481t  
 Nasal septal perforation, nasal  
   abnormalities (DDx) . . . . . 481t  
 National Committee for Clinical  
   Laboratory Standards.  
   see Clinical and Laboratory  
   Standards Institute (CLSI)  
 Naturally/intrinsically resistant  
   pathogens . . . . . 2  
 Nebcin . . . . . 705  
 NebuPent . . . . . 666  
 Necator americanus, hookworm . . . . . 278  
 Neck abnormalities (DDx)  
   bull neck . . . . . 483t  
   enlarged greater auricular  
     nerve . . . . . 483t  
   jugular vein tenderness . . . . . 483t  
   neck sinus tract . . . . . 483t  
   superior vena cava syndrome . . . . . 483t



- Neck infections, deep .....46
- Neck sinus tract, neck abnormalities  
(DDx).....483t
- Necrotizing abscesses .....135
- Necrotizing fasciitis .....134
- Neisseria gonorrhoeae* (GC) .....187
- cervicitis .....114
- conjunctivitis .....32
- endometritis .....113
- PID .....113
- post-exposure prophylaxis .....360t
- salpingitis .....113
- septic abortion .....113
- septic arthritis .....122
- tuboovarian abscess .....113
- urethritis .....114
- Neisseria meningitidis* .....187t, 214t
- acute bacterial meningitis .....18
- chronic medical prophylaxis .....364t
- post-exposure prophylaxis .....358t
- sepsis .....155
- travel prophylaxis .....373t
- Nelfinavir (NFV) .....653
- Nematodes (roundworms),  
  intestinal .....278
- Neonatal Group B streptococcal  
(GBS) infection (primary prevention),  
  chronic medical prophylaxis .....365t
- Nephrotic syndrome, general  
  appearance abnormalities (DDx) .....476t
- Neuroborreliosis, meningitis .....24
- Neurobrucellosis .....22
- Neurocysticercosis .....262
- meningitis .....22
- Neurological abnormalities (DDx)
- anisocoria .....494t
- Bell's palsy .....494t
- cranial nerve abnormalities,  
    unilateral/bilateral .....494t
- flaccid paralysis (acute) .....494t
- general muscle rigidity .....493t
- Guillain-Barré syndrome .....494t
- hemiplegia/hemiparesis .....495t
- mental confusion/encephalopathy  
    (acute) .....493t
- mononeuritis multiplex .....494t
- nuchal rigidity .....493t
- optic nerve atrophy .....494t
- papilledema .....494t
- transient deafness .....493t
- transverse myelitis .....494t
- Neurosyphilis .....119, 120
- Nevirapine (NVP) .....654
- NFV (nelfinavir) .....653
- Nitazoxanide .....656
- Nitrite positive, urinary  
  abnormalities (DDx) .....502t
- Nitrofurantoin .....657
- pearls & pitfalls .....516
- susceptibility profiles .....198t
- Nizoral .....631
- Nocardia asteroides* .....217t
- pulmonary nocardiosis .....271
- Nocardia brasiliensis* .....217t
- Nocardia* sp. .....188t
- cavitary infiltrates (thick-walled) .....466
- cerebral nocardiosis .....261
- cutaneous nocardiosis .....284
- transplant infections .....164
- unilateral ill-defined infiltrates  
    with effusion .....443
- unilateral ill-defined infiltrates  
    without effusion .....438
- Nocardiosis, cutaneous .....284, 286
- Nodular arm/leg lesions, extremity  
  abnormalities (DDx) .....492t
- Nodular lymphangitis, extremity  
  abnormalities (DDx) .....490t
- Nodular/pustular candidiasis .....290
- Nodules, skin .....289
- Non-aeruginosa *Pseudomonas*,  
  urosepsis .....112
- Non-cholera vibrios, travel prophylaxis .....372t
- Non-fermentative GNBs,  
  CAPD-associated peritonitis .....101
- Non-gonococcal urethritis (NGU),  
  post-exposure prophylaxis .....360t
- Norvir (ritonavir) .....686
- Norwalk virus, travel prophylaxis .....372t

- Nose tip gangrene, nasal abnormalities (DDx) ..... 481t
- Nosocomial pneumonia, bilateral infiltrates without effusion ..... 452
- Nosocomial pneumonia (NP), ..... 68
- Noxafil ..... 672
- Nuchal rigidity, neurological abnormalities (DDx) ..... 493t
- Nutritionally-variant streptococci (NVS), endocarditis ..... 71
- Nursing home-acquired pneumonia (NHAP) ..... 68
- NVP (nevirapine) ..... 654
- Nystatin, pediatric dose ..... 420
- O**-ccipital adenopathy, lymph node abnormalities (DDx) ..... 488t
- Ochrobactrum anthropi (CDC group Vd) ..... 227t
- Oflox ..... 658
- Ofloxacin ..... 658
- pearls & pitfalls ..... 515
- susceptibility profiles ..... 198t
- Olysio ..... 98t, 346t
- Omnicef ..... 558
- Onchocerca volvulus, cutaneous onchocerciasis ..... 290, 294
- Onchocerciasis, cutaneous ..... 290
- Onychomycosis ..... 148
- OPAT ..... 11
- Opisthorchiasis ..... 273, 274
- Opisthorchis viverrini ..... 273
- Opportunistic infections. *see* HIV infection
- Optic nerve atrophy, neurological abnormalities (DDx) ..... 494t
- Oral anaerobes ..... 49
- brain abscess ..... 27
- community-acquired pneumonia ..... 53
- deep neck infections ..... 46
- dental infections ..... 46
- gastric perforation ..... 85
- human bite wounds ..... 144, 411
- lung abscess/empyema ..... 67
- mastoiditis ..... 39
- suppurative parotitis ..... 40
- Oral anti-pseudomonal penicillins, pearls & pitfalls ..... 509
- Oral anti-staphylococcal penicillins, pearls & pitfalls ..... 509
- Oral first-generation cephalosporins, pearls & pitfalls ..... 509
- Oral second-generation cephalosporins pearls & pitfalls ..... 510
- Oral third-generation cephalosporins pearls & pitfalls ..... 511
- Oral ulcers, solitary/multiple, mouth abnormalities (DDx) ..... 482t
- Orbactiv ..... 659
- Orbital cellulitis, pediatric ..... 391
- Organ transplants
- bacteremia or candidemia ..... 160
- bone marrow ..... 160
- brain abscess/mass lesion ..... 164
- encephalitis/meningitis ..... 164
- hepatitis, viral ..... 166
- infections ..... 160
- pneumonias ..... 165
- solid organ ..... 161
- Oritavancin ..... 659
- pearls & pitfalls ..... 519
- susceptibility profiles ..... 198t
- Oritavancin (Orbactiv) ..... 659
- Oropharyngeal candidiasis, HIV ..... 336
- Orthopedic prosthetic implant surgery, surgical prophylaxis ..... 357t
- Orthopedic surgery (open fracture), surgical prophylaxis ..... 357t
- Oseltamavir ..... 660
- Osteochondritis, pediatric ..... 409
- Osteomyelitis ..... 129
- acute, pediatric ..... 409
- chronic ..... 130, 139
- Otitis externa ..... 37
- pediatric ..... 393
- Otitis media
- acute ..... 38
- pediatric ..... 393
- Oxacillin ..... 661
- pediatric dose ..... 420

- P**-cepacia, surgical prophylaxis ..... 357t
- P. maltophilia, surgical prophylaxis ..... 357t
- Pacemaker infections ..... 79, 82
- Painless purple palm/sole lesions, extremity abnormalities (DDx) ..... 490t
- Palatal perforation, mouth abnormalities (DDx) ..... 483t
- Palatal petechiae, mouth abnormalities (DDx) ..... 482t
- Palatal vesicles, mouth abnormalities (DDx) ..... 483t
- Palm lesions, painless, extremity abnormalities (DDx) ..... 490t
- Palm rashes, extremity abnormalities (DDx) ..... 492t
- Palpable purpura, extremity abnormalities (DDx) ..... 492t
- Pancreas transplants, transplant prophylaxis (BMT/SOT) ..... 368t
- Pancreatic abscess ..... 95
- Pancreatic pseudocyst, infected ..... 95
- Pancreatitis  
acute ..... 94  
edematous ..... 94  
hemorrhagic/necrotizing ..... 94
- Pancytopenia, platelet abnormalities (DDx) ..... 498t
- Papilledema, neurological abnormalities (DDx) ..... 494t
- Papular axillary lesions, extremity abnormality (DDx) ..... 491t
- Papules, skin ..... 289
- Paracoccidioides brasiliensis, pulmonary paracoccidioidomycosis ..... 267
- Paragonimiasis, cavity infiltrates (thin-walled) ..... 471
- Paragonimus westermani  
cerebral paragonimiasis (lung fluke) ..... 262  
pulmonary paragonimiasis ..... 265
- Parenteral first-generation cephalosporins pearls & pitfalls ..... 509
- Parenteral second-generation cephalosporins pearls & pitfalls ..... 510
- Parenteral third-generation cephalosporins pearls & pitfalls ..... 510
- Parenteral fourth-generation cephalosporins pearls & pitfalls ..... 511
- Paritaprevir/ritonavir/ombitasvir/dasabuvir ..... 98, 346t
- Parotid enlargement, head abnormalities (DDx) ..... 478t
- Parotitis, suppurative ..... 40
- Parvo B19, meningitis ..... 21
- Pasteurella multocida ..... 227t  
animal bite wounds ..... 144  
septic arthritis ..... 123
- PCP  
HIV, prophylaxis ..... 319t  
prophylaxis in opportunistic infections ..... 366t
- Peak serum levels, explanation ..... 521
- Pediatric infectious diseases ..... 386  
bone and joint ..... 407  
CNS ..... 387  
dosing ..... 414  
gastrointestinal ..... 403  
HEENT ..... 391  
lower respiratory ..... 396  
skin and soft tissue ..... 411  
vascular ..... 402
- Pediculus humanus var. capitis, head lice ..... 150
- Pediculus humanus var. corporis, body lice ..... 150
- Pegasis ..... 662
- Peg-Intron ..... 663
- Pegylated interferon  
alfa-2a ..... 662  
alfa-2b ..... 663
- Peliosis hepatitis ..... 291
- Pelvic inflammatory disease (PID) ..... 113
- Pelvic (OB/GYN) surgery, surgical prophylaxis ..... 356t
- Pelvic peritonitis/abscess ..... 100, 101

- Penicillin G .....664  
 pediatric dose .....420
- Penicillin  
 pearls & pitfalls .....508  
 susceptibility profiles .....186t
- Penicillin V .....665  
 pediatric dose .....421
- Penicillium, cutaneous .....289, 292
- Penicillium marneffeii .....245t  
 cutaneous penicillium .....289
- Pentam 300 .....666
- Pentamidine .....666
- Peptic ulcer disease .....85
- Peptococcus .....236t
- Peptostreptococcus .....236t
- Peramivir .....668
- Perianal warts .....118
- Pericardial effusion, heart  
 abnormalities (DDx) .....485t
- Pericarditis .....78
- Periorbital cellulitis .....29  
 pediatric .....391
- Periorbital edema, unilateral/bilateral,  
 eye abnormalities (DDx) .....479t
- Peripheral vascular disease (PVD),  
 non-diabetics with, osteomyelitis .....132
- Perirectal abscess .....151
- Perirectal fistula, genitourinary  
 abnormalities (DDx) .....489t
- Perirectal ulcer, genitourinary  
 abnormalities (DDx) .....489t
- Peritonitis  
 CAPD-associated .....101, 102  
 chronic tuberculosis .....101, 102  
 spontaneous bacterial .....101, 102
- Periumbilical nodule, lymph node  
 abnormalities (DDx) .....488t
- Periumbilical purpura  
 (thumbprint sign), abdominal  
 abnormalities (DDx) .....486t
- Persistent fevers, antibiotics for .....15
- Pertussis .....56  
 adult immunization .....380t  
 community-acquired pneumonia .....56  
 pediatric .....399
- post-exposure prophylaxis .....359t
- travel prophylaxis .....375t
- pH  
 pleural fluid .....503t  
 urinary .....502t
- Pharyngitis .....40  
 bacterial .....40  
 membranous .....40  
 other .....41  
 pediatric .....395  
 viral .....40
- Pneumonia .....134, 136
- Phialophora verrucosa,  
 chromomycosis .....289
- Phthirus pubis, pubic lice .....150, 151
- Pilonidal cysts, infected .....141, 142
- Pinworms .....279, 280
- Piperacillin .....669  
 pearls & pitfalls .....512  
 susceptibility profiles .....186t  
 pediatric dose .....421
- Piperacillin/tazobactam .....669  
 pearls & pitfalls .....512  
 susceptibility profiles .....186t  
 pediatric dose .....421
- Pipracil .....669
- Plague  
 pneumonic .....176  
 bioterrorist .....174  
 post-exposure prophylaxis .....363t
- Plantar hyperkeratoses, extremity  
 abnormalities (DDx) .....493t
- Plasma protein binding,  
 explanation .....522
- Plasmodium falciparum,  
 malaria .....259
- Plasmodium knowlesi, malaria .....259
- Plasmodium malariae, malaria .....259
- Plasmodium ovale, malaria .....259
- Plasmodium vivax, malaria .....259
- Platelet abnormalities (DDx)  
 pancytopenia .....498t  
 thrombocytopenia .....498t  
 thrombocytosis .....498t

- Plesiomonas, travel  
prophylaxis ..... 372t
- Plesiomonas shigelloides ..... 228t
- Pleural effusion profiles in various  
diseases (DDx)  
  eosinophils ..... 504  
  glucose ..... 504  
  RBCs (DDx) ..... 504  
  WBCs (DDx) ..... 504
- Pleural fluid abnormalities (DDx)  
  amylase ..... 504t  
  blood ..... 503t  
  brown ..... 503t  
  extracellular debris ..... 504t  
  glucose, decreased ..... 504t  
  lymphocytes ..... 504t  
  other ..... 505t  
  pH ..... 503t  
  PMNs ..... 504t  
  protein ..... 504t  
  rheumatoid factor (RF) ..... 505t  
  yellow-green ..... 503t  
  yellow-whitish ..... 503t
- Pneumatoceles, cavitory infiltrates  
(thin-walled) ..... 471
- Pneumococcus, adult  
immunization ..... 381t
- Pneumocystis (carinii) jiroveci  
  pneumonia (PCP)  
    bilateral infiltrates without  
      effusion ..... 450  
    community-acquired pneumonia ..... 55  
    HIV ..... 325
- Pneumonia  
  alcoholics ..... 54  
  Aspergillus ..... 55  
  aspiration ..... 53  
    unilateral focal segmental/lobar  
      infiltrate without effusion ..... 430  
    unilateral ill-defined infiltrates  
      without effusion ..... 439  
  atypical pathogens, non-zoonotic ..... 51  
  atypical pathogens, zoonotic ..... 51  
  bacterial pathogens ..... 50  
    bronchiectasis ..... 54  
    chickenpox ..... 52  
    chronic steroid use ..... 55  
    community-acquired ..... 49  
      pediatric ..... 396  
    cystic fibrosis ..... 54  
    HIV  
      Aspergillus ..... 323  
      bacterial ..... 324  
      CAP ..... 65  
      M. tuberculosis ..... 326  
      P. (carinii) jiroveci (PCP) ..... 302  
    hospital-acquired (HAP) ..... 68  
    influenza ..... 62  
    MERS ..... 62  
    M. avium-intracellulare (MAI) ..... 66  
    M. kansasii ..... 66  
    M. pneumoniae ..... 57  
    MSSA/MRSA ..... 64t, 65  
    M. tuberculosis (TB) ..... 66  
    non-zoonotic atypical,  
      diagnostic features ..... 59  
    nosocomial (NP) ..... 68  
    nursing home-acquired (NHAP) ..... 68  
    organ transplants ..... 67  
    SARS ..... 62  
    S. aureus ..... 64t, 65  
    transplant infections ..... 165  
    tuberculosis ..... 53  
    unknown pathogen ..... 49  
    ventilator-associated (VAP) ..... 69  
    zoonotic atypical ..... 60
- Pneumonic plague ..... 174, 176  
  post-exposure prophylaxis ..... 363t
- Pneumothorax, spontaneous,  
  chest abnormalities (DDx) ..... 484t
- Polymox ..... 533
- Polymyxin B ..... 671
- Polymyxin B/Colistin  
  pearls & pitfalls ..... 520  
  susceptibility profiles ..... 198t
- Pork tapeworm ..... 281
- Posaconazole ..... 672
- Positive blood cultures vs. bacteremia  
(MSSA, MRSA, CoNS) ..... 13t

- Posterior cervical adenopathy,  
unilateral/bilateral, lymph node  
abnormalities (DDx) ..... 488t
- Post-exposure prophylaxis ..... 385t
- anthrax (inhalation/cutaneous) ..... 363t
- avian influenza ..... 359t
- chancroid ..... 360t
- diphtheria ..... 359t
- gonorrhea ..... 360t
- hepatitis A (HAV) ..... 361t
- hepatitis B (HBV) ..... 361t
- hepatitis C (HCV) ..... 361t
- Lyme disease ..... 361t
- meningitis ..... 358t
- non-gonococcal urethritis (NGU) ..... 360t
- pertussis ..... 359t
- pneumonic plague ..... 363t
- rabies ..... 362t
- Rocky Mountain spotted fever ..... 361t
- smallpox ..... 363t
- swine influenza ..... 359t
- syphilis ..... 360t
- tuberculosis ..... 359t
- tularemia pneumonia ..... 363t
- varicella (chickenpox) ..... 360t
- viral hemorrhagic fever ..... 363t
- viral influenza ..... 358t
- zoonotic diseases  
(plague, anthrax) ..... 362t
- Preauricular adenopathy, lymph node  
abnormalities (DDx) ..... 488t
- Pregnancy  
asymptomatic bacteriuria in ..... 364t  
safety in ..... 523
- Prevotella sp. .... 240t
- Prezista (darunavir) ..... 591
- Primaquine ..... 674
- Primary amebic meningoencephalitis  
(PAM) ..... 23
- Primaxin ..... 623
- Procalcitonin levels (PCT), elevated,  
serum test abnormalities (DDx) ..... 499t
- Progressive multifocal  
encephalopathy (PML), HIV ..... 330
- Proloprim ..... 707
- Prophylaxis, antibiotic  
chronic medical ..... 364t  
endocarditis ..... 369t  
HIV  
opportunistic infections ..... 366t  
pre-exposure (PrEP) ..... 366t  
post-exposure ..... 366t  
malaria ..... 375t  
post-exposure ..... 358t  
surgical ..... 354t  
tetanus ..... 377t  
transplant (BMT/SOT) ..... 367t  
travel ..... 371t
- Propionibacterium acnes ..... 239t
- Prostaphlin ..... 661
- Prostate enlargement/tenderness,  
genitourinary abnormalities (DDx) ..... 489t
- Prostatic abscess ..... 110
- Prostatitis ..... 110
- Prosthetic valve endocarditis (PVE) ..... 76
- Protein, elevation, pleural fluid  
abnormalities (DDx) ..... 504t
- Proteinuria, urinary  
abnormalities (DDx) ..... 502t
- Proteus mirabilis, vulgaris ..... 228t  
cellulitis ..... 133  
decubitus ulcers ..... 137  
diabetic foot ulcers ..... 138  
osteomyelitis ..... 130  
skin abscesses/infected cysts ..... 141
- Prototheca, cutaneous ..... 289, 292
- Prototheca wickerhamii,  
cutaneous prototheca ..... 289
- Protozoa, intestinal ..... 276
- Providencia sp. .... 229t
- Pseudallescheria boydii, pulmonary  
pseudallescheriasis ..... 269
- Pseudomonas aeruginosa ..... 229t  
acute bacterial meningitis ..... 18  
bone marrow transplant infection ..... 160  
brain abscess ..... 27  
burn wounds ..... 145  
cavitary infiltrates (thick-walled) ..... 464  
community-acquired pneumonia ..... 54  
endocarditis ..... 75

- epididymitis .....106
- external otitis .....37
- febrile neutropenia .....158
- HIV infection.....324
- keratitis.....31
- mastoiditis .....39
- pneumonia, nosocomial.....68
- sepsis .....152
- septic arthritis .....122
- surgical prophylaxis.....357t
- urosepsis .....112
- Pseudomonas* (*Chryseomonas*)
- luteola (CDC group Ve-1) .....229t
- Pseudomonas* (*Flavimonas*)
- oryzihabitans (CDC group Ve-2).....230t
- Psittacosis*
- bilateral infiltrates without
- effusion .....451
- unilateral focal segmental/lobar
- infiltrate without effusion.....428
- unilateral ill-defined infiltrates
- without effusion.....438
- Pubic lice .....150, 151
- Pulmonary actinomycosis .....268, 270
- Pulmonary alveolar proteinosis (PAP),
- bilateral infiltrates without
- effusion .....457
- Pulmonary amebiasis.....265
- Pulmonary aspergillosis .....268
- Pulmonary blastomycosis .....267, 269
- Pulmonary coccidioidomycosis....268, 271
- Pulmonary coin lesions.....266
- Pulmonary cryptococcosis .....269, 272
- Pulmonary cystic lesions/masses .....265
- Pulmonary embolus/infarct
- unilateral focal segmental/lobar
- infiltrate with effusion .....434
- unilateral focal segmental/lobar
- infiltrate without effusion.....429
- unilateral ill-defined infiltrates with
- effusion .....445
- Pulmonary hemorrhage
- bilateral infiltrates without effusion...453
- unilateral ill-defined infiltrates
- without effusion.....442
- Pulmonary histoplasmosis.....267, 269
- Pulmonary infiltrates/mass lesions.....267
- Pulmonary nocardiosis .....268
- Pulmonary paracoccidioidomycosis
- (South American blastomycosis).....267
- Pulmonary pseudallescheriasis .....269
- Pulmonary sporotrichosis .....268
- Pulmonary zygomycosis
- (*mucor-mycosis*).....269
- Purple nodules, extremity
- abnormalities (DDx) .....492t
- Purple nose tip, nasal
- abnormalities (DDx) .....481t
- Purulent CSF with negative
- gram stain, CSF
- abnormalities (DDx) .....505t
- Pyelonephritis
- acute.....107
- chronic.....108
- Pyoderma abscesses, severe,
- pediatric .....411
- Pyogenic lung abscess, cavitary
- infiltrates (thick-walled) .....463
- Pyomyositis .....136
- Pyrazinamide .....675
- pediatric dose .....421
- Pyrimethamine.....676
- Pyuria, urinary abnormalities (DDx) ...508t
- PZA .....675
- Q**-fever
- bilateral infiltrates
- without effusion.....452
- community-acquired pneumonia.....51
- endocarditis.....72
- unilateral focal segmental/lobar
- infiltrate without effusion.....428
- unilateral ill-defined infiltrates
- without effusion.....438
- zoonotic atypical pneumonia.....61
- Quinine sulfate.....676
- Quinolones
- pearls & pitfalls .....516
- susceptibility profiles .....198t

- Quinupristin/dalfopristin .....677  
 pearls & pitfalls .....517  
 susceptibility profiles .....198t  
 pediatric dose .....421
- R**-ickettsia, post-exposure  
 prophylaxis .....361t
- Rabies  
 post-exposure  
 prophylaxis .....362t  
 travel prophylaxis .....374t
- Radiation pneumonitis, unilateral focal  
 segmental/lobar infiltrate  
 with effusion .....435
- RAL (raltegravir) .....678
- Raltegravir (RAL) .....678
- Rapid influenza diagnostic tests (RIDTs),  
 negative, swine influenza  
 pneumonia in hospitalized adults .....64
- Rapivab .....668
- Rash  
 below waist, general appearance  
 abnormalities (DDx) .....477t  
 migratory, extremity  
 abnormalities (DDx) .....492t  
 palms/soles, extremity  
 abnormalities (DDx) .....492t
- Rat bite fever (Streptobacillus  
 moniliformis, spirillum minus) .....257
- RBC abnormalities (DDx)  
 anemia (acute) .....497t  
 hemophagocytosis  
 (erythrophagocytosis) .....497t  
 Howell-Jolly bodies .....497t  
 RBC inclusions .....497t  
 schistocytes (microangiopathic  
 hemolytic anemia) .....497t  
 spherocytes .....497t  
 target cells .....497t
- RBC inclusions, RBC  
 abnormalities (DDx) .....497t
- RBCs in CSF, CSF  
 abnormalities (DDx) .....505t
- RBCs (per mm<sup>3</sup>), pleural  
 effusion profiles in various  
 diseases (DDx) .....504t
- Rebetol .....680
- Relapsing fever (B, recurrentis),  
 louse-borne and tick-borne .....257, 234t
- Relative bradycardia  
 defined .....14  
 fever curves (DDx) .....475t  
 temperature-pulse relationships .....14t
- Relative lymphocytosis,  
 WBC abnormalities (DDx) .....495t
- Relative lymphopenia,  
 WBC abnormalities (DDx) .....495t
- Relative tachycardia,  
 fever curves (DDx) .....475t
- Renal abscess .....109  
 intrarenal/perinephric .....109
- Renal failure, chronic  
 bilateral infiltrates with  
 effusion .....460  
 bilateral infiltrates without  
 effusion .....454
- Renal insufficiency,  
 dosing strategies in .....4, 5t
- Renal route of elimination .....5t
- Renal TB .....108
- Rescriptor (delavirdine) .....593
- Resistance, antibiotic .....2, 207
- Respiratory tract infections,  
 lower, empiric therapy .....48
- Respiratory viruses  
 epiglottitis .....47  
 pharyngitis .....40
- Retinitis. *see* Chorioretinitis
- Retrovir (zidovudine) .....718
- Reyataz (atazanavir) .....546
- Rheumatic test abnormalities (DDx)  
 angiotensin-converting enzyme  
 levels (ACE), elevated .....501t  
 anti-nuclear antibody titers (ANA),  
 elevated .....501t  
 anti-smooth muscle antibodies  
 (ASM), elevated .....501t  
 c-ANCA, elevated .....501t  
 double stranded DNA (DS-DNA),  
 elevated .....501t  
 rheumatoid factors (RF),  
 elevated .....501t



- Rheumatoid factor (RF)  
 elevation, rheumatic test  
 abnormalities (DDx) ..... 501t  
 pleural fluid abnormalities (DDx) ..... 505t
- Rheumatoid nodules, cavitory  
 infiltrates (thick-walled) ..... 469
- Rhizopus  
 cutaneous zygomycosis ..... 290  
 pulmonary zygomycosis ..... 269
- Rhodococcus equi ..... 217  
 cavitory infiltrates (thick-walled) ..... 470  
 unilateral focal segmental/lobar  
 infiltrate with effusion ..... 434
- Ribavirin ..... 680
- Rickettsia africae ..... 294
- Rickettsia akari ..... 294
- Rickettsia conorii ..... 294
- Rickettsialpox ..... 294
- Rickettsia parkeri ..... 294
- Rickettsia prowazekii ..... 294
- Rickettsia rickettsii (tick borne), Rocky  
 Mountain spotted fever (RMSF) ..... 294
- Rickettsia tsutsugamushi ..... 294
- Rickettsia typhi ..... 294
- Rifabutin ..... 681
- Rifadin ..... 682
- Rifampin ..... 682  
 pediatric dose ..... 421
- Rifapentine ..... 359
- Right lower quadrant tenderness,  
 abdominal abnormalities (DDx) ..... 486t
- Right upper quadrant mass,  
 abdominal abnormalities (DDx) ..... 486t
- Right upper quadrant tenderness,  
 abdominal abnormalities (DDx) ..... 486t
- Right upper quadrant tympany,  
 abdominal abnormalities (DDx) ..... 486t
- Rilpivirine (RPV) ..... 684
- Rimactane ..... 682
- Rimantadine ..... 685
- Ringworm  
 body ..... 147  
 scalp ..... 147
- Ritonavir (RTV) ..... 686
- Rocephin ..... 572
- Rocky Mountain spotted  
 fever (RMSF) ..... 294  
 post-exposure prophylaxis ..... 361t
- Rose spots, abdominal  
 abnormalities (DDx) ..... 485t
- Rotavirus, travel prophylaxis ..... 372t
- Roundworms, intestinal ..... 278
- Routes of elimination  
 (Renal and hepatic) ..... 5
- RPV (rilpivirine) ..... 684
- RSV  
 bilateral infiltrates  
 without effusion ..... 449  
 transplant infections ..... 163t
- RTV (ritonavir) ..... 686
- Rubella, adult immunization ..... 381t
- S**-agalactiae, bacterial cystitis ..... 103
- Safety in pregnancy,  
 explanation ..... 523
- Safety profile, antibiotics ..... 4
- Salmonella sp. .... 230  
 septic arthritis ..... 125  
 travel prophylaxis ..... 372t  
 typhi, non-typhi ..... 230t  
 HIV/AIDS ..... 337  
 typhi, travel prophylaxis ..... 373t  
 typhi/paratyphi, typhoid  
 (enteric) fever ..... 88  
 Vibrio sp. infectious diarrhea ..... 86
- Salpingitis ..... 113
- Saltwater wounds ..... 145
- Saquinavir (SQV) ..... 687
- Sarcoidosis  
 bilateral infiltrates with  
 effusion ..... 461  
 bilateral infiltrates  
 without effusion ..... 454
- Sarcoptes scabiei, scabies ..... 150
- SARS ..... 62  
 bilateral infiltrates  
 without effusion ..... 449
- S aureus (MSSA/MRSA) pneumonia ..... 65

- Scabies .....150
- Scalp nodules, head  
  abnormalities (DDx) .....478t
- Scedosporium apiospermum,  
  pulmonary pseudallescheriasis .....269
- Schistocytes (microangiopathic  
  hemolytic anemia),  
  RBC abnormalities (DDx).....497t
- Schistosoma mansoni,  
  japonicum.....275
- Schistosomiasis .....275
- Scleral nodules, eye  
  abnormalities (DDx) .....480t
- Scleroderma (skin lesions), extremity  
  abnormalities (DDx) .....492t
- Scrofula .....285
- Scrotal enlargement,  
  genitourinary abnormalities (DDx) ...489t
- Scrub (chigger/mite-borne)  
  typhus .....294
- Seborrheic dermatitis,  
  severe, head abnormalities  
  (DDx).....477t
- Selzentry (maraviroc).....642
- Sepsis .....151  
  asplenia/hyposplenia .....155  
  candidemia .....153  
  intraabdominal/pelvic source.....154  
  IV line .....156  
  lung source .....152  
  miliary BCG .....155  
  miliary TB .....155  
  overwhelming, with purpura .....155  
  steroids, chronic .....155  
  unknown source .....151  
  urosepsis .....157
- Septic abortion .....113
- Septic arthritis .....121  
  acute .....121  
  animal bite .....123  
  fungal .....123  
  pediatric .....407  
  TB .....123
- Septic pelvic thrombophlebitis....101, 102
- Septic pulmonary emboli, cavitary  
  infiltrates (thick-walled) .....465
- Septic shock .....151
- Septic thrombophlebitis .....82
- Septtra .....707
- Sequestered lung,  
  cavitory infiltrates (thin-walled) .....473
- Seromycin .....586
- Serpiginous lesions,  
  subcutaneous .....288
- Serratia marcescens .....230  
  acute bacterial meningitis .....19  
  endocarditis .....75  
  pneumonia .....68  
  sepsis .....152  
  urosepsis .....112
- Serum half-life (normal/ESRD),  
  explanation .....522
- Serum test abnormalities (DDx)  
  elevated cold agglutinins .....499t  
  elevated ferritin levels,  
    (acute/chronic) .....499t  
  erythrocyte sedimentation  
    rate (ESR) .....498t  
  lactate dehydrogenase  
    (LDH) .....499t  
  monoclonal gammopathy .....499t  
  monospot (false +) .....499t  
  procalcitonin levels (PCT) .....499t  
  SPEP (monoclonal, polyclonal  
    gammopathy) .....498t
- Serum transaminases  
  (SGOT/SGPT),  
  liver test abnormalities (DDx) .....500t
- Severe acute respiratory syndrome  
  (SARS) .....62
- Sexually transmitted diseases (STDs)  
  empiric therapy .....114
- Shigella sp. ....231  
  dysentery .....91  
  HIV .....338  
  travel prophylaxis .....372t
- Shingles .....142

- Shin hyperpigmentation, extremity  
abnormalities (DDx) ..... 492t
- Shock, general appearance  
abnormalities (DDx) ..... 476t
- Shoulder tenderness, chest  
abnormalities (DDx) ..... 484t
- Shunt infections, CNS,  
acute bacterial meningitis ..... 21
- Side effects, adverse/Simeprevir ..... 523
- Simeprevir ..... 98t, 346t
- Single fever spikes, fever  
curves (DDx) ..... 475t
- Sinusitis  
acute bacterial ..... 30  
chronic bacterial ..... 31  
pediatric ..... 392
- Sirturo ..... 53
- Sivextro ..... 691
- Skin  
boils ..... 140  
fungal infections ..... 147  
infections, empiric therapy ..... 132  
infestations (body lice,  
head lice, pubic lice, scabies) ..... 150  
other lesions ..... 297  
papules/nodules/abscesses ..... 289  
parasites, fungi,  
unusual organisms ..... 283  
pustules ..... 140  
rickettsia ..... 294  
subcutaneous lesions ..... 283  
subcutaneous serpiginous lesions ..... 288  
ulcers ..... 137  
decubitus ulcers ..... 137  
diabetic foot ulcers ..... 138  
ischemic foot ulcers ..... 139  
ulcers/skin tracts/abscesses ..... 284  
vesicles/bullae ..... 142
- Sleeping sickness ..... 255
- Smallpox ..... 177  
bioterrorist ..... 173  
post-exposure prophylaxis ..... 363t
- Sofosbuvir ..... 98, 346
- Soft tissue crepitation, general  
appearance abnormalities (DDx) ..... 476t
- Soft tissue infections,  
empiric therapy ..... 132
- Sole lesions, painless, extremity  
abnormalities (DDx) ..... 490t
- Sole rashes, extremity  
abnormalities (DDx) ..... 492t
- Solid organ transplant  
(SOT), transplant infections ..... 161
- Sovaldi ..... 98t, 346t
- Specific gravity, decreased,  
urinary abnormalities (DDx) ..... 502t
- Spectam ..... 689
- Spectinomycin ..... 689
- Spectracef ..... 559
- SPEP (monoclonal, polyclonal  
gammopathy), (DDx) ..... 498t
- Sphenoid sinus, acute  
bacterial sinusitis ..... 31
- Spherocytes, RBC  
abnormalities (DDx) ..... 497t
- Spinal tenderness, back  
abnormalities (DDx) ..... 484t
- Spirillum minus (rat bite fever) ..... 234t, 257
- Spirochetes  
aerobic ..... 234t  
blood ..... 257
- Splenomegaly, abdominal  
abnormalities (DDx) ..... 487t
- Splinter hemorrhages, extremity  
abnormalities (DDx) ..... 490t
- Spontaneous bacterial  
peritonitis (SBP) ..... 102
- Sporanox ..... 629
- Sporothrix schenckii  
chronic septic arthritis ..... 126  
cutaneous sporotrichosis ..... 290  
fungal arthritis ..... 123  
pulmonary sporotrichosis ..... 268
- Sporotrichosis  
cavitary infiltrates (thick-walled) ..... 466  
cavitary infiltrates (thin-walled) ..... 471  
cutaneous ..... 290
- Sputum gram stain, clinical use ..... 248t
- Squamous cell carcinoma, cavitary  
infiltrates (thick-walled) ..... 468

- SQV (saquinavir) .....687
- St. Louis encephalitis (St.LE) .....25
- Staphylococcus aureus .....208
- acute bacterial meningitis .....19
- animal bite wounds .....144
- bacteremia vs + blood cultures .....13t
- bone marrow transplant
- infection .....160
- brain abscess .....27
- burn wounds .....145
- CAPD-associated peritonitis .....102
- cavitary infiltrates (thick-walled) .....464
- complicated skin/skin
- structure tissue infections .....134
- decubitus ulcers .....137
- diabetic foot ulcers .....138
- endocarditis .....76
- acute bacterial (ABE) .....76
- prosthetic valve (PVE) .....77
- endophthalmitis .....35
- human bite wounds .....144
- infected joint prosthesis .....128
- ischemic foot ulcers .....139
- IV line (CVC) infections .....81
- keratitis .....31
- lung abscess/empyema .....67
- mastoiditis .....39
- osteomyelitis .....129
- pacemaker infections .....80
- pericarditis .....78
- pneumonia (community acquired) .....65
- positive BCs vs. bacteremia .....13t
- renal abscess .....109
- sepsis .....151
- septic arthritis .....121
- skin abscesses/infected cysts .....140
- solid organ transplant .....161
- suppurative parotitis .....40
- surgical prophylaxis .....355t, 356t, 357t
- toxic shock syndrome .....169
- transplant infections .....160
- vascular graft infections .....82
- Staphylococcus epidermidis (CoNS) ...209t
- brain abscess .....27
- CAPD-associated peritonitis .....101
- CNS shunt infection .....19
- endocarditis .....76
- infected joint prosthesis .....128
- pacemaker infection .....80
- sepsis .....152
- surgical prophylaxis .....355t
- Staphylococcus lugdunensis .....209t
- Staphylococcus saprophyticus .....210t
- cystitis .....103
- dysuria-pyuria syndrome .....103
- Stavudine (d4T) .....689
- Stenotrophomonas maltophilia .....231t
- community-acquired pneumonia .....54
- Sternal tenderness, chest
- abnormalities (DDx) .....484t
- Steroid therapy, chronic,
- pneumonia .....67
- Stomach, upper small bowel
- surgery, surgical prophylaxis .....356t
- Stomatitis .....44
- Stool, parasites, fungi,
- unusual organisms in .....276
- Streptobacillus moniliformis .....231t
- septic arthritis .....123
- Streptococcus anginosus (S. milleri)
- group (S. intermedius, S. anginosus,
- S. constellatus) .....213
- Streptococcus (bovis)
- gallolyticus .....213t
- endocarditis .....71
- Streptococci. *see* Group
- A streptococci; Group
- B streptococci; Groups C, F, G
- streptococci; Nutritionally-variant
- streptococci (NVS)
- deep neck infections .....46
- dental infections .....46
- endophthalmitis .....35
- Streptococcus pneumoniae .....213t
- acute bacterial exacerbation of
- chronic bronchitis (AECB) .....48
- acute bacterial meningitis .....18
- acute otitis media .....38
- adult immunization .....381t
- brain abscess .....27

chronic medical prophylaxis	364t	Suprax	562
CLSI susceptibility breakpoints	8t	Surgical prophylaxis	
community-acquired		arthroscopy	357t
pneumonia	49	biliary tract surgery	355t
conjunctivitis	32	cardiac valve replacement	
epiglottitis	47	surgery	355t
HIV infection	322t	CNS shunt (VP/VA) placement,	
prophylaxis	319	craniotomy, open CNS	
intraabdominal or pelvic		trauma	355t
peritonitis/abscess	101	coronary artery bypass	
keratitis	31	graft (CABG) surgery	355t
lung abscess/empyema	67	distal small bowel,	
mastoiditis	39	colon surgery	356t
pericarditis	78	factors affecting efficacy of	354t
prophylaxis of opportunistic		hepatic surgery	356t
infections	319t	orthopedic prosthetic implant	
sepsis	152	surgery (total hip/knee	
susceptibility breakpoints	8	replacement)	357t
unilateral focal segmental/lobar		orthopedic surgery	
infiltrate without effusion	426	(open fracture)	357t
Streptomycin	690	pelvic (OB/GYN) surgery	356t
Stribild [elvitegravir (EVG) + combicostat		stomach, upper small bowel	
(COBI) + emtricitabine (FTC) +		surgery	356t
tenofovir (TDF)]	602	thoracic (non-cardiac) surgery	355t
Strongyloides, bilateral		TURP, cytoscopy	357t
infiltrates with effusion	459	urological implant surgery	357t
Strongyloides stercoralis	278	Surgical therapy,	
transplant infections	165	inadequate, antibiotics and	15
Strongyloidiasis	278	Susceptibility testing	
Subacute bacterial		microbiology and	6
endocarditis (SBE)	70	<i>in vitro</i> vs. <i>in vivo</i> activity	7
Subconjunctival hemorrhage, eye		Sustiva	599
abnormalities (DDx)	479t	Swimmers ear	37
Subcutaneous serpiginous		Swimming pool granuloma	285
lesions	288	Swine influenza (H <sub>1</sub> N <sub>1</sub> )	65
Subdural empyema	27	community-acquired pneumonia	51
Superior vena cava syndrome,		post-exposure prophylaxis	359t
neck abnormalities (DDx)	483t	Symmetrel	531
Suppurating inguinal		Syndrome of inappropriate antidiuretic	
adenopathy	118	hormone (SIADH), encephalitis	26
Suppurative parotitis	40	Synercid	677
Suppurative pericarditis	78	Synergistic gangrene	135
Supraclavicular adenopathy,		Syphilis	119
lymph node		latent	120
abnormalities (DDx)	488t		

- neurosyphilis .....119  
 post-exposure prophylaxis .....360t  
 primary .....119  
 secondary .....120  
 tertiary .....120  
 Systemic lupus erythematosus (SLE)  
 bilateral infiltrates with effusion .....460  
 unilateral ill-defined infiltrates  
 without effusion .....442
- T**-achycardia, heart abnormalities  
 (DDx) .....485t
- Taenia multiceps, cerebral  
 coenurosis .....263
- Taenia saginata, beef tapeworm .....281
- Taenia solium, neurocysticercosis .....262
- Taenia solium, pork tapeworm .....281
- Tamiflu .....660
- Tapeworms, intestinal .....281
- Target cells, RBC  
 abnormalities (DDx) .....497t
- Tazicef .....568
- Tazidime .....568
- Tazocin .....669
- TB osteomyelitis .....132
- TDF (tenofovir) .....695
- Tedizolid .....691  
 pearls & pitfalls .....517  
 susceptibility profiles .....198t
- Teflaro .....567
- Teichoic acid antibody  
 titers (TAA) .....600t
- Telavancin .....693  
 pearls & pitfalls .....519  
 susceptibility profiles .....198t
- Telithromycin .....694
- Temporal muscle wasting, head  
 abnormalities (DDx) .....477t
- Tender fingertips, extremity  
 abnormalities (DDx) .....490t
- Tenofovir disoproxil  
 fumarate .....695
- Tenosynovitis, extremity  
 abnormality (DDx) .....491t
- Terbinafine .....697
- Tetanus .....170, 172  
 adult immunization .....381t  
 travel prophylaxis .....375t
- Tetanus prophylaxis in routine  
 wound management  
 more than/equal to 3 doses .....377t  
 unknown or less than 3 doses .....377t
- Tetracyclines .....698  
 pearls & pitfalls .....512  
 susceptibility profiles .....186t
- Thigh tenderness (bilateral),  
 extremity abnormality (DDx) .....491t
- Thoracic surgery, surgical  
 prophylaxis .....355t
- 3TC (lamivudine) .....632
- 3TC/ZDV (Combivir) .....633
- Thrombocytopenia, platelet  
 abnormalities (DDx) .....498t
- Thrombocytosis, platelet  
 abnormalities (DDx) .....498t
- Thrombophlebitis  
 extremity abnormality (DDx) .....491t  
 intracranial suppurative .....27  
 septic .....81  
 septic pelvic .....101
- Thrush .....43  
 HIV .....336
- Ticar .....699
- Ticarcillin .....699  
 pediatric dose .....421
- Tivicay (dolutegravir) .....595
- Ticarcillin/clavulanate .....699  
 pearls & pitfalls .....512  
 susceptibility profiles .....186t  
 pediatric dose .....421
- Trichophyton mentagrophytes,  
 tinea corporis .....147
- Tick-borne relapsing fever (TBRF) .....257
- Tick typhus fevers .....294
- Tigecycline .....700  
 pearls & pitfalls .....518  
 susceptibility profiles .....198t
- Time-dependent kinetics,  
 antibiotics with .....2
- Timentin .....699
- Tinea capitis (scalp ringworm) .....149

- Tinea corporis (body ringworm) . . . . .147
- Tinea cruris (jock itch) . . . . .147
- Tinea pedis . . . . .147
- Tinea versicolor (pityriasis) . . . . .148
- Tinidazole, pediatric dose . . . . .421
- Tipranavir (TPV) . . . . .704
- Tissue penetration,  
antibiotic selection and . . . . . 2
- TMP . . . . .707
- TMP-SMX . . . . .707
- pearls & pitfalls . . . . .515
- susceptibility profiles . . . . .186t
- Tobramycin . . . . .705
- pearls & pitfalls . . . . .513
- susceptibility profiles . . . . .198t
- pediatric dose . . . . .421
- Tongue tenderness,  
mouth abnormalities (DDx) . . . . .482t
- Tongue ulcers, mouth  
abnormalities (DDx) . . . . .482t
- Tonsillar membranes, mouth  
abnormalities (DDx) . . . . .483t
- Tonsillar ulcers, mouth  
abnormalities (DDx) . . . . .483t
- Toxic shock syndrome (TSS) . . . . .169
- Toxin-mediated infectious diseases . . . . .169
- botulism . . . . .170
- diphtheria . . . . .170
- tetanus . . . . .170
- toxic shock syndrome (TSS) . . . . .169
- Toxocarasis, meningitis . . . . .22
- Toxoplasma gondii  
    cerebral toxoplasmosis . . . . .262
- chorioretinitis . . . . .34
- encephalitis . . . . .25
- HIV . . . . .331
- HIV, prophylaxis . . . . .319t
- transplant infections . . . . .164
- Toxoplasmosis  
    HIV, prophylaxis . . . . .320
- meningitis . . . . .22
- prophylaxis of opportunistic  
        infections . . . . .366t
- TPV (tipranavir) . . . . .704
- Trachipleistophora,  
    microsporidia enteritis . . . . .335
- Trachoma conjunctivitis . . . . .33
- Transplant prophylaxis  
    (BMT/SOT) . . . . .367t
- Aspergillus sp. . . . .368t
- Candida sp. . . . .368t
- cytomegalovirus (CMV) . . . . .368t
- Herpes simplex (HSV) . . . . .367t
- PCP . . . . .368t
- toxoplasmosis . . . . .368t
- varicella zoster virus (VZV) . . . . .367t
- Transplants. *see* Organ transplants
- Transverse myelitis, neurological  
    abnormalities (DDx) . . . . .494t
- Trapezius tenderness,  
    chest abnormalities (DDx) . . . . .484t
- Traveler's diarrhea,  
    travel prophylaxis . . . . .373t
- Travel prophylaxis . . . . .371t
- general infectious disease travel  
        precautions . . . . .371t
- hepatitis A (HAV) . . . . .373t
- Japanese encephalitis (JE) . . . . .374t
- meningococcal meningitis . . . . .373t
- rabies . . . . .374t
- tetanus, diphtheria, pertussis . . . . .375t
- traveler's diarrhea . . . . .373t
- typhoid fever . . . . .373t
- yellow fever . . . . .374t
- Travel vaccines, HIV . . . . .323
- Trecator . . . . .611
- Trematodes (flukes/flatworms),  
    intestinal . . . . .282
- Treponema pallidum  
    meningitis . . . . .22
- post-exposure prophylaxis . . . . .360t
- primary syphilis . . . . .117
- syphilis . . . . .119, 120
- Trichinella spiralis, trichinosis . . . . .298
- Trichinosis . . . . .298
- Trichomonas vaginalis,  
    urethritis/cervicitis . . . . .114
- Trichophyton mentagrophytes,  
    onychomycosis . . . . .148

- Trichophyton rubrum  
  onychomycosis ..... 148  
  tinea corporis ..... 147  
Trichosporon beigeli, trichosporon ..... 289  
Trichostrongyliasis ..... 278  
Trichostrongylus orientalis ..... 278, 280  
Trichuris trichiura ..... 279  
Trimethoprim ..... 707  
Trimethoprim-sulfamethoxazole ..... 707  
Trimox ..... 533  
Trimpex ..... 707  
Trismus, mouth abnormalities  
  (DDx) ..... 482t  
Triumeq ..... 314t  
Trizivir [abacavir (ABC) + lamivudine  
  (3TC) + zidovudine (ZDV)] ..... 528  
Trobicin ..... 689  
Tropheryma whippelli,  
  Whipple's disease ..... 282  
Truvada [emtricitabine (FTC) +  
  tenofovir (TDF)] ..... 604  
Trypanosoma brucei  
  cruzi ..... 255, 256  
  gambiense ..... 255, 256  
  rhodesiense ..... 255, 256  
Trypanosoma brucei cruzi,  
  Chagas' disease ..... 273  
Trypanosomes in blood ..... 255  
Trypanosomiasis  
  American ..... 255  
  East African ..... 255  
  West African ..... 255  
Tsutsugamushi fever ..... 294  
Tuberculosis. *see* Mycobacterium  
  tuberculosis  
Tuberculosis  
  endophthalmitis ..... 35  
Tuboovarian abscess ..... 113  
Tularemia pneumonia ..... 173  
  bioterrorist ..... 176  
  post-exposure  
  prophylaxis ..... 363t  
  unilateral focal segmental/lobar  
  infiltrate with effusion ..... 432  
TURP, surgical prophylaxis ..... 357t
- Tyagil ..... 700  
Typhoid fever ..... 89  
  travel prophylaxis ..... 373t  
Typhus infections  
  epidemic ..... 295  
  flying squirrel ..... 294  
  murine ..... 296  
  scrub ..... 296  
  tick typhus fevers ..... 296  
T20 (enfuvirtide) ..... 606
- U**  
Ulcer-node syndromes, lymph  
  node abnormalities (DDx) ..... 488t  
Ulcers  
  Buruli ..... 285  
  foot/leg ..... 491t  
  frenal ..... 482t  
  genital ..... 117  
  mouth ..... 44  
  oral ..... 482t  
  peptic ..... 85  
  perirectal ..... 489t  
  skin ..... 137  
  tongue ..... 482t  
  tonsillar ..... 483t  
Ultra ..... 621  
Ultracef ..... 556  
Unasyn ..... 542  
Unipen ..... 652  
Unsafe water (fecally-contaminated),  
  infectious disease travel  
  precautions ..... 371t  
Ureaplasma urealyticum  
  post-exposure prophylaxis ..... 360t  
  urethritis/cervicitis ..... 114  
Urethritis  
  gonococcal ..... 114  
  non-gonococcal ..... 114  
Urex ..... 642  
Urinary abnormalities (DDx)  
  chyluria ..... 503t  
  eosinophiluria ..... 503t  
  hematuria, gross/microscopic ..... 502t  
  leukocyte esterase ..... 502t



- myoglobinuria (2° to rhabdomyolysis).....503t
- nitrite positive .....502t
- proteinuria .....502t
- pyuria .....501t
- specific gravity, decreased .....502t
- urinary pH, alkaline/acidic .....502t
- Urinary pH, alkaline/acidic, urinary abnormalities (DDx) .....502t
- Urinary tract infections (UTIs), chronic medical prophylaxis .....364t
- Urine gram stain, clinical use .....248t
- Urological implant surgery, surgical prophylaxis .....357t
- Urosepsis .....111
- community-acquired .....111
- urologic procedure .....112
- Urticaria, general appearance abnormalities (DDx) .....477t
- USFDA use-in-pregnancy letter code .....524t
- Usual dose, explanation .....521
- Uveitis, eye abnormalities (DDx) .....480t
- Uvular edema, mouth abnormalities (DDx) .....483t
- V**-accinations. (see Adult immunizations)
- Immunizations
- Vaginosis .....115
- Vaginitis .....115
- Vaginitis, Candida, in HIV/AIDS .....338
- Valacyclovir .....709
- Valcyte .....710
- Valganciclovir .....710
- Valtrex .....709
- Vancocin .....712
- Vancomycin .....712
- pearls & pitfalls .....516
- susceptibility profiles .....198t
- pediatric dose .....422
- Vancomycin-resistant enterococci (VRE) .....3
- Vantin .....565
- Varicella (VZV) chickenpox adult immunization .....382t
- post-exposure prophylaxis .....360t
- Varicella zoster virus (VZV). (see also Herpes zoster)
- chickenpox .....142
- herpes zoster .....288
- HIV .....343
- Variola virus post-exposure prophylaxis .....363t
- smallpox .....174
- Vascular graft infections .....82
- Venezuelan equine encephalitis (VEE) (VAP) .....69
- Verrucous foot/leg lesions, extremity abnormality (DDx) .....491t
- Verrucous hand/arm lesions, extremity abnormality (DDx) .....490t
- Vesicles, skin .....288
- Vfend .....714
- Vibativ .....693
- Vibramycin .....597
- Vibra-tabs .....597
- Vibrio alginolyticus .....232t
- Vibrio cholerae .....232t
- travel prophylaxis .....372t
- Vibrio parahaemolyticus .....232t
- Vibrio sp. saltwater wounds .....145
- Vibrio vulnificus .....232t
- saltwater wounds .....145
- Videx (didanosine) .....594
- Viekira Pak .....98t, 346t
- Vincent's angina .....44
- Viracept (nelfinavir) .....653
- Viral conjunctivitis .....33
- Viral esophagitis .....85
- Viral hemorrhagic fever, post-exposure prophylaxis .....363t
- Viral hepatitis .....97
- Viral influenza (human seasonal), post-exposure prophylaxis .....358t
- Viral pharyngitis .....42

- Viramune (nevirapine) .....654  
 Viread (tenofovir) .....695  
 Viridans streptococci endocarditis  
   prosthetic, subacute (SBE) ..... 70  
 Viridans  
   prosthetic valve endocarditis ..... 77  
   streptococci .....213t  
   bone marrow transplant infection ....160  
 Visceral leishmaniasis (kala-azar) .....275  
 Vistide .....577  
 Vitiligo, general appearance  
   abnormalities (DDx) .....477t  
 Volume of distribution, explanation ....522  
 Voriconazole .....714  
 VRE. *see* Vancomycin-resistant  
   enterococci (VRE)  
 VZV  
   bilateral infiltrates without effusion ....449  
   community-acquired pneumonia .....52  
   conjunctivitis .....33  
   encephalitis .....25  
   HIV, prophylaxis .....319  
   meningitis .....21  
   ophthalmicus .....33  
   post-exposure prophylaxis .....360t  
   prophylaxis of opportunistic  
     infections .....366t
- W**-arts, genital/perianal .....118  
 Watery eyes, eye abnormalities  
   (DDx) .....480t  
 WBC abnormalities (DDx)  
   atypical lymphocytes .....496t  
   basophilia .....496t  
   eosinophilia .....496t  
   immunoblasts .....496t  
   leukocytosis .....495t  
   leukopenia .....495t  
   lymphocytosis .....495t  
   monocytosis .....496t  
   relative lymphocytosis, .....495t  
   relative lymphopenia .....495t  
   WBC inclusions .....496t  
   WBC inclusions, WBC  
     abnormalities (DDx) .....496t  
   WBCs pleural  
     effusion profiles in various  
       diseases (DDx) .....504t  
   WBC type/glucose, clinical use .....247t  
 Wegener's granulomatosis  
   bilateral infiltrates with effusion .....461  
   bilateral infiltrates  
     without effusion .....453  
     cavitary infiltrates (thick-walled) .....468  
 West African trypanosomiasis .....255  
 Western equine encephalitis (WEE) .....25  
 West Nile encephalitis (WNE) .....25  
   meningitis .....21  
 Wheezing, diffuse, chest  
   abnormalities (DDx) .....484t  
 Whipple's disease .....282  
 Whipworm .....279  
 Wound infections .....144  
   animal bite .....144  
   burns .....145  
   cat scratch disease (CSD) .....144  
   freshwater .....145  
   human bite .....144  
   saltwater .....145  
 Wrist swelling, extremity  
   abnormality (DDx) .....491t  
 Wuchereria bancrofti .....254  
 Wymox .....533
- Y**-easts/fungi, clinical significance,  
   pending susceptibility testing .....242t  
 Yellow fever .....178  
   travel prophylaxis .....374t  
 Yellow/whitish pleural fluid  
   abnormalities (DDx) .....503t  
 Yersinia  
   infectious diarrhea .....86  
   travel prophylaxis .....373t  
 Yersinia enterocolitica .....232t  
 Yersinia pestis (plague) .....233t  
   bioterrorist, pneumonic plague .....174  
   post-exposure prophylaxis .....362t

<b>Z</b> -alcitabine (ddc).....	717	Zoonotic atypical pneumonias .....	60
ZDV (zidovudine).....	718	diagnostic features .....	61
Zerbaxa .....	572	Zoonotic diseases,	
Zerit (stavudine).....	689	post-exposure prophylaxis.....	362t
Ziagen (ABC).....	526	Zosyn .....	669
Zidovudine (ZDV).....	528	Zovirax.....	529
Zinacef.....	574	Zygomycosis, cutaneous .....	290
Zithromax.....	549	Zyvox .....	637

---


## **COLOR ATLAS**

### ***CSF, Sputum, and Urine Gram Stains***


Paul E. Schoch, PhD  
Edward J. Bottone, PhD  
Daniel Caplivski, MD

---


CSF GRAM STAINS




Staphylococcus aureus




Listeria monocytogenes




H. influenzae



Neisseria meningitidis




Klebsiella pneumoniae




Streptococcus pneumoniae


**SPUTUM GRAM STAINS**




*Streptococcus pneumoniae*




*Klebsiella pneumoniae*




*Haemophilus influenzae*



*Staphylococcus aureus*




*Moraxella catarrhalis*




*Pseudomonas aeruginosa*

## URINE GRAM STAINS




E. coli



Group B streptococci



Enterococci



Pseudomonas aeruginosa

---

# **COLOR ATLAS**


## ***Fungal Stains***

Daniel Caplivski, MD  
Edward J. Bottone, PhD


---




## FUNGAL STAINS




Actinomyces (lung biopsy)




Candida albicans (blood)




Alternaria (skin biopsy)




Candida albicans with  
pseudohyphae (blood)




Aspergillus (lung biopsy)




Candida as fungal ball (lung biopsy)




Candida (liver biopsy)




Fusarium (skin biopsy)




Cryptococcus (lung biopsy)




Histoplasma capsulatum (lung biopsy)




Cunninghamella (lung biopsy)




Malassezia (blood)




Mucormycosis (bone marrow)




Pneumocystis (lung biopsy)




Nocardia (lung biopsy)



Rhizomucor (sinus biopsy)



Paracoccidioidomycosis (bone biopsy)



Scedosporium (lung biopsy)